Supporting Information

Palladium-Catalyzed Reaction of Aryl Bromides with 7-Hydroxy-1,3-Dienes

Ming-Chang P. Yeh,^{*} Wen-Cheng Tsao and Ling-Hsien Tu

Department of Chemistry, National Taiwan Normal University, 88 Ding-Jou Road,
Section 4, Taipei 117, Taiwan

- (I) Analytical Data of Starting 7-Hydroxy-1,3-Dienes**
- (II) ¹H and ¹³C NMR Spectra for 2b, 2c, 2g, 2h, 2j, 2k, 2m and 2o**
- (III) X-ray Diffraction Analysis of 2c**
- (IV) CIF Format of X-ray Diffraction Analysis of 2c**

(I) Analytical Data of Starting 7-Hydroxy-1,3-Dienes

1-(2,4-Cyclohexadienyl)-2-methyl-2-propanol (1a)

IR (CH₂Cl₂) 3578, 3028, 2930, 1714, 1603, 1495, 1453, 1282, 1254 cm⁻¹; ¹H NMR (200 MHz, CDCl₃) δ 5.82 (m, 4 H), 2.39 (m, 2 H), 2.28 (m, 1 H), 2.05 (dd, *J* = 14.1, 6.6 Hz, 1 H), 1.50 (dd, *J* = 14.4, 5.6 Hz, 1 H), 1.21 (s, 6 H); ¹³C NMR (50 MHz, CDCl₃) δ 132.45, 125.77, 123.85, 123.18, 70.92, 47.93, 30.19, 29.82, 29.68, 29.15; MS (EI) *m/e* 152.2 (M⁺, 4), 135.2 (14), 134.2 (25), 95.1 (29), 91.1 (76), 79.1 (100), 59.1 (94); HRMS (EI) *m/e* calcd for C₁₀H₁₆O 152.1201, found 152.1208.

2-(2,4-Cyclohexadienyl)-1,1-diphenyl-1-ethanol (1b)

IR (CH₂Cl₂) 3575, 3034, 2928, 2861, 1714, 1601, 1495, 1455, 1365, 1284, 1251 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.39 (m, 4 H), 7.28 (m, 4 H), 7.19 (m, 2 H), 5.81 (m, 2 H), 5.77 (m, 1 H), 5.62 (m, 1 H), 2.52 (ddd, *J* = 16.3, 8.9, 1.8 Hz, 1 H), 2.35 (m, 2 H), 2.18 (m, 1 H), 1.96 (m, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ 147.29, 147.04, 132.27, 128.11, 128.08, 126.79, 126.77, 125.98, 125.84, 124.06, 123.46, 78.46, 46.25, 29.99, 28.71; MS (EI) *m/e* 276.3 (M⁺, 0.1), 183.2 (33), 181.2 (18), 180.2 (96), 165.1 (26), 105.1 (100), 84.0 (54), 77.1 (80); HRMS (EI) *m/e* calcd for C₂₀H₂₀O 276.1514, found 276.1512.

2-Benzyl-1-(2,4-cyclohexadienyl)-3-phenyl-2-propanol (1c)

IR (CH₂Cl₂) 3574, 3032, 2927, 2860, 1712, 1602, 1495, 1454, 1369, 1280, 1253 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 10 H), 5.84 (m, 2 H), 5.76 (m, 2 H), 2.79 (m, 4 H), 2.67 (m, 1 H), 2.33 (m, 1 H), 1.93 (m, 1 H), 1.68 (dd, *J* = 14.55, 6.95 Hz, 1 H), 1.45 (dd, *J* = 14.55, 5.6 Hz, 1 H); ¹³C NMR (125 MHz, CDCl₃) δ 137.2, 137.2, 132.6, 130.8, 130.7, 128.2, 128.1, 128.1, 126.5, 126.5, 126.1, 124.1, 123.7, 74.6, 46.2, 46.1, 43.6, 30.5, 28.9.

(5Z)-2-Methyl-5,7-octadien-2-ol (1d)

IR (CH₂Cl₂) 3575, 3026, 2930, 1713, 1605, 1495, 1452, 1280, 1256 cm⁻¹; ¹H NMR (200 MHz, CDCl₃) δ 6.66 (m, 1 H), 6.01 (t, *J* = 10.8 Hz, 1 H), 5.47 (q, *J* = 7.6 Hz, 1 H), 5.15 (dd, *J* = 21.6, 17.2 Hz, 2 H), 2.30 (q, *J* = 7.8 Hz, 2 H), 1.52 (m, 2 H), 1.25 (s, 6 H); ¹³C NMR (50 MHz, CDCl₃) δ 132.27, 131.83, 128.86, 116.59, 70.34, 43.18, 28.83, 22.57; MS (EI) *m/e* 140.2 (M⁺, 13), 125.2 (100), 107.2 (23), 91.1 (16), 84.1

(58), 83.1 (34), 69.1 (48), 55.1 (33); HRMS (EI) m/e calcd for $C_9H_{16}O$ 140.1201, found 140.1208.

2-Methyl-5,7-nonadien-2-ol (1e)

IR (CH_2Cl_2) 3580, 3031, 2929, 2861, 1711, 1602, 1494, 1451, 1366, 1281, 1255 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 6.32 (m, 1 H), 5.92 (d, $J = 10.9$ Hz, 1 H), 5.66 (m, 1 H), 5.27 (m, 1 H), 2.23 (dd, $J = 16.4, 7.7$ Hz, 2 H), 1.76 (d, $J = 6.8$ Hz, 3 H), 1.53 (m, 2 H), 1.22 (s, 6 H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 129.41, 129.24, 128.63, 126.73, 70.88, 43.52, 29.18, 22.72, 18.19.

(II) ^1H and ^{13}C NMR Spectra for 2b, 2c, 2g, 2h, 2j, 2k, 2m and 2o

(III) X-ray Diffraction Analysis of 2c

1. ORTEP diagram
2. Table 1. Crystal data and structure refinement
3. Atomic Coordinates
4. Bond Lengths and Angles
5. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$)
6. Anisotropic Thermal Parameters in CIF Format.

1. ORTEP Diagram

(±)-(3*aS*,5*S*,7*aS*)-5-(4-Methoxyphenyl)-2,2-dimethyl-2,3,3*a*,4,5,7*a*-hexahydrobenz
o[*b*]furan(2c).

2c

2. Table 1. Crystal data and structure refinement for 2c.

Identification code	3508	
Empirical formula	C17 H22 O2	
Formula weight	258.35	
Temperature	200(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	P 21 c n	
Unit cell dimensions	a = 5.7550(2) Å	= 90°.
	b = 14.2250(5) Å	= 90°.
	c = 17.5650(8) Å	= 90°.
Volume	1437.96(10) Å ³	
Z	4	
Density (calculated)	1.193 Mg/m ³	
Absorption coefficient	0.076 mm ⁻¹	
F(000)	560	
Crystal size	0.26 x 0.14 x 0.1 mm ³	
Theta range for data collection	2.32 to 25.01°.	
Index ranges	-5<=h<=6, -15<=k<=16, -18<=l<=20	
Reflections collected	7793	
Independent reflections	2432 [R(int) = 0.0672]	
Completeness to theta = 25.01°	99.9 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	1.0171 and 0.948	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2432 / 1 / 173	
Goodness-of-fit on F ²	0.971	
Final R indices [I>2sigma(I)]	R1 = 0.0449, wR2 = 0.0982	
R indices (all data)	R1 = 0.0805, wR2 = 0.1175	
Absolute structure parameter	1(2)	
Extinction coefficient	0.016(3)	
Largest diff. peak and hole	0.141 and -0.133 e.Å ⁻³	

3. Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$)for **2c**. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U_{ij} tensor.

	x	y	z	$U(\text{eq})$
O(1)	7808(4)	4212(2)	-1292(1)	53(1)
O(2)	2903(3)	3638(1)	3312(1)	43(1)
C(1)	6047(7)	4794(2)	-1606(2)	58(1)
C(2)	7443(5)	3862(2)	-569(2)	41(1)
C(3)	5496(6)	4046(2)	-139(2)	47(1)
C(4)	5293(6)	3652(2)	576(2)	48(1)
C(5)	7001(5)	3084(2)	882(2)	39(1)
C(6)	8934(5)	2905(2)	432(2)	42(1)
C(7)	9147(6)	3284(2)	-284(2)	45(1)
C(8)	6761(5)	2693(2)	1681(2)	41(1)
C(9)	4428(6)	2251(2)	1812(2)	42(1)
C(10)	3251(5)	2316(2)	2459(2)	45(1)
C(11)	4198(5)	2794(2)	3158(2)	40(1)
C(12)	6706(5)	3122(2)	3092(2)	38(1)
C(13)	7204(5)	3451(2)	2285(2)	42(1)
C(14)	6822(6)	3902(2)	3692(2)	46(1)
C(15)	4280(6)	4179(2)	3853(2)	43(1)
C(16)	3754(7)	5205(2)	3684(2)	60(1)
C(17)	3543(6)	3928(3)	4652(2)	57(1)

4. Table 3. Bond lengths [Å] and angles [°] for 2c.

O(1)-C(2)	1.380(3)
O(1)-C(1)	1.419(4)
O(2)-C(11)	1.439(3)
O(2)-C(15)	1.456(3)
C(2)-C(7)	1.373(4)
C(2)-C(3)	1.377(5)
C(3)-C(4)	1.379(4)
C(4)-C(5)	1.382(4)
C(5)-C(6)	1.388(4)
C(5)-C(8)	1.516(4)
C(6)-C(7)	1.374(4)
C(8)-C(9)	1.500(4)
C(8)-C(13)	1.536(4)
C(9)-C(10)	1.326(4)
C(10)-C(11)	1.505(4)
C(11)-C(12)	1.521(4)
C(12)-C(13)	1.520(4)
C(12)-C(14)	1.531(4)
C(14)-C(15)	1.541(4)
C(15)-C(17)	1.510(4)
C(15)-C(16)	1.521(4)
C(2)-O(1)-C(1)	117.4(2)
C(11)-O(2)-C(15)	106.4(2)
C(7)-C(2)-C(3)	119.7(3)
C(7)-C(2)-O(1)	116.3(3)
C(3)-C(2)-O(1)	124.0(3)
C(4)-C(3)-C(2)	119.4(3)
C(3)-C(4)-C(5)	122.1(3)
C(4)-C(5)-C(6)	117.1(3)
C(4)-C(5)-C(8)	120.7(3)
C(6)-C(5)-C(8)	122.2(3)
C(7)-C(6)-C(5)	121.4(3)
C(6)-C(7)-C(2)	120.3(3)
C(9)-C(8)-C(5)	112.2(2)
C(9)-C(8)-C(13)	109.6(2)

C(5)-C(8)-C(13)	111.5(2)
C(10)-C(9)-C(8)	124.1(3)
C(9)-C(10)-C(11)	123.0(3)
O(2)-C(11)-C(10)	110.1(2)
O(2)-C(11)-C(12)	104.5(2)
C(10)-C(11)-C(12)	114.9(3)
C(11)-C(12)-C(13)	110.1(2)
C(11)-C(12)-C(14)	102.2(2)
C(13)-C(12)-C(14)	114.2(2)
C(12)-C(13)-C(8)	113.4(2)
C(12)-C(14)-C(15)	105.7(2)
O(2)-C(15)-C(17)	109.2(3)
O(2)-C(15)-C(16)	105.7(2)
C(17)-C(15)-C(16)	110.7(3)
O(2)-C(15)-C(14)	105.2(2)
C(17)-C(15)-C(14)	112.1(3)
C(16)-C(15)-C(14)	113.5(3)

Symmetry transformations used to generate equivalent atoms:

5. Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 2c.

The anisotropic displacement factor exponent takes the form: $-2 \left[h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12} \right]$

	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
O(1)	65(2)	52(1)	43(1)	4(1)	0(1)	9(1)
O(2)	34(1)	46(1)	48(1)	-7(1)	-4(1)	5(1)
C(1)	76(3)	49(2)	49(2)	7(2)	-7(2)	3(2)
C(2)	49(2)	37(2)	35(2)	-3(1)	-2(2)	1(1)
C(3)	46(2)	50(2)	46(2)	3(2)	-5(2)	11(2)
C(4)	44(2)	55(2)	44(2)	-2(2)	5(2)	11(2)
C(5)	44(2)	37(2)	37(2)	-4(1)	-3(1)	7(1)
C(6)	41(2)	43(2)	44(2)	-5(1)	-4(2)	6(2)
C(7)	44(2)	50(2)	41(2)	0(2)	6(2)	6(2)
C(8)	40(2)	41(2)	43(2)	-5(1)	-2(1)	6(2)
C(9)	49(2)	39(2)	38(2)	-6(1)	-7(2)	1(2)
C(10)	39(2)	41(2)	55(2)	-5(2)	-7(2)	-3(1)
C(11)	37(2)	41(2)	41(2)	1(1)	1(1)	2(2)
C(12)	33(2)	42(2)	40(2)	-1(1)	-4(1)	4(1)
C(13)	38(2)	46(2)	41(2)	2(1)	-1(1)	-2(1)
C(14)	43(2)	55(2)	40(2)	-2(2)	-3(2)	-3(2)
C(15)	47(2)	43(2)	40(2)	-6(1)	-5(2)	1(2)
C(16)	70(2)	50(2)	61(2)	-3(2)	-15(2)	1(2)
C(17)	60(2)	70(2)	42(2)	-5(2)	3(2)	-4(2)

IV. CIF Format of X-ray Diffraction Analysis of Compound 2c

```
#=====
data_global
#=====
```

1. SUBMISSION DETAILS

_publ_contact_author #Name and address of author for correspondence

Professor Ming-Chang P. Yeh
Department of Chemistry
National Taiwan Normal University
88, Ding-Jou Road, Section 4
Taipei, Taiwan

_publ_contact_author_phone 886-2-29309080
_publ_contact_author_fax 886-6-29324249
_publ_contact_author_email cheyeh@scc.ntnu.edu.tw

_publ_requested_journal *Organometallics.*
_publ_requested_coeditor_name ?

_publ_contact_letter
;
? #Insert text if wanted
;

```
#=====
```

2. PROCESSING SUMMARY (IUCr Office Use Only)

_journal_date_recd_electronic ?
_journal_date_to_coeditor ?
_journal_date_from_coeditor ?

_journal_date_accepted ?
 _journal_date_printers_first ?
 _journal_date_printers_final ?
 _journal_date_proofs_out ?
 _journal_date_proofs_in ?
 _journal_coeditor_name ?
 _journal_coeditor_code ?
 _journal_coeditor_notes
 ;?
 ;
 _journal_techeditor_code ?
 _journal_techeditor_notes
 ;?
 ;
 _journal_coden_ASTM ?
 _journal_name_full ?
 _journal_year ?
 _journal_volume ?
 _journal_issue ?
 _journal_page_first ?
 _journal_page_last ?
 _journal_suppl_publ_number ?
 _journal_suppl_publ_pages ?

#=====

 =====

3. TITLE AND AUTHOR LIST

(±)-(3aS,5S,7aS)-5-(4-Methoxyphenyl)-2,2-dimethyl-2,3,3a,4,5,7a-hexahydro-
 benzo[*b*]furan(2c).

2c

Yeh, Ming-Chang P. *

Tsao, Wen-Cheng

Tu, Ling-Hsien

Department of Chemistry, National Taiwan Normal University, 88 Ding-Jou Road,
Section 4, Taipei 117, Taiwan

#

4. TEXT

_publ_section_abstract

;

? #Insert appropriate text here if wanted.

;

_publ_section_comment

;

? #Insert appropriate text here if wanted.

;

_publ_section_experimental

;

? #Insert appropriate text here if wanted.

;

_publ_section_references

;

Gabe, E.J., Le Page, Y., Charland, J.-P., Lee, F.L. and White, P.S.
(1989) J. Appl. Cryst., 22, 384-387.

Flack, H., (1983) Acta Cryst., A39, 876-881.

International Tables for X-ray Crystallography, Vol. IV, (1974)
Kynoch Press, Birmingham, England.

Johnson, C.K., (1976) ORTEP - A Fortran Thermal Ellipsoid Plot
Program, Technical Report ORNL-5138, Oak Ridge National Lab., U.S.A.

Larson, A.C., (1970) p.293, Crystallographic Computing, Munksgaard,
Copenhagen.

Le Page, Y., (1988) J. Appl. Cryst., 21, 983-984.

Le Page, Y. and Gabe, E.J., (1979) J. Appl. Cryst., 12, 464-466.

Rogers, D., (1981) Acta Cryst., A37, 734-741.

? #Insert any other references here

;

_publ_section_figure_captions

;

? #Insert figure captions here if wanted

;

_publ_section_acknowledgements

;

? #Insert any acknowledgements here.

;

=====

If more than one structure is reported, sections 5-10 should be filled in
per structure. For each data set, replace the ? in the data_? line below
by a unique identifier.

data_?

=====

5. CHEMICAL DATA

_audit_creation_method SHELXL-97

_chemical_name_systematic

;

**(±)-(3a*S*,5*S*,7a*S*)-5-(4-Methoxyphenyl)-2,2-dimethyl-2,3,3a,4,5,7a-hexahydrobenz
o[*b*]furan(2c).**

?

;

_chemical_name_common ?

_chemical_melting_point 76-77 °C ?

_chemical_formula_moiety 'C17 H12 O2'

_chemical_formula_sum 'C17 H22 O2'

_chemical_formula_weight 258.36

loop_

_atom_type_symbol

_atom_type_description

_atom_type_scatter_dispersion_real

_atom_type_scatter_dispersion_imag

_atom_type_scatter_source

'C' 'C' 0.0033 0.0016

'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

'H' 'H' 0.0000 0.0000

'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

'O' 'O' 0.0106 0.0060

'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

CRYSTAL DATA

_symmetry_cell_setting orthorhombic

_symmetry_space_group_name_H-M 'P 21 c n'

loop_

_symmetry_equiv_pos_as_xyz

'x, y, z'

'x+1/2, -y, -z'

'x, -y+1/2, z+1/2'

'x+1/2, y+1/2, -z+1/2'

_cell_length_a	5.7550(2)
_cell_length_b	14.2250(5)
_cell_length_c	17.5650(8)
_cell_angle_alpha	90.00
_cell_angle_beta	90.00
_cell_angle_gamma	90.00
_cell_volume	1437.96(10)
_cell_formula_units_Z	4
_cell_measurement_temperature	200(2)
_cell_measurement_reflns_used	7670
_cell_measurement_theta_min	2.037
_cell_measurement_theta_max	25.028

_exptl_crystal_description	prism
_exptl_crystal_colour	colorless
_exptl_crystal_size_max	0.26
_exptl_crystal_size_mid	0.14
_exptl_crystal_size_min	0.1
_exptl_crystal_density_meas	?
_exptl_crystal_density_diffm	1.193
_exptl_crystal_density_method	'not measured'
_exptl_crystal_F_000	560


```

_exptl_absorpt_coefficient_mu    0.076
_exptl_absorpt_correction_type    multi-scan
_exptl_absorpt_correction_T_min    0.948
_exptl_absorpt_correction_T_max    1.0171
_exptl_absorpt_process_details

```

```
;
```

7. EXPERIMENTAL DATA

```

_exptl_special_details
_diffm_ambient_temperature    200(2)
_diffm_radiation_wavelength    0.71073
_diffm_radiation_type    MoK $\alpha$ 
_diffm_radiation_source    'fine-focus sealed tube'
_diffm_radiation_monochromator    graphite
_diffm_measurement_device_type    KappaCCD
_diffm_measurement_method    'CCD rotation images, thick slices'
_diffm_detector_area_resol_mean    ?
_diffm_standards_number    ?
_diffm_standards_interval_count    ?
_diffm_standards_interval_time    ?
_diffm_standards_decay_%    ?
_diffm_reflns_number    7793
_diffm_reflns_av_R_equivalents    0.0672
_diffm_reflns_av_sigmaI/netI    0.0749
_diffm_reflns_limit_h_min    -5
_diffm_reflns_limit_h_max    6
_diffm_reflns_limit_k_min    -15
_diffm_reflns_limit_k_max    16
_diffm_reflns_limit_l_min    -18
_diffm_reflns_limit_l_max    20
_diffm_reflns_theta_min    2.32
_diffm_reflns_theta_max    25.01
_reflns_number_total    2432
_reflns_number_gt    1712
_reflns_threshold_expression    >2sigma(I)

_computing_data_collection    'Collect (Nonius BV, 1997-2000)'
_computing_cell_refinement    'HKL Scalepack (Otwinowski & Minor 1997)'

```

```

_computing_data_reduction      'HKL Denzo and Scalepack (Otwinowski & Minor 19'
_computing_structure_solution  'SHELXS-97 (Sheldrick, 1997)'
_computing_structure_refinement 'SHELXL-97 (Sheldrick, 1997)'
_computing_molecular_graphics  'Ortep-3 for Windows (Farrugia, 1997)'
_computing_publication_material 'WinGX publication routines (Farrugia, 1999)'

```

_#8. REFINEMENT DATA

```
;
```

Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

```
;
```

```

_refine_ls_structure_factor_coef  Fsqd
_refine_ls_matrix_type            full
_refine_ls_weighting_scheme       calc
_refine_ls_weighting_details
'calc w=1/[\s^2*(Fo^2)+(0.0546P)^2+0.1923P] where P=(Fo^2+2Fc^2)/3'
_atom_sites_solution_primary      direct
_atom_sites_solution_secondary    difmap
_atom_sites_solution_hydrogens    geom
_refine_ls_hydrogen_treatment     mixed
_refine_ls_extinction_method       SHELXL
_refine_ls_extinction_coef         0.016(3)
_refine_ls_extinction_expression
'Fc^*=kFc[1+0.001xFc^2\^3/sin(2\q)]^-1/4^'
_refine_ls_abs_structure_details
'Flack H D (1983), Acta Cryst. A39, 876-881'
_refine_ls_abs_structure_Flack     1(2)
_refine_ls_number_reflns           2432
_refine_ls_number_parameters       173
_refine_ls_number_restraints       1
_refine_ls_R_factor_all             0.0805
_refine_ls_R_factor_gt             0.0449

```

_refine_ls_wR_factor_ref	0.1175
_refine_ls_wR_factor_gt	0.0982
_refine_ls_goodness_of_fit_ref	0.971
_refine_ls_restrained_S_all	0.971
_refine_ls_shift/su_max	0.001
_refine_ls_shift/su_mean	0.000

9. ATOMIC COORDINATES AND THERMAL PARAMETERS

loop_

```

_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_occupancy
_atom_site_symmetry_multiplicity
_atom_site_calc_flag
_atom_site_refinement_flags
_atom_site_disorder_assembly
_atom_site_disorder_group

```

O1 O 0.7808(4) 0.42122(15) -0.12916(11) 0.0532(6) Uani 1 1 d ...

O2 O 0.2903(3) 0.36384(14) 0.33121(11) 0.0425(5) Uani 1 1 d ...

C1 C 0.6047(7) 0.4794(2) -0.16055(18) 0.0581(10) Uani 1 1 d ...

H1A H 0.6510 0.4985 -0.2107 0.070 Uiso 1 1 d R ..

H1B H 0.5847 0.5340 -0.1290 0.070 Uiso 1 1 d R ..

H1C H 0.4608 0.4454 -0.1633 0.070 Uiso 1 1 d R ..

C2 C 0.7443(5) 0.3862(2) -0.05686(16) 0.0405(7) Uani 1 1 d ...

C3 C 0.5496(6) 0.4046(2) -0.01389(17) 0.0471(8) Uani 1 1 d ...

H3 H 0.4287 0.4429 -0.0353 0.057 Uiso 1 1 d R ..

C4 C 0.5293(6) 0.3652(2) 0.05755(17) 0.0476(8) Uani 1 1 d ...

H4 H 0.3949 0.3798 0.0877 0.057 Uiso 1 1 d R ..

C5 C 0.7001(5) 0.3084(2) 0.08818(16) 0.0392(7) Uani 1 1 d ...

C6 C 0.8934(5) 0.2905(2) 0.04318(17) 0.0424(8) Uani 1 1 d ...

H6 H 1.0139 0.2501 0.0622 0.051 Uiso 1 1 d R ..

C7 C 0.9147(6) 0.3284(2) -0.02842(16) 0.0450(8) Uani 1 1 d ...

H7 H 1.0495 0.3163 -0.0591 0.054 Uiso 1 1 d R ..

C8 C 0.6761(5) 0.2693(2) 0.16805(16) 0.0412(7) Uani 1 1 d . . .
 H8 H 0.7907 0.2212 0.1754 0.049 Uiso 1 1 d R . .
 C9 C 0.4428(6) 0.2251(2) 0.18124(17) 0.0419(7) Uani 1 1 d . . .
 H9 H 0.3705 0.1906 0.1406 0.050 Uiso 1 1 d R . .
 C10 C 0.3251(5) 0.2316(2) 0.24589(18) 0.0447(8) Uani 1 1 d . . .
 H10 H 0.1721 0.2048 0.2470 0.054 Uiso 1 1 d R . .
 C11 C 0.4198(5) 0.2794(2) 0.31575(17) 0.0396(7) Uani 1 1 d . . .
 H11 H 0.4055 0.2348 0.3564 0.047 Uiso 1 1 d R . .
 C12 C 0.6706(5) 0.3122(2) 0.30922(16) 0.0382(7) Uani 1 1 d . . .
 H12 H 0.7778 0.2627 0.3213 0.046 Uiso 1 1 d R . .
 C13 C 0.7204(5) 0.3451(2) 0.22851(16) 0.0415(8) Uani 1 1 d . . .
 H13A H 0.8790 0.3656 0.2256 0.050 Uiso 1 1 d R . .
 H13B H 0.6190 0.3972 0.2179 0.050 Uiso 1 1 d R . .
 C14 C 0.6822(6) 0.3902(2) 0.36916(17) 0.0460(8) Uani 1 1 d . . .
 H14A H 0.7584 0.3681 0.4144 0.055 Uiso 1 1 d R . .
 H14B H 0.7687 0.4416 0.3480 0.055 Uiso 1 1 d R . .
 C15 C 0.4280(6) 0.4179(2) 0.38527(16) 0.0433(8) Uani 1 1 d . . .
 C16 C 0.3754(7) 0.5205(2) 0.36837(19) 0.0600(10) Uani 1 1 d . . .
 H16A H 0.4611 0.5600 0.4027 0.072 Uiso 1 1 d R . .
 H16B H 0.2120 0.5312 0.3751 0.072 Uiso 1 1 d R . .
 H16C H 0.4182 0.5349 0.3169 0.072 Uiso 1 1 d R . .
 C17 C 0.3543(6) 0.3928(3) 0.46521(17) 0.0571(9) Uani 1 1 d . . .
 H17A H 0.4422 0.4283 0.5016 0.069 Uiso 1 1 d R . .
 H17B H 0.3820 0.3269 0.4729 0.069 Uiso 1 1 d R . .
 H17C H 0.1918 0.4057 0.4717 0.069 Uiso 1 1 d R . .

loop_

_atom_site_aniso_label
 _atom_site_aniso_U_11
 _atom_site_aniso_U_22
 _atom_site_aniso_U_33
 _atom_site_aniso_U_23
 _atom_site_aniso_U_13
 _atom_site_aniso_U_12

O1 0.0648(15) 0.0516(13) 0.0432(13) 0.0040(10) -0.0004(12) 0.0088(12)
 O2 0.0336(10) 0.0456(12) 0.0483(12) -0.0075(9) -0.0044(10) 0.0046(10)
 C1 0.076(3) 0.049(2) 0.049(2) 0.0068(16) -0.0074(18) 0.0030(18)
 C2 0.049(2) 0.0372(17) 0.0349(17) -0.0031(13) -0.0015(15) 0.0005(14)

C3 0.0456(18) 0.050(2) 0.0462(19) 0.0026(15) -0.0047(17) 0.0110(16)
 C4 0.044(2) 0.055(2) 0.044(2) -0.0016(16) 0.0053(16) 0.0110(16)
 C5 0.0438(18) 0.0372(16) 0.0366(18) -0.0042(13) -0.0028(14) 0.0073(14)
 C6 0.0407(18) 0.0431(18) 0.0435(19) -0.0046(14) -0.0042(15) 0.0059(15)
 C7 0.0435(18) 0.0501(19) 0.0414(18) -0.0002(15) 0.0060(16) 0.0059(16)
 C8 0.0398(18) 0.0405(18) 0.0433(18) -0.0045(14) -0.0021(14) 0.0056(15)
 C9 0.0489(19) 0.0392(17) 0.0377(18) -0.0061(14) -0.0068(16) 0.0005(15)
 C10 0.0385(18) 0.0410(18) 0.055(2) -0.0048(15) -0.0075(15) -0.0026(14)
 C11 0.0372(17) 0.0406(17) 0.0408(17) 0.0010(14) 0.0005(14) 0.0025(15)
 C12 0.0333(16) 0.0416(17) 0.0399(18) -0.0005(14) -0.0041(13) 0.0042(14)
 C13 0.0376(18) 0.0458(18) 0.0412(18) 0.0015(14) -0.0009(14) -0.0016(14)
 C14 0.0427(18) 0.055(2) 0.0400(18) -0.0020(15) -0.0030(15) -0.0029(16)
 C15 0.0473(19) 0.0431(19) 0.0395(18) -0.0056(14) -0.0050(15) 0.0009(16)
 C16 0.070(2) 0.050(2) 0.061(2) -0.0034(17) -0.0151(18) 0.0011(18)
 C17 0.060(2) 0.070(2) 0.0423(19) -0.0052(17) 0.0033(16) -0.0040(17)

10. MOLECULAR GEOMETRY

;

All esds (except the esd in the dihedral angle between two l.s. planes)
 are estimated using the full covariance matrix. The cell esds are taken
 into account individually in the estimation of esds in distances, angles
 and torsion angles; correlations between esds in cell parameters are only
 used when they are defined by crystal symmetry. An approximate (isotropic)
 treatment of cell esds is used for estimating esds involving l.s. planes.

;

loop_

_geom_bond_atom_site_label_1

_geom_bond_atom_site_label_2

_geom_bond_distance

_geom_bond_site_symmetry_2

_geom_bond_publ_flag

O1 C2 1.380(3) . ?

O1 C1 1.419(4) . ?

O2 C11 1.439(3) . ?

O2 C15 1.456(3) . ?

C2 C7 1.373(4) . ?

C2 C3 1.377(5) . ?

C3 C4 1.379(4) . ?
 C4 C5 1.382(4) . ?
 C5 C6 1.388(4) . ?
 C5 C8 1.516(4) . ?
 C6 C7 1.374(4) . ?
 C8 C9 1.500(4) . ?
 C8 C13 1.536(4) . ?
 C9 C10 1.326(4) . ?
 C10 C11 1.505(4) . ?
 C11 C12 1.521(4) . ?
 C12 C13 1.520(4) . ?
 C12 C14 1.531(4) . ?
 C14 C15 1.541(4) . ?
 C15 C17 1.510(4) . ?
 C15 C16 1.521(4) . ?

loop_

_geom_angle_atom_site_label_1
 _geom_angle_atom_site_label_2
 _geom_angle_atom_site_label_3
 _geom_angle
 _geom_angle_site_symmetry_1
 _geom_angle_site_symmetry_3
 _geom_angle_publ_flag

C2 O1 C1 117.4(2) . . ?
 C11 O2 C15 106.4(2) . . ?
 C7 C2 C3 119.7(3) . . ?
 C7 C2 O1 116.3(3) . . ?
 C3 C2 O1 124.0(3) . . ?
 C4 C3 C2 119.4(3) . . ?
 C3 C4 C5 122.1(3) . . ?
 C4 C5 C6 117.1(3) . . ?
 C4 C5 C8 120.7(3) . . ?
 C6 C5 C8 122.2(3) . . ?
 C7 C6 C5 121.4(3) . . ?
 C6 C7 C2 120.3(3) . . ?
 C9 C8 C5 112.2(2) . . ?
 C9 C8 C13 109.6(2) . . ?

C5 C8 C13 111.5(2) . . ?
 C10 C9 C8 124.1(3) . . ?
 C9 C10 C11 123.0(3) . . ?
 O2 C11 C10 110.1(2) . . ?
 O2 C11 C12 104.5(2) . . ?
 C10 C11 C12 114.9(3) . . ?
 C11 C12 C13 110.1(2) . . ?
 C11 C12 C14 102.2(2) . . ?
 C13 C12 C14 114.2(2) . . ?
 C12 C13 C8 113.4(2) . . ?
 C12 C14 C15 105.7(2) . . ?
 O2 C15 C17 109.2(3) . . ?
 O2 C15 C16 105.7(2) . . ?
 C17 C15 C16 110.7(3) . . ?
 O2 C15 C14 105.2(2) . . ?
 C17 C15 C14 112.1(3) . . ?
 C16 C15 C14 113.5(3) . . ?

_diffm_measured_fraction_theta_max 0.999

_diffm_reflms_theta_full 25.01

_diffm_measured_fraction_theta_full 0.999

_refine_diff_density_max 0.141

_refine_diff_density_min -0.133

_refine_diff_density_rms 0.034

#==**END**