Palladium(II)-Catalyzed Enantioselective Aerobic Dialkoxylation of 2-Propenylphenols: A Pronounced Effect of Copper Additives on Enantioselectivity

Yang Zhang and Matthew S. Sigman*

Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, UT 84112-8500

Supporting Information

Table of Contents

General Information	S2
Preparation of 2-propenylphenol Derivatives	S2
Dialkoxylation of 2-propenylphenol using Pd[(-)-Sparteine]Cl ₂ and CuCl ₂	S2
Dialkoxylation of 2-propenylphenol using Pd[(-)-Sparteine]Cl ₂ without CuCl ₂	S2
Preparation of Quinoline Oxazoline Derivatives	S2
Screening of Chiral Ligands in the Dialkoxylation of 2-Propenylphenol	S3
Study of Copper Effect on the Enantioselectivity of the Dialkoxylation of 2-Propenylphenol	S3
Preparation of 1-((1S,2S)-1,2-dimethoxypropyl)-2-methoxybenzene (Determination of Product Absolute Configuration)	S4-S5
General Procedure of Enantioselective Dialkoxylation of 2-Propenylphenol	S6
Characterization Data	S6
Chiral Separations	S7-S18
¹ H- and ¹³ C-NMR Spectra	S18-S30
References	S31

General Information Dichloromethane and dichloroethane were distilled from CaH_2 . Toluene was dried before use by passing through a column of activated alumina. Methanol was distilled from magnesium methoxide. THF was distilled from sodium benzophenone ketyl. The 3 Å molecular sieves were powdered and activated by putting under vacuum in a flask and heating with a Bunsen burner (ca. 1 minute). ^1H -NMR spectra were obtained at 300 MHz and referenced to the CHCl_3 singlet at 7.26 ppm. ^{13}C -NMR spectra were obtained at 75 MHz and referenced to the center line of the CDCl_3 triplet at 77.2 ppm. All melting points were recorded on an electrothermal melting point apparatus. IR spectra were recorded using Bruker Tensor 27. MS data were obtained on a Micromass Quattro II. GC/MS data were obtained on an HP 5890 (EI).

Preparation of 2-propenylphenol Derivatives:¹ To a 250 mL flame dried round bottom flask equipped with a stir bar was placed 2.56 g of ethyltriphenylphosphonium bromide (6.9 mmol, 2.3 equiv.) and 40 mL of toluene under a nitrogen atmosphere. To this was added a solution of 785 mg KO^tBu (7 mmol, 2.33 equiv.) in 10 mL of THF dropwise via cannulation. The reaction mixture slowly turned a deep red color over 4 hours. The mixture was cooled to -78 °C and 3 mmol of aldehyde (1 equiv.), dissolved in 8 mL of toluene, was added dropwise via cannulation. The mixture was allowed to slowly warm to ambient temperature and stirred 12 hours then quenched with 20 mL of saturated NH_4Cl solution. The mixture was diluted with an additional 40 mL of H_2O and extracted 3 times with 20 mL each of diethyl ether. The organic extracts were combined, washed once with 15 mL brine, dried over MgSO_4 , filtered, and the solvent removed in vacuo. The product was purified by bulb to bulb distillation. The ^1H -NMR spectra of the products were compared to previously reported ^1H -NMR spectra.²

Dialkoxylation of 2-propenylphenol using $\text{Pd}[(+)\text{-Sparteine}]\text{Cl}_2$ and CuCl_2 : To a 5 mL side-arm round bottom flask equipped with a stir bar was added 4.1 mg $\text{Pd}[(+)\text{-sparteine}]\text{Cl}_2$ (0.01 mmol, 0.1 equiv.), 2.6 mg of CuCl_2 (0.02 mmol, 0.2 equiv.), 50 mg of freshly activated, crushed 3 Å molecular sieves, and 800 μL methanol. A three-way joint fitted with a balloon of O_2 was attached. The mixture was stirred for 20 minutes at room temperature under balloon of O_2 . To the reaction mixture, a solution of 0.1 mmol of 2-propenyl phenol and 5 μL 5-nonanone internal standard dissolved in 200 μL methanol was added dropwise. Aliquot of the reaction was taken periodically. The aliquot was passed through a short silica plug eluting with 2 mL of EtOAc. The sample was then analyzed by GC and referenced against a time zero sample. The response factors of 2-propenyl phenol, 2-(1,2-dimethoxypropyl)phenol were determined to be 1.06 : 1 by ^1H -NMR and GC. The enantiomeric excess of the 2-(1,2-dimethoxypropyl)phenol product were determined by GC using a chiral GTA column (100 °C isothermal).

Dialkoxylation of 2-propenylphenol using $\text{Pd}[(+)\text{-Sparteine}]\text{Cl}_2$ without CuCl_2 : To a 5 mL side-arm round bottom flask equipped with a stir bar was added 4.1 mg $\text{Pd}[(+)\text{-sparteine}]\text{Cl}_2$ (0.01 mmol, 0.1 equiv.), 50 mg of freshly activated, crushed 3 Å molecular sieves, and 800 μL methanol. A three-way joint fitted with a balloon of O_2 was attached. The mixture was stirred for 20 minutes at room temperature under balloon of O_2 . To the reaction mixture, a solution of 0.1 mmol of 2-propenyl phenol and 5 μL 5-nonanone internal standard dissolved in 200 μL methanol was added dropwise. Aliquots of the reaction were taken periodically. The aliquot was passed through a short silica plug eluting with 2 mL of EtOAc. The sample was then analyzed by GC and referenced against a time zero sample. The enantiomeric excess of the 2-(1,2-dimethoxypropyl)phenol product were determined by GC using a chiral GTA column (100 °C isothermal).

Preparation of Quinoline Oxazoline Derivatives: To a 50 mL flame dried round bottom flask equipped with a stir bar was placed 0.346g of quinaldic acid (2 mmol, 1 equiv.) and 18 mL of dichloromethane under a nitrogen atmosphere. Cooled to -5 °C in an ice bath. To this was added 330 μL of n-methylmorpholine (3 mmol, 1.5 equiv.) dropwise via a syringe. After 15 minutes, 301 μL of isobutylchloroformate (2.3 mmol, 1.15 equiv.) was added dropwise over a period of 2 minutes. Stirred for 30 minutes at -5 °C, 2.4 mmol of amino alcohol (1.2 equiv. (S)-valinol, (S)-tert-leucinol, (S)-2-Phenylglycinol and (R)-2-Amino-3-phenyl-1-propanol respectively) and 253 μL of n-methylmorpholine (2.3 mmol, 1.15 equiv.) in dichloromethane was added dropwise via a syringe. The reaction solution was warmed up to room temperature after 1 hour and stirred for 12 hours. Diluting the reaction solution with 50 mL dichloromethane and washed once with 2 mL of 1 M HCl, 10 mL of H_2O , 10 mL of brine, dried over

Na₂SO₄, filtered and the solvent removed in vacuo. The peptide products were crystallized in Hexane and Et₂O or purified with flash silica-gel column chromatography with Hexane/EtOAc as eluent.

(R)-N-(1-hydroxy-3-phenylpropan-2-yl)quinoline-2-carboxamide: 88% yield of a clear viscous oil. R_f = 0.3 w/1:1 EtOAc/Hexanes. ¹H-NMR (300 MHz CDCl₃) δ = 8.50-8.60 (d, J = 8.0 Hz, 1H), 8.21 (s, 1H), 8.00-8.06 (d, J = 8.2 Hz, 1H), 7.68-7.87 (m, 2H), 7.50-7.62 (m, 1H), 7.18-7.37 (m, 5H), 4.36-4.50 (m, 1H), 3.70-3.92 (m, 2H), 3.39-3.46 (m, 1H), 3.03-3.10 (d, J = 7.1 Hz, 2H), 2.18 (s, 1H); ¹³C-NMR (75 MHz, CDCl₃) δ = 165.3, 150.2, 146.7, 137.9, 137.7, 130.3, 129.9, 129.6, 129.4, 128.9, 128.1, 127.9, 126.8, 118.9, 64.3, 54.5, 37.5. MS (Quattro II) m/z (M)⁺ calcd. 306.1 obsd. 306.3.

To a 50 mL flame dried round bottom flask equipped with a stir bar was placed 0.571 g tosyl chloride (2 mmol, 1.5 equiv.), 0.036 g DMAP (0.3 mmol, 0.1 equiv.), 2 mmol of quinoline peptide, and 20 mL of dichloroethane under a nitrogen atmosphere. To this was added 1123 μ L of triethylamine dropwise via a syringe. The mixture was stirred at room temperature for 3 hours, then heated to reflux for 12 hours. The reaction solution was cooled to room temperature. The reaction solution was diluted with 50 mL dichloromethane and washed once with 10 mL of saturated NaHCO₃, 10 mL of H₂O, 10 mL of brine, dried over Na₂SO₄, filtered and the solvent removed in vacuo. The quinoline oxazoline products were crystallized in Hexane and Et₂O or purified with flash silica-gel column chromatography with Hexane/EtOAc as eluent. The ¹H-NMR spectra of the products were compared to previously reported ¹H-NMR spectra.³

(R)-4-benzyl-2-(quinolin-2-yl)-4,5-dihydrooxazole: 82% yield of a white solid, R_f = 0.5 w/1:1 EtOAc/Hexanes, MP = 59-60°C. IR: 3419, 2923, 2361, 1595, 1457, 1371, 1123, 1087, 841, 766. ¹H-NMR (300 MHz CDCl₃) δ = 8.10-8.37 (m, 3H), 7.66-7.83 (m, 2H), 7.51-7.58 (m, 1H), 7.10-7.40 (m, 5H), 4.62-4.74 (m, 1H), 4.44-4.52 (t, J = 9.5 Hz, 1H), 4.25-4.32 (t, J = 9.5 Hz, 1H), 3.26-3.34 (dd, J = 13.4 Hz, J = 5.4 Hz, 1H), 2.72-2.82 (dd, J = 13.4 Hz, J = 5.4 Hz, 1H); ¹³C-NMR (75 MHz, CDCl₃) δ = 163.9, 147.8, 147.4, 138.1, 137.0, 130.6, 129.4, 128.9, 128.7, 128.2, 128.0, 126.7, 121.1, 73.2, 68.6, 41.9; MS (Quattro II) m/z (M)⁺ calcd. 288.4 obsd. 288.9.

Screening of Chiral Ligands in the Dialkoxylation of 2-Propenylphenol: To a 5 mL side-arm round bottle flask equipped with a stir bar was added 2.6 mg Pd(MeCN)₂Cl₂ (0.01 mmol, 0.1 equiv.), 0.012 mmol chiral ligand (0.12 equiv) and 800 μ L methanol, and stirred for 20 minutes. Freshly activated, crushed 3Å molecular sieves was added (50 mg), then a three-way joint fitted with a balloon of O₂ was attached and the apparatus was evacuated and refilled with oxygen three times. The mixture was stirred vigorously for 20 min at room temperature under O₂. At this time, a solution of 0.1 mmol of 2-propenylphenol and 4 μ L 5-nonanone internal standard dissolved in 200 μ L methanol was added dropwise. Aliquots of the reaction were taken periodically. The aliquot was passed through a short silica plug eluting with 2 mL of EtOAc. The sample was then analyzed by GC and referenced against a time zero sample. The enantiomeric excess of the 2-(1,2-dimethoxypropyl)phenol product were determined by GC using a chiral GTA column (100°C isothermal).

Study of Copper Effect on the Enantioselectivity of the Dialkoxylation of 2-Propenylphenol: To four individual 5 mL side-arm round bottle flasks was added 2.6 mg Pd(MeCN)₂Cl₂ (0.01 mmol, 0.1 equiv.), 2.9 mg iPr-quinox ligand (0.012 mmol, 0.12 equiv), various amount of CuCl₂ (0.0025 mmol to 0.02 mmol, 0.025 equiv. to 0.2 equiv), and 800 μ L methanol, respectively. Stirred for 20 minutes. 50 mg of freshly activated, crushed 3Å molecular sieves was added to each of the flask. The flasks were attached to a four neck cow. A three-way joint fitted with a balloon of O₂ was attached and the apparatus was evacuated and refilled with oxygen three times. The mixture was stirred vigorously for 20 minutes at room temperature under O₂. At this time, a solution of 0.1 mmol of 2-propenyl phenol and 5 μ L 5-nonanone internal standard dissolved in 200 μ L methanol was added dropwise to each of the flasks. Aliquots of the reaction were taken periodically. The aliquot was passed through a short silica plug eluting with 2 mL of EtOAc. The sample was then analyzed using a GC and referenced against a time zero sample. The enantiomeric excess of the 2-(1,2-dimethoxypropyl)phenol product were determined by GC using a chiral GTA column (100°C isothermal).

Determination of the Absolute Configuration of Product 1: Preparation of 1-((1*S*,2*S*)-1,2-dimethoxypropyl)-2-methoxybenzene.

To a stirred suspension of 830 mg of AD-mix α in 4 mL of H_2O and 3 mL of $t\text{BuOH}$ in a 25 mL round bottom flask at 0°C was added 90 mg of 1-methoxy-2-(prop-1-en-1-yl)benzene (0.6 mmol, 1 equiv.) in 1 mL $t\text{BuOH}$. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 hours. At this time, 880 mg of Na_2SO_3 (6.6 mmol, 11 equiv.) was added and the reaction mixture was stirred for 1 hour. The mixture was diluted with 30 mL of H_2O and extracted three times with 10 mL of EtOAc. The organic layers were combined and dried over Na_2SO_4 , filtered, and solvent removed in vacuo. The dihydroxylated product **1y** was purified by column chromatography eluting with 40% EtOAc/hexanes. The resulting product was dissolved in 3 mL of THF and 80.8 mg of NaH (60% in mineral oil) (2.02 mmol, 4 equiv.) was added in portions. The mixture was allowed to stir for 10 minutes then 252 μL of MeI (4.04 mmol, 8 equiv.) was added dropwise and the mixture was stirred for 3 hours followed by quenching with 25 mL of H_2O . The aqueous layer was extracted three times w/ 10 mL of EtOAc. The combined organic layers were dried over Na_2SO_4 , filtered, and solvent removed. The product **1z** was purified by column chromatography eluting with 2-10% Et₂O/hexanes afforded a clean oil product (45mg): $[\alpha]_D^{+105.6}$ (c 0.25, CHCl_3). The $^1\text{H-NMR}$ spectra of the products were compared to previously reported $^1\text{H-NMR}$ spectra.^{2a} The absolute configuration was assigned to be (*S,S*) by applying the mnemonic for styrene derivatives developed by Sharpless and others.⁴

Compound **1a** (Table 2, entry 1) was methylated (NaH, MeI) and the retention time of product **1z** was compared to the retention time of the product **1z** prepared using the above method (see following HPLC traces).

Chiral separation condition: HPLC analysis (chiral OD, hexanes/*i*PrOH = 98/2, flow rate = 0.5 mL/min).

a. Racemic **1z** prepared by methylation of racemic **1a**.

b. Product **1z** prepared by enantioselective dialkoxylation using (*R*)-Bn-quinox followed by methylation.

c. Product **1z** prepared by Sharpless dihydroxylation followed by methylation of **1y**.

General Procedure of Enantioselective Dialkoxylation of 2-Propenylphenol: To a flame dried 100 mL schlenk flask equipped with a stir bar was added 13.0 mg Pd(MeCN)₂Cl₂ (0.05 mmol, 0.1 equiv.), 17.3 mg (R)-Bn-quinox ligand (0.06 mmol, 0.12 equiv), and 4 mL methanol, and stirred vigorously for 20 minutes. Freshly activated, crushed 3Å molecular sieves (250 mg) was added. A three-way joint fitted with a balloon of O₂ was attached and the apparatus was evacuated and refilled with oxygen three times. The mixture was stirred vigorously for 20 minutes at room temperature under O₂. At this time, a solution of 0.5 mmol of propenylphenol dissolved in 1 mL methanol was added dropwise. After 48 hours, the reaction mixture was passed through a short plug of silica eluting with 50 mL of 1:1 Et₂O/Hexanes. The solvent was removed in vacuo and the residue was purified via flash silica-gel column chromatography (dry loaded; 1:10 to 1:5 EtOAc/hexanes was used as eluent). The ¹H-NMR spectra of the products **1a**, **3a**, **4a**, **5a**, **9a** and **10a** were compared to previously reported ¹H-NMR spectra.^{2a}

Characterization Data:

2-((1S,2S)-1,2-dimethoxypropyl)-4-methoxyphenol (2a). Clear oil, *R*_f=0.3, 1:3 EtOAc/Hexanes, clear oil, IR: 3375, 2978, 2936, 2831, 2361, 1623, 1587, 1510, 1459, 1107, 1037, 959, 839, 734; ¹H-NMR (500 Mhz CDCl₃): δ 7.55 (s, 1H), 6.72-6.83 (m, 2H), 6.58-6.65 (m, 1H), 4.23-4.28 (d, *J* = 4.9 Hz, 1H), 3.77 (s, 3H), 3.70-3.75 (m, 1H), 3.40 (s, 3H), 3.36 (s, 3H), 1.03-1.08 (d, *J* = 6.3 Hz, 3H); ¹³C-NMR (75 MHz CDCl₃): δ 153.2, 149.4, 124.4, 118.1, 114.7, 114.3, 86.1, 80.2, 58.0, 57.1, 55.9, 15.2; MS (E.I.) *m/z* (M)⁺ calcd. 226.1 obsd. 226.1.

2,4-di-tert-butyl-6-((1R,2R)-1,2-dimethoxypropyl)phenol (6a). Clear oil, *R*_f = 0.45, 1:3 EtOAc/Hexanes, IR: 3343, 2958, 2828, 1743, 1604, 1481, 1460, 1420, 1244, 1087, 912, 825, 769, 702; ¹H-NMR (500 Mhz CDCl₃): δ 8.00 (s, 1H), 7.21-7.23 (d, *J* = 2.5 Hz, 1H), 6.83-6.86 (d, *J* = 2.5 Hz, 1H), 4.20-4.26 (d, *J* = 6.0 Hz, 1H), 3.70-3.80 (m, 1H), 3.42 (s, 3H), 3.9 (s, 3H), 1.41 (s, 9H), 1.28 (s, 9H), 0.98-1.00 (t, *J* = 6.4 Hz, 3H); ¹³C-NMR (75 MHz CDCl₃): δ 152.3, 141.2, 136.6, 124.3, 123.6, 122.1, 88.8, 79.6, 57.8, 57.5, 31.8, 29.8, 15.8; MS (E.I.) *m/z* (M)⁺ calcd. 308.2 obsd. 308.1.

2-((1S,2S)-1,2-dimethoxypropyl)-5-methoxyphenol (7a). Clear oil, *R*_f = 0.3, 1:3 EtOAc/Hexanes, IR: 3381, 262, 2923, 2361, 1597, 1418, 1262, 1071, 801, 734; ¹H-NMR (500 Mhz CDCl₃): δ 8.10 (s, 1H), 6.88-6.95 (d, *J* = 8.2 Hz, 1H), 6.37-6.49 (m, 2H), 4.14-4.20 (d, *J* = 4.7 Hz, 1H), 3.76 (s, 3H), 3.70-3.73 (m, 1H), 3.40 (s, 3H), 3.34 (s, 3H), 1.07-1.11 (d, *J* = 6.3 Hz, 3H); ¹³C-NMR (75 MHz CDCl₃): δ 160.9, 156.9, 130.5, 115.6, 106.1, 102.7, 87.3, 80.4, 57.7, 57.3, 55.4, 15.5; MS (E.I.) *m/z* (M)⁺ calcd. 226.1 obsd. 226.1.

2-((1S,2S)-1,2-dimethoxybutyl)phenol (8a). Clear oil, *R*_f = 0.45, 1:5 EtOAc/Hexanes, IR: 3358, 2969, 2936, 1604, 1487, 1458, 1362, 1285, 1241, 1141, 1082, 929, 851, 755; ¹H-NMR (500 Mhz CDCl₃): δ 8.07 (s, 1H), 7.17-7.24 (m, 1H), 6.98-7.02 (m, 1H), 6.80-6.92 (m, 2H), 4.23-4.28 (d, *J* = 4.4 Hz, 1H), 3.43 (s, 3H), 3.33-3.41 (m, 4H), 1.41-1.58 (m, 2H), 0.88-0.97 (t, *J* = 7.4 Hz, 3H); ¹³C-NMR (75 MHz CDCl₃): δ 155.9, 129.6, 123.2, 119.8, 117.3, 86.7, 86.1, 59.4, 57.9, 23.9, 10.2; MS (E.I.) *m/z* (M)⁺ calcd. 210.1 obsd. 210.1.

Chiral Separations and GC traces: All chiral separations were performed on GC with a chiral GTA column except product **6a** (HPLC, *R,R*-Whelk-OI, hexanes/*i*PrOH = 99.5/0.5, flow rate = 0.5 mL/min). The racemic samples of **2a**, **5a**, **7a**, and **8a** were prepared utilizing racemic *i*Pr-quinox ligand following the standard enantioselective dialkoxylation procedure. The racemic samples of others in Table 1 were obtained from previous work.^{2a}

Table 1. Chiral separation of dialkoxylation products.

Compound	Condition	Retention Times
R = H 1a 100°C, isothermal	anti: 25.4 (major) and 26.4 min syn: 27.7 and 29.7 (major) min
	R = OMe 2a 115°C, isothermal	anti: 84.0 (major) and 85.8 min syn: 95.9 and 99.8 (major) min
	R = Me 3a 100°C, isothermal	anti: 34.0 (major) and 35.0 min syn: 37.4 and 38.4 (major) min
	R = Cl 4a 130°C, isothermal	anti: 16.2 (major) and 16.7 min syn: 18.0 and 18.5 (major) min
5a 100°C, isothermal	anti: 24.3, 25.5 (major) min syn: 27.2 (major), 30.1 min
6a (<i>R,R</i>) Whelk-O1 column	syn: 11.18 (major), 13.02 min
7a 120°C, isothermal	anti: 39.9 (major) and 41.5 min syn: 43.8 and 45.4 (major) min
8a 100°C, isothermal	anti: 59.9 (major) and 62.3 min syn: 66.3 and 68.6 (major) min
9a 95°C, isothermal	anti: 46.4 (major) and 49.1 min syn: 55.1 and 57.6 (major) min
10a 100°C, isothermal	syn: 55.2, 57.5 (major) min anti: 89.0 (major), 92.3 min
11a 110°C, isothermal	23.1 and 24.5 min

a. GC spectrum of racemic product **1a**

b. GC spectrum of product **1a** in Table 2, entry 1. (anti: 85% ee, syn: 85% ee)

a. GC spectrum of racemic product **2a**

b. GC spectrum of product **2a** in Table 2, entry 2. (anti: 89% ee, syn: 92% ee)

a. GC spectrum of racemic product **3a**

b. GC spectrum of product **3a** in Table 2, entry 3. (anti: 90% ee, syn: 91% ee)

a. GC spectrum of racemic product **4a**

b. GC spectrum of product **4a** in Table 2, entry 4. (anti: 85% ee, syn: 86% ee)

a. GC spectrum of racemic product **5a**

b. GC spectrum of product **5a** prepared with *S*-*i*PrQuinox in Table 2 (entry 5). (anti: 73% ee, syn: 83% ee)

a. GC spectrum of racemic product **6a**

b. GC spectrum of product **6a** prepared with *S*-*i*PrQuinox in Table 2 (entry 6). (63% ee)

a. GC spectrum of racemic product **7a**.

b. GC spectrum of product **7a** in Table 2 (entry 6). (anti: 85% ee, syn: 84% ee)

a. GC spectrum of racemic product **8a**.

b. GC spectrum of product **8a** in Table 2 (entry 7). (anti: 81% ee, syn: 82% ee)

a. GC spectrum of racemic product **9a**.

b. GC spectrum of product **9a** in Table 2 (entry 8). (anti: 78% ee, syn: 83% ee)

a. GC spectrum of racemic product **10a**.

b. GC spectrum of product **10a** in Table 2 (entry 9). (anti: 82% ee, syn: 81% ee)

a. GC spectrum of racemic product **11a**.

b. GC spectrum of product **11a** prepared using *R*-BnQuinox ligand.

Reference:

1. Adapted from: Johannes, C. W.; Visser, M. S.; Weatherhead, G. S.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1998**, *120*, 8340.
2. a) Schultz, M. J.; Sigman, M. S. *J. Am. Chem. Soc.* **2006**, *128*, 1461. b) Jimenez, G. L.; Alvarez, C.M.; Munoz, D. M.; Rodriguez, G. I. *Chem. Commun.* **2005**, *21*, 2689. c) Alberola, A.; Ortega, A. G.; Pedrosa, R.; Bragado, J. P.; Amo, J. R. *J. Chem. Soc. Perkin Trans. 1* **1983**, 1209. d) Nishinaga, A.; Iwasaki, H.; Shimizu, T.; Toyoda, Y.; Matsuura, T. *J. Org. Chem.* **1986**, *51*, 2257.
3. Chelucci, G.; Medici, S.; Saba, A. *Tetrahedron: Asymmetry* **1999**, *10*, 543.
4. The absolute configuration can be readily assigned by applying the rule developed by Sharpless for styrene derivatives. See a) Sharpless, K. B.; Amberg, W.; Bennani, Y.; Grispino, G. A.; Hartung, J.; Jeong, K.; Kwong, H.; Morikawa, K.; Wang, Z.; Wu, D.; Zhang, X. *J. Org. Chem.* **1992**, *57*, 2766. b) Spivey, A. C.; Scoriah, N.; Thorpe, S. J. *J. Chem. Educ.* **1999**, *76*, 655. c) Wipf, P.; Hopkins, C. R. *J. Org. Chem.* **2001**, *66*, 3133.