Supporting Information for

Palladium(II)-Catalyzed Carboxylation and Hydroxyl-Carboxylation of Arenes via C–H Bond Cleavage Using Formic Acid as a Carbonyl Source

Fumitoshi Shibahara, Shinsuke Kinoshita, Kyoko Nozaki*

General Method

All manipulations involving air- and/or moisture-sensitive compounds were carried out using the standard Schlenk technique under argon purified by passing through a hot column packed with BASF catalyst R3-11. NMR spectra were recorded in deuteriochloroform on a Varian Mercury 200 (^1H 200 MHz; ^{13}C 50 MHz) or JEOL JNM-ECP500 (^1H 500 MHz; ^{13}C 125 MHz) spectrometers. Size exclusion chromatography (SEC) analysis was carried out using a GL Sciences instrument (HPLC pump PU610, LC column oven MODEL556) equipped with Shodex SE-61 RI detector, SIC GPC board and two columns (Shodex KF-804L) using polystyrene as a standard and tetrahydrofuran as an eluent at 40 °C at 1 mL/min.

All the solvents used for reactions were distilled under argon after drying over an appropriate drying reagent. Most of reagents were purchased from Aldrich Chemical Co. or Kanto Chemical Co., Inc. and were used without further purification unless otherwise specified.

Carboxylation of arenes 1a-h at 80 °C in trifluoroacetic acid (TFA)

In a 5-mL test tube with a screw cap equipped with a magnetic stirring bar, benzene (**1a**, 0.89 mL, 10.0 mmol), HCOOH (1.00 mL, 26 mmol), Pd(tfa)₂ (6.7 mg, 0.020 mmol), K₂S₂O₈ (1.62 g, 6.0 mmol) and TFA (2.0 mL) were placed. The mixture was stirred at 80 °C for 70 h. After addition of water (30 mL), the mixture was extracted

by ethyl acetate (20 mL x 3) and the combined organic layers were dried over MgSO₄. After evaporation of the solvent, the mixture was purified by silica-gel column chromatography (Hex:*i*-PrOH = 5:1) to give benzoic acid (**2a**, 0.26 g, 2.2 mmol, TON = 106). The other arenes **1b-h** were allowed to react under the same conditions. The *ortho:meta:para* ratios were determined by ¹H NMR. The following compounds are commercially available from Aldrich and the structures were confirmed by comparison of their ¹H NMR spectra: **2a**, *o*-**2b**, *m*-**2b**, *p*-**2b**, *p*-**2c**, *p*-**2d**, *o*-**2f**, *m*-**2f**, and *p*-**2f**. *m*-**2d** was reported in S. Berger. *Tetrahedron* **1976**, 32, 2451. Products from **1e** were obtained as hydroxybenzoic acids because the ester of acetic acid was hydrolyzed either under the reaction conditions or at the stage of aqueous work up. Both of *o*- and *p*-hydroxybenzoic acids are commercially available from Aldrich.

Carboxylation of biphenyl (**1h**) at 80 °C in TFA.

Biphenyl (**1h**, 10 mmol) was treated under the same conditions as mentioned above. Purification by silica-gel column chromatography (Hex:AcOEt = 5:1) gave biphenylcarboxylic acids (**2h**, R_f 0.08, 479 mg, 2.4 mmol, TON = 120, 24% yield from biphenyl). The *meta:para* ratio of 1:4 was determined by ¹H NMR. Benzo[c]chromen-6-one (R_f 0.40, 26 mg, 0.13 mmol, 1%), 9-fluorenone (R_f 0.55, 12 mg, 0.070 mmol, 0.7%), and 4-hydroxybiphenyl (R_f 0.30, 15 mg, 0.09 mmol, 9%) were obtained as byproducts and biphenyl was recovered in 62%. 4-Biphenylcarboxylic acid, 9-fluorenone, and 4-hydroxybiphenyl are available from Aldrich and the structures were confirmed by comparison of their ¹H and ¹³C NMR. The following compounds were reported in literature. 3- Biphenylcarboxylic acid: E. Nicholas, S. M. Resouly *Tetrahedron* **1999**, 55, 11889. Benzo[c]chromen-6-one: T. Murai.; H. Togo.; M. Yokoyama. *J.Chem.Soc.,Perkin Trans. I* **1999**, 1713.

Hydroxy-Carboxylation of biphenyl (**1h**) at 50°C in $\text{TFA}/\text{CH}_2\text{Cl}_2$.

A mixture of biphenyl (0.31 g, 2.0 mmol), HCOOH (0.50 mL, 13 mmol), $\text{Pd}(\text{tfa})_2$ (0.033 g, 0.100 mmol), $\text{K}_2\text{S}_2\text{O}_8$ (1.62 g, 6.0 mmol), CH_2Cl_2 (1.0 mL) and TFA (2.0 mL) was stirred at 50°C for 48 h in a 5-mL test tube equipped with a screw cap. After aqueous work-up and removal of solvents, the crude product was isolated by silica-gel column chromatography (Hex:*i*-PrOH = 10:1) to give hydroxy-biphenylcarboxylic acid (**3**) as the major product in 45% yield (193 g, 0.90 mmol, R_f 0.33 at Hex:*i*-PrOH = 5:1). The regioisomer ratio of [4'-hydroxy-3-biphenylcarboxylic acid (meta-isomer)]/[4'-hydroxy-4-biphenylcarboxylic acid (para-isomer)] was determined by ^1H NMR to be 17/83. A minor isolated product was 4,4'-dihydroxybiphenyl obtained in 2% yield (7.0 mg, 0.040 mmol, of R_f 0.38 at Hex:*i*-PrOH = 5:1). Biphenyl was completely consumed and a complex mixture of other unidentified byproducts was obtained. 4-Hydroxy4'-carboxylbiphenyl and 4,4'-dihydroxybiphenyl are available from Aldrich and the structures were confirmed by comparison of their ^1H and ^{13}C NMR charts with literature data.

Characterization of the Byproducts with Higher Molecular Weight.

A mixture of biphenyl (0.31 g, 2.0 mmol), HCOOH (0.50 mL, 13 mmol), Pd(tfa)₂ (0.033 g, 0.100 mmol), K₂S₂O₈ (1.62 g, 6.0 mmol), CH₂Cl₂ (1.0 mL) and TFA (2.0 mL) was stirred 50 °C for 48 h in 5-mL test tube in the same way as mentioned above. After filtration through a pad of Celite, the solids were washed with ethyl acetate. After evaporation of the solvents, the crude product was subjected to SEC analysis using polystyrene standard (M_w 402, M_w/M_n = 1.26).

Attempted Hydroxy-Carboxylation of biphenyl using CO as a Carbonyl source.

In a 20 mL schlenk tube equipped with a magnetic stirring bar, a mixture of biphenyl (0.31 g, 2.0 mmol), Pd(tfa)₂ (0.033 g, 0.100 mmol), and K₂S₂O₈ (1.62 g, 6.0 mmol) in CH₂Cl₂ (1.0 mL) and TFA (2.0 mL) was stirred at 50 °C for 48 h under atmospheric pressure of CO (balloon). After the same work-up as mentioned above, the major products were biphenylcarboxylic acids **2h** obtained in 31% yield (123 mg, 0.62 mmol, *meta:para* = 13:87), accompanied by a byproduct 9-fluorenone in 6% yield (21 mg, 0.12 mmol). Biphenyl (**1h**) was recovered in 46% (0.14 g, 0.91 mmol).

Attempted Hydroxy-Carboxylation of biphenyl using CO in the Presence of Water.

The reaction procedure mentioned above was repeated in the presence of H₂O (54 µL, 3.0 mmol). Biphenylcarboxylic acids **2h** were obtained in 36% yield and biphenyl (**1h**) was recovered in 57%.

Control experiment with 4-biphenylcarboxylic acid (4**), with 4-hydroxybiphenyl (**5**), 4-biphenyl α,α,α -trifluoroacetate (**6a**), and 4-biphenyl formate (**6b**).**

When 4-biphenylcarboxylic acid (**4**, 0.40 g, 2.0 mmol) was subjected to the same reaction conditions as those for the hydroxylation-carboxylation of biphenyl, quantitative recovery of **4** was confirmed by ^1H NMR analysis.

In contrast, treatment of 4-hydroxybiphenyl (**5**, 0.34 g, 2.0 mmol) with HCOOH (0.5 mL, 13 mmol), $\text{Pd}(\text{tfa})_2$ (0.033 g, 0.100 mmol), $\text{K}_2\text{S}_2\text{O}_8$ (1.62 g, 6.0 mmol), CH_2Cl_2 (1.0 mL) and TFA (2.0 mL) at 50 °C for 48 h. After aqueous work up, hydroxy-biphenylcarboxylic acid (**3**) was obtained in 6% yield (26 mg, 0.12 mmol).

Treatment of 4-biphenyl α,α,α -trifluoroacetate (**6a**, 0.53 g, 2.0 mmol) by the same procedure gave **3** (*meta:para* = 29:71) in 52% yield accompanied by a byproduct 4,4'-dihydroxybiphenyl (2% yield). From 4-biphenyl formate (**6b**, 0.40 g, 2.0 mmol), **3** (*meta:para* = 23:77) and 4,4'-dihydroxybiphenyl were obtained in 62% and 4%, respectively.

About the temperature dependency of the hydroxy-carboxylation: The reaction at 80 °C.

A mixture of biphenyl (0.31 g, 2.0 mmol), HCOOH (0.50 mL, 13 mmol), $\text{Pd}(\text{tfa})_2$ (0.033 g, 0.100 mmol), $\text{K}_2\text{S}_2\text{O}_8$ (1.62 g, 6.0 mmol), 1,2-dichloroethane (1.0 mL) and TFA (2.0 mL) was stirred at 80 °C for 48 h. Standard work-up and purification provided 40% of **2h**. No trace of **3** was detected.

Hydroxylation-Carboxylation of Benzene

A mixture of benzene (178 μL , 2.0 mmol), HCOOH (1.00 mL, 26 mmol), $\text{Pd}(\text{tfa})_2$ (0.033 g, 0.100 mmol), $\text{K}_2\text{S}_2\text{O}_8$ (2.2 g, 8.0 mmol), CH_2Cl_2 (1.0 mL) and TFA (2.0 mL) was stirred at 40 °C for 91 h in a 5-mL test tube equipped with a screw cap. After

aqueous work-up and removal of the organic solvents in vacuo, the crude product was isolated by silica-gel column chromatography (Hex:*i*-PrOH = 10:1) to give benzoic acid (41 mg 0.34 mmol, Rf 0.38), 2-hydroxybenzoic acid (6 mg 0.04 mmol, Rf 0.35), and 4-hydroxybenzoic acid (23 mg 0.16 mmol, Rf 0.10). Benzoic acid, 2-hydroxybenzoic acid, and 4-hydroxybenzoic acid are available from Aldrich and the structures were confirmed by comparison of their ^1H and ^{13}C NMR charts to authentic samples.

About the stability of formic acid under acidic conditions.

A mixture of formic acid (0.20 mL), trifluoroacetic acid (0.8 mL), dichloroethane (50 μL , internal standard) and CD_2Cl_2 (0.5 mL) in an NMR tube was heated at 50 $^\circ\text{C}$ and ^1H NMR was recorded after 0, 25, 40, and 80 h. During the time, the integration ratio of dichloroethane and formic proton was unchanged. Addition of $\text{Pd}(\text{tfa})_2$ to this sample resulted in the immediate reduction of Pd(II) to give black powders of palladium.