Paxdaphnidines A and B, Novel Penta- and Tetracyclic Alkaloids
from *Daphniphyllum paxianum*

Zha-Jun Zhan, Sheng-Ping Yang, and Jian-Min Yue*

State Key Laboratory of Drug Research, Institute of Materia Medica, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, 555 Zu Chong Zhi Road, Zhangjiang Hi-Tech Park, Shanghai, 201203, P. R. China

jmyue@mail.shcnc.ac.cn.

Supporting Information

Figure S2. ¹H NMR spectrum of paxdaphnidineA (**1**) in CD₃OD.

Figure S3. ¹³C NMR spectrum of paxdaphnidineA (**1**) in CD₃OD.

Figure S4. ¹³C NMR spectrum of paxdaphnidineA (**1**) in CDCl₃.

Figure S5. ¹H-¹H COSY spectrum of paxdaphnidineA (**1**) in CD₃OD.

Figure S6. HSQC spectrum of paxdaphnidineA (**1**) in CD₃OD (20-50 ppm).

Figure S7. HSQC spectrum of paxdaphnidineA (**1**) in CD₃OD (50-58 ppm).

Figure S8. HMBC spectrum of paxdaphnidineA (**1**) in CD₃OD.

Figure S9. NOESY spectrum of paxdaphnidineA (**1**) in CD₃OD.

Figure S10. EI MS spectrum of paxdaphnidineA (**1**).

Figure S11. ¹H NMR spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S12. ¹³C NMR spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S13. ¹H-¹H COSY spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S14. HSQC spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S15. HMBC spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S16. NOESY spectrum of paxdaphnidine B (**2**) in CD₃OD.

Figure S17. EI MS spectrum of paxdaphnidine B (**2**).

General Procedures S18.

Solvent: cd3od
Ambient temperature
Mercury-400Bb "SIN400"

Relax. delay 1.500 sec
COSY 90-45
Acq. time 0.171 sec
Width 6009.6 Hz
2D Width 6009.6 Hz
8 repetitions
312 increments
OBSERVED F2, 400.1587622 MHz
DATA PROCESSING
Sd time 0.085 sec
F1 DATA PROCESSING
Sd time 0.044 sec
F1 size 2048 x 2048
Total time 2 hr, 2 min, 37 sec

paxdaphnidine A (1)
H-H COSY in CD₃OD

Solvent: CD3OD
Ambient temperature
User: 11-12-86 "SIMM400"
Mercury-400BB

Relax. delay 1.000 sec
Acq. time 0.170 sec
Width 6009.6 Hz
2D Width 20000.0 Hz
16 repetitions
2 x 200 increments
OBSERVE H1, 400.1587697 MHz
DECOUPLE C13, 100.6275247 MHz
Power 54 db
on during acquisition
off during delay
GARP-1 modulated
DATA PROCESSING
F2 axis apodization 0.079 sec
F1 axis apodization 0.013 sec
Gain 2.000
F2 size 2048
F1 size 2048
Total time 2 hr, 18 min, 4 sec

Solvent: CD3OD
Ambient temperature
User: 11-12-86 "SIMM400"
Mercury-400B8

Relax. delay 1.000 sec
Acq. time 0.170 sec
Width 6009.6 Hz
2D Width 20000.0 Hz
2D repetitions
2 x 100 increments
OSERVE 11.000 587697 MHz
DECOUPLE C13, 100.6275247 MHz
Power 54 dB
on during acquisition
off during delay
GARP-1 modulated
DATA PROCESSING
Gauss apodization 0.079 sec
F1 DATA PROCESSING
Gauss apodization 0.013 sec
FT size 2048 x 2048
Total time 2 hr, 18 min, 4 sec

S9

SPEC: 131282
Samp: Dap1
Comm: 70ev/IST:200C/R:1000
Mode: EI +VE +LMR BSCAN (EXP) UP LR NRM
Oper: SIMM/MAT95/WANG J.
Base: 357.3
Norm: 357.3
Peak: 1000.00 mmu

13-Aug-03
Elapse: 01:00.2
Start : 15:26:27
8

Inlet :
Masses: 50 > 750
#peaks: 579

Dap2 1H CD3OD 03-12-11

```
Solvent: cd3od
Ambient temperature
Mercury-400BB "SIMM400"
```

```
Relax. delay 1.500 sec
COSY 90-45
```

COSY 90-45
Acq. time 0.171 sec

Acq. time 0.171 sec
Width 6009.6 Hz

Width	6009.6 Hz
2D Width	6009.6 Hz

2D width 6009.6 Hz
8 repetitions8 repetitions
512 increments

```
512 increments
OBSERVE H1 200 15
```

OBSERVE H1, 400.1587644 MHZ
DATA PROCESSING

DATA PROCESSING

Sq. sine bell 0.085 sec

```
04.316 0.000 sec
F1 DATA PROCESSING
04.317 0.040 sec
```

```

11 DATA PROCROSSING
Sq. sine bel1 0.043 sec

```

sq: slice belt 0.043 sec
FT size 2048 x 2048

FI size 2048 x 2048
Total time 2 hr, 2 min, 37 sec

DATE: 0000000000

Solvent: CD3OD
Ambient temperature
User: 11-12-86 "SIMM400"
Mercury-400BB

Relax. delay 1.000 sec
Acq. time 0.170 sec
Width 6009.6 Hz
2D Width 20000.0 Hz
8 repetitions
2 x 200 increments
OBSERVE H1, 400.1587687 MHz
DECUPLE C13, 100.6275247 MHz
Power 34 dB
of during acquisition
GARP-1 modulated
DATA PROCESSING
Gauss apodization 0.079 sec
F1 DATA PROCESSING
Gauss apodization 0.014 sec
FT size 2048 x 2048
Total time 1 hr, 10 min, 39 sec

S15

S17

SPEC: 131281
Samp: Dap2
Comm: 70eV/IST:200C/R:1000
Mode: EI +VE +LMR BSCAN (EXP) UP LR NRM
Oper: SIMM/MAT95/WANG J.
Base: 260.2
Norm: 260.2
Peak: 1000.00 mmu

13-Aug-03
Elapse: 07:34.7
Start : 15:15:20
73

Inlet :
Masses: 50 > 750
#peaks: 766

General Procedures. Melting point was recorded on Fisher-Johns melting point apparatus and uncorrected. Optical rotations were determined on a Perkin-Elmer 341 polarimeter. IR spectra were measured on a Perkin-Elmer 577 spectrometer. NMR spectra were made on a Bruker AM-400 and Bruker AM-600 spectrometer with TMS as internal standard. Standard pulse sequences were employed to obtain ^1H - ^1H COSY, HMBC, HSQC and NOESY. EIMS (70 eV) was carried out on a Finnigan MAT 95 mass spectrometer instrument. All solvents used were of analytical grade (Shanghai Chemical Plant). Silica gel (230–400 mesh) was used for column chromatography, and pre-coated silica gel GF₂₅₄ plates (Qingdao Haiyang Chemical Plant) were used for TLC. Sephadex LH-20 was also used for column chromatograph.