Pd-*m*BDPP-Catalyzed Regioselective Internal Arylation of Electron-rich Olefins by Aryl Halides

*Shifang Liu,^a Neil Berry^a, Nick Thomson,^b Alan Pettman,^b Zeynab Hyder,^a Jun Mo^a and
Jianliang Xiao^{a*}*

^a Liverpool Centre for Materials and Catalysis

Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, UK

^b Pfizer Global Research and Development, Chemical Research and Development

Department, Ramsgate Road, Sandwich, Kent CT13 9NJ, UK

Supporting Information

Contents

1. Materials and apparatus	2
2. General procedure for the Heck arylation	2
3. Characterization of products	3
4. Methodology of DFT calculations	5
5. Results of DFT calculations	6
6. References	10
7. Sample NMR spectra	11

1. Materials and apparatus

All chemicals used were purchased from Aldrich or Lancaster and used as received. 1-Butyl-3-methylimidazolium hexafluorophosphate ([bmim][PF₆]) was prepared according to the literature method,¹ and were dried *in vacuo* at 70 °C for 5 h before use. The reactions were all carried out under a nitrogen atmosphere. The ¹H and ¹³C NMR spectra were recorded on a Gemini 400 spectrometer at 400 MHz (¹H) and 100 MHz (¹³C) in ppm with reference to TMS internal standard in CDCl₃. Mass spectra were obtained by chemical ionization (CI). All the arylation products were satisfactorily characterized by ¹H and ¹³C NMR, MS, and HRMS and by comparison of their NMR spectra with available literature data.² The ligand *meso*-2,4-bis(diphenylphosphino)pentane (*m*BDPP) was synthesized according to a literature procedure.³

2. General procedure for the Heck arylation

Specific procedures are given for the arylation of 1-bromonaphthalene (**1b**) by *n*-butyl vinyl ether (**2**) in DMSO; similar procedures were followed for other coupling reactions. An oven-dried, screw-capped reaction tube containing a stir bar was charged with **1b** (1.0 mmol), Pd(OAc)₂ (2 mol%), *m*BDPP (4 mol%), DMSO (1 mL) under nitrogen at room temperature. Following degassing three times, **2** (2.0 mmol) and NEt₃ (1.5 mmol) were injected sequentially under nitrogen. The mixture was heated to 115 °C and stirred at this temperature for 36 h. The reaction mixture was then cooled to room temperature and acidified by addition of 10 mL of 20% HCl. The product was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic layer was washed with water until neutrality, dried (MgSO₄), filtered, and concentrated *in vacuo*. A small sample was taken for NMR analysis to determine the conversion. The arylated ketone **5b** was isolated in 94% yield (160.7 mg,

0.94 mmol) as a white powder out of the crude product by flash chromatography on silica gel using a mixture of ethyl acetate and hexane (1:3 to 1:0) as eluant.

3. Characterization of products

Characterization of products 5a – 5m: compounds **5a -5m** were characterized by ^1H and ^{13}C NMR, MS, and HRMS, and all data were compared with the available data in literature.²

2-Acetonaphthone (5a): ^1H NMR (CDCl_3 , δ_{ppm}): 8.46 (s, 1H), 8.03 (dd, $J = 8.58$, 1.75 Hz, 1H), 7.95 (d, $J = 8.58$ Hz, 1H), 7.88 (dd, $J = 6.20$, 5.56 Hz, 2H), 7.60-7.55 (m, 2H), 2.71 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 198.5, 136.0, 134.9, 132.9, 130.6, 129.9, 128.9, 128.8, 128.2, 127.2, 124.3, 27.1. MS: CI, m/z 188 $[\text{M}+\text{NH}_4]^+$ (91), 171 $[\text{M}+\text{H}]^+$ (100), 170 M^+ (22), 155 $[\text{M}-\text{CH}_3]^+$ (22); HRMS, Calcd for $\text{C}_{12}\text{H}_{11}\text{O}$: 171.0810; Found 171.0809.

1-Acetonaphthone (5b) ^1H NMR (CDCl_3 , δ_{ppm}): 8.74 (d, $J = 8.75$ Hz, 1H), 7.92 (dt, $J = 7.84$, 1.60 Hz, 3H), 7.60 (dt, $J = 7.84$, 1.43 Hz, 1H), 7.53 (m, 2H), 2.74 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 202.5, 135.9, 134.4, 133.4, 130.6, 129.1, 128.8, 128.5, 126.8, 126.4, 124.7, 30.4. MS: CI, m/z 188 $[\text{M}+\text{NH}_4]^+$ (100), 171 $[\text{M}+\text{H}]^+$ (63), 170 M^+ (10), 155 $[\text{M}-\text{CH}_3]^+$ (10); HRMS, Calcd for $\text{C}_{12}\text{H}_{11}\text{O}$: 171.0810; Found 171.0809.

4'-Methylacetophenone (5c) ^1H NMR (CDCl_3 , δ_{ppm}): 7.86 (d, $J = 8.26$ Hz, 2H), 7.25 (d, $J = 8.26$ Hz, 2H), 2.64 (s, 3H), 2.41 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 198.3, 144.3, 135.1, 129.6, 128.8, 26.9, 22.0. MS: CI, m/z 152 $[\text{M}+\text{NH}_4]^+$ (100), 135 $[\text{M}+\text{H}]^+$ (97), 119 $[\text{M}-\text{CH}_3]^+$ (25); HRMS: Calcd for $\text{C}_9\text{H}_{11}\text{O}$: 135.0810; Found 135.0809.

3'-Methylacetophenone (5d) ^1H NMR (CDCl_3 , δ_{ppm}) : 7.78 (m, 2H), 7.37 (m, 2H), 2.60 (s, 3H), 2.42 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 197.0, 136.9, 135.7, 132.4, 127.4, 127.0, 124.2, 25.2, 19.9. MS: CI, m/z 152 $[\text{M}+\text{NH}_4]^+$ (100), 135 $[\text{M}+\text{H}]^+$ (30), 134 M^+ (<2), 119

$[M-CH_3]^+$ (10); HRMS, Calcd for $C_9H_{11}O$: 135.0810; Found 135.0808.

Acetophenone (5e) 1H NMR ($CDCl_3$, δ_{ppm}): 7.98 (d, $J = 7.16$ Hz, 2H), 7.55 (t, $J = 7.16$ Hz, 1H), 7.47 (t, $J = 7.31$ Hz, 2H), 2.62 (s, 3H). ^{13}C NMR ($CDCl_3$, δ_{ppm}): 198.6, 137.5, 133.5, 128.9, 128.7, 27.0. MS: CI, m/z 138 $[M+NH_4]^+$ (100), 121 $[M+H]^+$ (34), 120 M^+ (5), 105 $[M-CH_3]^+$ (62); HRMS, Calcd for C_8H_9O : 121.0653; Found 121.0655.

4'-Fluoroacetophenone (5f) 1H NMR ($CDCl_3$, δ_{ppm}): 7.99 (m, 2H), 7.16 (t, $J = 8.68$ Hz, 2H), 2.60 (s, 3H). ^{13}C NMR ($CDCl_3$, δ_{ppm}): 197.3, 166.6 (d, $J_{CF} = 254.50$ Hz), 134.4, 131.9 (d, $J_{CF} = 9.46$ Hz), 116.5 (d, $J_{CF} = 22.13$ Hz), 27.3. MS: CI, m/z 156 $[M+NH_4]^+$ (100), 139 $[M+H]^+$ (10), 138 M^+ (27), 123 $[M-CH_3]^+$ (78); HRMS, Calcd for $C_8H_{11}FON$: 156.0825, Found 156.0826.

4'-Methoxyacetophenone (5g) 1H NMR ($CDCl_3$, δ_{ppm}): 7.94 (d, $J = 8.90$ Hz, 2H), 6.93 (d, $J = 8.90$ Hz, 2H), 3.87 (s, 3H), 2.56 (s, 3H). ^{13}C NMR ($CDCl_3$, δ_{ppm}): 197.1, 163.8, 130.9, 130.7, 114.0, 55.8, 26.7. MS: CI, m/z 168 $[M+NH_4]^+$ (38), 151 $[M+H]^+$ (100), 135 $[M-CH_3]^+$ (16); HRMS, Calcd for $C_9H_{11}O_2$: 151.0759; Found 151.0755.

3'-Methoxyacetophenone (5h) 1H NMR ($CDCl_3$, δ_{ppm}): 7.53 (d, $J = 7.63$ Hz, 1H), 7.48 (dd, $J = 3.02, 2.54$ Hz, 1H), 7.37 (t, $J = 7.95$ Hz, 1H), 7.12 (ddd, $J = 8.11, 2.70, 0.96$ Hz, 1H), 3.86 (s, 3H), 2.60 (s, 3H). ^{13}C NMR ($CDCl_3$, δ_{ppm}): 198.3, 160.1, 138.8, 129.9, 121.5, 120.0, 112.7, 55.8, 27.1. MS: CI, m/z 168 $[M+NH_4]^+$ (100), 151 $[M+H]^+$ (35), 150 M^+ (11), 135 $[M-CH_3]^+$ (8); HRMS, Calcd for $C_9H_{11}O_2$: 151.0759; Found 151.0759.

2'-Methoxyacetophenone (5i) 1H NMR ($CDCl_3$, δ_{ppm}): 7.73 (dd, $J = 7.63, 1.91$ Hz, 1H), 7.47 (ddd, $J = 7.63, 1.91, 0.80$ Hz, 1H), 6.98 (q, $J = 7.63$ Hz, 2H), 3.91 (s, 3H), 2.62 (s, 3H). ^{13}C NMR ($CDCl_3$, δ_{ppm}): 200.3, 159.3, 134.1, 131.8, 120.9, 120.5, 111.9, 55.8, 32.2. MS: CI, m/z 151 $[M+H]^+$ (100), 150 M^+ (11), 135 $[M-CH_3]^+$ (19); HRMS, Calcd for $C_9H_{11}O_2$: 151.0759; Found 151.0759.

4'-Acetylbenzonitrile (5j) ^1H NMR (CDCl_3 , δ_{ppm}): 8.05 (d, $J = 8.58$ Hz, 2H), 7.79 (d, $J = 8.58$ Hz, 2H), 2.66 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 195.9, 140.3, 132.9, 129.1, 118.3, 116.8, 27.2. MS: CI, m/z 163 $[\text{M}+\text{NH}_4]^+$ (100), 145 M^+ (10), 130 $[\text{M}-\text{CH}_3]^+$ (38); HRMS, Calcd for $\text{C}_9\text{H}_{11}\text{N}_2\text{O}$: 163.0871, Found 163.0873.

4'-Acetylbenzaldehyde (5k) ^1H NMR (CDCl_3 , δ_{ppm}): 10.11 (s, 1H), 8.12 (d, $J = 6.60$ Hz, 2H), 8.01 (d, $J = 6.60$ Hz, 2H), 2.66 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 197.8, 191.9, 141.6, 139.4, 130.2, 129.2, 27.3. MS: CI, m/z 166 $[\text{M}+\text{NH}_4]^+$ (30), 148 M^+ (100), 133 $[\text{M}-\text{CH}_3]^+$ (100); HRMS, Calcd for $\text{C}_9\text{H}_{12}\text{O}_2\text{N}$: 166.0868, Found 166.0861.

1,4-Diacetylbenzene (5l) ^1H NMR (CDCl_3 , δ_{ppm}): 8.04 (s, 4H), 2.65 (s, 6H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 197.8, 140.5, 128.8, 27.2. MS: CI, m/z 180 $[\text{M}+\text{NH}_4]^+$ (100), 163 $[\text{M}+\text{H}]^+$ (5), 162 M^+ (7), 147 $[\text{M}-\text{CH}_3]^+$ (21); HRMS, Calcd for $\text{C}_{10}\text{H}_{14}\text{O}_2\text{N}$: 180.1024; Found 180.1019.

Methyl-4-acetylbenzoate (5m) ^1H NMR (CDCl_3 , δ_{ppm}): 8.13 (d, $J = 8.26$ Hz, 2H), 8.01 (d, $J = 8.27$ Hz, 2H), 3.96 (s, 3H), 2.65 (s, 3H). ^{13}C NMR (CDCl_3 , δ_{ppm}): 197.9, 166.6, 140.6, 134.2, 130.2, 128.6, 52.8, 27.2. MS: CI, m/z 196 $[\text{M}+\text{NH}_4]^+$ (100), 179 $[\text{M}+\text{H}]^+$ (16), 178 M^+ (8), 163 $[\text{M}-\text{CH}_3]^+$ (83); HRMS, Calcd for $\text{C}_{10}\text{H}_{14}\text{O}_3\text{N}$: 196.0974; Found 196.0971.

4. Methodology of DFT calculations

All calculations were performed on a PC using either Scientific Linux 3.0.3⁴ or Windows XP⁵ operating systems. All structures were completely optimized at the density functional theory (DFT) level by using the B3LYP functional as implemented in GAMESS⁶ or PCGAMESS.⁷ For all optimizations the 6-31* basis set was used for all atoms excluding Pd. Relativistic effects for Pd were addressed by using SBKJC effective core potential (ECP) together with the SBKJC⁶ basis set. Charges were calculated by

fitting charges which reproduced the charge, dipole and quadrupole of the molecule.⁶ Calculations were performed in parallel using Linux cluster in the Physics department of Liverpool University. Structures were visualized using Molekel.⁸

5. Results of DFT calculations

Optimized geometry of [PdBr(Ph)(DPPP)] in the singlet ground state in the gas phase

Selected parameters:

Energy (au)		-4656.9298691834
Distance (Å) (Pd14-Br25)		2.493
Distance (Å) (P7-Pd14)		2.439
Distance (Å) (P15-Pd14)		2.324
Distance (Å) (Pd14-C19)		2.061
Angle(°) (C19-Pd14-Br25)		86.758
Angle(°) (Br25-Pd14-P7)		90.582
Angle(°) (C19-Pd14-P15)		87.516
Angle(°) (P7-Pd14-P15)		95.273
Charge	P7	0.1222
	P15	0.3557
	Pd14	-0.3066
	C19	0.1429
	Br25	-0.3468

Coordinates of [PdBr(Ph)(DPPP)] in xyz format in Angstroms

68

0.0000000

C	-3.03029	2.11277	-1.34979
C	-3.33347	0.75754	-1.15071
C	-4.65939	0.32281	-1.29049
C	-5.66432	1.22809	-1.63462
C	-5.35383	2.57370	-1.84217
C	-4.03645	3.01366	-1.69995
P	-1.95759	-0.40768	-0.77748
C	-2.72808	-1.77052	0.20247
C	-3.10698	-3.00520	-0.34844
C	-3.67423	-3.99877	0.45378
C	-3.87324	-3.77141	1.81546
C	-3.49861	-2.54656	2.37331
C	-2.92464	-1.55418	1.57934
Pd	0.02368	0.56920	0.25718
P	1.54668	-0.61279	-1.03982
C	0.90946	-1.29118	-2.66359
C	-0.42887	-2.04983	-2.59340
C	-1.66882	-1.14382	-2.46703
C	1.63627	1.48251	1.15843
C	2.35978	0.88558	2.19829
C	3.42656	1.55941	2.80569
C	3.78415	2.83933	2.38083
C	3.06192	3.44488	1.34994
C	1.99618	2.77231	0.74461
Br	-1.40220	1.77634	1.90755
C	3.10305	0.22223	-1.56568
C	4.27379	0.11853	-0.79860
C	5.42650	0.81165	-1.16860
C	5.42829	1.62127	-2.30521
C	4.26504	1.74379	-3.06641
C	3.10985	1.05434	-2.69702
C	2.08431	-2.15469	-0.18201
C	1.45521	-2.52566	1.01570
C	1.80011	-3.71793	1.65718
C	2.77888	-4.54831	1.11120
C	3.41395	-4.18635	-0.08035
C	3.06851	-2.99889	-0.72428
H	1.43053	3.27107	-0.03937
H	3.32209	4.44842	1.01839
H	4.61120	3.36421	2.85359
H	3.97234	1.08073	3.61701
H	2.09388	-0.10669	2.55437
H	2.21188	1.18374	-3.29455
H	4.25197	2.38014	-3.94757
H	6.32740	2.16090	-2.59105
H	6.32236	0.72234	-0.55994
H	4.28318	-0.49176	0.09791
H	0.69672	-1.87441	1.44367
H	1.30375	-3.99190	2.58417
H	3.05004	-5.47430	1.61180
H	4.18017	-4.82843	-0.50733
H	3.57929	-2.72296	-1.64342
H	1.68946	-1.94892	-3.06452
H	0.81179	-0.45333	-3.36349

H	-0.40441	-2.79138	-1.78491
H	-0.52543	-2.62511	-3.52382
H	-2.56993	-1.67625	-2.79277
H	-1.57032	-0.28485	-3.14313
H	-2.96638	-3.20988	-1.40515
H	-3.95966	-4.94934	0.00986
H	-4.31579	-4.54446	2.43873
H	-3.64837	-2.35995	3.43363
H	-2.62832	-0.60411	2.01771
H	-4.91165	-0.72001	-1.11746
H	-6.69028	0.88253	-1.73441
H	-6.13849	3.27889	-2.10465
H	-3.79215	4.06276	-1.84483
H	-2.01177	2.46337	-1.21051

Optimized geometry of [PdBr(Ph)(*m*BDPP)] in the singlet ground state in the gas phase

Selected parameters:

Energy (au)		-4723.6778492809
Distance (Å) (Pd18-Br19)		2.508
Distance (Å) (P7-Pd18)		2.421
Distance (Å) (P17-Pd18)		2.541
Distance (Å) (Pd18-C32)		2.033
Angle (°)(C32-Pd18- Br19)		88.081
Angle (°)(Br19-Pd18- P17)		90.152
Angle (°)(C32-Pd18-P7)		87.398
Angle(°)(P17-Pd18-P7)		94.411
Charge	P7	0.0453
	P17	-0.2355
	Pd18	-0.0155
	C32	0.1034
	Br19	-0.4859

Coordinates of [PdBr(Ph)(*m*BDPP)] in xyz format in Angstroms

74

0.0000000

C	3.26131	-2.93578	-0.56020
C	2.13282	-2.19595	-0.20017
C	1.35019	-2.65357	0.85747
C	1.67447	-3.82389	1.52796
C	2.79301	-4.54691	1.15690
C	3.58886	-4.09845	0.11272
P	1.60145	-0.67569	-1.08675
C	3.13452	0.21906	-1.56656
C	4.27033	0.18849	-0.76233
C	5.39121	0.93083	-1.09394
C	5.39463	1.72352	-2.22823
C	4.26430	1.77953	-3.02596
C	3.14353	1.03788	-2.69472
C	0.92815	-1.29267	-2.72553
C	-0.43079	-2.01649	-2.59181
C	-1.68183	-1.10593	-2.53388
P	-2.02339	-0.44042	-0.82344
Pd	0.03361	0.57156	0.27253
Br	-1.43075	1.84117	1.86407
C	-3.38666	0.74886	-1.12752
C	-3.05704	2.07495	-1.39431
C	-4.04429	3.00282	-1.67957
C	-5.37509	2.62061	-1.68423
C	-5.71673	1.30767	-1.40317
C	-4.72886	0.37753	-1.12677
C	-2.75731	-1.77227	0.21674
C	-2.78905	-3.11943	-0.13272
C	-3.30559	-4.06905	0.73689
C	-3.79781	-3.68645	1.97102
C	-3.76067	-2.34973	2.33783
C	-3.24109	-1.40147	1.47522
C	1.64089	1.43414	1.17055
C	2.32750	0.82321	2.21577
C	3.39988	1.45040	2.83836
C	3.80836	2.70691	2.42744
C	3.12640	3.33422	1.39556
C	2.05659	2.70581	0.77904
H	1.52939	3.22579	-0.00261
H	3.42335	4.31871	1.07567
H	4.63657	3.19646	2.90983
H	3.90867	0.95656	3.64932
H	2.02403	-0.14764	2.56805
H	2.27743	1.11460	-3.32531
H	4.24991	2.40194	-3.90262
H	6.26507	2.30115	-2.48260
H	6.25501	0.89652	-0.45529
H	4.28039	-0.39366	0.13766
H	0.48942	-2.09438	1.17400
H	1.05663	-4.15714	2.34179
H	3.05025	-5.45041	1.68009
H	4.46422	-4.65223	-0.17556
H	3.89574	-2.59981	-1.35793
C	1.90557	-2.17333	-3.51863

H	0.75858	-0.38975	-3.30300
H	-0.39846	-2.70875	-1.75702
H	-0.53520	-2.63890	-3.47522
C	-2.86706	-1.71994	-3.29595
H	-1.43463	-0.18767	-3.06207
H	-2.41167	-3.45186	-1.07841
H	-3.32253	-5.10343	0.44298
H	-4.20158	-4.42162	2.64409
H	-4.12698	-2.04191	3.30045
H	-3.18822	-0.37587	1.78716
H	-5.00930	-0.63487	-0.90255
H	-6.74934	1.00770	-1.39427
H	-6.14286	3.34374	-1.89432
H	-3.77392	4.02453	-1.87583
H	-2.03253	2.39436	-1.35078
H	1.51760	-2.32610	-4.52142
H	2.88544	-1.72262	-3.61128
H	2.01939	-3.14782	-3.05903
H	-2.58329	-1.87252	-4.33354
H	-3.17424	-2.67883	-2.89946
H	-3.72655	-1.06330	-3.28612

6. References

- (1) Fuller, J.; Carlin, R. T.; de Long, H. C.; Haworth, D. *J. Chem. Soc., Chem. Commun.* **1994**, 299.
- (2) Mo, J.; Xu, L.; Xiao, J. *J. Am. Chem. Soc.* **2005**, 127, 751.
- (3) Bakos, J.; Tóth, I.; Hell, B.; Markó, L. *J. Organomet. Chem.* **1985**, 279, 23.
- (4) <https://www.scientificlinux.org/>.
- (5) <http://www.windows.com>.
- (6) Schmidt, M. W.; Baldrige, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. H.; Koseki, S.; Matsunaga, N.; Nguyen, K. A.; Su, S. J.; Windus, T. L.; Dupuis, M.; Montgomery, J. A. *J. Comput. Chem.* **1993**, 14, 1347.
- (7) Granovsky, A. A. www <http://classic.chem.msu.su/gran/gamess/index.html>.
- (8) Portmann, S.; Lüthi, H. P. *CHIMIA*, **2000**, 54, 766.

7. Sample NMR spectra

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR **5b**

^1H NMR

^{13}C NMR **5c**

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

51

^1H NMR

^{13}C NMR

5m

^1H NMR

^{13}C NMR