S1

Pd on Surface Modified Ferrites: A Complete Facile Magnetically Recoverable Catalyst for Hydrogenation Reactions

Debanjan Guin, Babita Baruwati, Sunkara V Manorama*

**Nanomaterials Laboratory, Inorganic and Physical Chemistry Division,
Indian Institute of Chemical Technology, Hyderabad 500 007**

manorama@iict.res.in

Synthesis Procedure & Characterization	S2- S3
FTIR spectra	S4
VSM plots	S5
Table for Fe ₃ O ₄ - DA- Pd catalyst	S6
GCMS spectra	S7-S14
¹ H and ¹³ C NMR of	S15
¹ H and ¹³ C NMR of	S16

Preparation of surface modified nanocrystalline ferrites: 1 g of the above-synthesized NiFe_2O_4 was taken in deionized water. 2 g of dopamine was added to the mixture and refluxed for 12 hrs at 100°C . The resulting product was precipitated using excess amount of acetone and centrifugation. It was then dried and used for Pd incorporation.

In case of Fe_3O_4 surface modification, 1 g of the above-synthesized product was taken in deionized water. 2 g of dopamine was added to the mixture and then ultrasonicated for 100 minutes. The resulting product was highly dispersed in water and was precipitated using excess acetone and centrifugation. In this case we intentionally avoided the heating process as heating in air may cause the oxidation of Fe_3O_4 to Fe_2O_3 .

Incorporation of Pd (0): 1 g of surface modified $\text{Fe}_3\text{O}_4/\text{NiFe}_2\text{O}_4$ nanoparticle was dispersed in water. NaPdCl_4 solution in water was added to the mixture to get 10 wt% of Pd. Diluted solution of hydrazine monohydrate was added dropwise to adjust the pH to 9. The reaction mixture was stirred overnight at room temperature and then allowed to settle. The product was washed several times with water, centrifuged and dried at room temperature.

Henceforth the catalysts are referred as $\text{NiFe}_2\text{O}_4\text{-DA-Pd}$ and $\text{Fe}_3\text{O}_4\text{-DA-Pd}$.

Characterization

X-Ray diffraction patterns of the powdered samples are recorded on a Rigaku Miniflex Table Top diffractometer using $\text{CuK}\alpha$ radiation ($1.5406 \text{ \AA}$) source. The intensity data were collected over a 2θ range of 2 to 80° using a step scan mode (0.06 degree per sec). The average crystallite size of the samples was estimated with the help of Scherrer equation using the FWHM applied to all major peaks.

The amount of Pd in the samples before and after reactions are determined with an IRIS Intrepid II XDL inductively coupled plasma, atomic emission spectroscopy (ICPAES) instrument. 2.5 mg of each sample was dissolved in 2.5 mL of concentrated aqua regia and the volume was adjusted to 25 mL. 5 mL of this solution was then further diluted to 25 mL. This diluted sample was then used for the AES analysis.

Transmission electron micrographs and SAED patterns are recorded on a Phillips Tecnai G² FEI F12 electron microscope. The powder samples are dispersed in ethanol by ultrasonication before loading onto a carbon coated copper grid and then allowed to dry at room temperature before recording the micrographs.

FTIR spectra was recorded on a Nicolet Nexus 670 Spectrometer equipped with DTGS KBr detector in the range $400\text{-}4000 \text{ cm}^{-1}$.

XPS measurements for the as synthesized samples were performed on a Kratos X-Ray photoelectron spectrometer. The X-ray excitation energy was 1253.6 eV ($\text{MgK}\alpha$) and the spectra were recorded with pass energy of 80 eV. The angle between the detector and the X-ray flux direction was constant and equal to 90° , while the measurements were made at an electron take off angle (angle between the surface plane and electron emission direction) of 70° .

The calibration of the spectrometer and the charging correction were verified by determining the binding energy of C 1s level with a graphite substrate (at 285 eV). The binding energies were compared with the reported data from literature to identify the chemical composition and the oxidation state.

GCMS spectra were recorded on a Thermo Finnigan TRACE DSQ MS Spectrophotometer using BP-1 (30x0.25x1.0) capillary column.

^1H NMR spectra have been recorded on a Bruker UX NMR/XWIN NMR 300 MHz spectrometer.

^{13}C NMR spectra have been recorded on a Bruker Avance 600 MHz spectrometer.

Magnetic hysteresis loops of the pure NiFe_2O_4 and Fe_3O_4 samples were recorded on a Lakeshore 7000 Vibrating sample magnetometer up to a field 1T.

S4

FTIR

FTIR spectrum of Dopamine

FTIR spectrum of Fe₃O₄ functionalized with dopamine

FTIR spectrum of NiFe₂O₄ functionalized with dopamine

VSM:

Magnetization Vs Applied field plot for the as synthesized NiFe_2O_4 nanoparticles showing the superparamagnetic behavior of the nanoparticles.

Magnetization Vs Applied field plot for the as synthesized Fe_3O_4 nanoparticles showing the superparamagnetic behavior of the nanoparticles.

S6

Table. Summary of the observations from the different reactions carried out on Fe₃O₄–DA-Pd catalyst.

entry	substrate	product	conversion (%)	isolated yield ^a (%)	time (min) etoh
1.			100	97	40
2.			100	98	40
3.			100	70	60
4.			100	98	60
5.			100	98	30
6.			100	98	40
7.			100	98	50
8.			100	99	35
9.			100	85	65
10.			100	95	30

Reactant amount 2 mmol, catalyst amount 0.025 g and solvent 10 mL respectively in each case.

^a yield after 5 reactions.

S7

GCMS

C:\Xcalibur\data\MANOH2

2/15/2007 5:16:12 PM

RT: 0.00 - 43.03

MANOH2 #494 RT: 10.10 AV: 1 NL: 6.00E7

T: + c Full ms [40.00-400.00]

S8

C:\Xcalibur\data\MANOH3

2/15/2007 6:16:02 PM

RT: 0.00 - 43.01

MANOH3 #357 RT: 8.40 AV: 1 NL: 3.85E7

T: + c Full ms [40.00-400.00]

S9

C:\Xcalibur\data\MANP-H4

2/16/2007 11:33:07 AM

RT: 0.00 - 43.01

Mol. Wt.:93

MANP-H4 #148 RT: 5.82 AV: 1 NL: 7.57E7

T: + c Full ms [40.00-400.00]

S10

C:\Xcalibur\data\MANO-H6

2/17/2007 10:27:08 AM

RT: 0.00 - 43.01

MANO-H6 #756 RT: 13.34 AV: 1 NL: 4.74E7

T: + c Full ms [40.00-400.00]

S11

C:\Xcalibur\data\MANO-H6A

2/19/2007 10:06:34 AM

RT: 0.00 - 43.01

MANO-H6A #756 RT: 13.34 AV: 1 NL: 4.59E7

T: + c Full ms [40.00-400.00]

S12

C:\Xcalibur\data\MANO-H8

2/17/2007 12:21:57 PM

RT: 0.00 - 43.01

MANO-H8 #761 RT: 13.40 AV: 1 NL: 6.19E7

T: + c Full ms [40.00-400.00]

S13

C:\Xcalibur\data\MANO-H9

2/17/2007 1:10:59 PM

RT: 0.00 - 43.03

MANO-H9 #1044 RT: 16.90 AV: 1 NL: 3.75E7

T: + c Full ms [40.00-400.00]

S14

C:\Xcalibur\data\MANO-H10

2/17/2007 2:03:10 PM

RT: 0.00 - 43.02

Mol. Wt.: 112

MANO-H10 #95 RT: 5.16 AV: 1 NL: 3.61E7

T: + c Full ms [40.00-400.00]

S15

^1H NMR spectra

^{13}C -H1-SVM-DEBA
 ^{13}C IN CDCl_3
AV-600

S16

13C-H5-SVM-DEBA
13C IN CDCL3
AV-600