Phase-Transfer Catalyzed Asymmetric Glycolate Alkylation

Merritt B. Andrus,* Erik J. Hicken, and Jeffrey C. Stephens

*Brigham Young University, Department of Chemistry and Biochemistry, C100 BNSN,
Provo, Utah 84602-5700*

mbandrus@chem.byu.edu

Supporting Information

Pg. 1-14	Experimentals
Pg. 14-17	X-ray data
Pg. 18-31	NMR data
Pg. 31-35	HPLC chromatograms

General Experimental

Air and water sensitive reactions were performed in flame-dried glassware under a nitrogen atmosphere. Air and moisture sensitive reagents were introduced via dry syringe or cannula. Methylene chloride, toluene, THF, diethyl ether, DMSO, DMF, methanol, triethylamine and acetonitrile were dried by passing through columns of activated alumina. Benzene and pyridine were stored over molecular sieves. Reagents were purchased from Aldrich and Lancaster. Flash chromatography was carried out using 60-230 mesh silica gel. Radial chromatography was performed using 1, 2, and 4 mm plates loaded with 230-400 mesh PF-254 gypsum bound silica. Analytical thin-layer chromatography (TLC) was performed with Merck silica gel 60 F₂₅₄, 0.25 mm pre-coated TLC plates. TLC plates were visualized using UV₂₅₄ and cerium molybdate with charring. All ¹H NMR spectra were obtained with either 200, 300 or 500 MHz Varian spectrometers using TMS (0.0 ppm) or chloroform (7.27 ppm) as an internal reference. Signals are reported as m (multiplet), s (singlet), d (doublet), t (triplet), q (quartet), bs (broad singlet), ABq (AB quartet) obs (obscured); and coupling constants are reported in hertz (Hz). ¹³C NMR were obtained with either 75 or 125 MHz Varian spectrometers using chloroform (77.2 ppm) as the internal standard. Infrared spectra were obtained on a Perkin Elmer FTIR instrument. Mass spectral data (HRMS, CI, EI, FAB) were obtained from the Brigham Young University mass spectrometry facility. Optical rotations were obtained using the sodium D line at ambient temperature. Low temperatures were maintained using an immersion cooler with a cooling probe placed in an acetone bath.

Catalyst Preparation

Park & Jew – ***O*(9)-allyl-*N*-2',3',4'-trifluorobenzylhydrocinchonidinium bromide (4)**. The catalyst was prepared according to the published procedure with the exceptions that in the workup of *N*-(2',3',4'-Trifluoro)benzylhydrocinchonidinium bromide the reaction mixture was diluted with a 1:40 ratio of MeOH :

diethyl ether to precipitate the catalyst. Also the purification of *O*(9)-allyl-*N*-2',3',4'-trifluorobenzylhydrocinchonidinium bromide **4** was accomplished via column chromatography (5% MeOH/CH₂Cl₂) rather than by recrystallization.

Substrate Preparation

Benzhydryloxy-acetic acid (5). To a flame dried round bottom flask was added benzhydrol (1.0 g, 5.42 mmol) and 54 ml of benzene. Then tetrabutylammonium hydrogensulfate (0.092 g, 0.27 mmol) was added with stirring followed by 10 mL of a 50% aq. (w/w) NaOH solution. The reaction was stirred for 15 min then ethyl bromoacetate (0.900 mL, 8.13 mmol) was added dropwise. The solution was allowed to stir at ambient temperature for 6 hours. The resulting thick white solution was then diluted with H₂O and hexanes, the layers were mixed and then separated. The aqueous layer was then acidified, while stirring vigorously, with 6M HCl until a pH of ~ 4 was obtained. Then 1M HCl was added until the pH was between 1.4-1.0, as monitored by pH 0 – 2.5 indicator strips. Next, the resulting white, cloudy solution was extracted with Et₂O (5 x 100 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated to provide 1.03 g (79%) of the title compound as a white powder. Observations by TLC and ¹H NMR concluded that the product was analytically pure and carried on to the next step. Data are: ¹H NMR (DMSO-*d*₆, 300 MHz) δ 12.76 (bs, 1H), 7.40-7.22 (m, 10H), 5.60 (s, 1H), 3.99 (s, 2H); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ 171.3, 141.7, 128.4, 127.5, 126.8, 82.1, 65.2.

2-Benzhydryloxy-1-(2,5-dimethoxyphenyl)-ethanone (1). To a flame dried round bottom flask was added benzhydryloxy-acetic acid **5** (1.94 g, 8.01 mmol) and 32 mL of CH₂Cl₂. The solution was cooled to 0°C and *N,O*-dimethylhydroxylamine hydrochloride (1.17 g, 12.02 mmol) was added in one portion followed by *N,N*-diisopropylethylamine (2.09 mL, 12.02 mmol). Then 4-(dimethylamino)pyridine (0.146 g, 1.20 mmol) was added followed by 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.70g, 12.02 mmol). The reaction was stirred at 0°C for 1 h then warmed to ambient temperature where it stirred for an additional 24 h. The solution was then diluted with CH₂Cl₂ and H₂O. The layers were mixed and separated, and the aqueous layer was extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layers were washed with an aqueous 1M H₃PO₄ solution, then with a saturated aqueous NaHCO₃ solution and finally with a saturated aqueous NaCl solution. The organic layer was then dried over MgSO₄, filtered and concentrated *in vacuo*. Dry THF (40.0 mL) was added to the crude mixture followed by cooling to -40°C. To a separate flame dried round bottom flask was added 1-bromo-2,5-dimethoxybenzene (2.41 mL, 16.02 mmol) and 40.0 mL of THF. The solution was cooled to -40°C and then, with stirring, *n*-BuLi (9.56 mL, 1.6 M in hexanes, 15.3 mmol)

was added dropwise. The solution was allowed to stir for 1 hr then added via cannula to the previously described, pre-cooled solution of the crude 2-benzhydryloxy-*N*-methoxy-*N*-methyl-acetamide solution. The resulting solution was allowed to stir for 15 min at -40°C then quenched by the addition of a 5% HCl/EtOH solution. The reaction was warmed to ambient temperature and the solution partitioned between a saturated aqueous NaCl solution and a 1:1 mixture of Et₂O:CH₂Cl₂. The layers were separated and the aqueous layer extracted with (1:1) Et₂O:CH₂Cl₂ (3 x 50 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated. The crude product was purified via column chromatography (10% EtOAc/hexanes) to produce 2.63 g (91%) of the desired compound as an off-white crystalline solid. Data are: TLC R_f = 0.23 (20% EtOAc/hexanes); ¹H NMR (CDCl₃, 300 MHz) δ 7.43–7.20 (m, 11H), 7.00 (dd, J = 3, 9Hz, 1H), 6.81 (d, J = 9Hz, 1H), 5.62 (s, 1H), 4.71 (s, 1H), 3.75 (s, 3H), 3.69 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 197.7, 153.7, 141.8, 128.5, 127.7, 127.5, 125.9, 121.1, 113.7, 113.1, 83.4, 75.1, 56.0, 55.9; mp = 82–86°C; HRMS (FAB⁺) found 385.1412 [M+Na]⁺, calcd 385.1410 for C₂₃H₂₂O₄Na; Anal. calcd for C₂₃H₂₂O₄: C, 76.22; H, 6.12. Found: C, 75.98; H, 6.09.

Alternative 2 Step Procedure for Substrate Synthesis

2-Benzhydryloxy-*N*-methoxy-*N*-methyl-acetamide. To a flame dried round bottom flask was added benzhydryloxy-acetic acid (6.39 g, 26.4 mmol) and 100 mL of CH₂Cl₂. The solution was cooled to 0°C and *N,O*-dimethylhydroxylamine hydrochloride (3.09 g, 31.7 mmol) was added in one portion followed by *N,N*-diisopropylethylamine (5.52 mL, 31.7 mmol). Then 4-(dimethylamino)pyridine (0.387 g, 3.17 mmol) was added followed by 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (5.06 g, 26.4 mmol). The reaction stirred at 0°C for 1 h then warmed to ambient temperature where it stirred for an additional 2.5 h. The solution was then diluted with 250 mL of CH₂Cl₂ and 200 mL of H₂O. The layers were mixed and separated, and the aqueous layer was extracted with CH₂Cl₂ (3 x 100 mL). The combined organic layers were washed with an aqueous 1M H₃PO₄ solution, then with a saturated aqueous NaHCO₃ solution and finally with a saturated aqueous NaCl solution. The organic layer was then dried over MgSO₄, filtered and concentrated *in vacuo*. Purification via flash column chromatography (30% EtOAc/hexanes) provided 5.32 g (71%) of the pure compound as a thick oil that became off-white crystals in cold conditions. Data are: TLC R_f = 0.17 (20% EtOAc/hexanes); ¹H NMR (CDCl₃, 500 MHz) δ 7.41–7.24 (m, 10H), 5.66 (s, 1H), 4.28 (s, 2H), 3.56 (s, 3H), 3.18 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz) δ 172.2, 141.6, 128.6, 127.8, 127.6, 83.6, 65.9, 61.6, 32.5; HRMS (FAB⁺) found 308.1266 [M+Na]⁺, calcd 308.1257 for C₁₇H₁₉NO₃Na.

2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-ethanone (1). To a flame dried round bottom flask was added 1-bromo-2,5-dimethoxybenzene (2.33 mL, 15.6 mmol) and 10.0 mL of THF. The solution was cooled to -

40°C and then, with stirring, *n*-BuLi (9.1 mL, 1.6 M in hexanes, 14.6 mmol) was added dropwise. The solution was allowed to stir for 1 hr then added via cannula to a pre-cooled solution of 2-benzhydryloxy-*N*-methoxy-*N*-methyl-acetamide (2.97 g, 10.4 mmol) in 10.4 mL of THF at -40°C. The resulting solution was allowed to stir for 15 min at -40°C then quenched by the addition of 30 mL of a 5% HCl/EtOH solution. The reaction was warmed to ambient temperature and the solution partitioned between a saturated aqueous NaCl solution and a 1:1 mixture of Et₂O:CH₂Cl₂. The layers were separated and the aqueous layer extracted with (1:1) Et₂O:CH₂Cl₂ (3 x 50 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated. The crude product was purified via column chromatography (10% EtOAc/hexanes) to produce 3.10 g (82%) of the desired compound as an off-white crystalline solid. Data are: TLC R_f = 0.23 (20% EtOAc/hexanes); ¹H NMR (CDCl₃, 300 MHz) δ 7.43-7.20 (m, 11H), 7.00 (dd, *J* = 3, 9 Hz, 1H), 6.81 (d, *J* = 9 Hz, 1H), 5.62 (s, 1H), 4.71 (s, 2H), 3.75 (s, 3H), 3.69 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 197.7, 153.7, 141.8, 128.5, 127.7, 127.5, 125.9, 121.1, 113.7, 113.1, 83.4, 75.1, 56.0, 55.9; mp = 82–86°C; HRMS (FAB⁺) found 385.1412 [M+Na]⁺, calcd 385.1410 for C₂₃H₂₂O₄Na; Anal. calcd for C₂₃H₂₂O₄: C, 76.22; H, 6.12. Found: C, 75.98; H, 6.09.

Representative Procedure for the Enantioselective Catalytic Phase Transfer Alkylation of 2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-ethanone.

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-3-phenyl-propan-1-one. To a flame dried round bottom flask was added 2-benzhydryloxy-1-(2,5-dimethoxy-phenyl)-ethanone **1** (0.10g, 0.276 mmol), *O*(9)-allyl-*N*-2',3',4'-trifluorobenzylhydrocinchonidinium bromide **4** (15.7 mg, 0.028 mmol), CH₂Cl₂ (1.4 mL) and hexane (1.4 mL). The solution was cooled to -35°C and then CsOH•H₂O (0.232g, 1.38 mmol) was added in one portion. The mixture stirred for 10 min at which time benzyl bromide (0.165 mL, 1.38 mmol) was added dropwise. The mixture stirred at -35°C for 13 h at which time the reaction was diluted with Et₂O (40 mL) and H₂O (15 mL). The layers were mixed and then separated and the organic layer was washed with H₂O (2 x 15 mL) followed by a saturated aqueous solution of NaCl, then dried over MgSO₄. The mixture was filtered, the solvent removed *in vacuo* and the crude residue purified by column chromatography (15% EtOAc/hexane) to afford 0.116 g (93%) of the desired compound as a colorless oil. Data are: TLC R_f = 0.35 (20% EtOAc/hexanes); [α]_D²³ -6.7° (c 2.14, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.32-7.07 (m, 14H), 7.00 (dd, *J* = 3.0, 9.0 Hz, 1H), 6.88-6.80 (m, 3H), 5.41 (s, 1H), 5.14 (dd, *J* = 3.0, 9.6 Hz, 1H), 3.76 (s, 3H), 3.58 (s, 3H), 3.06 (dd, *J* = 2.7, 13.8 Hz, 1H), 2.88 (dd, *J* = 9.6, 14.1 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 201.2, 153.9, 152.7, 142.7, 141.4, 138.5, 129.9, 128.29, 128.27, 127.8, 127.5, 127.4, 127.3, 127.2, 126.5, 120.5, 114.3, 113.4, 82.8, 82.5, 56.1, 56.0, 39.1; HRMS (EI⁺) found 452.1982 M⁺, calcd 452.1988 for C₃₀H₂₈O₄; Anal. calcd for C₃₀H₂₈O₄: C, 79.62; H, 6.24. Found: C, 79.59; H, 6.24. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 7.3 min, *R* (minor) 6.4 min, 93 : 7 er). The absolute configuration was determined by elaboration of the product to known compounds described below.

A larger scale reaction employing ketone **1** (1.00g, 2.76 mmol) and catalyst **4** (0.155g, 0.276 mmol) with identical reaction conditions and reagents provided the title compound as a colorless oil, 1.180g (94%) with matching enantioselectivity. The combined aqueous layers were extracted with CH₂Cl₂ (3 x 50 mL). The combined organic layers were then dried over MgSO₄, filtered and concentrated to provide 0.102g

(71%) of catalyst **4** with the majority of the Br⁻ counterion exchanged for Cl⁻ due to the saturated aqueous NaCl solution contained in the combined aqueous layers.

**Physical Data for Enantioselective Phase Transfer
Alkylation Products 2 From Table 2**

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-4-methyl-pent-4-en-1-one. Following purification via chromatography the product was obtained in 78% yield as a colorless oil. TLC R_f = 0.36 (20% EtOAc/hexanes); $[\alpha]_D^{23}$ -26.5° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.42-7.16 (m, 11H), 6.98 (dd, J = 3.3, 9.0 Hz, 1H), 6.79 (d, J = 9.0 Hz, 1H), 5.52 (s, 1H), 5.12 (dd, J = 4.8, 7.5 Hz, 1H), 4.82-4.77 (m, 2H), 3.75 (s, 3H), 3.55 (s, 3H), 2.40-2.35 (m, 2H), 1.61 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.5, 153.8, 152.6, 144.0, 142.8, 141.8, 141.7, 128.6, 128.4, 128.3, 128.2, 127.8, 127.7, 127.5, 127.3, 127.2, 126.7, 120.3, 114.3, 113.9, 113.2, 82.2, 80.1, 56.0, 55.9, 41.1, 22.5; HRMS (FAB⁺) found 439.1878 [M+Na]⁺, calcd 439.1880 for C₂₇H₂₈O₄Na. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 5.9 min, *R* (minor) 5.4 min, 94 : 6 er).

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-pent-4-en-1-one. Following purification via chromatography the product was obtained in 83% yield as a colorless oil. TLC R_f = 0.31 (20% EtOAc/hexanes); $[\alpha]_D^{23}$ -10.8° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.40-7.16 (m, 10H), 7.12 (d, J = 3.0 Hz, 1H), 6.98 (dd, J = 3.0, 9.0 Hz, 1H), 6.79 (d, J = 9.0 Hz, 1H), 5.95-5.81 (m, 1H), 5.53 (s, 1H), 5.05-4.97 (m, 3H), 3.75 (s, 3H), 3.55 (s, 3H), 2.55-2.32 (m, 2H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.2, 153.8, 152.6, 142.7, 141.8, 134.5, 128.6, 128.5, 128.2, 128.1, 127.8, 127.7, 127.4, 127.3, 126.7, 120.1, 117.3, 114.2, 113.2, 82.3, 81.4, 56.0, 55.9, 37.2; HRMS (FAB⁺) found 425.1715 [M+Na]⁺, calcd 425.1723 for C₂₆H₂₆O₄Na; Anal. calcd for C₂₆H₂₆O₄: C, 77.59; H, 6.51. Found: C, 77.61; H, 6.57. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 0.5 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 12.6 min, *R* (minor) 11.2 min, 91 : 9 er).

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-5-(trimethyl-silanyl)-pent-4-yn-1-one.

Following purification via chromatography the product was obtained in 88% yield as a colorless oil. TLC R_f = 0.34 (20% EtOAc/hexanes); $[\alpha]_D^{23} +26.3^\circ$ (c 1.95, CHCl₃); ¹H NMR (CDCl₃, 500 MHz) δ 7.46-7.28 (m, 10H), 7.15 (d, J = 3.0 Hz, 1H), 7.01 (dd, J = 3.0, 9.0 Hz, 1H), 6.81 (d, J = 9.0 Hz, 1H), 5.65 (s, 1H), 5.16 (dd, J = 5.5, 7.0 Hz, 1H), 3.78 (s, 3H), 3.58 (s, 3H), 2.71-2.69 (m, 2H), 0.16 (s, 9H); ¹³C NMR (CDCl₃, 125 MHz) δ 201.1, 153.8, 152.8, 142.3, 141.5, 128.5, 128.3, 127.8, 127.7, 127.6, 127.5, 127.4, 120.5, 114.1, 113.2, 103.4, 86.5, 82.6, 80.0, 56.0, 55.9, 24.6, 0.19; HRMS (EI⁺) found 472.2078 M⁺, calcd 472.2070 for C₂₉H₃₂O₄. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% IPA/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 4.7 min, *R* (minor) 6.0 min, 91 : 9 er).

Following purification via chromatography the product was obtained in 89% yield as a colorless oil. TLC R_f = 0.23 (20% EtOAc/hexanes); $[\alpha]_D^{23} +10.3^\circ$ (c 0.93, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.45-7.19 (m, 10H), 7.12 (d, J = 3.3 Hz, 1H), 6.97 (dd, J = 3.0, 9.0 Hz, 1H), 6.78 (d, J = 9.0 Hz, 1H), 5.58 (s, 1H), 5.06 (dd, J = 5.1, 6.9 Hz, 1H), 3.75 (s, 3H), 3.57 (s, 3H), 2.59-2.56 (m, 2H), 1.72 (t, J = 2.7 Hz, 3H); ¹³C NMR (CDCl₃, 125 MHz) δ 201.7, 153.9, 152.7, 142.4, 141.7, 128.5, 128.4, 128.2, 128.0, 127.9, 127.8, 127.7, 127.6, 120.2, 114.3, 113.2, 82.8, 80.8, 75.5, 56.2, 56.0, 23.5, 3.8; HRMS (FAB⁺) found 437.1725 [M+Na]⁺, calcd 437.1723 for C₂₇H₂₆O₄Na. Anal. calcd for C₂₇H₂₆O₄: C, 78.24; H, 6.32. Found: C, 78.24; H, 6.17. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 0.5 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 16.8 min, *R* (minor) 15.7 min, 90 : 10 er).

(2*S*)-2-Benzhydryloxy-3-(4-*tert*-butyl-phenyl)-1-(2,5-dimethoxy-phenyl)-propan-1-one.

Following purification via chromatography the product was obtained in 96% yield as a colorless oil. TLC R_f = 0.34 (20% EtOAc/hexanes); $[\alpha]_D^{23} +18.7^\circ$ (c 1.26, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.33-6.97 (m, 14H), 6.87-6.79 (m, 3H), 5.41 (s, 1H), 5.12 (dd, J = 3.3, 9.6 Hz, 1H), 3.75 (s, 3H), 3.58 (s, 3H), 3.02 (dd, J = 3.3, 13.8 Hz, 1H), 2.86 (dd, J = 9.6, 13.8 Hz, 1H), 1.35 (s, 9H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.4, 153.9, 152.7, 149.3, 142.8, 141.5, 135.3, 129.6, 128.3, 128.2, 127.8, 127.4, 127.3, 125.1, 120.4, 114.3, 113.4, 83.0, 82.6, 56.0, 55.9, 38.6, 34.6, 31.7; HRMS (FAB⁺) found 531.2530 [M+Na]⁺, calcd 531.2506 for C₃₄H₃₆O₄Na. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 0.5 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 11.2 min, *R* (minor) 10.5 min, 92 : 8 er).

(2S)-2-Benzhydryloxy-3-biphenyl-2-yl-1-(2,5-dimethoxy-phenyl)-propan-1-one. Following purification via chromatography the product was obtained in 99% yield as a colorless oil. TLC R_f = 0.27 (20% EtOAc/hexanes); $[\alpha]_D^{23}$ -58.9° (c 1.35, CHCl₃); ¹H NMR (CDCl₃, 500 MHz) δ 7.35-7.11 (m, 15H), 7.00-6.98 (m, 2H), 6.88-6.76 (m, 4H), 6.62 (d, J = 9.0 Hz, 1H), 5.33 (s, 1H), 4.87-4.85 (m, 1H), 3.68 (s, 3H), 3.36 (s, 3H), 3.12 (dd, J = 3.0, 14.0 Hz, 1H), 2.94 (dd, J = 10.0, 14.0 Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz) δ 203.9, 153.5, 151.9, 142.7, 142.5, 141.5, 141.4, 135.3, 131.3, 130.3, 129.4, 128.3, 128.1, 128.0, 127.7, 127.4, 127.3, 126.7, 126.5, 119.1, 113.8, 112.8, 82.3, 81.8, 55.9, 55.7, 36.1; HRMS (FAB⁺) found 551.2198 [M+Na]⁺, calcd 551.2193 for C₃₆H₃₂O₄Na. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 7.5 min, *R* (minor) 6.8 min, 95 : 5 er).

(2S)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-3-naphthalen-2-yl-propan-1-one. Following purification via chromatography the product was obtained in 91% yield as a colorless oil. TLC R_f = 0.3 (2:1:7, CH₂Cl₂:Et₂O:hexanes); $[\alpha]_D^{23}$ +40.6° (c 1.1, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.83-7.69 (m, 3H), 7.58 (s, 1H), 7.45-7.41 (m, 2H), 7.33-6.79 (m, 14H), 5.43 (s, 1H), 5.25 (dd, J = 3.3, 9.3 Hz, 1H), 3.72 (s, 3H), 3.59 (s, 3H), 3.22 (dd, J = 3.0, 14.1 Hz, 1H), 3.05 (dd, J = 9.3, 13.8 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.1, 153.9, 152.7, 142.6, 141.2, 135.9, 133.6, 132.4, 128.5, 128.3, 128.2, 128.1, 127.8, 127.7, 127.6, 127.4, 127.3, 127.2, 126.0, 125.5, 120.5, 114.3, 113.4, 82.7, 82.6, 56.1, 55.9, 39.2; HRMS (FAB⁺) found 525.2048 [M+Na]⁺, calcd 525.2036 for C₃₄H₃₀O₄Na. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 9.9 min, *R* (minor) 8.9 min, 92 : 8 er).

(2S)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-5,9-dimethyl-deca-4,8-dien-1-one. Following purification via chromatography the product was obtained in 80% yield as a colorless oil. TLC R_f = 0.39

(20% EtOAc/hexanes); $[\alpha]_{\text{D}}^{23} +9.5^\circ$ (c 1.0, CHCl_3); ^1H NMR (CDCl_3 , 300 MHz) δ 7.41-7.18 (m, 10H), 7.10 (d, $J = 3.0$ Hz, 1H), 6.96 (dd, $J = 3.0, 9.0$ Hz, 1H), 6.78 (d, $J = 9.0$ Hz, 1H), 5.52 (s, 1H), 5.22-5.17 (m, 1H), 5.11-5.07 (m, 1H), 4.97 (t, $J = 6.0$ Hz, 1H), 3.74 (s, 3H), 3.54 (s, 3H), 2.42 (t, $J = 6.3$ Hz, 2H), 2.06-1.92 (m, 4H), 1.67 (d, $J = 0.9$ Hz, 3H), 1.59 (d, $J = 0.9$ Hz, 3H), 1.49 (d, $J = 1.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 202.9, 153.8, 152.5, 142.8, 141.9, 137.6, 131.5, 128.4, 128.2, 128.0, 127.7, 127.4, 127.3, 124.5, 119.9, 119.8, 114.2, 113.2, 82.3, 81.9, 56.0, 55.9, 39.9, 31.7, 26.8, 25.8, 17.9, 16.4; HRMS (FAB^+) found 521.2676 $[\text{M}+\text{Na}]^+$, calcd 521.2662 for $\text{C}_{33}\text{H}_{38}\text{O}_4\text{Na}$. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 0.5 mL/min, 23°C , $\lambda = 254$ nm, retention times: *S* (major) 10.1 min, *R* (minor) 9.1 min, 92 : 8 er).

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-3-(2-methoxy-5-nitro-phenyl)-propan-1-one.

Following purification via chromatography the product was obtained in 91% yield as a white solid. TLC $R_f = 0.2$ (30% EtOAc/hexanes); $[\alpha]_{\text{D}}^{23} -35.6^\circ$ (c 1.2, CHCl_3); ^1H NMR (CDCl_3 , 300 MHz) δ 8.09 (dd, $J = 2.7, 9.0$ Hz, 1H), 8.01 (d, $J = 3.0$ Hz, 1H), 7.31-7.10 (m, 9H), 6.99-6.93 (m, 3H), 6.79-6.75 (m, 2H), 5.42 (s, 1H), 5.25 (dd, $J = 5.4, 8.4$ Hz, 1H), 3.75 (s, 3H), 3.69 (s, 3H), 3.58 (s, 3H), 3.13-2.99 (m, 2H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 202.2, 162.8, 153.8, 152.5, 142.3, 141.4, 141.1, 128.3, 128.2, 127.9, 127.7, 127.3, 127.1, 127.0, 124.2, 120.0, 114.4, 113.0, 109.7, 82.4, 80.3, 56.0, 55.9, 33.1; HRMS (FAB^+) found 550.1844 $[\text{M}+\text{Na}]^+$, calcd 550.1836 for $\text{C}_{31}\text{H}_{29}\text{NO}_7\text{Na}$. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 2.0 mL/min, 23°C , $\lambda = 254$ nm, retention times: *S* (major) 12.2 min, *R* (minor) 15.8 min, 94 : 6 er).

(2*S*)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-hept-4-en-1-one. Following purification via chromatography the product was obtained in 85% yield as a colorless oil. TLC $R_f = 0.35$ (20% EtOAc/hexanes); $[\alpha]_{\text{D}}^{23} +1.0^\circ$ (c 1.8, CHCl_3); ^1H NMR (CDCl_3 , 300 MHz) δ 7.41-7.18 (m, 10H), 7.10 (d, $J = 3.3$ Hz, 1H), 6.97 (dd, $J = 3.3, 9.0$ Hz, 1H), 6.79 (d, $J = 9.0$ Hz, 1H), 5.53 (s, 1H), 5.46-5.43 (m, 2H), 4.96 (dd, $J = 4.2, 7.5$ Hz, 1H), 3.75 (s, 3H), 3.55 (s, 3H), 2.48-2.30 (m, 2H), 2.02-1.94 (m, 2H), 0.94 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 202.6, 153.8, 152.5, 142.7, 141.8, 135.0, 134.1, 128.4, 128.2, 128.1, 127.7, 127.4, 127.3, 124.8, 119.9, 114.2, 113.2, 82.3, 82.0, 56.0, 55.9, 36.1, 25.7, 13.8; HRMS (FAB^+) found 453.2040 $[\text{M}+\text{Na}]^+$, calcd 453.2036 for $\text{C}_{28}\text{H}_{30}\text{O}_4\text{Na}$. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 10% EtOH/hexane, 0.5 mL/min, 23°C , $\lambda = 254$ nm, retention times: *S* (major) 12.2 min, *R* (minor) 12.9 min, 91 : 9 er).

Enolate Trapping with Ethyl Triflate. Following purification via chromatography the product **13** was obtained in 26% yield as a colorless oil. TLC R_f = 0.45 (30% Et₂O/hexanes); ¹H NMR (CDCl₃, 300 MHz) δ 7.44-7.26 (m, 10H), 7.05 (d, J = 3.0 Hz, 1H), 6.81 (s, 1 H), 6.75 (d, J = 9.0 Hz, 1 H), 6.66 (dd, J = 3.0, 9.0 Hz, 1 H), 5.81 (s, 1H), 3.94 (q, J = 7.2 Hz, 2H), 3.75 (s, 3H), 3.63 (s, 3H), 1.35 (t, J = 6.9 Hz, 3H); nOe (500 MHz) irradiation of the signal at 7.05 ppm gave a 0.5% enhancement of the vinyl proton at 6.81 ppm; ¹³C NMR (CDCl₃, 125 MHz) δ 153.9, 150.8, 141.6, 137.5, 135.7, 128.6, 127.9, 127.3, 125.3, 113.7, 112.8, 112.1, 85.6, 67.1, 56.3, 55.9, 15.9; HRMS (FAB⁺) found 413.1724 [M+Na]⁺, calcd 413.1723 for C₂₅H₂₆O₄Na.

Product Elaboration and Determination of the Absolute Configuration.

(2S)-1-(2,5-Dimethoxy-phenyl)-2-hydroxy-3-phenylpropan-1-one (6). To a flame dried round bottom flask was added (3S)-2-benzhydryloxy-1-(2,5-dimethoxy-phenyl)-3-phenylpropan-1-one **2** (0.625 g, 1.38 mmol) and 27 mL of CH₂Cl₂. The solution was cooled to -78°C and then TiCl₄ was added dropwise (0.75 mL, 1.0 M CH₂Cl₂). The resulting dark red/orange solution was stirred at -78°C for 15 min, at which time additional TiCl₄ was added dropwise (0.65 mL, 1.0 M CH₂Cl₂). The reaction was stirred for an another 15 min, then the reaction was quenched by the addition of a saturated aqueous NaHCO₃ solution (25 mL). The layers were separated and the aqueous layer was washed with Et₂O (2 x 25 mL). The combined organic layers were washed with H₂O followed by a saturated aqueous NaCl solution, then dried over MgSO₄, filtered and concentrated *in vacuo*. Chromatography (radial, 4mm plate, 20% EtOAc/hexanes) afforded the title compound, 0.361 g (91%), as a colorless oil. Data are: TLC R_f = 0.33 (20% EtOAc/hexanes); [α]_D²³ - 42.2° (c 1.15, CHCl₃) ¹H NMR (CDCl₃, 300 MHz) δ 7.34 (d, J = 3.3 Hz, 1H), 7.28-7.08 (m, 6H), 6.93 (d, J = 9.0 Hz, 1H), 5.42-5.36 (m, 1H), 3.87 (s, 3H), 3.86 (obs m, 1H), 3.78 (s, 3H), 3.14 (dd, J = 3.6, 14.1 Hz, 1H), 2.70 (dd, J = 7.5, 14.1 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.0, 153.8, 153.4, 137.8, 129.5, 128.3, 126.6, 124.4, 121.9, 114.7, 113.3, 77.6, 56.1, 55.9, 40.8; HRMS (EI⁺) found 286.1200 M⁺, calcd 286.1205 for C₁₇H₁₈O₄.

(2S)-2-Hydroxy-3-phenyl-propionic acid 2,5-dimethoxy-phenyl ester (7). To a flame dried round bottom flask containing activated 4 angstrom molecular sieves and K₂CO₃ (0.097 g, 0.70 mmol) was added trans-N,N-bis(*p*-toluenesulfonyl)-1,2-cyclohexanediamine (0.148 g, 0.35 mmol), followed by CH₂Cl₂ (2.5 mL).

The mixture was cooled to 0°C and then SnCl₄ (0.350 mL, 1.0 M in CH₂Cl₂) was added followed by bis(trimethylsilyl) peroxide (0.150 mL, 0.70 mmol). The mixture was stirred at 0°C for 15 minutes, then (2*S*)-1-(2,5-Dimethoxy-phenyl)-2-hydroxy-3-phenyl-propan-1-one **6** (0.100 g, 0.35 mmol) was added dropwise as a CH₂Cl₂ (3.5 mL) solution. The reaction mixture was stirred at 0°C for 1.5 h, at which time it was quenched by the addition of a saturated aqueous NaHCO₃ solution (9 mL) followed by the addition of a saturated aqueous Na₂S₂O₃ solution (9 mL). The mixture stirred for 0.5 h at ambient temperature then diluted with CH₂Cl₂ (10 mL). The layers were mixed and separated; then the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layers were then washed with H₂O followed by a saturated aqueous NaCl solution, then dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by column chromatography (50% Et₂O/hexanes) to provide 0.083 g (79%) of the title compound as a white solid. The product was put into a round bottom flask, then dissolved in a minimal amount of warm diethyl ether. The flask was then capped loosely and allowed to cool to ambient temperature where it sat for ~5 h. The remaining solution was then decanted from the needle-like crystals and the residual crystals were washed quickly with cold diethyl ether. The crystals were then analyzed by chiral HPLC which revealed the product was enantiomerically enriched to >99% ee. Data are: TLC R_f = 0.23 (50% Et₂O/hexanes); [α]_D²³ – 15.9° (c 0.17, CHCl₃); mp = 128-130°C; ¹H NMR (CDCl₃, 300 MHz) δ 7.37-7.25 (m, 5H), 6.91 (d, *J* = 9.0 Hz, 1H), 6.76 (dd, *J* = 3.3, 9.3 Hz, 1H), 6.57 (d, *J* = 3.0 Hz, 1H), 4.76-4.70 (m, 1H), 3.77 (s, 3H), 3.74 (s, 3H), 3.34 (dd, *J* = 4.5, 14.1 Hz, 1H), 3.15 (dd, *J* = 7.2, 14.1 Hz, 1H), 2.77-2.74 (m, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 172.5, 153.9, 145.3, 139.9, 136.6, 129.9, 128.7, 127.2, 113.6, 112.1, 109.4, 71.5, 56.6, 56.0, 40.7; HRMS (EI⁺) found 302.1155 M⁺, calcd 302.1154 for C₁₇H₁₉O₅. The enantioselectivity was determined by chiral HPLC (DAICEL Chiralpack AD column, 20% IPA/hexane, 1.0 mL/min, 23°C, λ = 254 nm, retention times: *S* (major) 19.0 min, *R* (minor) 17.2 min, >99 : 1 er).

Benzoic acid-(1*S*)-1-benzyl-2-(2,5-dimethoxy-phenyl)-2-oxo-ethyl ester (8**).** To a flame dried round bottom flask was added benzoyl chloride (0.200 mL, 1.73 mmol), 6.0 mL of CHCl₃, and pyridine (0.465 mL, 5.75 mmol). The solution was cooled to 0°C, then (2*S*)-1-(2,5-Dimethoxy-phenyl)-2-hydroxy-3-phenyl-propan-1-one **6** (0.330g, 1.15 mmol) was added slowly as a solution in CHCl₃ (2.0 mL). The reaction was stirred at 0°C for 2.0 h, then quenched by the addition of H₂O (30 mL). The layers were separated and the aqueous layer was extracted with CHCl₃ (2 x 15 mL). The combined organic layers were washed with a saturated aqueous NH₄Cl solution, then dried over MgSO₄, filtered and concentrated. The title compound was obtained after radial chromatography (4mm plate, gradient 10 – 20% EtOAc/hexanes) as a white solid, 0.407 g (90%), which could be recrystallized from CH₂Cl₂. Data are: TLC R_f = 0.37 (30% EtOAc/hexanes); FTIR (film) 3088-2836, 1714, 1669, 1600, 1582, 1495, 1463, 1313, 1276, 1115, 1039, 991 cm⁻¹; [α]_D²³ +13.6° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 8.04-8.01 (m, 2H), 7.57-7.51 (m, 1H), 7.44-7.20 (m, 8H), 7.10 (dd, *J* = 3.3, 9.0 Hz, 1H), 6.95 (d, *J* = 9.3 Hz, 1H), 6.41 (dd, *J* = 3.0, 9.3 Hz, 1H), 3.93 (s, 3H), 3.79 (s, 3H), 3.37 (dd, *J* = 3.3, 14.4 Hz, 1H), 3.13 (dd, *J* = 9.3, 14.4 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 196.3, 166.3, 154.0, 153.4, 137.6, 133.2, 130.1, 130.0, 129.5, 128.6, 128.5, 126.9, 125.5, 121.9, 114.7, 113.4, 80.1, 56.4, 56.0, 36.9; HRMS (EI⁺) found 390.1459 M⁺, calcd 390.1467 for C₂₄H₂₂O₅.

Benzoic acid-(1*S*)-1-(2,5-dimethoxy-phenoxycarbonyl)-2-phenyl-ethyl ester (9). To a flame dried round bottom flask was added activated 4 angstrom molecular sieves (0.200 g), trans-*N,N*-bis(*p*-toluenesulfonyl)-1,2-cyclohexanediamine (0.026 g, 0.062 mmol), K₂CO₃ (0.034 g, 0.246 mmol) and 1.25 mL of CH₂Cl₂. The mixture was cooled to 0°C and SnCl₄ (0.062 mL, 1.0 M in CH₂Cl₂) was added followed by bis(trimethylsilyl) peroxide (0.066 mL, 0.308 mmol). The mixture was stirred at 0°C for 15 minutes, then benzoic acid-(1*S*)-1-benzyl-2-(2,5-dimethoxy-phenyl)-2-oxo-ethyl ester **8** (0.048 g, 0.123 mmol) was added as a CH₂Cl₂ solution (3.0 mL). The reaction stirred at 0°C for 1.0 h at which time the reaction was quenched by the addition of a saturated aqueous NaHCO₃ (6 mL) followed by a saturated aqueous Na₂S₂O₃ solution (6 mL). The mixture was then stirred at ambient temperature for 30 minutes and diluted with 10 mL of CH₂Cl₂ and the layers separated. The aqueous phase was extracted with EtOAc (2 x 20 mL) and the combined organic layers were washed with H₂O followed by a saturated aqueous NaCl solution. The mixture was then filtered, concentrated and purified via flash column chromatography (30% EtOAc/hexanes) to provide 0.037 g (74%) of the title compound as an oil. Data are: TLC R_f = 0.37 (30% Et₂O/hexanes); [α]_D²³ +13.6° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 8.07-8.04 (m, 2H), 7.58-7.53 (m, 1H), 7.45-7.24 (m, 7H), 6.89 (d, *J* = 9.3 Hz, 1H), 6.73 (dd, *J* = 3.0, 9.0 Hz, 1H), 6.61 (d, *J* = 3.3 Hz, 1H), 5.72-5.67 (m, 1H), 3.76 (s, 3H), 3.74 (s, 3H), 3.54 (dd, *J* = 4.2, 14.4 Hz, 1H), 3.43 (dd, *J* = 9.0, 14.4 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 167.9, 166.0, 153.9, 145.4, 139.9, 136.3, 133.6, 130.1, 129.8, 129.5, 128.8, 128.6, 127.3, 113.8, 112.2, 109.3, 73.6, 56.8, 56.0, 37.8; HRMS (EI⁺) found 406.1418 M⁺, calcd 406.1416 for C₂₄H₂₂O₆.

(2*S*)-2-Hydroxy-3-phenyl-propionic acid methyl ester (3). To a flame dried round bottom flask containing benzoic acid-(1*S*)-1-(2,5-dimethoxy-phenoxycarbonyl)-2-phenyl-ethyl ester **9** (0.023 g, 0.057 mmol) was added 0.57 mL of MeOH and 0.57 mL of THF. The solution was cooled to 0°C and then a freshly prepared NaOMe/MeOH solution (1.19 mL, 0.1 M) was added. The solution stirred at 0°C for 1 h then at ambient temperature for 13 h at which time the reaction was quenched by the addition of a saturated aqueous NH₄Cl solution (0.5 mL) and further diluted with 10 mL of H₂O. The resulting solution was extracted with Et₂O (3 x 10 mL) and the combined organic layers were dried over MgSO₄, filtered and concentrated. The crude residue was purified via flash column chromatography (40% Et₂O/hexanes) to provide 0.0084 g (82%) of the title compound as an oil that slowly became a white solid. Data are: TLC R_f = 0.21 (40% Et₂O/hexanes); [α]_D²³ -8.8° (c 0.4, CHCl₃), lit. [α]_D²⁴ = -6.8° (c 1.39, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 7.35-7.21 (m, 5H), 4.50-4.45 (m, 1H), 3.79 (s, 3H), 3.14 (dd, *J* = 4.5, 13.8 Hz, 1H), 2.98 (dd, *J* = 6.9, 13.8 Hz, 1H), 2.70 (d, *J* = 6.3 Hz, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 174.8, 136.5, 129.7, 128.7, 127.2, 71.5, 52.7, 40.8; product

verification and absolute configuration were obtained by comparison to: Yoshikawa, N.; Yamada, Y. M. A.; Das, J.; Sasai, H.; Shibasaki, M. *J. Am. Chem. Soc.* **1999**, *121*, 4168.

Benzoic acid-(1S)-1-methoxycarbonyl-2-phenyl-ethyl ester (12). To a round bottom flask containing benzoic acid-(1S)-1-(2,5-dimethoxy-phenoxy-carbonyl)-2-phenyl-ethyl ester **9** (0.012 g, 0.0295 mmol) was added 0.30 mL of MeOH and 0.30 mL of THF. The solution was cooled to 0°C and then a freshly prepared solution of NaOMe/MeOH (0.006 mL, 0.1M) was added. The solution was then warmed to ambient temperature and stirred for 1 hour, then an additional 0.06 mL of the NaOMe/MeOH solution was added and the solution stirred for 16 h. The reaction was then quenched by the addition of a saturated aqueous NH₄Cl solution (0.25 mL) and then further diluted with H₂O (5 mL). The solution was then extracted with Et₂O (3 x 10 mL) and the combined organic layers were dried over MgSO₄, filtered and concentrated. Purification of the product via preparative TLC (0.25 mm precoated plate, 50% Et₂O/hexanes) produced 0.007 g (81%) of the title compound as a colorless oil. Data are: TLC R_f = 0.45 (40% Et₂O/hexanes); [α]_D²³ -35.5° (c 0.45, MeOH), lit. [α]_D²⁰ = -40.2° (c 1.85, MeOH); ¹H NMR (CDCl₃, 500 MHz) δ 8.03-8.01 (m, 2H), 7.58-7.55 (m, 1H), 7.45-7.42 (m, 2H), 7.33-7.29 (m, 3H), 7.26-7.24 (m, 1H), 5.45 (dd, *J* = 5.0, 8.5 Hz, 1H), 3.74 (s, 3H), 3.31 (dd, *J* = 5.0, 14.0 Hz, 1H), 3.26 (dd, *J* = 8.0, 14.0 Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz) δ 170.3, 166.1, 136.2, 133.6, 130.1, 129.6, 129.5, 128.8, 128.6, 127.3, 73.7, 52.6, 37.8; product verification and absolute configuration were obtained by comparison to: Burk, M. J.; Kalberg, C. S.; Pizzano, A. *J. Am. Chem. Soc.* **1998**, *120*, 4345.

(2S)-1-(2,5-Dimethoxy-phenyl)-2-hydroxy-hept-4-en-1-one. To a round bottom flask containing (2S)-2-Benzhydryloxy-1-(2,5-dimethoxy-phenyl)-hept-4-en-1-one (0.189 g, 0.439 mmol) was added CH₂Cl₂ (8.8 mL). The solution was then cooled to -78°C and TiCl₄ (0.44 mL, 1.0 M CH₂Cl₂) was added dropwise. The resulting dark red/orange solution was stirred at -78°C for 30 min. The reaction was then quenched by the addition of a saturated aqueous NaHCO₃ solution (10 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 15 mL). The combined organic layers were washed with H₂O followed by a saturated aqueous NaCl solution, then dried over MgSO₄, filtered and concentrated *in vacuo*. Chromatography (radial, 1mm plate, 20% EtOAc/hexanes) afforded the title compound, 0.103g (89%), as a colorless oil. Data are: TLC R_f = 0.23 (20% EtOAc/hexanes); [α]_D²³ -64.5° (c 1.0, CHCl₃) ¹H NMR (CDCl₃, 300 MHz) δ 7.31 (d, *J* = 3.3 Hz, 1H), 7.09 (dd, *J* = 3.0, 9.0 Hz, 1H), 6.93 (d, *J* = 9.3 Hz, 1H), 5.42-5.39 (m, 2H), 5.21-5.16 (m, 1H), 3.88 (s, 3H), 3.82 (obs d, *J* = 6.3 Hz, 1H), 3.80 (s, 3H), 2.57-2.49 (m, 1H), 2.23-2.14 (m, 1H), 2.02-1.93 (m, 2H), 0.93 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 202.7, 153.8, 153.3,

135.7, 124.8, 123.7, 121.5, 114.6, 113.2, 76.8, 56.1, 55.9, 37.8, 25.8, 13.9; HRMS (EI⁺) found 287.1250 M⁺, calcd 287.1254 for C₁₅H₂₀O₄Na.

(1S)-Benzoic acid 1-(2,5-dimethoxy-benzoyl)-hex-3-enyl ester (10). To a round bottom flask containing (2S)-1-(2,5-Dimethoxy-phenyl)-2-hydroxy-hept-4-en-1-one (0.073 g, 0.276 mmol) was added 1.4 mL of CHCl₃ and pyridine (0.112 mL, 1.38 mmol). The solution was cooled to 0°C, then benzoyl chloride (0.048 mL, 0.414 mmol) was added slowly. The reaction was stirred at 0°C for 2.5 h, then quenched by the addition of H₂O (10 mL). The layers were separated and the aqueous layer was extracted with CHCl₃ (2 x 10 mL). The combined organic layers were washed with a saturated aqueous NH₄Cl solution, then dried over MgSO₄, filtered and concentrated. The title compound was obtained after radial chromatography (1mm plate, 20% EtOAc/hexanes) as a white solid, 0.094 g (92%). Data are: mp = 68-72°C; [α]_D²³ +25.2° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 8.11-8.08 (m, 2H), 7.59-7.54 (m, 1H), 7.47-7.42 (m, 2H), 7.38 (d, *J* = 3.3 Hz, 1H), 7.08 (dd, *J* = 3.0, 9.0 Hz, 1H), 6.92 (d, *J* = 9.3 Hz, 1H), 6.25-6.21 (m, 1H), 5.63-5.47 (m, 2H), 3.89 (s, 3H), 3.79 (s, 3H), 2.76-2.70 (m, 1H), 2.62-2.53 (m, 1H), 2.06-1.97 (m, 2H), 0.94 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 196.5, 166.3, 153.8, 153.2, 136.0, 133.2, 130.2, 129.9, 128.5, 125.7, 123.7, 121.4, 114.5, 113.2, 79.3, 56.3, 55.9, 34.2, 25.8, 13.9.

(1S)-Benzoic acid 1-(2,5-dimethoxy-phenoxy-carbonyl)-hex-3-enyl ester (11). To a flame dried round bottom flask was added activated 4 angstrom molecular sieves (0.100 g), trans-N,N-bis(*p*-toluenesulfonyl)-1,2-cyclohexanediamine (0.027 g, 0.065 mmol), K₂CO₃ (0.036 g, 0.260 mmol) and 1.25 mL of CH₂Cl₂. The mixture was cooled to 0°C and SnCl₄ (0.065 mL, 1.0 M in CH₂Cl₂) was added followed by bis(trimethylsilyl) peroxide (0.070 mL, 0.325 mmol). The mixture was stirred at 0°C for 5 minutes, then (1S)-Benzoic acid 1-(2,5-dimethoxy-benzoyl)-hex-3-enyl ester **10** (0.048 g, 0.130 mmol) was added as a CH₂Cl₂ solution (2.0 mL). The reaction stirred at 0°C for 20 min, at which time the reaction was quenched by the addition of a saturated aqueous NaHCO₃ (5 mL) followed by a saturated aqueous Na₂S₂O₃ solution (5 mL). The mixture was stirred at ambient temperature for 30 minutes then diluted with 10 mL of CH₂Cl₂ and the layers separated. The aqueous phase was extracted with CH₂Cl₂ (2 x 20 mL) and the combined organic layers were washed with H₂O followed by a saturated aqueous NaCl solution. The mixture was then filtered, concentrated and purified via flash column chromatography (30% Et₂O/hexanes) to provide 0.037 g (74%) of the title compound as a colorless oil. Data are: [α]_D²³ -11.1° (c 1.0, CHCl₃); ¹H NMR (CDCl₃, 300 MHz) δ 8.14-8.11 (m, 2H), 7.62-7.57 (m, 1H), 7.49-7.44 (m, 2H), 6.90 (d, *J* = 9.0 Hz, 1H), 6.75 (dd, *J* = 3.0, 9.0 Hz, 1H), 6.70 (d, *J* = 3.0 Hz, 1H), 5.82-5.72 (m, 1H), 5.65-5.50 (m, 2H), 3.78 (s, 3H), 3.76 (s, 3H), 2.90-2.86 (m,

2H), 2.08 (m, 2H), 1.00 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 168.1, 166.1, 153.9, 145.4, 140.0, 137.1, 133.5, 130.1, 129.7, 128.6, 122.5, 113.7, 112.1, 109.4, 72.6, 56.8, 56.0, 34.9, 25.9, 13.9.

X-ray structure determination of intermediate 8.

Table 1. Crystal data and structure refinement for cmpd#8

Identification code	EJH24CU	
Empirical formula	C ₂₄ H ₂₂ O ₅	
Formula weight	390.2	
Temperature	100(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	P2(1)	
Unit cell dimensions	a = 7.1604(3) Å	$\alpha = 90^\circ$.
	b = 12.9607(5) Å	$\beta = 104.743(2)^\circ$.
	c = 11.3097(5) Å	$\gamma = 90^\circ$.
Volume	1015.03(7) Å ³	
Z	2	
Density (calculated)	1.277 Mg/m ³	
Absorption coefficient	0.727 mm ⁻¹	
F(000)	412	
Crystal size	0.18 x 0.15 x 0.10 mm ³	
Theta range for data collection	4.04 to 55.78°.	
Index ranges	-7 ≤ h ≤ 7, -13 ≤ k ≤ 13, -11 ≤ l ≤ 11	
Reflections collected	7137	
Independent reflections	2512 [R(int) = 0.0285]	
Completeness to theta = 55.78°	97.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9301 and 0.8833	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2503 / 1 / 265	
Goodness-of-fit on F ²	1.153	
Final R indices [I > 2σ(I)]	R1 = 0.0362, wR2 = 0.0795	
R indices (all data)	R1 = 0.0362, wR2 = 0.0796	
Absolute structure parameter	0.05(13)	
Extinction coefficient	0.083(3)	
Largest diff. peak and hole	0.225 and -0.310 e.Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for EJH24Cu. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
C(1)	6906(2)	1498(1)	5742(2)	30(1)
C(2)	6840(2)	548(1)	6295(2)	34(1)
C(3)	6568(2)	-353(1)	5608(2)	39(1)
C(4)	6376(2)	-314(2)	4361(2)	42(1)
C(5)	6408(2)	631(2)	3796(2)	40(1)
C(6)	6669(2)	1527(1)	4486(2)	34(1)
C(7)	7251(2)	2483(1)	6481(2)	29(1)
C(8)	9393(2)	2774(1)	6876(2)	27(1)
C(9)	12273(2)	1996(1)	8002(1)	26(1)
C(10)	13141(2)	1145(1)	8843(1)	28(1)
C(11)	12035(2)	363(1)	9172(2)	32(1)
C(12)	12936(3)	-428(1)	9927(2)	38(1)
C(13)	14924(3)	-446(2)	10353(2)	43(1)
C(14)	16026(3)	333(2)	10039(2)	44(1)
C(15)	15148(2)	1127(1)	9282(2)	36(1)
C(16)	9660(2)	3826(1)	7503(2)	27(1)
C(17)	9076(2)	4791(1)	6776(1)	27(1)
C(18)	8978(2)	5687(1)	7467(2)	29(1)
C(19)	8521(2)	6628(1)	6908(2)	30(1)
C(20)	8161(2)	6699(1)	5647(2)	32(1)
C(21)	8230(2)	5838(1)	4952(2)	30(1)
C(22)	8669(2)	4876(1)	5498(2)	29(1)
C(23)	8399(2)	4108(1)	3537(1)	34(1)
C(24)	8517(4)	7471(2)	8784(2)	52(1)
O(1)	10327(1)	1991(1)	7731(1)	28(1)
O(2)	13192(2)	2619(1)	7587(1)	31(1)
O(3)	10275(2)	3866(1)	8610(1)	32(1)
O(4)	8728(2)	4001(1)	4832(1)	32(1)
O(5)	8360(2)	7535(1)	7509(1)	39(1)

Table 3. Bond lengths [Å] and angles [°] for EJH24Cu.

C(1)-C(2)	1.387(2)
C(1)-C(6)	1.388(2)
C(1)-C(7)	1.510(2)
C(2)-C(3)	1.389(3)
C(3)-C(4)	1.382(3)
C(4)-C(5)	1.385(3)
C(5)-C(6)	1.385(3)
C(7)-C(8)	1.532(2)
C(8)-O(1)	1.4438(19)
C(8)-C(16)	1.526(2)
C(9)-O(2)	1.209(2)
C(9)-O(1)	1.3484(19)
C(9)-C(10)	1.485(2)
C(10)-C(11)	1.394(2)
C(10)-C(15)	1.397(2)
C(11)-C(12)	1.384(3)
C(12)-C(13)	1.382(3)
C(13)-C(14)	1.383(3)
C(14)-C(15)	1.384(2)
C(16)-O(3)	1.2173(19)

C(16)-C(17)	1.497(2)
C(17)-C(22)	1.404(2)
C(17)-C(18)	1.411(2)
C(18)-C(19)	1.375(2)
C(19)-O(5)	1.377(2)
C(19)-C(20)	1.386(2)
C(20)-C(21)	1.372(2)
C(21)-C(22)	1.392(2)
C(22)-O(4)	1.368(2)
C(23)-O(4)	1.4305(19)
C(24)-O(5)	1.420(2)
C(2)-C(1)-C(6)	118.43(15)
C(2)-C(1)-C(7)	121.37(14)
C(6)-C(1)-C(7)	120.20(14)
C(1)-C(2)-C(3)	120.71(16)
C(4)-C(3)-C(2)	120.18(18)
C(3)-C(4)-C(5)	119.67(17)
C(4)-C(5)-C(6)	119.79(16)
C(5)-C(6)-C(1)	121.19(16)
C(1)-C(7)-C(8)	112.37(12)
O(1)-C(8)-C(16)	109.59(12)
O(1)-C(8)-C(7)	106.32(12)
C(16)-C(8)-C(7)	110.74(13)
O(2)-C(9)-O(1)	123.22(14)
O(2)-C(9)-C(10)	124.35(14)
O(1)-C(9)-C(10)	112.43(14)
C(11)-C(10)-C(15)	119.76(14)
C(11)-C(10)-C(9)	122.51(14)
C(15)-C(10)-C(9)	117.70(14)
C(12)-C(11)-C(10)	119.74(15)
C(13)-C(12)-C(11)	120.33(17)
C(12)-C(13)-C(14)	120.17(17)
C(13)-C(14)-C(15)	120.22(17)
C(14)-C(15)-C(10)	119.78(17)
O(3)-C(16)-C(17)	120.49(14)
O(3)-C(16)-C(8)	119.03(13)
C(17)-C(16)-C(8)	120.41(13)
C(22)-C(17)-C(18)	118.62(14)
C(22)-C(17)-C(16)	125.97(14)
C(18)-C(17)-C(16)	115.39(13)
C(19)-C(18)-C(17)	121.01(15)
C(18)-C(19)-O(5)	124.79(14)
C(18)-C(19)-C(20)	119.45(15)
O(5)-C(19)-C(20)	115.76(14)
C(21)-C(20)-C(19)	120.72(14)
C(20)-C(21)-C(22)	120.78(14)
O(4)-C(22)-C(21)	122.30(14)
O(4)-C(22)-C(17)	118.30(13)
C(21)-C(22)-C(17)	119.39(14)
C(9)-O(1)-C(8)	115.14(12)
C(22)-O(4)-C(23)	117.57(13)
C(19)-O(5)-C(24)	117.19(13)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for EJH24Cu. The anisotropic

displacement factor exponent takes the form: $-2\pi^2[h^2 a^{*2}U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
C(1)	22(1)	31(1)	35(1)	0(1)	5(1)	1(1)
C(2)	30(1)	34(1)	35(1)	-1(1)	3(1)	0(1)
C(3)	31(1)	32(1)	52(1)	-2(1)	7(1)	-1(1)
C(4)	30(1)	42(1)	54(1)	-18(1)	11(1)	-8(1)
C(5)	32(1)	50(1)	37(1)	-9(1)	9(1)	-7(1)
C(6)	30(1)	35(1)	36(1)	0(1)	5(1)	-5(1)
C(7)	30(1)	28(1)	29(1)	1(1)	7(1)	3(1)
C(8)	30(1)	25(1)	25(1)	3(1)	5(1)	3(1)
C(9)	31(1)	26(1)	22(1)	-4(1)	5(1)	0(1)
C(10)	35(1)	28(1)	22(1)	-2(1)	7(1)	4(1)
C(11)	36(1)	30(1)	28(1)	-1(1)	7(1)	1(1)
C(12)	51(1)	33(1)	32(1)	4(1)	15(1)	3(1)
C(13)	50(1)	40(1)	34(1)	8(1)	5(1)	13(1)
C(14)	39(1)	47(1)	41(1)	8(1)	2(1)	10(1)
C(15)	37(1)	37(1)	31(1)	4(1)	5(1)	2(1)
C(16)	25(1)	29(1)	27(1)	2(1)	6(1)	0(1)
C(17)	26(1)	29(1)	26(1)	2(1)	6(1)	-1(1)
C(18)	30(1)	32(1)	24(1)	-1(1)	6(1)	-2(1)
C(19)	33(1)	25(1)	34(1)	0(1)	10(1)	-1(1)
C(20)	33(1)	29(1)	36(1)	7(1)	12(1)	2(1)
C(21)	30(1)	34(1)	25(1)	4(1)	6(1)	0(1)
C(22)	26(1)	30(1)	31(1)	-1(1)	7(1)	-2(1)
C(23)	40(1)	39(1)	24(1)	0(1)	8(1)	1(1)
C(24)	88(2)	34(1)	37(1)	-4(1)	18(1)	8(1)
O(1)	30(1)	27(1)	26(1)	3(1)	5(1)	2(1)
O(2)	31(1)	29(1)	31(1)	1(1)	6(1)	-1(1)
O(3)	38(1)	30(1)	26(1)	0(1)	4(1)	2(1)
O(4)	43(1)	30(1)	23(1)	0(1)	8(1)	0(1)
O(5)	60(1)	26(1)	34(1)	0(1)	16(1)	3(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for EJH24Cu.

	x	y	z	U(eq)
H(2)	6979	514	7134	41
H(3)	6514	-985	5987	47
H(4)	6226	-919	3905	50
H(5)	6255	665	2955	47
H(6)	6686	2160	4101	41
H(7A)	6766	2399	7202	35
H(7B)	6534	3039	5996	35
H(8)	9940	2782	6165	33
H(11)	10697	372	8885	38
H(12)	12200	-950	10148	45
H(13)	15521	-983	10853	51
H(14)	17362	323	10337	52
H(15)	15893	1647	9066	43
H(18)	9227	5639	8313	35
H(20)	7869	7336	5268	38

H(21)	7980	5899	4107	36
H(23A)	7096	4333	3194	52
H(23B)	8601	3456	3187	52
H(23C)	9279	4609	3360	52
H(24A)	7548	7010	8927	79
H(24B)	8335	8144	9092	79
H(24C)	9774	7217	9194	79

NMR Spectra

Chiral HPLC Chromatograms

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	6.416	N	4253	61324	6.961
2	7.370	N	44235	819563	93.038
Total Area				880887	99.999

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	6.256	N1	30376	468403	48.821
2	7.160	N2	27087	491013	51.178
Total Area				959416	99.999

Data:

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	15.740	*N1	4196	129223	9.865
2	16.846	*N2	32556	1180675	90.134
Total Area				1309898	99.999

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	16.045	N1	20624	666222	48.944
2	17.191	N2	19162	694955	51.055
Total Area				1361177	99.999

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	4.735	N	49972	570242	90.769
2	5.981	N	4112	57992	9.230
Total Area				628234	99.999

Data:

Analysis: Channel A

Peak No.	Time	Type	Height(μ V)	Area(μ V-sec)	Area%
1	4.736	N	25329	285607	50.492
2	5.986	N	19101	280038	49.507
Total Area				565645	99.999