Supporting Information

Phosphoric Acid-Promoted Synthesis of 4-Acylpyrroles and Dipyrrolylketones from Mixed Anhydrides

Cory S. Beshara and Alison Thompson*

Department of Chemistry, Dalhousie University, Halifax, Nova Scotia,

Canada B3H 4J3 alison.thompson@dal.ca

Table of Contents

<i>General Experimental Procedures</i>	<i>2</i>
<i>Non-duplicate pyrroles synthesized via Knight's method</i>	<i>2</i>
4-(4-Ethoxy-benzoyl)-3,5-dimethyl-1H-pyrrole-2-carboxylic acid ethyl ester (4d)	2
¹³ C NMR spectra.....	3
¹³ C NMR spectrum of 4b	3
¹³ C NMR spectrum of 4c	4
¹³ C NMR spectrum of 4d	5
¹³ C NMR spectrum of 4e	6
¹³ C NMR spectrum of 4f	7
¹³ C NMR spectrum of 4g	8
¹³ C NMR spectrum of 4h	9
¹³ C NMR spectrum of 4i	10
¹³ C NMR spectrum of 7	11
¹³ C NMR spectrum of 8	12
¹³ C NMR spectrum of 9	13
<i>References.....</i>	<i>14</i>

General Experimental Procedures

All acids and TFAA were used as purchased. Reagent-grade acetonitrile was purchased from Caledon and used without further purification. Pyrroles **1**,¹ **3**,² and **10**³ were made following known literature procedures. Pyrrole **2** is a known compound.⁴ The syntheses of **5**³ and **6**⁵ were achieved by utilizing hydrogenolytic conditions⁶ from the corresponding benzyl esters: 3,5-dimethyl-1*H*-pyrrole-2,4-dicarboxylic acid 2-benzyl ester 4-ethyl ester⁷ (yields **5**) and 4-acetyl-3,5-dimethyl-1*H*-pyrrole-2-carboxylic acid benzyl ester⁸ (yields **6**). Melting points are uncorrected. ¹H and ¹³C NMR spectra were obtained at 500 and 125 MHz, respectively. Chemical shifts (δ) are referenced to TMS at 0 ppm and all coupling constants are reported in hertz. Silica gel used for column flash chromatography was silica gel 60, 230-400 mesh. TLC was performed using aluminum-backed plates and visualized under UV light and a vanillin in ethanol dip. Electrospray mass spectra were obtained using an ion trap instrument. All reported R_f values are as eluted with 3:7 ethyl acetate:hexanes.

Non-duplicate pyrroles synthesized *via* Knight's method

4-(4-Ethoxy-benzoyl)-3,5-dimethyl-1*H*-pyrrole-2-carboxylic acid ethyl ester (**4d**)

Isolation of the title compound as a an off white solid (0.126 g, 67%). mp: 183-185°C; δ_{H} (CDCl₃): 1.37 (3H, t, J 7), 1.44 (3H, t, J 7), 2.25 (3H, s), 2.26 (3H, s), 4.10 (2H, q, J 7), 4.34 (2H, q, J 7), 6.92 (2H, d, J 5), 7.74 (2H, d, J 5), 9.84 (1H, bs); δ_{C} (CDCl₃): 12.6, 13.5, 14.7, 15.0, 60.6, 64.0, 114.3, 118.6, 123.8, 129.2, 132.0, 132.9, 136.6, 162.4, 162.8, 193.4; *m/z* 652.9 (dimer as Na adduct); R_f 0.30.

^{13}C NMR spectra ^{13}C NMR spectrum of 4b

^{13}C NMR spectrum of 4c

^{13}C NMR spectrum of 4d

^{13}C NMR spectrum of 4e

^{13}C NMR spectrum of 4f

^{13}C NMR spectrum of 4g

^{13}C NMR spectrum of 4h

^{13}C NMR spectrum of 4i

^{13}C NMR spectrum of 7

^{13}C NMR spectrum of 8

^{13}C NMR spectrum of 9

References

- (1) Smith, K. M.; Miura, M.; Tabb, H. D. *J. Org. Chem.* **1983**, *48*, 4779-4781.
- (2) Thompson, A.; Alattar, Y.; Beshara, C. S.; Burley, R. K.; Cameron, T. S.; Robertson, K. N. *J. Heterocycl. Chem.* **2004**, *41*, 777-781.
- (3) Cordell, G. A. *J. Org. Chem.* **1975**, *40*, 3161-3169.
- (4) Omote, M.; Ando, A.; Sato, K.; Kumadaki, I. *Tetrahedron* **2001**, *57*, 8085-8094.
- (5) Badger, G. M.; Harris, R. L. N.; Jones, R. A. *Aust. J. Chem.* **1964**, *17*, 987-1001.
- (6) Thompson, A.; Dolphin, D. *J. Org. Chem.* **2000**, *65*, 7870-7877.
- (7) Rezzano, I.; Buldain, G.; Frydman, B. *J. Org. Chem.* **1982**, *47*, 3059-3063.
- (8) Awruch, J.; Frydman, B. *Tetrahedron* **1986**, *42*, 4137-4146.