Supporting Information

Photoinduced Diastereoselective Addition of Perfluoroalkyl Iodides to Acrylic Acid Derivatives for the Synthesis of Fluorinated Amino Acids

Tomoko Yajima* and Hajime Nagano

Department of Chemistry, Faculty of Science, Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo
112-8610, Japan

Contents

Experiment Procedure and Characterization Data	S1
¹ H-NMR and ¹³ C-NMR Spectra	S10

General methods.

All reactions were performed with a high-pressure Hg lamp (Toshiba, 460W). ¹H NMR spectra were recorded on a JEOL AL-400 (400 MHz) or EX-400 (400 MHz) spectrometer with CDCl₃ as the solvent and tetramethylsilane as an internal standard unless otherwise noted. ¹³C NMR spectra were recorded on the instruments operating at 100.5 MHz with CDCl₃ as the solvent and internal standard (δ 77.0). IR spectra were taken on a SHIMADZU FTIR-8700 spectrometer. Mass spectra (EI⁺) were obtained on a JEOL JMS-700 mass spectrometer. Precoated Merck Kieselgel 60 F254 and Kanto silica gel 60 (spherical neutral) were used for thin layer chromatography and flash chromatography, respectively. (1*R*,2*S*,5*R*)-Menthol and phenethyl alcohol were purchased from Kanto Chemical Co. Ethyl acrylate and (*R*)-4-benzyl-2-oxazolidinone were purchased from Wako Pure Chemical Co. Pantolactone, (-)-8-phenylmenthyl and (-)-10.2-camphorsultam were purchased from Aldrich. Perfluoro-*n*-hexyl iodide, heptafluoro-*i*-propyl iodide and pentafluoroethyl iodide were purchased from Daikin Chemical Sales LTD. Heptafluoro-*n*-propyl iodide was purchased from Tokyo Chemical Industry Co. Trifluoromethyl iodide was purchased from Tosoh F-tech inc.

General procedure for iodoperfluoroalkylation.

In a Pyrex glass tube were placed olefin (0.2 mmol), perfluoroalkyl iodide (2.0 mmol) and CH₂Cl₂ (5 mL). Then Na₂S₂O₃ (1 mmol, 158 mg) and water (1 mL) were added to the mixture. After sealing the tube, the mixture was mixed by shaking and then irradiated with a Hg lamp at room temperature. After the reaction was completed, the mixture was extracted with CH₂Cl₂. The extract was washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to afford pure product.

(1*R*,2*S*,5*R*)-2-Isopropyl-5-methylcyclohexyl 2-iodo-3-perfluorohexylpropanoate (2) (50:50 diastereomer mixture)

White powder; mp. 56 °C (from CHCl₃); $[\alpha]_D^{24}$ -21.4 (*c* 1.22, CHCl₃); ¹H NMR (400 MHz) δ 0.74 (d, *J* = 6.8 Hz, 3H, CH₃), 0.77 (d, *J* = 6.8 Hz, 3H, CH₃), 0.87–0.93 (m, 14H, CH₃×4, cyclohexyl), 0.95–1.10 (m, 4H, cyclohexyl), 1.46 (m, 4H, cyclohexyl), 1.70 (m, 4H, cyclohexyl), 1.85–2.07 (m, 4H, cyclohexyl), 3.35 (m, 2H, C₆F₁₃CHH×2), 3.69 (m, 2H, C₆F₁₃CHH×2), 4.57 (dd, *J* = 3.2, 10.8 Hz, 1H, OCH), 4.58 (dd, *J* = 3.2, 10.8 Hz, 1H, OCH), 4.70 (dt, *J* = 4.4, 10.4 Hz, 1H, ICH), 4.77 (dt, *J* = 4.4, 10.4 Hz, 1H, ICH); ¹³C NMR (100 MHz) δ 5.7, 6.2, 15.9, 16.0, 20.7, 20.8, 21.98, 22.00, 23.2, 23.4, 25.9, 26.1, 31.3, 31.4, 34.18, 34.19, 38.4 (t, *J*_{C-F} = 22.7 Hz, C₆F₁₃C), 38.6 (t, *J*_{C-F} = 22.7 Hz, C₆F₁₃C), 39.4, 40.0, 47.0, 47.1, 76.5, 76.6, 169.5, 169.7; IR (neat, cm⁻¹) 2961, 1737, 1461, 1238, 1147, 962, 700, 656, 519; MS *m/z* 501 (M⁺ – C₁₀H₁₉O (menthyloxy) 3%), 473 (17), 139 (25), 138 (100); HRMS calcd for C₉H₃F₁₃IO (M⁺ – C₁₀H₁₉O) 500.9021, found 500.9007.

Dehydroiodination of 2: (1*R*,2*S*,5*R*)-2-Isopropyl-5-methylcyclohexyl (2*E*)-3-perfluorohexyl-2-propenoate (3)

To a solution of **2** (50 mg, 0.08 mmol) in CH₂Cl₂ (1.0 mL) was added DBU (21 μL, 0.8 mmol). After stirring for 3 h, the reaction mixture was concentrated under reduced pressure. The residue was chromatographed on silica gel to afford pure product **3** (40.0 mg, 99%).

Colorless oil; $[\alpha]_D^{23}$ -34.0 (*c* 0.80, CHCl₃); ¹H NMR (400 MHz) δ 0.78 (d, *J* = 6.8 Hz, 3H, CH₃), 0.88 (m, 1H, cyclohexyl), 0.89 (d, *J* = 6.8 Hz, 3H, CH₃), 0.93 (d, *J* = 6.8 Hz, 3H, CH₃), 1.07 (m, 2H, cyclohexyl), 1.45 (m, 1H, cyclohexyl), 1.52 (m, 1H, cyclohexyl), 1.71 (m, 2H, cyclohexyl), 1.84 (dsep, *J* = 2.8, 6.8 Hz, 1H, (CH₃)₂CH), 2.04 (m, 1H, cyclohexyl), 4.82 (dt, *J* = 4.0, 11.2 Hz, 1H, OCH), 6.53 (dt, *J* = 16.4, 2.0 Hz, 1H, COCH=), 6.82 (dt, *J* = 16.4, 12.0 Hz, 1H, C₆F₁₃CH=); ¹³C NMR (100 MHz) δ 16.4, 20.7, 22.0, 23.5, 26.4, 31.5, 34.2, 40.7, 47.0, 76.0, 130.3 (t, *J*_{C-F} = 23.1 Hz, C₆F₁₃C), 131.2 (t, *J*_{C-F} = 7.4 Hz, C₆F₁₃C), 163.1; IR (neat, cm⁻¹) 2961, 1731, 1241, 1200, 1146, 977; MS *m/z* 513 (M⁺ – CH₃, 1%), 373 (M⁺ – C₁₀H₁₉O (menthyloxy), 100); HRMS calcd for C₁₈H₁₈F₁₃O₂ (M⁺ – CH₃) 513.1099, found 513.1056.

1-Phenylethyl 2-iodo-3-perfluorohexylpropanoate (5) (50:50 diastereomer mixture)

Colorless oil; ¹H NMR (400 MHz) δ 1.57 (d, *J* = 6.3 Hz, 3H, CH₃), 1.58 (d, *J* = 6.3 Hz, 3H, CH₃), 2.68 (m, 2H, C₆F₁₃CHH×2), 3.31 (m, 2H, C₆F₁₃CHH×2), 4.62 (q, *J* = 6.3 Hz, 1H, PhCH), 4.65 (q, *J* = 6.3 Hz, 1H, PhCH), 5.92 (m, 2H, ICH×2), 7.32 (m, 10H, arom); ¹³C NMR (100 MHz) δ 5.7, 5.9, 21.0, 22.0, 38.4 (t, *J*_{C-F} = 25.7 Hz, C₆F₁₃C), 38.6 (t, *J*_{C-F} = 25.7 Hz, C₆F₁₃C), 74.6, 74.7, 126.0, 126.2, 128.2, 128.3, 128.4, 128.5, 140.0, 140.1, 169.1, 169.2; IR (neat, cm⁻¹) 2987, 1739, 1239, 1199, 1146, 698; MS *m/z* 495

($M^+ - I$, 75%), 121 (51), 105 (100); HRMS calcd for $C_{17}H_{12}O_2F_{13}$ ($M^+ - I$) 495.0630, found 495.0641.

4,4-Dimethyl-2-oxotetrahydrofuran-3-yl 2-iodo-3-perfluorohexylpropanoate (7)

(60:40 diastereomer mixture)

Colorless oil; IR (neat, cm^{-1}) 1799, 1755, 1241, 1203, 1146; MS m/z 630 (M^+ , 42%), 503 ($M^+ - I$, 18), 500 (31), 473 (85), 373 (80), 86 (100); HRMS calcd for $C_{15}H_{12}O_4F_{13}I$ (M^+) 629.9573, found 629.9601. **Major diastereomer.** 1H NMR (400 MHz) δ 1.21 (s, 3H, CH_3), 1.27 (s, 3H, CH_3), 2.78 (m, 1H, $C_6F_{13}CHH$), 3.33 (m, 1H, $C_6F_{13}CHH$), 4.07 (m, 2H, OCH_2), 4.82 (dd, $J = 4.4, 9.6$ Hz, 1H, ICH), 5.42 (s, 1H, OCH); ^{13}C NMR (100 MHz) δ 3.2 (t, $J_{C-F} = 7.8$ Hz, $C_6F_{13}CH_2C$), 19.7, 22.90, 38.0 (t, $J_{C-F} = 23.4$ Hz, $C_6F_{13}C$), 40.8, 76.1, 76.2, 169.54, 170.9. **Minor diastereomer.** 1H NMR (400 MHz) δ 1.15 (s, 3H, CH_3), 1.23 (s, 3H, CH_3), 2.78 (m, 1H, $C_6F_{13}CHH$), 3.33 (m, 1H, $C_6F_{13}CHH$), 4.07 (m, 2H, OCH_2), 4.71 (dd, $J = 3.2, 9.6$ Hz, 1H, ICH), 5.38 (s, 1H, OCH); ^{13}C NMR (100 MHz) δ 3.4 (t, $J_{C-F} = 7.8$ Hz, $C_6F_{13}CH_2C$), 19.7, 22.88, 38.7 (t, $J_{C-F} = 23.4$ Hz, $C_6F_{13}C$), 40.5, 76.0, 76.3, 169.48, 171.0.

(1R,2S,5R)-5-Methyl-2-(1-methyl-1-phenylethyl)cyclohexyl 2-iodo-3-perfluorohexylpropanoate (9)

(24:76 diastereomer mixture)

Colorless oil; IR (neat, cm^{-1}) 2961, 1733, 1241, 1205, 1146, 701; MS m/z 613 ($M^+ - C_9H_{11}$, 8%), 605 ($M^+ - I$, 8), 473 (25), 437 (13), 214 (31), 119 (100), 91 (69); HRMS calcd for $C_{16}H_{15}F_{13}IO_2$ ($M^+ - C_9H_{11}$) 612.9909, found 612.9958; $[\alpha]^{24}_D +18.0$ (c 1.22, $CHCl_3$). **Major diastereomer.** 1H NMR (400 MHz) δ 0.90 (d, $J = 7.2$ Hz, 3H, CH_3), 0.97 (m, 1H, cyclohexyl), 1.14 (m, 1H, cyclohexyl), 1.20 (s, 3H, CH_3), 1.28 (m, 1H, cyclohexyl), 1.29 (s, 3H, CH_3), 1.48 (m, 1H, cyclohexyl), 1.68 (m, 1H, cyclohexyl), 1.80 (m, 1H, cyclohexyl), 1.92 (m, 1H, cyclohexyl), 2.10 (m, 1H, cyclohexyl), 2.51 (m, 1H, $C_6F_{13}CHH$), 3.03 (m, 1H, $C_6F_{13}CHH$), 3.77 (dd, $J = 5.2, 8.0$ Hz, 1H, ICH), 4.80 (dt, $J = 4.4, 10.8$ Hz, 1H, OCH), 7.12–7.32 (m, 5H, arom); ^{13}C NMR (100 MHz) δ 5.3, 21.8, 23.6, 26.4, 28.7, 31.1, 34.6, 37.1 (t, $J_{C-F} = 19.8$ Hz, $C_6F_{13}C$), 39.4, 41.66, 50.5, 76.7, 125.09, 125.19, 127.82, 151.6, 169.2. **Minor diastereomer.** 1H NMR (400 MHz) δ 0.86 (d, $J = 7.2$ Hz, 3H, CH_3), 0.97 (m, 1H, cyclohexyl), 1.14 (m, 1H, cyclohexyl), 1.18 (s, 3H, CH_3), 1.28 (m, 1H, cyclohexyl), 1.27 (s, 3H, CH_3), 1.48 (m, 1H, cyclohexyl), 1.68 (m, 1H, cyclohexyl), 1.80 (m, 1H, cyclohexyl), 1.92 (m, 1H, cyclohexyl), 2.10 (m, 1H, cyclohexyl), 2.51 (m, 1H, $C_6F_{13}CHH$), 3.03 (m, 1H, $C_6F_{13}CHH$), 4.19 (dd, $J = 3.2, 10.0$ Hz, 1H, ICH), 4.77 (dt, $J = 4.4, 10.8$ Hz, 1H, OCH), 7.12–7.32 (m, 5H, arom); ^{13}C NMR (100 MHz) δ 7.4, 21.7, 25.5, 27.1, 28.4, 31.3, 34.5, 36.9 (t, $J_{C-F} = 19.8$ Hz, $C_6F_{13}C$), 39.5, 41.64, 50.3, 78.3, 125.12, 125.6, 127.79, 150.7, 169.4.

(4S)-3-(2-Iodo-3-perfluorohexylpropanoyl)-4-benzyl-2-oxazolidinone (11)

Less polar diastereomer. White powder; mp. 96 °C (from CHCl₃); [α]_D²³ -26.5 (*c* 0.68, CHCl₃); ¹H NMR (400 MHz) δ 2.77 (dd, *J* = 9.7, 13.2 Hz, 1H, OCHH), 2.78 (m, 1H, C₆F₁₃CHH), 3.32 (dd, *J* = 3.4, 13.2 Hz, 1H, OCHH), 3.59 (m, 1H, C₆F₁₃CHH), 4.23 (dd, *J* = 3.4, 7.6 Hz, 1H, PhCHH), 4.28 (t, *J* = 7.6 Hz, 1H, PhCHH), 4.75 (ddt, *J* = 7.6, 9.7, 3.4 Hz, 1H, NCH), 6.15 (dd, *J* = 2.9, 10.7 Hz, 1H, ICH), 7.25–7.36 (m, 5H, arom); ¹³C NMR (100 MHz) δ 5.8, 36.4, 37.3 (t, *J*_{C-F} = 21.5 Hz, C₆F₁₃C), 55.3, 66.3, 127.4, 129.0, 129.4, 134.5, 152.2, 169.3. IR (KBr, cm⁻¹) 2956, 1771, 1714, 1399, 1357, 1183, 1145, 1103, 744, 704, 513, 447; MS *m/z* 677 (M⁺, 15%), 550 (M⁺ – I, 100), 501 (34), 473 (89), 133 (62), 117 (57); HRMS calcd for C₁₉H₁₃F₁₃INO₃ (M⁺) 676.9732, found 676.9717. **More polar diastereomer.** Colorless oil; [α]_D²³ -25.7 (*c* 0.49, CHCl₃); ¹H NMR (400 MHz) δ 2.74 (dd, *J* = 9.6, 13.2 Hz, 1H, OCHH), 2.80 (m, 1H, C₆F₁₃CHH), 3.32 (dd, *J* = 3.2, 13.2 Hz, 1H, OCHH), 3.60 (m, 1H, C₆F₁₃CHH), 4.22 (dd, *J* = 3.2, 8.8 Hz, 1H, PhCHH), 4.28 (t, *J* = 8.8 Hz, 1H, PhCHH), 4.70 (m, 1H, NCH), 6.18 (dd, *J* = 3.2, 9.6 Hz, 1H, ICH), 7.21–7.38 (m, 5H, arom); ¹³C NMR (100 MHz) δ 5.0, 37.7, 37.8 (t, *J*_{C-F} = 20.6 Hz, C₆F₁₃C), 55.8, 66.3, 127.4, 129.0, 129.3, 134.7, 152.3, 169.5; IR (KBr, cm⁻¹) 3032, 1784, 1700, 1394, 1354, 1238, 1209, 1100, 707; MS *m/z* 677 (M⁺, 11%), 550 (M⁺ – I, 93), 501 (25), 473 (62), 133 (48), 91 (100); HRMS calcd for C₁₉H₁₃F₁₃INO₃ (M⁺) 676.9732, found 676.9688.

***N*-{(2*R*)-2-Iodo-3-perfluorohexylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (13-major isomer)**

Plate crystals; mp. 147 °C (from CH₂Cl₂-hexane 1:1); [α]_D²¹ -42.6 (*c* 0.97, CHCl₃); ¹H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.21 (s, 3H, CH₃), 1.40 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.73 (m, 1H, C₆F₁₃CHH), 3.46 (m, 1H, C₆F₁₃CHH), 3.48 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.54 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.99 (t, *J* = 6.8 Hz, 1H, NCH), 5.31 (m, 1H, ICH); ¹³C NMR (100 MHz) δ 6.9, 19.9, 20.6, 26.5, 32.7, 37.0, 37.4 (t, *J*_{C-F} = 20.7 Hz, C₆F₁₃C), 44.3, 47.9, 49.0, 52.8, 64.9, 168.2; IR (KBr, cm⁻¹) 2977, 1695, 1329, 1238, 1146, 1065, 776, 699, 534; MS *m/z* 715 (M⁺, 6%), 588 (M⁺ – I, 21), 524 (72), 473 (39), 134 (100); HRMS calcd for C₁₉H₁₉F₁₃INO₃S (M⁺) 714.9923, found 714.9927.

***N*-{(2*S*)-2-Iodo-3-perfluorohexylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (13-minor isomer)**

Plate crystals; mp. 203 °C (from CH₂Cl₂-hexane 1:1); [α]_D²¹ -37.6 (*c* 0.92, CHCl₃); ¹H NMR (400 MHz) δ 0.97 (s, 3H, CH₃), 1.09 (s, 3H, CH₃), 1.40 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.79 (m, 1H, C₆F₁₃CHH), 3.44 (m, 1H, C₆F₁₃CHH), 3.49 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.53 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.86 (dd, *J* = 5.4, 7.3 Hz, 1H, NCH), 5.20 (dd, *J* = 2.4, 12.7 Hz, 1H, ICH); ¹³C NMR (100 MHz) δ 6.6, 19.9, 20.1, 26.4, 32.9, 37.8, 39.8 (t, *J*_{C-F}

= 19.0 Hz, C₆F₁₃C), 44.5, 47.9, 48.9, 52.5, 65.8, 167.5; IR (KBr, cm⁻¹) 2999, 1679, 1334, 1241, 1147, 1065, 846, 772, 710, 528; MS *m/z* 715 (M⁺, 14%), 651 (6), 588 (M⁺ – I, 23), 524 (86), 473 (43), 404 (25), 134 (100); HRMS calcd for C₁₉H₁₉F₁₃INO₃S (M⁺) 714.9923, found 714.9890.

***N*-{(2*R*)-2-Iodo-3-perfluorohexylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (14-major isomer)**

Plate crystals; mp. 125 °C (from CH₂Cl₂-hexane 1:1); [α]_D²¹ -55.3 (c 1.39, CHCl₃); ¹H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.21 (s, 3H, CH₃), 1.40 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.71 (m, 1H, C₆F₁₃CHH), 3.46 (m, 1H, C₆F₁₃CHH), 3.48 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.54 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.99 (t, *J* = 6.0 Hz, 1H, NCH), 5.31 (m, 1H, ICH); ¹³C NMR (100 MHz) δ 6.8, 19.9, 20.6, 26.5, 32.7, 37.0, 37.1 (t, *J*_{C-F} = 11.9 Hz, C₆F₁₃C), 44.3, 47.9, 48.9, 52.8, 64.9, 168.2; IR (KBr, cm⁻¹) 2980, 1693, 1294, 1225, 1137, 1118, 1060, 960, 758, 721, 535; MS *m/z* 565 (M⁺, 8%), 438 (M⁺ – I, 22), 374 (69), 323 (56), 134 (100); HRMS calcd for C₁₆H₁₉F₇INO₃S (M⁺) 565.0019, found 565.0035.

***N*-{(2*S*)-2-Iodo-3-perfluoropropylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (14-minor isomer)**

Plate crystals; mp. 180 °C (from CH₂Cl₂-hexane 1:1); [α]_D²⁵ -42.9 (c 0.06, CHCl₃); ¹H NMR (400 MHz) δ 0.97 (s, 3H, CH₃), 1.09 (s, 3H, CH₃), 1.40 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.52 (m, 1H, C₆F₁₃CHH), 2.78 (m, 1H, C₆F₁₃CHH), 3.50 (d, *J* = 13.4 Hz, 1H, SO₂CHH), 3.54 (d, *J* = 13.4 Hz, 1H, SO₂CHH), 3.87 (t, *J* = 8.4 Hz, 1H, NCH), 5.21 (m, 1H, ICH); ¹³C NMR (100 MHz) δ 6.6, 19.9, 20.8, 26.4, 32.9, 37.8, 38.3 (t, *J*_{C-F} = 13.2 Hz, C₆F₁₃C), 44.5, 47.8, 48.8, 52.5, 65.8, 167.4; IR (KBr, cm⁻¹) 3002, 1681, 1335, 1229, 1069, 1085, 1067, 962, 772, 532; MS *m/z* 565 (M⁺, 19%), 438 (M⁺ – I, 23), 374 (82), 323 (59), 254 (25), 134 (100); HRMS calcd for C₁₆H₁₉F₇INO₃S (M⁺) 565.0019, found 565.0037.

***N*-{(2*R*)-2-Iodo-3-perfluoroisopropylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (15-major isomer)**

Plate crystals; mp. 147 °C (from CH₂Cl₂-hexane 1:1); [α]_D²¹ -64.6 (c 1.12, CHCl₃); ¹H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.20 (s, 3H, CH₃), 1.40 (m, 2H, CH₂), 1.94 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.71 (dd, *J* = 16.0, 22.4 Hz, 1H, C₆F₁₃CHH), 3.47 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.53 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.57 (m, 1H, C₆F₁₃CHH), 3.99 (t, *J* = 6.4 Hz, 1H, NCH), 5.30 (m, 1H, ICH); ¹³C NMR (100 MHz) δ 6.7, 19.9, 20.6, 26.5, 32.7, 34.4 (t, *J*_{C-F} = 10.0 Hz, C₆F₁₃C), 36.9, 44.3, 47.9, 48.9, 52.8, 65.0, 168.2; IR (KBr, cm⁻¹) 2985, 1695, 1329, 1302, 1232, 1172, 1155,

1136, 1049, 775, 719, 544; MS m/z 565 (M^+ , 7%), 438 ($M^+ - I$, 18), 374 (57), 323 (42), 134 (100); HRMS calcd for $C_{16}H_{19}F_7INO_3S$ (M^+) 565.0019, found 565.0026.

***N*-{(2*S*)-2-Iodo-3-perfluoroisopropylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (15-minor isomer)**

White powder; mp. 128 °C (from $CHCl_3$); $[\alpha]_D^{21}$ -51.6 (c 0.43, $CHCl_3$); 1H NMR (400 MHz) δ 0.97 (s, 3H, CH_3), 1.08 (s, 3H, CH_3), 1.41 (m, 2H, CH_2), 1.92 (m, 3H, CH_2 , CH), 2.09 (m, 2H, CH_2), 2.74 (dd, J = 15.1, 28.8 Hz, 1H, $C_6F_{13}CHH$), 3.47 (d, J = 14.2 Hz, 1H, SO_2CHH), 3.50 (d, J = 14.2 Hz, 1H, SO_2CHH), 3.61 (m, 1H, $C_6F_{13}CHH$), 3.84 (dd, J = 5.6, 7.8 Hz, 1H, NCH), 5.20 (m, 1H, ICH); ^{13}C NMR (100 MHz) δ 8.3, 18.2, 20.0, 26.5, 32.9, 36.7 (d, J_{C-F} = 18.1 Hz, $C_6F_{13}C$), 37.8, 44.4, 47.9, 48.8, 52.5, 65.8, 167.4; IR (KBr, cm^{-1}) 2967, 1681, 1337, 1227, 1137, 989, 771, 531; MS m/z 565 (M^+ , 6%), 438 ($M^+ - I$, 11), 374 (30), 323 (20), 119 (100); HRMS calcd for $C_{16}H_{19}F_7INO_3S$ (M^+) 565.0019, found 565.0027.

***N*-{(2*R*)-2-Iodo-3-perfluoroethylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (16-major isomer)**

Needle crystals; mp. 153 °C (from CH_2Cl_2); $[\alpha]_D^{21}$ -57.9 (c 0.94, $CHCl_3$); 1H NMR (400 MHz) δ 0.99 (s, 3H, CH_3), 1.21 (s, 3H, CH_3), 1.44 (m, 2H, CH_2), 1.92 (m, 3H, CH_2 , CH), 2.05 (m, 2H, CH_2), 2.69 (m, 1H, $C_6F_{13}CHH$), 3.44 (m, 1H, $C_6F_{13}CHH$), 3.47 (d, J = 13.6 Hz, 1H, SO_2CHH), 3.54 (d, J = 13.6 Hz, 1H, SO_2CHH), 3.99 (m, 1H, NCH), 5.29 (m, 1H, ICH); ^{13}C NMR (100 MHz) δ 6.8, 19.9, 20.6, 26.5, 32.7, 37.1, 37.2 (t, J_{C-F} = 16.9 Hz, $C_6F_{13}C$), 44.3, 47.9, 49.0, 52.8, 64.9, 168.2; IR (KBr, cm^{-1}) 2968, 1687, 1407, 1336, 1219, 1192, 1138, 1113, 1066, 769, 531; MS m/z 515 (M^+ , 14%), 388 ($M^+ - I$, 29), 324 (93), 273 (66), 241 (48), 173 (33), 134 (100); HRMS calcd for $C_{15}H_{19}F_5INO_3S$ (M^+) 515.0050, found 515.0079.

***N*-{(2*S*)-2-Iodo-3-perfluoroethylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (16-minor isomer)**

1H NMR (400 MHz) δ 0.97 (s, 3H, CH_3), 1.10 (s, 3H, CH_3), 1.40 (m, 2H, CH_2), 1.90 (m, 3H, CH_2 , CH), 2.10 (m, 2H, CH_2), 2.74 (m, 1H, $C_6F_{13}CHH$), 3.40 (m, 1H, $C_6F_{13}CHH$), 3.49 (d, J = 13.6 Hz, 1H, SO_2CHH), 3.53 (d, J = 13.6 Hz, 1H, SO_2CHH), 3.86 (dd, J = 5.2, 7.6 Hz, 1H, NCH), 5.19 (dd, J = 2.4, 11.6 Hz, 1H, ICH); ^{13}C NMR (100 MHz) δ 6.5, 20.0, 20.3, 26.4, 33.0, 37.8, 39.5 (t, J_{C-F} = 19.8 Hz, $C_6F_{13}C$), 44.5, 47.8, 48.9, 52.5, 65.8, 168.2; MS m/z 515 (M^+ , 8%), 388 ($M^+ - I$, 14), 324 (43), 273 (30), 173 (41), 134 (100); HRMS calcd for $C_{15}H_{19}F_5INO_3S$ (M^+) 515.0050, found 515.0068.

The minor isomer of **16** contained a small portion of *N*-(3-perfluoroethylpropanoyl)-(1*S*,2*R*,4*R*)-bornane-10,2-sultam as an inseparable side product. To identify the structure of the side product, **16** (major and minor mixture) was reduced by radical method.

***N*-(3-perfluoroethylpropanoyl)-(1*S*,2*R*,4*R*)-bornane-10,2-sultam:** To a solution of **16** (major/minor = 80:20 mixture, 35 mg, 0.7 mmol) in dry toluene (1.0 mL) were added *n*-Bu₃SnH (38 μ L, 0.14 mmol) and Et₃B (1.06 mol/L in hexane; 70 μ L, 0.07 mmol) at 0 °C. The mixture was stirred at 0 °C for 6 h. KF and water were added and the reaction mixture was stirred at room temperature for 10 h. After filtration, the solvent was evaporated under reduced pressure. The residue was chromatographed on silica gel to afford pure product (10.1 mg, 38%).

¹H NMR (400 MHz) δ 0.98 (s, 3H, CH₃), 1.15 (s, 3H, CH₃), 1.38 (m, 2H, CH₂), 1.90 (m, 3H, CH₂, CH), 2.11 (m, 2H, CH₂), 2.47 (m, 2H, C₆F₁₃CH₂), 3.04 (t, *J* = 7.6 Hz, 2H), 3.46 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.53 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.87 (dd, 1H, *J* = 5.2, 7.2 Hz NCH); ¹³C NMR (100 MHz) δ 19.9, 20.8, 25.4 (t, *J*_{C-F} = 19.0 Hz, C₆F₁₃C), 26.5, 26.8, 32.9, 38.3, 44.6, 47.9, 48.7, 52.9, 65.3, 168.8.

Procedure for iodotrifluoromethylation.

In a pressure-resistant Pyrex glass tube were placed olefin (0.2 mmol) and CH₂Cl₂ (5 mL). Then the mixture was cooled to -30°C, and trifluoromethyl iodide (2.0 mmol), Na₂S₂O₃ (1 mmol, 158 mg) and water (1 mL) were added to the mixture. After sealing the tube, the mixture was mixed by shaking and then irradiated with a Hg lamp at room temperature. After the reaction was complete, the mixture was extracted with CH₂Cl₂. The extract was washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to afford pure product.

N-{(2*R*)-2-Iodo-3-trifluoromethylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (17-major isomer)

Needle crystals; mp. 183 °C (from CH₂Cl₂); [α]_D²³ -72.1 (*c* 0.28, CHCl₃); ¹H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.21 (s, 3H, CH₃), 1.37 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.78 (m, 1H, C₆F₁₃CHH), 3.35 (m, 1H, C₆F₁₃CHH), 3.47 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.53 (d, *J* = 13.6 Hz, 1H, SO₂CHH), 3.98 (m, 1H, NCH), 5.21 (m, 1H, ICH); ¹³C NMR (100 MHz) δ 7.8, 19.9, 20.6, 26.5, 32.7, 37.0, 39.9 (q, *J*_{C-F} = 29.7 Hz, CF₃C), 44.4, 47.9, 48.9, 52.9, 64.9, 124.8 (q, *J*_{C-F} = 277.2 Hz, CF₃), 168.2; IR (KBr, cm⁻¹) 2945, 1689, 1395, 1354, 1329, 1293, 1259, 1156, 1085, 1061, 769; MS *m/z* 465 (M⁺, 29%), 401 (9), 338 (M⁺ - I, 17), 274 (53), 223 (37), 134 (100); HRMS calcd for C₁₄H₁₉F₃INO₃S (M⁺) 465.0096, found 465.0082.

***N*-{(2*R*)-2-Iodo-3-trifluoromethylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (17-minor isomer)**

The minor isomer of **17** was inseparable with starting material **12**. ^1H NMR (400 MHz) δ 0.97 (s, 3H, CH₃), 1.10 (s, 3H, CH₃), 1.37 (m, 2H, CH₂), 1.92 (m, 3H, CH₂, CH), 2.08 (m, 2H, CH₂), 2.75 (m, 1H, C₆F₁₃CHH), 3.36 (m, 1H, C₆F₁₃CHH), 3.47 (d, J = 13.6 Hz, 1H, SO₂CHH), 3.53 (d, J = 13.6 Hz, 1H, SO₂CHH), 3.85 (dd, 1H, J = 5.2, 7.6 Hz, NCH), 5.10 (dd, 1H, J = 2.8, 11.6 Hz, ICH); ^{13}C NMR (100 MHz) δ 7.3, 20.0, 20.6, 26.5, 33.0, 37.9, 40.1 (q, $J_{\text{C-F}}$ = 29.7 Hz, CF₃C), 44.5, 47.9, 48.8, 52.6, 65.8, 124.8 (q, $J_{\text{C-F}}$ = 277.2 Hz, CF₃), 167.4.

Azide displacement of **14 (major): *N*-{(2*S*)-2-Azido-3-perfluoropropylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (**18**)**

To a solution of **14** (95 mg, 0.17 mmol) in CH₂Cl₂ (1.0 mL) were added NaN₃ (110 mg, 1.7 mmol) and DMF (1.0 mL). After stirring for 1 h, the reaction mixture was concentrated under reduced pressure. The residue was chromatographed on silica gel to afford pure product **18** (65.4 mg, 82%).

Needle crystals; mp. 87 °C (from CH₂Cl₂); $[\alpha]_{\text{D}}^{21}$ -43.9 (c 1.17, CHCl₃); ^1H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.10 (s, 3H, CH₃), 1.42 (m, 2H, CH₂), 1.93 (m, 3H, CH₂, CH), 2.09 (m, 2H, CH₂), 2.56 (m, 1H, C₆F₁₃CHH), 2.84 (m, 1H, C₆F₁₃CHH), 3.53 (d, J = 14.1 Hz, 1H, SO₂CHH), 3.57 (d, J = 14.1 Hz, 1H, SO₂CHH), 3.99 (t, J = 6.8 Hz, 1H, NCH), 4.80 (t, J = 6.4 Hz, 1H, N₃CH); ^{13}C NMR (100 MHz) δ 19.9, 20.4, 26.4, 32.80, 32.83 (t, $J_{\text{C-F}}$ = 19.8 Hz, C₆F₁₃C), 37.7, 44.5, 47.9, 48.9, 52.9, 54.4, 65.3, 166.8; IR (neat, cm⁻¹) 2966, 2117, 1700, 1340, 1239, 1139, 1117; MS m/z 374 ($\text{M}^+ - \text{N}_2$, SO₂, 6%), 242 (68), 210 (40), 135 (100), 93 (72); HRMS calcd for C₁₆H₁₉F₇NO ($\text{M}^+ - \text{N}_2$, SO₂) 374.1338, found 374.1355.

Azide displacement of **17 (major): *N*-{(2*S*)-2-Azido-3-perfluoroethylpropanoyl}-(1*S*,2*R*,4*R*)-bornane-10,2-sultam (**19**)**

Compound **19** was prepared following the procedure described for the preparation of **18** except for the use of **16** instead of **14** (88%).

Needle crystals; mp. 98 °C (from CH₂Cl₂); $[\alpha]_{\text{D}}^{25}$ -38.2 (c 0.57, CHCl₃); ^1H NMR (400 MHz) δ 0.99 (s, 3H, CH₃), 1.11 (s, 3H, CH₃), 1.42 (m, 2H, CH₂), 1.96 (m, 3H, CH₂, CH), 2.10 (m, 2H, CH₂), 2.53 (m, 1H, C₆F₁₃CHH), 2.79 (m, 1H, C₆F₁₃CHH), 3.52 (d, J = 14.1 Hz, 1H, SO₂CHH), 3.56 (d, J = 14.1 Hz, 1H, SO₂CHH), 3.99 (t, J = 6.4 Hz, 1H, NCH), 4.78 (t, J = 6.8 Hz, 1H, N₃CH); ^{13}C NMR (100 MHz) δ 19.9, 20.5, 26.4, 32.89 (t, $J_{\text{C-F}}$ = 18.2 Hz, C₆F₁₃C), 32.92, 37.7, 44.5, 48.0, 48.9, 52.9, 54.4, 65.3, 166.8; IR (neat, cm⁻¹) 2997, 2110, 1702, 1684, 1396, 1333, 1199, 1141; MS m/z 324 ($\text{M}^+ - \text{N}_2$, SO₂, 8%), 242 (46), 160 (30), 135 (100), 93 (90); HRMS calcd for C₁₆H₁₉F₇NO ($\text{M}^+ - \text{N}_2$,

SO₂) 324.1337, found 324.1379.

Hydrolysis and hydrogenation of **18**: (*S*)-4, 4, 5, 5, 6, 6, 6-Heptafluoronorleucine (**20**)

To a solution of **18** (90 mg, 0.18 mmol) in THF (1.0 mL) were added LiOH-H₂O (15 mg, 0.36 mmol) and water (1.0 mL). After stirring for 1 h, the reaction mixture was evaporated to remove THF and the residue was extracted with CH₂Cl₂. The organic layer contained (1*S*)-(-)-2,10-bornane sultam which was recovered after evaporation. The aqueous layer was acidified to pH 1 by the addition of 10% HCl and then extracted with CH₂Cl₂. The organic layer was concentrated under reduced pressure to give a crude product.

To a solution of the crude carboxylic acid in water-acetic acid (3:1, 1.5 mL) was added 10% Pd on carbon (8 mg, 5 mol%). The mixture was stirred under H₂ atmosphere for 20 h. The reaction mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give **20** as a white solid, which was recrystallized from water to give white powder.

White powder; mp. 230 °C (decomp.); [α]_D²¹ +2.7 (*c* 0.15, 1N HCl); ¹H NMR (400 MHz, 5% NaOD in D₂O) δ 1.84 (m, 1H, C₃F₇CHH), 2.07 (m, 1H, C₃F₇CHH), 3.10 (t, *J* = 6.8 Hz, 1H, CH); ¹³C NMR (100 MHz, 5% NaOD in D₂O) δ 34.9 (t, *J*_{C-F} = 21.4 Hz, C₃F₇C), 49.9, 180.2; IR (KBr, cm⁻¹) 3034, 1665, 1588, 1405, 1360, 1179, 1115, 929, 687; MS *m/z* 212 (*M*⁺ – CO₂H, 100%), 192 (33), 74 (22); HRMS calcd for C₅H₅F₇N (*M*⁺ – CO₂H) 212.0310, found 212.0313.

Hydrolysis and hydrogenation of **19**: (*S*)-4,4,5,5,5-Pentafluoronorvaline (**21**)

Compound **21** was prepared following the procedure described for **20** except for the use of **19** instead of **18** (63%).

White powder; mp. 230 °C (decomp.); [α]_D²¹ +2.5 (*c* 0.16, 1N HCl); ¹H NMR (400 MHz, 5% NaOD in D₂O) δ 1.92 (m, 1H, C₂F₅CHH), 2.19 (m, 1H, C₂F₅CHH), 3.22 (t, *J* = 6.0 Hz, 1H, CH); ¹³C NMR (100 MHz, 5% NaOD in D₂O) δ 35.0 (t, *J*_{C-F} = 18.2 Hz, C₂F₅C), 50.0, 180.3; IR (KBr, cm⁻¹) 3012, 1671, 1590, 1410, 1341, 1188, 1096, 1022, 980, 681; MS *m/z* 162 (*M*⁺ – CO₂H, 100%), 142 (13); HRMS calcd for C₄H₅F₅N (*M*⁺ – CO₂H) 162.0342, found 162.0339.

11 (More polar)

C:\IMMR data\yajima\899\820c.als
can-c6

C:\1H-NMR\y4j\lmg\1130.d1s
c3-amino 5% NaODaq.