Platinum(II) Catalyzed 1,6-Diene Cycloisomerizations:
Turnover in the Absence of β -Hydride Elimination
Supporting Information

William D. Kerber, Jeong Hwan Koh, and Michel R. Gagné

Department of Chemistry, University of North Carolina at Chapel Hill,

Chapel Hill, North Carolina, 27599-3290

Table of Contents

1. Experimental Procedures
 - 1.1 Synthesis of Dienes
 - 1.2 Catalytic Cycloisomerizations
 - 1.3 Deuterium Labeling Experiments
2. 2D NMR Experiments
 - 2.1 Relative Stereochemistry of **3a**
 - 2.2 Structural Determination of **4a**

1. Experimental Procedures

General: Catalytic cycloisomerization experiments were performed in a dinitrogen-filled MBraun Labmaster 100 glovebox. Synthetic procedures were performed under dinitrogen using standard Schlenk techniques. CH_2Cl_2 and THF were sparged with dry argon and passed through a column of activated alumina. For catalytic experiments,

CH₂Cl₂ was further freeze-pump-thaw degassed. DMSO and NEt₃ were distilled from CaH₂. Hept-1-en-6-yne,¹ benzyl bromides,² bis(η⁵-cyclopentadienyl)-di(methyl-d₃) titanium,³ [(PNP)Pt(ethylene)](BF₄)₂,⁴ (triphos)PtI₂,⁵ and Ag⁺-impregnated silica⁶ were prepared according to literature procedures. All other chemicals were used as received. NMR spectra were recorded on a Bruker Avance 400 MHz or Varian Inova 600 MHz spectrometer; chemical shifts are given in ppm and are referenced to residual solvent resonances. GC was performed on a HP 6890 with an HP-5 column. GC/MS was performed on an Agilent 5973 with an HP-5 column. High-resolution mass spectrometry was performed by the Mass Spectrometry Service Laboratory at the University of Minnesota.

1.1 Synthesis of Dienes 2a-2d

***Trans*-1-Methoxy-2-(3-methyl-octa-2,7-dienyl)-benzene⁷ (2a) Representative Procedure:** To a solution of 1.11 g AlMe₃ (15.4 mmol) in 12.0 mL CH₂Cl₂ was added 1.10 g Cp₂ZrCl₂ (3.7 mmol) and the resulting lemon-yellow solution was stirred for 10 min at RT. A solution of 0.74 g hept-1-en-6-yne (7.5 mmol) in 7.0 mL CH₂Cl₂ was then added and stirred at RT for 24 h. The reaction mixture was then cannula transferred to a solution of 1.35 g 2-methoxybenzyl bromide (6.7 mmol) and 400 mg Pd(PPh₃)₄ (0.35 mmol) in 40 mL THF at 0°C, raised to RT and stirred for 24 h. The reaction was quenched at 0°C by slow addition of 40 mL 3N HCl and the aqueous phase was extracted with ether. The combined organic extracts were washed with sat. NaHCO₃, sat. NaCl, dried over MgSO₄ and concentrated under vacuum. The crude material was purified by flash chromatography in 99:1 hexanes:EtOAc. Yielded 1.22 g of a colorless oil (78%); ¹H NMR: (400 MHz, CDCl₃) δ 7.17 (m, 2H), 6.86 (m, 2H), 5.81 (m, 1H), 5.31 (td, 1H, J = 7.2, 1.2 Hz), 4.95 (m, 2H), 3.82 (s, 3H), 3.32 (d, 2H, J = 7.2 Hz), 2.02 (m, 4H), 1.68 (s, 3H), 1.51 (m, 2H); ¹³C NMR: (100 MHz, CDCl₃) δ 157.3, 139.0, 135.9, 130.0, 129.2,

126.8, 122.6, 120.4, 114.3, 110.2, 55.3, 39.2, 33.4, 28.2, 27.2, 15.9; HR ESI m/z = 253.1552 (253.1563 calculated for $C_{16}H_{22}ONa$).

***Trans*-1-Methoxy-3-(3-methyl-octa-2,7-dienyl)-benzene (2b):** A procedure similar to **2a** was followed. The crude material purified by flash chromatography in 49:1 hexanes:EtOAc. Yielded 0.99 g of a colorless oil (69%); 1H NMR: (400 MHz, $CDCl_3$) δ 7.11 (t, 2H, J = 7.6 Hz), 6.80 (m, 3H), 5.86 (m, 1H), 5.39 (t, 1H, J = 7.2 Hz), 5.01 (m, 2H), 3.82 (s, 3H), 3.38 (d, 2H, J = 6.8 Hz), 2.08 (m, 4H), 1.75 (s, 3H), 1.58 (m, 2H); ^{13}C NMR: (100 MHz, $CDCl_3$) δ 159.6, 143.4, 138.8, 136.2, 129.2, 122.9, 120.7, 114.4, 114.0, 110.1, 55.0, 39.0, 34.2, 33.3, 27.1, 16.0.; LR EI m/z = 230.2 (calculated 230.17 for $C_{16}H_{22}O$)

***Trans*-1-Methoxy-4-(3-methyl-octa-2,7-dienyl)-benzene (2c):** A procedure similar to **2a** was followed. The crude material purified by flash chromatography in 49:1 hexanes:EtOAc. Yielded 0.76 g of a colorless oil (67%); 1H NMR: (400 MHz, $CDCl_3$) δ 7.08 (d, 2H J = 8.8 Hz), 6.81 (d, 2H, J = 8.8 Hz), 5.80 (m, 1H), 5.31 (td, 1H, J = 7.2, 1.2 Hz), 4.96 (m, 2H), 3.77 (s, 3H), 3.28 (d, 2H, J = 7.2 Hz), 2.02 (m, 4H), 1.68 (s, 3H), 1.51 (m, 2H); ^{13}C NMR: (100 MHz, $CDCl_3$) δ 157.7, 138.9, 135.8, 133.8, 129.1, 123.6, 114.4, 113.7, 55.3, 39.1, 33.4, 33.3, 27.2, 16.0; HR ESI m/z = 253.1564 (253.1563 calculated for $C_{16}H_{22}ONa$).

***Trans*-(3-Methyl-octa-2,7-dienyl)-benzene (2d):** A procedure similar to **2a** was followed. The crude material purified by flash chromatography in 199:1 hexanes:EtOAc. Yielded 0.87 g of a colorless oil (43%); 1H NMR: (400 MHz, $CDCl_3$) δ 7.26 (m, 2H), 7.17 (m, 3H), 5.81 (m, 1H), 5.24 (td, 1H, J = 7.2, 1.2 Hz), 4.97 (m, 2H), 3.35 (d, 2H, J = 7.2 Hz), 2.03 (m, 4H), 1.70 (s, 3H), 1.51 (m, 2H); ^{13}C NMR: (100 MHz, $CDCl_3$) δ 141.8, 138.9, 136.1, 128.3, 128.3, 125.7, 123.1, 114.4, 39.1, 34.2, 33.7, 27.2, 16.1; HR EI m/z = 200.1566 (calculated 200.1565 for $C_{15}H_{20}O$)

1.2 Catalytic Cycloisomerizations

***Trans*-1-(2-Methoxy-benzyl)-2-methyl-bicyclo[4.1.0]heptane (3a) Representative Procedure:** To a solution of 9.8 mg (triphos)PtI₂ (10 μ mol) in 2.0 mL CH_2Cl_2 was added 1.5 μ L acetone (20 μ mol) and 6.8 mg AgSbF₆ (20 μ mol) and stirred in the dark for 30 min. The heterogeneous mixture was sub-micron filtered onto 23 mg *trans*-1-Methoxy-

2-(3-methyl-octa-2,7-dienyl)-benzene **2a** (100 μ mol) and 30 μ L N-methyl-diphenylamine (1 μ mol; 0.0260 M in CH_2Cl_2). The reaction was stirred at 22°C until all starting material had been consumed (GC, 16 h). The reaction was filtered through a plug of silica, CH_2Cl_2 eluent, and concentrated under reduced pressure. The crude material was purified by flash chromatography on Ag^+ -impregnated silica in 400:1 hexanes:EtOAc to yield 19 mg (83%) of a colorless oil. >95:5 d.r. by ^1H NMR; ^1H NMR: (600 MHz, CDCl_3) δ 7.24 (d, 1H, J = 7.2 Hz), 7.16 (m, 1H), 6.89 (m, 1H), 6.83 (d, 1H, J = 7.8 Hz), 3.79 (s, 3H), 3.18 (d, 1H, J = 14.4 Hz), 2.08 (d, 1H, J = 14.4 Hz), 1.97 (m, 1H), 1.67 (m, 1H), 1.33 (m, 2H), 1.27 (m, J = 1H), 0.99 (m, 2H), 0.94 (d, 3H, J = 6.6 Hz), 0.64 (m, 1H), 0.27 (dd, 1H, J = 9.0, 4.2 Hz), 0.10 (t, 1H, J = 4.8 Hz); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.9, 130.4, 128.9, 126.8, 120.1, 110.1, 55.2, 37.3, 30.0, 29.9, 24.7, 24.5, 22.0, 19.4, 18.5, 13.8; HR ESI m/z = 253.1563 (253.1563 calculated for $\text{C}_{16}\text{H}_{22}\text{ONa}$)

Trans-1-(4-Methoxy-benzyl)-2-methyl-bicyclo[4.1.0]heptane (3c): A procedure similar to **3a** was followed. The crude material flashed on Ag^+ -impregnated silica⁶ in 200:1 hexanes:EtOAc. Yielded 15 mg (65%) of a colorless oil. >95:5 d.r. by ^1H NMR; ^1H NMR: (400 MHz, CDCl_3) δ 7.11 (d, 2H, J = 8.4 Hz), 6.80 (d, 2H, J = 8.8 Hz), 3.78 (s, 3H), 2.94 (d, 1H, J = 14.4 Hz), 2.07 (d, 1H, J = 14.4 Hz), 1.99 (m, 1H), 1.62 (m, 1H), 1.38 (m, 3H), 0.99 (m, 2H), 0.89 (d, 3H, J = 6.4 Hz), 0.61 (m, 1H), 0.28 (dd, 1H, J = 4.6, 9.0 Hz), 0.10 (t, 1H, J = 5.0 Hz); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.8, 132.7, 130.1, 113.4, 55.2, 43.4, 29.7, 29.4, 25.7, 24.5, 22.0, 19.1, 18.6, 13.6; LR EI m/z = 230.2 (230.17 calculated for $\text{C}_{16}\text{H}_{22}\text{O}$)

[2-(1-Methyl-cyclohexenyl)-ethyl]-benzene (5d) Representative Procedure: To a solution of 50.2 mg (3-Methyl-octa-2,7-dienyl)-benzene **2d** (0.251 mmol) and 21.9 mg Pt(II) dication **1b** (0.025 mmol, 10 mol%) in 5.0 mL CH_2Cl_2 was added 0.055 mL (0.013 mmol, 0.5 mol%) of a Ph_2NMe solution (0.0225 M in CH_2Cl_2) and stirred at 22°C until all starting material had been consumed (GC, 12 hours). The solution was filtered through silica plug, eluting with CH_2Cl_2 and concentrated under reduced pressure. The crude material flashed on Ag^+ -impregnated silica⁶, 200:1 hexanes/EtOAc eluent. Yielded 34 mg (68%) of a colorless oil, obtained as a 2.1:1.0 mixture of cyclohexene regioisomers (GC, major is cyclohex-3-ene); ^1H NMR: (400 MHz, CDCl_3) δ 7.28 (m, 2H major, 2H minor), 7.14 (m, 3H major, 3H minor), 5.62 (m, 2H major, 1H minor), 5.47

(m, 1H minor), 2.57 (m, 2H major, 2H minor), 2.05 (m, 2H major, 2H minor), 1.81-1.96 (m, 2H major, 2H minor), 1.40-1.66 (m, 4H major, 4H minor), 1.02 (s, 3H minor), 0.97 (s, 3H major); ^{13}C NMR: (100 MHz, CDCl_3) δ 143.6 (major), 143.5 (minor), 136.4 (minor), 128.3 (2C major, 2C minor), 128.3 (2C major, 2C minor), 126.1 (major), 125.7 (major), 125.5 (major, 2C minor), 45.1, 43.8, 37.7, 34.7, 34.4, 33.3, 31.4, 30.7, 30.2, 27.3, 25.2, 24.5, 22.7, 19.3; HR EI m/z = 200.1577 (Calculated 200.1565 for $\text{C}_{15}\text{H}_{20}$).

1-Methoxy-2-[2-(1-methyl-cyclohex-3-enyl)-ethyl]-benzene (5a): A procedure similar to **5d** was followed. The crude product purified by flash chromatography on Ag^+ -impregnated silica, 200:1 hexanes/EtOAc eluent, to yield 23 mg (68%) of a colorless oil, as a 2.1:1.0 mixture of cyclohexene regioisomers (GC, major is cyclohex-3-ene); ^1H NMR: (600 MHz, CDCl_3) δ 7.13 (m, 2H major, 2H minor), 6.80 (m, 2H major, 2H minor), 5.61 (m, 2H major, 1H minor), 5.48 (m, 1H minor), 3.80 (s, 3H major, 3H minor), 2.57 (m, 2H major, 2H minor), 2.04 (m, 2H major, 2H minor), 1.78-1.94 (m, 2H major, 2H minor), 1.39-1.66 (m, 4H major, 4H minor), 1.02 (s, 3H minor), 0.97 (s, 3H major); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.4 (major, minor), 136.8 (minor), 131.9 (major), 131.8 (minor), 129.5 (major, minor), 126.7 (major, minor), 126.1 (major), 125.8 (major), 125.3 (minor), 120.4 (major, minor), 110.2 (major, minor), 55.3 (major, minor), 42.9, 41.7, 37.8, 34.5, 34.4, 32.2, 31.4, 27.2, 25.3, 24.9, 24.5, 24.4, 22.7, 19.3; LR EI m/z = 230.2 (calculated 230.17 for $\text{C}_{16}\text{H}_{22}\text{O}$).

1-Methoxy-3-[2-(1-methyl-cyclohex-3-enyl)-ethyl]-benzene (5b): A procedure similar to **5d** was followed. The crude product purified by flash chromatography on Ag^+ -impregnated silica, 100:1 hexanes/EtOAc eluent, to yield 10 mg (70%) of a colorless oil as a 2.1:1.0 mixture of cyclohexene regioisomers (GC, major is cyclohex-3-ene); ^1H NMR: (400 MHz, CDCl_3) δ 7.18 (m, 1H major, 1H minor), 6.73 (m, 3H major, 3H minor), 5.63 (m, 2H major, 1H minor), 5.44 (m, 1H minor), 3.78 (s, 3H major, 3H minor), 2.55 (m, 2H major, 2H minor), 2.02 (m, 2H major, 2H minor), 1.76-1.97 (m, 2H major, 2H minor), 1.38-1.64 (m, 4H major, 4H minor), 0.99 (s, 3H minor), 0.95 (s, 3H major); ^{13}C NMR: (100 MHz, CDCl_3) δ 159.6 (major, minor), 145.2 (major, minor), 136.4 (minor), 129.2 (major, minor), 126.1 (major), 125.7 (major, minor), 120.8 (major, minor), 114.2 (major), 114.1 (minor), 110.8 (minor), 110.7 (minor), 55.1 (major, minor),

44.9, 43.7, 37.6, 34.7, 34.4, 33.3, 31.4, 30.8, 30.3, 27.4, 25.2, 24.5, 22.7, 19.3; LR EI m/z = 230.2 (calculated 230.17 for $C_{16}H_{22}O$).

1-Methoxy-4-[2-(1-methyl-cyclohex-3-enyl)-ethyl]-benzene (5c): A procedure similar to **5d** was followed. The crude product purified by flash chromatography on Ag^+ -impregnated silica, 100:1 hexanes/EtOAc eluent, to yield 9 mg (64%) of a colorless oil, as a 2.1:1.0 mixture of cyclohexene regioisomers (GC, major is cyclohex-3-ene); 1H NMR: (400 MHz, $CDCl_3$) δ 7.08 (d, 2H major, 2H minor, J = 8.4 Hz), 6.80 (d, 2H major, 2H minor, J = 8.4 Hz), 5.61 (m, 2H major, 1H minor), 5.42 (m, 1H minor), 3.77 (s, 3H major, 3H minor), 2.51 (m, 2H major, 2H minor), 2.02 (m, 2H major, 2H minor), 1.74-1.94 (m, 2H major, 2H minor), 1.39-1.62 (m, 4H major, 4H minor), 1.00 (s, 3H minor), 0.95 (s, 3H major); ^{13}C NMR: (100 MHz, $CDCl_3$) δ 157.6 (major, minor), 136.5 (minor), 135.6 (major, minor), 129.1 (2C major, 2C minor), 126.1 (major), 125.7 (major), 125.6 (minor), 113.7 (2C major, 2C minor), 55.3 (major, minor), 45.3, 44.1, 37.7, 34.8, 34.4, 33.3, 31.3, 29.7, 29.2, 27.4, 25.2, 24.5, 22.7, 19.3; LR EI m/z = 230.2 (calculated 230.17 for $C_{16}H_{22}O$).

Figure 1 Synthesis of Labeled Substrates

1.3 Deuterium Labeling Experiments

The synthesis of deuterium labeled derivatives of **2a** is outlined in Fig.1. The results of cycloisomerization of these substrates by **1a** is shown in Fig. 2.

7-(2-Methoxy-phenyl)-5-methyl-hept-5-en-1-ol (6): To a solution of 3.83 g AlMe_3 (53.1 mmol) in 30 mL CH_2Cl_2 was added 2.59 g Cp_2ZrCl_2 (8.9 mmol) and the resulting lemon-yellow solution was stirred for 10 min at RT and cooled to 0°C . A solution of 2.0 mL hex-5-yn-1-ol (17.7 mmol) in 14 mL CH_2Cl_2 was then added and stirred at RT for 24 h. The reaction mixture was then cannula transferred to a solution of 3.56 g 2-methoxybenzyl bromide (17.7 mmol) and 1.02 g $\text{Pd}(\text{PPh}_3)_4$ (0.88 mmol) in 88 mL THF at 0°C . Raised to RT and stirred for 24 hours. The reaction was quenched at 0°C by slow addition of 120 mL 3N HCl and the aqueous phase was extracted with ether. The combined organic extracts were washed with sat. NaHCO_3 , sat. NaCl, dried over MgSO_4 and concentrated under vacuum. The crude material was purified by flash chromatography in 1.5:1 hexanes:EtOAc to yield 4.08 g of a yellow oil (93%). ^1H NMR: (400 MHz, CDCl_3) δ 7.14 (m, 2H), 6.85 (m, 2H), 5.30 (td, 1H, $J = 7.2, 1.2$ Hz), 3.80 (s, 3H), 3.63 (t, 2H, $J = 6.4$ Hz), 3.31 (d, 2H, $J = 7.2$ Hz), 2.03 (t, 2H, $J = 7.2$ Hz), 1.68 (s, 3H), 1.50 (m, 4H); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.3, 135.8, 130.0, 129.2, 126.8, 122.6, 120.4, 110.1, 63.0, 55.3, 39.4, 32.4, 28.2, 24.0, 15.9; HR ESI $m/z = 257.1509$ (257.1512 calculated for $\text{C}_{15}\text{H}_{22}\text{O}_2\text{Na}$)

7-(2-Methoxy-phenyl)-5-methyl-hept-5-enal (7): A solution of 0.365 mL oxalyl chloride (4.18 mmol) in 8.3 mL CH_2Cl_2 was cooled to -78°C and 0.590 mL DMSO (8.31 mmol) was added dropwise via syringe. After stirring 30 min at this temperature, 865 mg 7-(2-Methoxy-phenyl)-5-methyl-hept-5-en-1-ol **6** (3.32 mmol) in 8.3 mL CH_2Cl_2 was added dropwise via cannula. The reaction was stirred for 30 min and 2.3 mL triethylamine (17 mmol) was added via syringe. The temperature was maintained for 60 min before warming to RT and stirring for 60 min. The reaction was quenched with 20 mL water and extracted with ether. The combined organic extracts were washed with sat. NaCl, dried over MgSO_4 and concentrated under vacuum. The crude material was purified by flash chromatography in 4:1 hexanes/EtOAc to yield 638 mg (83%) of a yellow oil. ^1H NMR: (400 MHz, CDCl_3) δ 9.74 (t, 1H, $J = 1.6$ Hz), 7.15 (m, 2H), 6.85

(m, 2H), 5.32 (td, 1H, J = 7.2, 1.2 Hz), 3.82 (s, 3H), 3.32 (d, 2H, J = 7.2 Hz), 2.38 (td, 2H, J = 7.4, 1.6 Hz), 2.05 (t, 2H, J = 7.2 Hz), 1.76 (m, 2H), 1.69 (s, 3H); ^{13}C NMR: (100 MHz, CDCl_3) δ 202.7, 157.2, 134.7, 129.7, 129.2, 126.9, 123.7, 120.3, 110.1, 55.2, 43.2, 38.8, 28.3, 20.2, 15.7; HR ESI m/z = 255.1365 (255.1356 calculated for $\text{C}_{15}\text{H}_{20}\text{O}_2\text{Na}$)

1-(8,8-Dideutero-3-methyl-octa-2,7-dienyl)-2-methoxy-benzene³ (8): To 273 mg 7-(2-Methoxy-phenyl)-5-methyl-hept-5-enal **7** (1.18 mmol) was added 6.9 mL of a 0.5 M solution of bis(η^5 -cyclopentadienyl)-di(methyl- d_3) titanium in THF (3.45 mmol). The reaction was heated to reflux in the dark until no aldehyde was present (TLC, 48 hours). The reaction was diluted with 20 mL ether and cooled to 0°C for several hours before filtering through celite. Polar impurities were removed by filtering through a plug of silica (4:1 hexanes/EtOAc) and the crude material was purified by flash chromatography in 200:1 hexanes/EtOAc to yield 41 mg (15%) of a colorless oil. The isotopic purity was >95% at the 8 position (^1H NMR); ^1H NMR: (400 MHz, CDCl_3) δ 7.14 (m, 2H), 6.85 (m, 2H), 5.79 (bm, 1H), 5.30 (td, 1H, J = 7.2, 1.2 Hz), 3.81 (s, 3H), 3.31 (d, 2H, J = 7.2 Hz), 2.01 (m, 4H), 1.67 (s, 3H), 1.50 (m, 2H); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.3, 138.8, 136.0, 130.1, 129.2, 126.8, 122.5, 120.4, 113.8 (m), 110.1, 55.3, 39.2, 33.3, 28.2, 27.2, 15.9; LR EI m/z = 232.2 (232.18 calculated for $\text{C}_{16}\text{H}_{20}\text{D}_2\text{O}$)

Trans-1-(2-deutero-3-methyl-octa-2,7-dienyl)-2-methoxy-benzene (9): To a solution of 2.06 g AlMe_3 (28.6 mmol) in 20 mL CH_2Cl_2 was added 2.09 g Cp_2ZrCl_2 (7.1 mmol) and the resulting lemon-yellow solution was stirred for 10 min at ambient temperature. A solution of 1.36 g 7-Deutero-hept-1-en-6-yne (14.3 mmol) in 15 mL CH_2Cl_2 was then added and stirred for 24 h. The reaction mixture was then cannula transferred to a solution of 2.87 g 2-methoxybenzyl bromide (14.3 mmol) and 0.83 g $\text{Pd}(\text{PPh}_3)_4$ (0.7 mmol) in 70 mL THF at 0°C. Raised to RT and stirred for 24 hours. The reaction was quenched at 0°C by slow addition of 70 mL 3N HCl and the aqueous phase was extracted with ether. The combined organic extracts were washed with sat. NaHCO_3 , sat. NaCl, dried over MgSO_4 and concentrated under vacuum. The crude material was purified by flash chromatography in 99:1 hexanes:EtOAc to yield 2.25 g of a colorless oil (68%). The isotopic purity was 90% D at the 2 position (^1H NMR); ^1H NMR: (400 MHz, CDCl_3) δ 7.17 (m, 2H), 6.87 (m, 2H), 5.81 (m, 1H), 4.97 (m, 2H), 3.82 (s, 3H), 3.33 (s, 2H), 2.03 (m, 4H), 1.69 (s, 3H), 1.52 (m, 2H); ^{13}C NMR: (100 MHz, CDCl_3) δ 157.3,

139.0, 135.8, 130.0, 129.2, 126.8, 122.2 (t, $J_{CD} = 23$ Hz), 120.4, 114.3, 110.1, 55.3, 39.1, 33.4, 28.1, 27.2, 15.9; LR EI $m/z = 231.2$ (231.17 calculated for $C_{16}H_{21}DO$)

Trans-1-(1,1-Dideutero-3-methyl-octa-2,7-dienyl)-2-methoxy-benzene (10): To a solution of 1.11 g $AlMe_3$ (15.4 mmol) in 12.0 mL CH_2Cl_2 was added 1.10 g Cp_2ZrCl_2 (3.7 mmol) and the resulting lemon-yellow solution was stirred for 10 min at RT. A solution of 0.74 g hept-1-en-6-yne (7.5 mmol) in 7.0 mL CH_2Cl_2 was then added and stirred at RT for 24 h. The reaction mixture was then cannula transferred to a solution of 1.60 g 1-(Bromo-dideutero-methyl)-2-methoxy-benzene (7.86 mmol) and 400 mg $Pd(PPh_3)_4$ (0.35 mmol) in 40 mL THF at $0^\circ C$. Raised to RT and stirred for 24 hours. The reaction was quenched at $0^\circ C$ by slow addition of 40 mL 3N HCl and the aqueous phase was extracted with ether. The combined organic extracts were washed with sat. $NaHCO_3$, sat. NaCl, dried over $MgSO_4$ and concentrated under vacuum. The crude material was purified by flash chromatography in 99:1 hexanes:EtOAc to yield 1.25 g of a colorless oil (72%). The isotopic purity was >95% at the 1 position (1H NMR); 1H NMR: (400 MHz, $CDCl_3$) δ 7.14 (m, 2H), 6.86 (m, 2H), 5.81 (m, 1H), 5.30 (bs, 1H), 4.97 (m, 2H), 3.82 (s, 3H), 2.02 (m, 4H), 1.69 (s, 3H), 1.51 (m, 2H); ^{13}C NMR: (100 MHz, $CDCl_3$) δ 157.3, 139.0, 135.9, 130.0, 129.2, 126.8, 122.4, 120.4, 114.3, 110.1, 55.3, 39.2, 33.4, 27.6 (p, $J_{CD} = 19$ Hz), 27.2, 15.9; LR EI $m/z = 232.2$ (232.18 calculated for $C_{16}H_{20}D_2O$)

Figure 2 Cycloisomerization of Labeled Substrates

7,7-Dideuterio-1-(2-methoxy-benzyl)-2-methyl-bicyclo[4.1.0]heptane (3e): A procedure similar to **2a** was followed. The crude material flashed on Ag⁺-impregnated silica in 100:1 hexanes:EtOAc to yield 20 mg (74%) of a colorless oil. >95:5 d.r. by ¹H NMR; ¹H NMR: (400 MHz, CDCl₃) δ 7.25 (m, 1H), 7.16 (m, 1H), 6.85 (m, 2H), 3.79 (s, 3H), 3.18 (d, 1H, J = 14.4 Hz), 2.07 (d, 1H, J = 14.4 Hz), 1.97 (m, 1H), 1.66 (m, 1H), 1.35 (m, 2H), 1.26 (m, 1H), 1.00 (m, 2H), 0.93 (d, 3H, J = 6.8 Hz), 0.66 (m, 1H); ¹³C NMR: (100 MHz, CDCl₃) δ 157.9, 130.4, 128.9, 126.8, 120.0, 110.2, 55.2, 37.2, 30.0, 29.9, 24.5, 24.4, 21.9, 19.4, 18.3, 12.9 (m); LR EI *m/z* = 232.2 (calculated 232.18 for C₁₆H₂₀D₂O)

Trans-1-(2-Methoxy-benzyl)-2-deuterio-2-methyl-bicyclo[4.1.0]heptane (3f): A procedure similar to **2a** was followed. The crude material flashed on Ag⁺-impregnated silica in 100:1 hexanes:EtOAc to yield 33 mg (80%) of a colorless oil. >95:5 d.r. by ¹H NMR; ¹H NMR: (400 MHz, CDCl₃) δ 7.24 (m, 1H), 7.16 (m, 1H), 6.90 (m, 1H), 6.86 (m, 1H), 3.79 (s, 3H), 3.19 (d, 1H, J = 14.8 Hz), 2.03 (d, 1H, J = 14.8 Hz), 1.98 (m, 1H), 1.32 (m, 3H), 0.99 (m, 2H), 0.95 (s, 3H), 0.66 (m, 1H), 0.30 (dd 1H J = 9.2, 4.4 Hz), 0.11 (t, 1H, J = 5.0 Hz); ¹³C NMR: (100 MHz, CDCl₃) δ 158.0, 130.5, 129.0, 126.9, 120.1, 110.3, 55.2, 37.4, 30.0, 29.6 (t, J_{C-D} = 76 Hz), 24.7, 24.6, 21.9, 19.4, 19.3, 13.8; LR EI *m/z* = 231.2 (231.17 calculated for C₁₆H₂₂O)

Trans-1-(α,α-dideuterio-2-methoxy-benzyl)-2-methyl-bicyclo[4.1.0]heptane (3g): Followed procedure **2a** to yield 30 mg (71%) of a colorless oil; >95:5 d.r. by ¹H NMR; ¹H NMR: (400 MHz, CDCl₃) δ 7.24 (dd, 1H, J = 7.2, 1.6 Hz), 7.17 (td, 1H, J = 7.6, 1.6 Hz), 6.86 (m, 2H), 3.80 (s, 3H), 1.97 (m, 1H), 1.67 (m, 1H), 1.33 (m, 3H), 0.99 (m, 2H), 0.94 (d, 3H, J = 6.8 Hz), 0.66 (m, 1H), 0.27 (dd, 1H, J = 9.2, 4.4 Hz), 0.09 (t, 1H, J = 4.8 Hz); ¹³C NMR: (100 MHz, CDCl₃) δ 157.9, 130.4, 128.9, 126.8, 120.0, 110.2, 55.2, 36.5 (m), 30.0, 29.9, 24.5 (2C), 22.0, 19.4, 18.4, 13.8; LR EI *m/z* = 232.2 (232.18 calculated for C₁₆H₂₀D₂O)

Figure 3 Calculated conformations of 1,2-dimethyl-bicyclo[4.1.0]heptane

2. 2D NMR Experiments

2.1 Stereochemical Determination of **3a**

The relative stereochemistry of **3a** was determined by a 2D NOESY experiment. The lowest energy conformations of the two possible stereoisomers of the related compound, 1,2-dimethyl-bicyclo[4.1.0]heptane, were obtained from a Monte-Carlo minimization (Spartan) at the 3-21G* energy level as shown in Fig. 1. Because a strong NOE was observed in **3a** between the cyclopropyl methylene and the methyl (and not the methine), the *trans* relative stereochemistry has been assigned. The relevant NOESY correlations are tabulated below, along with ^1H and COSY data. The minor (*cis*) isomer could be distinctly observed in the crude ^1H NMR of **3a**. The ^1H NMR spectrum of **3c** matches the *trans* isomer of **3a**, and so **3c** has been assigned the *trans* relative stereochemistry as well.

2.2 Structural Determination of **4a**

Because isomerization product **4** was obtained as an inseparable mixture of isomers, definitive assignment of structure by 2D NMR experiments alone was difficult (COSY, HMQC, HMBC). When **4a** was hydrogenated however, a single compound was obtained (eq. 1). This product was fully characterized by NMR

¹H NMR Chemical Shifts (ppm) for 2a

Position	δ ¹ H
1	--
2	1.67
3	1.33, 0.64 ^a
4	1.27, 0.99 ^a
5	1.97, 1.33 ^a
6	0.99
7a	0.1
7b	0.27
8	0.94
9	3.18, 2.08 ^a
10	--
11	--
12	6.83
13	7.16
14	6.89
15	7.24
16	3.79

(a) diastereotopic methylene

¹H/¹H Correlations (COSY) for 2a

Proton	Cross Peaks
H2	H3, H8
H3	H2, H3 ^a , H4
H4	H3, H4 ^a , H5 ^b
H5	H4 ^b , H5 ^a , H6
H6	H5, H7
H7a	H6, H7b
H7b	H6, H7a
H8	H2
H9	H9 ^a
H12	H13
H13	H12, H14
H14	H13, H15
H15	H14

(a) geminal coupling (b) weak

Selected NOESY Correlations for 2a

Proton	Cross Peaks
H2	H3, H4 ^a , H8, H9 ^a
H3	H2, H3 ^b , H4, H7a, H8
H4	H2 ^a , H3, H4 ^b , H5, H6 ^a
H5	H4, H5 ^b , H6, H7a ^a
H6	H4 ^a , H5, H7b
H7a	H3, H5 ^a , H7b, H8,
H7b	H6, H7a, H9
H8	H2, H3 ^a , H7a, H9
H9	H2, H7b, H8, H9 ^b

(a) weak (b) diastereotopic methylene

2a

Once the carbon skeleton of **4a** was certain, the assignment of major and minor regioisomers was easily determined. 2D NMR data for **11** is summarized below.

1-Methoxy-2-[2-(1-methyl-cyclohexyl)-ethyl]-benzene (11): To a solution of 6.0 mg (0.026 mmol) 1-Methoxy-2-[2-(1-methyl-cyclohexenyl)-ethyl]-benzene (**4a**; mixture of regioisomeric alkenes) in 1.0 mL MeOH was added 5.5 mg Pd/C (0.003 mmol Pd). The

solution was saturated with H₂ at 0°C and raised to 25°C, stirring under H₂. After complete reaction (GC, 5 h), the solution was filtered through celite and concentrated under reduced pressure to yield 5 mg (83%) of a colorless oil; ¹H NMR: (600 MHz, CDCl₃) δ 7.13 (m, 2H), 6.86 (m, 1H), 6.82 (d, 1H, J = 7.8 Hz), 3.81 (s, 3H), 2.53 (m, 2H), 1.45 (m, 6H), 1.33 (m, 4H), 1.27 (m, 2H), 0.94 (s, 3H); ¹³C NMR: (100 MHz, CDCl₃) δ 157.4, 132.3, 129.6, 126.6, 120.4, 110.3, 55.3, 41.9, 37.8, 32.9, 26.6, 25.2, 24.2, 22.1; LR EI *m/z* = 232.2 (calculated 232.18 for C₁₆H₂₄O)

¹H and ¹³C NMR Chemical Shifts (ppm) for 10

Position	δ ¹ H	δ ¹³ C
1	--	157.4
2	6.82	110.3
3	7.13	126.6
4	6.86	120.4
5	7.13	129.6
6	--	132.3
7	3.81	55.3
8	2.53	24.2
9	1.45	41.9
10	--	32.9
11	1.27, 1.33 ^a	37.8
12	1.45 ^a	22.1
13	1.33, 1.45	26.6
14	0.94	25.2

(a) diastereotopic methylene

One-Bond ¹³C/¹H Correlations (HMQC) for 10

Carbon	Cross Peak
C1	--
C2	H2
C3	H3
C4	C4
C5	C5
C6	--
C7	H7
C8	C8
C9	H9
C10	--
C11	H11
C12	H12
C14	H14

Long-range (Mainly Three-bond) ¹³C/¹H Correlations (HMBC) for 10

Carbon	Cross Peaks
C1	H2 ^a , H3, H5, H8
C2 ^b	H3
C3	H5
C4	H2
C5	H3, H4 ^a , C8
C6	H2, H4, H8, H9
C7	none
C8 ^c	H9, H11 ^a
C9	H8, H11 ^a , H14
C10	H8 ^a , H9, H11, H12 ^a , H14
C11	H9 ^a , H11', H12 ^a , H13, H14
C12	H11, H12', H13
C13	H11, H12
C14	C9, C11, C12 ^a

(a) weak (b) H4 not observed (c) H5 not observed

-
- ¹ Negishi, E., Holmes, S.J., Tour, J.M., Miller, J.A., Cederbaum, F.E., Swanson, D.R., Takahashi, T. *J. Am. Chem. Soc.* **1989**, *111*, 3336.
- ² (a) Kelley, J.L., Linn, J.A., Selway, J.W.T. *J. Med. Chem.* **1989**, *32*, 1757. (b) Hussain, A., Parrick, J. *Tett. Lett.* **1983**, *24*, 609.
- ³ Petasis, N. A.; Bzowej, E. I. *J. Am. Chem. Soc.* **1990**, *112*, 6392.
- ⁴ Hahn, C.; Morvillo, P.; Herdtweck, E.; Vitagliano, A. *Organomet.* **2002**, *21*, 1807.
- ⁵ Fernandez, D., Sevilano, P., Garcia-Seijo, M.I., Castineiras, A., Janosi, L., Berente, Z., Kollar, L., Garcia-Fernandez, M.E. *Inorg.Chim. Act.* **2001**, *312*, 40.
- ⁶ Tong-Shuang, L.; Ji-Tai, L.; Hui-Zhang, L. *J. Chrom. A.* **1995**, *715*, 372.
- ⁷ Negishi, E.; Matsushita, H.; Okukado, N. *Tett. Lett.* **1981**, *22*, 2715.