Potassium Hydroxide-Mediated Novel Rearrangement of 2-Alkylsulfonyl-2-arylsulfonyl-1-phenylethanones to 1-aryl-2-(arylsulfonylmethanesulfonyl)ethanones

Jong Kwan Bin, Jae Sang Lee and Kyongtae Kim*

School of Chemistry and Molecular Engineering, Seoul National University, Seoul 151-742, Korea

kkim@plaza.snu.ac.kr

2,2-Bis(ethanesulfonyl)-1-phenylethanone 7b

: white solid; mp 142-143 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 2993, 2937, 1671, 1594, 1336, 1313, 1280, 1187, 1149, 1122, 973, 856, 840, 792, 761, 682, and 586 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (t, *J* = 7.5 Hz, 6H), 3.70 (m, 4H), 6.21 (s, 1H), 7.52-8.00 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 5.68, 48.5, 80.2, 129.2, 129.4, 135.2, 136.3, 184.5. Anal. Calcd for C₁₂H₁₆O₅S₂: C, 47.35; H, 5.30; S, 21.07. Found: C, 47.37; H, 5.32; S, 21.12.

1-Phenyl-2,2-bis(propane-1-sulfonyl)ethanone 7c

: white solid; mp 138-140 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 2992, 1667, 1311, 1225, 1178, 1134, 1112, 960, 853, 734, 578, and 534 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.03 (t, *J* = 7.4 Hz), 1.96 (m, 4H), 3.74 (m, 4H), 6.14 (s, 1H), 7.42-7.97 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 13.1, 25.6, 55.3, 81.3, 129.2, 129.4, 133.7, 137.9, 186.2. Anal. Calcd for C₁₄H₂₀O₅S₂: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.53; H, 6.06; S, 19.21.

1-Phenyl-2,2-bis(propane-2-sulfonyl)ethanone 7d

: white solid; mp 138-139 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3002, 2940, 1677, 1594, 1448, 1392, 1322, 1305, 1272, 1203, 1182, 1162, 1132, 1116, 1054, 975, 881, 784, 696, 684, 582, and 543 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.43 (d, *J* = 6.2 Hz), 1.55 (d, *J* = 6.2 Hz), 4.26 (m, 2H), 6.30 (s, 1H), 7.52-8.01 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 16.0, 17.3, 56.1, 78.8, 129.6, 129.7, 135.5, 136.8, 185.0. Anal. Calcd for C₁₄H₂₀O₅S₂: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.58; H, 6.11; S, 19.30.

2,2-Bis(butane-1-sulfonyl)-1-phenylethanone 7e

: white solid; mp 131-132 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 2964, 2942, 1675, 1594, 1467, 1326, 1274, 1195, 1180, 1137, 1122, 1099, 1078, 998, 910, 840, 732, 684, 580, and 530 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.99 (t, *J* = 7.3 Hz, 6H), 1.54 (m, 4H), 1.91

(m, 4H), 3.69 (m, 4H), 6.17 (s, 1H), 7.52- 7.99 (m, 5H); ^{13}C NMR(75 MHz, CDCl_3) δ 14.0, 22.1, 23.4, 53.8, 81.5, 129.6, 129.7, 135.5, 136.7, 185.0. Anal. Calcd for $\text{C}_{16}\text{H}_{24}\text{O}_5\text{S}_2$: C, 53.31; H, 6.71; S, 17.79. Found: C, 53.28; H, 6.74; S, 17.82.

2,2-Bis(methanesulfonyl)-1-(*m*-tolyl)ethanone 7f

: white solid; mp 151-152 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3029, 2992, 1677, 1581, 1388, 1326, 1212, 1080, 1023, 1008, 984, 840, 682, and 464 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.46 (s, 3H), 3.44 (s, 6H), 6.15 (s, 1H), 7.42-7.81 (m, 4H); ^{13}C NMR (75 MHz, CDCl_3) δ 21.8, 42.2, 82.8, 127.1, 129.5, 136.5, 139.7, 185.0. Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_5\text{S}_2$: C, 45.50; H, 4.68; S, 22.09. Found: C, 45.52; H, 4.64; S, 22.05.

2,2-Bis(ethanesulfonyl)-1-(*p*-Bromophenyl)ethanone 7g

: white solid; mp 190-192 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3089, 2942, 1677, 1581, 1562, 1396, 1326, 1284, 1070, 1043, 1008, 964, 840, 727, 682, and 482 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.49 (t, $J = 7.4$ Hz, 6H), 3.66 (m, 4H), 6.12 (s, 1H), 7.68 (d, $J = 8.7$ Hz, 2H), 7.84 (d, $J = 8.7$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 48.5, 80.3, 130.7, 131.0, 132.6, 135.1 183.7. Anal. Calcd for $\text{C}_{12}\text{H}_{12}\text{BrO}_5\text{S}_2$: C, 37.60; H, 3.94; S, 16.73. Found: C, 37.62; H, 3.95; S, 16.77.

2,2-Bis(ethanesulfonyl)-1-(naphthalen-1-yl)ethanone 7h

pale yellow liquid; IR (neat) 2960, 1668, 1598, 1337, 1313, 1221, 1142, 1104, 926, 868, 728, 684 and 569 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.51 (t, $J = 7.4$ Hz, 6H), 3.76 (m, 4H), 6.23 (s, 1H), 7.57-8.62 (m, 7H); ^{13}C NMR (75 MHz, CDCl_3) δ 5.59, 48.5, 83.5, 124.2, 125.4, 125.9, 127.3, 128.6, 129.2, 129.8, 133.9, 134.9, 135.1, 189.0. Anal. Calcd for $\text{C}_{16}\text{H}_{18}\text{O}_5\text{S}_2$: C, 54.22; H, 5.12; S, 18.09. Found: C, 54.12; H, 5.21; S, 18.01.

2,2-Bis(methanesulfonyl)-1-(phenyl)propan-1-one 7i

: white solid; mp 186-187 °C (CH_2Cl_2 – *n*-hexane); IR (KBr) 2928, 1664, 1324, 1305, 1270, 1184, 1136, 1113, 967, 851, 752, 678, 579, and 483 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.14 (s, 3H), 3.00 (s, 6H), 7.32-7.80 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ 7.3, 38.4, 82.4, 129.3, 129.7, 135.6, 137.1, 185.7. Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_5\text{S}_2$: C, 45.50; H, 4.86; S, 22.09. Found: C, 45.63; H, 4.77; S, 22.03.

2,2-Bis(methanesulfonyl)-1-(phenyl)butan-1-one 7j

: white solid; mp 132-134 °C (CH_2Cl_2 – *n*-hexane); IR (KBr) 3024, 2944, 1667, 1587, 1328, 1302, 1222, 1139, 1084, 963, 771, 755, 694, 544, 512, and 474 cm^{-1} ; ^1H NMR

(300 MHz, CDCl₃) δ 0.95 (t, J = 7.5 Hz, 3H), 2.87 (q, J = 7.5 Hz, 2H), 3.34 (s, 6H), 7.45-7.91 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 9.4, 22.0, 41.8, 96.4, 128.5, 129.1, 133.3, 138.9, 192.9. Anal. Calcd for C₁₂H₁₆O₅S₂: C, 47.35; H, 5.30; S, 21.07. Found: C, 47.25; H, 5.31; S, 21.01.

2-Benzenesulfonyl-2-methanesulfonyl-1-phenylethanone 7k

: white solid; mp 180-182 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3062, 2938, 1677, 1587, 1446, 1324, 1272, 1151, 1083, 966, 842, 742, 678, 638, 574, and 505 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.49 (s, 3H), 6.27 (s, 1H), 7.45-7.98 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 41.2, 84.12, 125.9, 129.1, 129.4, 135.1, 135.3, 136.1, 137.6, 185.3. Anal. Calcd for C₁₅H₁₄O₅S₂: C, 53.24; H, 4.17; S, 18.95. Found: C, 53.20; H, 4.21; S, 18.93.

2-Benzenesulfonyl-2-ethanesulfonyl-1-phenylethanone 7l

: white solid; mp 158-159 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2987, 2942, 1679, 1594, 1448, 1334, 1272, 1159, 1081, 1047, 998, 970, 912, 838, 750, 684, 640, 584, and 541 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (t, J = 7.5 Hz, 3H), 3.80 (m, 2H), 6.30 (s, 1H), 7.46-7.96 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 5.1, 48.0, 83.9, 129.1, 129.2, 129.3, 129.9, 135.0, 135.3, 136.1, 137.5, 185.1. Anal. Calcd for C₁₆H₁₆O₅S₂: C, 54.53; H, 4.58; S, 18.20. Found: C, 54.55; H, 4.43; S, 18.17.

2-Benzenesulfonyl-2-(propane-2-sulfonyl)-1-phenylethanone 7m

: white solid; mp 154-156 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3064, 2983, 2940, 1685, 1594, 1448, 1332, 1268, 1159, 1130, 1079, 1054, 968, 750, 684, and 590 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.43 (d, J = 6.9, 3H), 1.56 (d, J = 6.9, 3H), 4.18 (m, 1H), 6.45 (s, 1H), 7.26-7.95 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 15.0, 16.8, 56.0, 81.2, 128.9, 129.0, 129.1, 129.4, 130.7, 134.9, 135.3, 136.2, 184.3. Anal. Calcd for C₁₇H₁₈O₅S₂: C, 55.72; H, 4.95; S, 17.50. Found: C, 55.69; H, 4.81; S, 17.54

2-Benzenesulfonyl-2-(butane-1-sulfonyl)-1-phenylethanone 7n

: pale yellow liquid, IR (neat) 3066, 2962, 2935, 2875, 1681, 1594, 1448, 1336, 1272, 1159, 1081, 998, 970, 752, 684, and 530 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.98 (t, J = 7.3 Hz, 3H), 1.53 (m, 2H), 1.90 (m, 2H), 3.77 (m, 2H), 6.35 (s, 1H), 7.43-7.93 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 13.6, 21.8, 22.5, 53.0, 84.3, 129.1, 129.2, 129.3, 129.9, 135.0, 135.2, 136.2, 137.6, 185.3. Anal. Calcd for C₁₈H₂₀O₅S₂: C, 56.82; H, 5.30; S, 16.86. Found: C, 56.82; H, 5.25; S, 16.84.

2-Benzenesulfonyl-1-phenyl-2-phenylmethanesulfonyl-ethanone 7o

: white solid; mp 182-183 °C (CH₂Cl₂ – *n*-hexane); IR(neat) 3064, 3006, 2919, 1681, 1594, 1448, 1407, 1340, 1272, 1189, 1159, 1132, 1081, 975, 840, 748, 682, 568, and 505 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 4.76 (d, *J* = 13.7 Hz, 1H) 5.19 (d, *J* = 13.7 Hz, 1H), 6.40 (s, 1H) 7.36-7.99 (m, 15H); ¹³C NMR (75 MHz, CDCl₃) δ 60.2, 81.2, 126.0, 129.1, 129.2, 129.3, 129.6, 130.5, 131.9, 135.0, 135.4, 135.9, 136.7, 184.5. Anal. Calcd for C₂₁H₁₈O₅S₂: C, 60.85; H, 4.38; S, 15.47. Found: C, 60.85; H, 4.31; S, 15.46.

2-Methanesulfonyl-1-phenyl-2-(*p*-toluenesulfonyl)-ethanone 7p

: white solid; mp 177-178 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3045, 3014, 2946, 1675, 1594, 1446, 1311, 1272, 1203, 1186, 1147, 1126, 1083, 960, 835, 811, 676, 649, 568, 509, and 478 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.44 (s, 3H), 3.47 (s, 3H), 6.18 (s, 1H), 7.34-7.87 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 21.8, 41.2, 84.4, 129.1, 129.2, 129.8, 130.1, 134.9, 135.0, 136.3, 146.8, 185.4. Anal. Calcd for C₁₆H₁₆O₅S₂: C, 54.53; H, 4.58; S, 18.20. Found: C, 54.55; H, 4.60; S, 18.25.

2-Ethanesulfonyl-1-phenyl-2-(*p*-toluenesulfonyl)-ethanone 7q

: white solid; mp 143-144 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3072, 2942, 2871, 1679, 1594, 1448, 1336, 1270, 1157, 1081, 1045, 970, 912, 838, 752, 684, 661, 578 and 541 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (t, *J* = 7.5 Hz, 3H), 2.44 (s, 3H), 3.70-3.89 (m, 2H), 6.28 (s, 1H), 7.35-7.89 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 5.1, 21.8, 47.9, 83.9, 129.1, 129.2, 129.9, 134.4, 135.0, 136.2, 146.8, 185.2. Anal. Calcd for C₁₇H₁₈O₅S₂: C, 55.72; H, 4.95; S, 17.50. Found: C, 55.70; H, 4.91; S, 17.41.

2-(Butane-1-sulfonyl)-1-phenyl-2-(*p*-toluenesulfonyl)-ethanone 7r

: white solid; mp 126-127 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2960, 2875, 1681, 1594, 1448, 1405, 1338, 1272, 1186, 1157, 1081, 970, 910, 838, 813, 752, 703, 684, 657, 578, and 543 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.99 (t, *J* = 7.3 Hz, 3H), 1.49-1.57 (m, 2H), 1.86-1.93 (m, 2H), 2.43 (s, 3H), 3.68-3.80 (m, 2H), 6.27 (s, 1H), 7.34-7.88 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 13.6, 21.8, 21.9, 22.5, 52.5, 84.3, 129.1, 129.2, 129.9, 134.5, 134.9, 136.2, 146.8, 185.3. Anal. Calcd for C₁₉H₂₂O₅S₂: C, 57.85; H, 5.62; S, 16.26. Found: C, 57.85; H, 5.52; S, 16.12.

1-Phenyl-2-phenylmethanesulfonyl-2-(*p*-toluenesulfonyl)-ethanone 7s

: white solid; mp 182-183 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3064, 2921, 1679, 1592, 1450, 1338, 1270, 1189, 1155, 1079, 973, 914, 840, 808, 746, 694, 653, 601, 559, and

507 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.42 (s, 3H), 4.75 (d, *J* = 13.7 Hz, 1H), 5.16 (d, *J* = 13.7 Hz, 1H), 6.38 (s, 1H), 7.32-7.86 (m, 14H); ¹³C NMR (75 MHz, CDCl₃) δ 21.8, 60.3, 81.3, 126.2, 128.9, 129.0, 129.2, 129.5, 129.7, 130.5, 131.8, 133.8, 134.8, 136.1, 146.8, 184.6. Anal. Calcd for C₂₂H₂₀O₅S₂: C, 61.66; H, 4.70; S, 14.97. Found: C, 61.66; H, 4.79; S, 14.87.

2-(*p*-Methoxybenzenesulfonyl)-1-phenyl-2-methanesulfonylethanone 7t

: white solid; mp 156-157 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3018, 2944, 2840, 1679, 1594, 1448, 1336, 1268, 1149, 1083, 1022, 962, 912, 838, 804, 752, 684, 647, 630, 572, and 532 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.47 (s, 3H), 3.88 (s, 3H), 6.22 (s, 1H), 7.01-7.90 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 41.1, 55.8, 84.3, 114.5, 128.6, 129.1, 129.2, 132.3, 135.0, 136.2, 165.0, 185.5. Anal. Calcd for C₁₆H₁₆O₆S₂: C, 52.16; H, 4.38; S, 17.41. Found: C, 52.16; H, 4.29; S, 17.38.

2-(*p*-Methoxybenzenesulfonyl)-1-phenyl-2-ethanesulfonylethanone 7u

: white solid; mp 142-142 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3072, 2944, 2840, 1681, 1592, 1496, 1336, 1268, 1151, 1083, 1047, 1022, 970, 838, 752, 684, 582, and 543 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.47 (t, *J* = 7.5 Hz, 3H), 3.68-3.84 (m, 2H), 3.88 (s, 3H), 6.28 (s, 1H), 6.99-7.89 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 5.2, 48.0, 55.8, 83.9, 114.4, 128.5, 129.1, 129.2, 132.4, 134.9, 136.2, 165.0, 185.3. Anal. Calcd for C₁₇H₁₈O₆S₂: C, 53.39; H, 4.74; S, 16.77. Found: C, 53.38; H, 4.59; S, 16.79.

2-(*p*-Methoxybenzenesulfonyl)- 2-(propane-2-sulfonyl)- 1-phenylethanone 7v

: white solid; mp 150-151 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2979, 2942, 1683, 1592, 1496, 1332, 1267, 1153, 1081, 1054, 1022, 970, 839, 804, 750, 686, 663, 588, 548, and 528 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.42 (d, *J* = 6.9 Hz, 3H), 1.55 (d, *J* = 6.9 Hz, 3H), 3.88 (s, 3H), 4.14 (m, 1H), 6.41 (s, 1H), 6.99-7.93 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 14.9, 16.9, 55.8, 55.9, 81.3, 114.1, 125.9, 129.1, 129.2, 133.3, 134.8, 136.6, 165.1, 184.6. Anal. Calcd for C₁₈H₂₀O₆S₂: C, 54.53; H, 5.08; S, 16.17. Found: C, 54.49; H, 5.18; S, 16.29.

2-(*p*-Methoxybenzenesulfonyl)-1-phenyl-2-phenylethanesulfonylethanone 7w

: white solid; mp 86-87 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3066, 2944, 2844, 1681, 1591, 1496, 1452, 1413, 1338, 1268, 1149, 1083, 1022, 971, 910, 840, 754, 696, 597, 566, and 511 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.87(s, 3H), 4.71 (d, *J* = 13.7 Hz, 1H), 5.16 (d, *J* = 13.7 Hz, 1H), 6.36 (s, 1H), 6.98-7.91 (m, 14H); ¹³C NMR (75 MHz, CDCl₃)

δ 55.8, 60.3, 81.0, 114.2, 126.3, 127.7, 128.9, 129.0, 129.2, 129.5, 131.8, 133.1, 134.8, 136.2, 165.1, 184.7. Anal. Calcd for $C_{22}H_{20}O_6S_2$: C, 59.44; H, 4.53; S, 14.43. Found: C, 59.54; H, 4.59; S, 14.43.

2-Methanesulfonylmethylsulfonyl-1-phenylethanone 8a

: white solid; mp 155-156 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 1670, 1292, 1120, 851, 688, and 496 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 3.29 (s, 3H), 4.97 (s, 2H), 5.06 (s, 2H), 7.53-7.97 (m, 5H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 42.9, 59.2, 70.6, 129.1, 129.6, 135.4, 135.5, 189.0. Anal. Calcd for $C_{10}H_{12}O_5S_2$: C, 43.46; H, 4.38; S, 23.21. Found: C, 43.53; H, 4.41; S, 23.14.

2-Ethanesulfonylmethylsulfonyl-1-phenylpropan-1-one 8b

: white solid; mp 134-136 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3056, 2960, 1668, 1312, 1165, 855, 567, 496 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 1.45 (t, $J = 7.4$ Hz, 3H), 1.78 (d, $J = 7.2$ Hz, 3H), 3.46 (q, $J = 7.4$ Hz, 2H), 4.56 (d, $J = 15$ Hz, 1H), 4.80 (d, $J = 15$ Hz, 1H), 5.54 (q, $J = 7.2$ Hz, 1H), 7.52-7.55 (m, 2H), 7.56-7.58 (m, 1H), 7.97-7.99 (m, 2H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 6.4, 12.3, 49.2, 63.9, 65.9, 129.0, 129.1, 134.7, 134.9, 193.1. Anal. Calcd for $C_{12}H_{16}O_5S_2$: C, 47.35; H, 5.30; S, 21.07. Found: C, 47.28; H, 5.24; S, 21.17.

1-Phenyl-2-(propane-1-sulfonylmethylsulfonyl)butan-1-one 8c

: white solid; mp 167-168 °C (CH_2Cl_2 – *n*-hexane); IR (KBr) 2928, 1670, 1315, 1215, 1168, 1004, 960, 855, 736, 567, and 512 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 0.92-0.96 (m, 6H), 1.90-1.94 (m, 2H), 2.20-2.29 (m, 2H), 3.44 (t, $J = 7.5$ Hz, 2H), 4.46 (d, $J = 15$ Hz, 1H), 4.60 (d, $J = 15$ Hz, 1H), 5.42 (dd, $J = 14, 4.6$ Hz, 1H), 7.50-8.01 (m, 5H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 12.6, 15.7, 25.2, 31.2, 53.7, 68.4, 71.2, 129.3, 129.5, 134.7, 134.8, 191.2. Anal. Calcd for $C_{14}H_{20}O_5S_2$: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.63; H, 6.00; S, 19.38.

2-Methyl-1-phenyl-2-(propane-2-sulfonylmethylsulfonyl)propan-1-one 8d

: white solid; mp 127-128 °C (CH_2Cl_2 – *n*-hexane); IR (KBr) 3056, 2960, 1668, 1312, 1120, 860, 727, 567, and 496 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 1.46 (s, 3H), 1.48 (s, 3H), 1.85 (s, 6H), 3.79-3.89 (m, 1H), 4.90 (s, 2H), 7.45-7.75 (m, 5H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 14.8, 14.9, 21.2, 53.2, 55.5, 63.2, 125.8, 127.9, 128.5, 132.5, 201.3. Anal. Calcd for $C_{14}H_{20}O_5S_2$: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.66; H, 6.01; S, 19.41.

2-(Butane-1-sulfonylmethylsulfonyl)-1-phenylpentan-1-one 8e

: white solid; mp 119-121 °C (CH₂Cl₂ – *n*-hexane); IR (KBr) 3056, 2960, 1668, 1312, 1248, 1165, 1047, 928, 892, 855, and 567 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.93-0.99 (m, 6H), 1.39-1.53 (m, 4H), 1.83-1.89 (m, 2H), 2.16-2.29 (m, 2H), 3.40 (t, *J* = 7.9 Hz, 2H), 4.57 (d, *J* = 15 Hz, 1H), 4.72 (d, *J* = 15 Hz, 1H), 5.50 (dd, *J* = 14, 4.7 Hz, 1H), 7.53-7.58 (m, 2H), 7.66-7.69 (m, 1H), 8.03-8.06 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 13.8, 14.1, 20.8, 21.9, 24.1, 30.6, 54.9, 67.0, 68.5, 129.4, 129.5, 135.1, 136.5, 193.2. Anal. Calcd for C₁₆H₂₄O₅S₂: C, 53.31; H, 6.71; S, 17.79. Found: C, 53.24; H, 6.67; S, 17.85.

2-Methanesulfonylmethylsulfonyl-1-(*m*-tolyl)ethanone 8f

: white solid; mp 171-173 °C (CH₂Cl₂ – *n*-hexane); IR (KBr) 3056, 2960, 1668, 1312, 1165, 855, 567, and 496 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.46 (s, 3H), 3.29 (s, 3H), 4.96 (s, 2H), 5.04 (s, 2H), 7.43-7.82 (m, 4H); ¹³C NMR (75 MHz, CDCl₃) 21.2, 42.5, 59.8, 71.2, 126.2, 128.7, 130.9, 132.0, 139.1, 141.5, 198.6. Anal. Calcd for C₁₁H₁₄O₅S₂: C, 45.50; H, 4.68; S, 22.09. Found: C, 45.44; H, 4.70; S, 22.15.

1-(*p*-Bromophenyl)-2-ethanesulfonylmethylsulfonyl-propan-1-one 8g

: white solid; mp 183-185 °C (CH₂Cl₂ – *n*-hexane); IR (KBr) 2928, 1674, 1226, 1115, and 490 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.47 (t, *J* = 7.5 Hz, 3H), 1.78 (d, *J* = 7.2 Hz, 3H), 3.44 (q, *J* = 7.5 Hz, 2H), 4.55 (d, *J* = 15 Hz, 1H), 4.76 (d, *J* = 15 Hz, 1H), 5.54 (q, *J* = 7.2 Hz, 1H), 7.70 (d, *J* = 8.7 Hz, 2H), 7.88 (d, *J* = 8.7 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 41.7, 58.6, 69.6, 127.5, 130.5, 131.7, 137.5, 192.4. Anal. Calcd for C₁₂H₁₂BrO₅S₂: C, 37.60; H, 3.94; S, 16.73. Found: C, 37.53; H, 3.85; S, 16.77.

2-Ethanesulfonylmethylsulfonyl-1-(naphthalen-2-yl)propan-1-one 8h

: white solid; mp 207-209 °C (CH₂Cl₂ – *n*-hexane); IR (KBr) 2960, 1668, 1445, 1315, 1221, 1142, 1004, 926, 858, 728, and 569 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (t, *J* = 7.5 Hz, 3H), 1.80 (d, *J* = 7.2 Hz, 3H), 3.46 (q, *J* = 7.5 Hz, 2H), 4.55 (s, *J* = 15 Hz, 1H), 4.85 (d, *J* = 15 Hz, 1H), 5.64 (q, *J* = 7.2 Hz, 1H), 7.56-8.78 (m, 7H); ¹³C NMR (75 MHz, CDCl₃) δ 5.9, 12.8, 50.6, 61.7, 65.9, 125.1, 126.1, 126.3, 128.0, 128.5, 129.1, 130.6, 133.4, 134.2, 135.6, 192.0. Anal. Calcd for C₁₆H₁₈O₅S₂: C, 54.22; H, 5.12; S, 18.09. Found: C, 54.26; H, 5.22; S, 18.01.

2-(1-Methanesulfonylethylsulfonyl)-1-phenylethanone 8i

: white solid; mp 130-132 °C (CH₂Cl₂ – *n*-hexane); IR (KBr) 3040, 2960, 1670, 1315, 1219, 1120, 976, 851, 752, 678, and, 481 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.67 (d, *J* = 7.1 Hz, 3H), 3.20 (s, 3H), 4.61 (q, *J* = 7.0 Hz, 1H), 5.14 (s, 2H), 7.50-7.99 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 5.7, 41.6, 57.3, 75.2, 129.2, 129.3, 135.6, 135.7, 192.6. Anal. Calcd for C₁₁H₁₄O₅S₂: C, 45.50; H, 4.86; S, 22.09. Found: C, 45.57; H, 4.81; S, 22.16.

2-(1-Methanesulfonylpropane-1-sulfonyl)-1-phenylethanone 8j

: pale yellow liquid; IR (neat) 2924, 2854, 1683, 1596, 1462, 1308, 1213, 1126, 913, 965, 909, 754, 687, 504, and 467 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.41 (t, *J* = 7.4 Hz, 3H), 2.25 (m, 1H), 2.49 (m, 1H), 2.50 (m, 1H), 3.15 (s, 3H), 4.87 (d, *J* = 16 Hz, 1H), 4.91 (t, *J* = 6.0 Hz, 1H), 5.38 (d, *J* = 16 Hz, 1H), 7.51-7.96 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 13.7, 18.3, 39.2, 60.3, 81.8, 129.1, 129.5, 135.3, 135.8, 189.4. Anal. Calcd for C₁₂H₁₆O₅S₂: C, 47.35; H, 5.30; S, 21.07. Found: C, 47.32; H, 5.28; S, 21.05.

2-Benzenesulfonylmethylsulfonyl-1-phenylpropan-1-one 8l

: white solid; mp 118-119 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2987, 2923, 1681, 1594, 1448, 1332, 1209, 1160, 1137, 1081, 995, 958, 836, 752, 684, and 505cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.78 (d, *J* = 7.2 Hz, 3H), 4.72 (d, *J* = 15 Hz, 1H), 4.95 (d, *J* = 15 Hz, 1H), 5.58 (q, *J* = 7.2 Hz, 1H), 7.53-8.02 (m 10H); ¹³C NMR (75 MHz, CDCl₃) δ 12.4, 64.6, 70.1, 128.9, 129.1, 129.2, 129.4, 134.7, 135.0, 135.1, 138.4, 193.4. Anal. Calcd for C₁₆H₁₆O₅S₂: C, 54.53; H, 4.58; S, 18.20. Found: C, 54.45; H, 4.47; S, 18.20.

2-Benzenesulfonylmethylsulfonyl-2-methyl-1-phenylpropan-1-one 8m

: white solid; mp 116-117 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2996, 2935, 1673, 1596, 1326, 1265, 1155, 1118, 1081, 964, 835, 752, and 534cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.76 (s, 6H), 5.01 (s, 2H), 7.39-8.06 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 21.3, 68.7, 74.7, 125.9, 128.0, 128.4, 129.1, 129.3, 132.4, 134.8, 136.9, 201.3. Anal. Calcd for C₁₇H₁₈O₅S₂: C, 55.72; H, 4.95; S, 17.50. Found: C, 55.68; H, 4.91; S, 17.52

2-Benzenesulfonylmethylsulfonyl-1-phenylpentan-1-one 8n

: white solid; mp 112-114 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3064, 2967, 2929, 2875, 1679, 1594, 1448, 1332, 1251, 1160, 1139, 1081, 960, 748, and 686cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.94 (t, *J* = 7.3 Hz, 3H), 1.24 (m, 2H), 2.22 (m, 2H), 4.76 (d, *J* = 15 Hz, 1H), 4.84 (d, *J* = 15 Hz, 1H), 5.43 (dd, *J* = 14.0, 4.62 Hz, 1H), 7.52-7.80 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ 13.8, 20.5, 30.2, 68.9, 69.9, 128.9, 129.1, 129.2, 129.4, 134.7,

135.0, 136.2, 138.3, 193.1. Anal. Calcd for $C_{18}H_{20}O_5S_2$: C, 56.82; H, 5.30; S, 16.86. Found: C, 56.81; H, 5.28; S, 16.74

2-Benzenesulfonylmethylsulfonyl-1,2-diphenylethanone 8o

: white solid; mp 182-183 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3064, 2991, 2923, 1683, 1594, 1492, 1448, 1336, 1282, 1207, 1139, 1191, 997, 910, 838, 732, 696, 647, 514, and $436cm^{-1}$; 1H NMR (300 MHz, $CDCl_3$) δ 4.59 (d, J = 15 Hz, 1H), 4.94 (d, J = 15 Hz, 1H), 6.83 (s, 1H), 7.42-7.98 (m, 15H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 70.3, 74.7, 126.5, 128.9, 129.0, 129.2, 129.4, 129.7, 130.4, 131.1, 134.4, 135.0, 135.1, 138.3, 190.6. Anal. Calcd for $C_{21}H_{18}O_5S_2$: C, 60.85; H, 4.38; S, 15.47. Found: C, 60.86; H, 4.28; S, 15.43.

1-Phenyl-2-(*p*-toluenesulfonylmethylsulfonyl)ethanone 8p

: white solid; mp 122-123 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3064, 2987, 2921, 1683, 1594, 1452, 1334, 1220, 1139, 1083, 991, 848, 808, 759, 688, 576, and $511cm^{-1}$; 1H NMR (300 MHz, $CDCl_3$) δ 2.47 (s, 3H), 5.02 (s, 2H), 5.07 (s, 2H), 7.41-7.97 (m, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 21.8, 59.7, 71.7, 128.8, 128.9, 129.2, 130.2, 134.9, 135.5, 146.4, 189.0. Anal. Calcd for $C_{16}H_{16}O_5S_2$: C, 54.53; H, 4.58; S, 18.20. Found: C, 54.59; H, 4.62; S, 18.15.

1-Phenyl-2-(*p*-toluenesulfonylmethylsulfonyl)propan-1-one 8q

: white solid; mp 112-114 °C (CH_2Cl_2 – *n*-hexane); IR (neat) 3064, 2987, 2973, 1683, 1594, 1450, 1334, 1207, 1186, 1160, 1137, 1083, 958, 838, 813, 761, 570, 538, and $509cm^{-1}$; 1H NMR (300 MHz, $CDCl_3$) δ 1.78 (d, J = 7.2 Hz, 3H), 2.46 (s, 3H), 4.69 (d, J = 15 Hz, 1H), 4.91 (d, J = 15 Hz, 1H), 5.59 (q, J = 7.2 Hz, 1H), 7.36-8.0 (m, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 12.3, 21.8, 64.3, 70.0, 128.9, 129.1, 129.2, 130.1, 134.7, 134.9, 135.2, 146.4, 193.3. Anal. Calcd for $C_{17}H_{18}O_5S_2$: C, 55.72; H, 4.95; S, 17.50. Found: C, 55.68; H, 4.89; S, 17.41.

1-Phenyl-2-(*p*-toluenesulfonylmethylsulfonyl)pentan-1-one 8r

: pale yellow liquid; IR (neat) 3066, 2956, 2925, 1681, 1594, 1448, 1334, 1274, 1159, 1081, 970, 838, 750, 719, 684, 584, 543, 530, and $507cm^{-1}$; 1H NMR (300 MHz, $CDCl_3$) δ 0.93 (t, J = 7.3 Hz, 3H), 1.35-1.45 (m, 2H), 2.13-2.25 (m, 2H), 2.43 (s, 3H), 4.76 (q, J = 15 Hz, 2H), 5.46 (q, J = 4.5 Hz, 1H), 7.35-8.04 (m, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 13.8, 20.5, 21.8, 30.2, 68.7, 69.9, 128.9, 129.1, 130.0, 134.7, 135.2, 136.2, 146.3, 193.0. Anal. Calcd for $C_{19}H_{22}O_5S_2$: C, 57.85; H, 5.62; S, 16.26. Found: C, 57.75; H, 5.59; S, 16.26.

1,2-Diphenyl-2-(*p*-toluenesulfonylmethylsulfonyl)ethanone 8s

: white solid; mp 188-189 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3064, 2921, 1679, 1592, 1450, 1338, 1270, 1189, 1155, 1079, 973, 914, 840, 808, 746, 694, 653, 602, 559, and 507 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.42 (s, 3H), 4.75 (d, *J* = 13.7 Hz, 1H), 5.16 (d, *J* = 13.7 Hz, 1H), 6.38 (s, 1H), 7.32-7.86 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 21.8, 60.3, 81.3, 126.2, 128.9, 129.0, 129.2, 129.5, 129.7, 130.5, 131.8, 133.8, 134.8, 136.1, 146.8, 184.6. Anal. Calcd for C₂₂H₂₀O₅S₂: C, 61.66; H, 4.70; S, 14.97. Found: C, 61.76; H, 4.78; S, 14.97.

2-(*p*-Methoxybenzenesulfonylmethylsulfonyl)-1-phenylethanone 8t

: white solid; mp 142-143 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3068, 2991, 2923, 1683, 1592, 1496, 1330, 1267, 1211, 1147, 1085, 1020, 910, 840, 757, 686, and 503cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.90 (s, 3H), 5.03 (s, 2H), 5.10 (s, 2H), 7.06-7.98 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 55.9, 59.6, 71.7, 114.7, 128.8, 129.2, 129.4, 131.3, 135.0, 135.3, 164.8, 189.0. Anal. Calcd for C₁₆H₁₆O₆S₂: C, 52.16; H, 4.38; S, 17.41. Found: C, 52.26; H, 4.36; S, 17.39.

2-(*p*-Methoxybenzenesulfonylmethylsulfonyl)-1-phenylpropan-1-one 8u

: pale yellow liquid; IR (neat) 3058, 2923, 2850, 1681, 1594, 1496, 1450, 1332, 1265, 1207, 1155, 1085, 1022, 954, 836, 804, 759, 572, and 516cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.58 (d, *J* = 7.5 Hz, 3H), 3.88 (s, 3H), 4.69 (d, *J* = 15 Hz, 1H), 4.90 (d, *J* = 15 Hz, 1H), 5.58 (q, *J* = 7.1 Hz, 1H), 7.01-8.02 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 12.3, 29.7, 55.8, 64.2, 70.1, 114.6, 128.3, 129.1, 129.4, 131.3, 134.6, 135.0, 164.7, 193.3. Anal. Calcd for C₁₇H₁₈O₆S₂: C, 53.39; H, 4.74; S, 16.77. Found: C, 53.40; H, 4.64; S, 16.78.

2-(*p*-Methoxybenzenesulfonylmethylsulfonyl)-2-methyl-1-phenylpropan-1-one 8v

: pale yellow liquid; IR (neat) 3064, 2996, 2933, 2846, 1673, 1594, 1577, 1498, 1463, 1415, 1328, 1265, 1213, 1151, 1118, 1085, 1022, 966, 836, 804, 6703, 609, 541 and 486cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.75 (s, 6H), 3.86 (s, 3H), 4.99 (s, 2H), 7.00-7.95 (m, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 21.2, 55.7, 68.9, 74.8, 114.4, 128.0, 128.5, 130.1, 131.5, 132.3, 137.0, 164.6, 201.1. Anal. Calcd for C₁₈H₂₀O₆S₂: C, 54.53; H, 5.08; S, 16.17. Found: C, 54.52; H, 5.21; S, 16.30.

2-(*p*-Methoxybenzenesulfonylmethylsulfonyl)-1,2-diphenylethanone 8w

: white solid; mp 108-109 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 3066, 2987, 2919, 1677, 1594, 1496, 1336, 1303, 1274, 1209, 1187, 1137, 1083, 1022, 85, 838, 771, and 698 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.88 (s, 3H), 4.57 (d, *J* = 15 Hz, 1H), 4.89 (d, *J* = 15 Hz, 1H), 6.85 (s, 1H), 7.00-7.98 (m, 14H); ¹³C NMR (75 MHz, CDCl₃) δ 55.8, 70.4, 74.4, 114.6, 126.5, 129.0, 129.2, 129.4, 129.4, 129.6, 130.4, 131.1, 131.3, 134.5, 135.1, 164.8, 190.5. Anal. Calcd for C₂₂H₂₀O₆S₂: C, 59.44; H, 4.53; S, 14.43. Found: C, 59.44; H, 4.50; S, 14.41.

2-Ethanesulfonylmethylsulfonyl-1-phenylpentan-1-one 8x

: white solid; mp 126-127 °C (CH₂Cl₂ – *n*-hexane); IR (neat) 2975, 2921, 2875, 1675, 1590, 1454, 1322, 1251, 1218, 1133, 1049, 952, 910, 852, 742, and 688 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.94 (t, *J* = 7.3 Hz, 3H), 1.37-1.46 (m, 5H), 2.15-2.27 (m, 2H), 3.38-3.45 (m, 2H), 4.56 (d, *J* = 15.3 Hz, 1H), 4.74 (d, *J* = 15.3 Hz, 1H), 4.97 (dd, *J* = 14, 4.7 Hz, 1H), 7.52-8.04 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 6.47, 13.7, 20.5, 30.2, 49.4, 65.8, 68.2, 129.1, 129.2, 134.8, 136.1, 192.9. Anal. Calcd for C₁₄H₂₀O₅S₂: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.48; H, 6.07; S, 19.21.

2-Butanesulfonylmethylsulfonyl-1-phenylpropan-1-one 8y

: pale yellow liquid; IR (neat) 2964, 2931, 2875, 1683, 1594, 1581, 1450, 1332, 1241, 1209, 1130, 1105, 1000, 958, 916, 846, 765, 728, and 686 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.96 (t, *J* = 7.3 Hz, 3H), 1.45-1.52 (m, 2H), 1.75-1.88 (m, 5H), 3.39 (t, *J* = 7.9 Hz, 2H), 4.59 (d, *J* = 15.4 Hz, 1H), 4.80 (d, *J* = 15.4 Hz, 1H), 5.57 (q, *J* = 7.2 Hz, 1H), 7.51-8.01 (m, 5H); ¹³C NMR (75 MHz, CDCl₃) δ 12.7, 13.9, 21.9, 24.0, 54.8, 64.2, 67.0, 129.4, 129.5, 135.1, 135.3, 193.5. Anal. Calcd for C₁₄H₂₀O₅S₂: C, 50.58; H, 6.06; S, 19.29. Found: C, 50.52; H, 6.11; S, 19.30.