SUPPORTING INFORMATION

A Practical Asymmetric Approach to Medium-Sized Carbocycles based on the Combination of Two Ru-Catalyzed Transformations and a Lewis Acid- Induced Cyclization

*Fernando López, Luis Castedo and José L. Mascareñas**

*Departamento de Química Orgánica, Universidad de Santiago de Compostela, 15782 Santiago
de Compostela, Spain.*

General Procedures: All dry solvents were freshly distilled under argon from the appropriate drying agent before use. THF and Et₂O were distilled from sodium/benzophenone. CH₂Cl₂ was distilled from CaH₂. All reactions were conducted in dry solvents under argon atmosphere unless otherwise stated. The ruthenium complexes [CpRu(CH₃CN)₃]PF₆¹ and (S,S)-TsDPEN [(S,S)-**7**]² were prepared according to published procedures. Thin-layer chromatography (TLC) was performed on silica gel plates and components were visualized by observation under UV light, or by treating the plates with a *p*-anisaldehyde or phosphomolybdic reagent followed by heating. Flash chromatography was performed on silica gel, unless otherwise stated. Drying was performed with anhydrous Na₂SO₄. Concentrations were carried out in a rotary evaporator. ¹H and ¹³C NMR spectra were recorded in CDCl₃, at 250 MHz and 62.9 MHz, respectively, and in some cases at 300 or 500 MHz (75.4 or 125.7 for ¹³C NMR). Carbon types were determined from DEPT ¹³C NMR experiments.

General procedure for the synthesis of ketones 6a-c (exemplified for 6a).³

4-(trimethylsilyl)-1-phenylbut-3-yn-2-one (6a). AlCl₃ (1.0 g, 7.56 mmol) over 5 min was added to a solution of 2-phenylacetyl chloride (1.17 g, 7.56 mmol) and 1,2-bis(trimethylsilyl)ethyne (1.35g, 7.94 mmol) in CH₂Cl₂ (25 mL) at 0 °C. After stirring for 15 min at that temperature, water (10 mL) and HCl (10% aq., 1 mL) were added. The organic phase was separated, extracted with CH₂Cl₂ and washed with NaHCO₃ (10% aq.) and brine. The combined organic phases were dried, filtered, concentrated and chromatographed (1% Et₂O/hexanes) to afford **6a** as a colorless oil [(1.37 g, 84%). R_f = 0.61 (10% EtOAc/hexanes)]; ¹H NMR δ 7.33 (m, 5H), 3.85 (s, 2H), 0.26 (s, 9H); ¹³C NMR δ 184.6 (C), 132.7 (C), 129.7 (CH), 128.5 (CH), 127.2 (CH), 101.5 (C), 99.9 (C), 51.8 (CH₂), -1.10 (CH₃); LRMS (CI, m/z): 217 (M⁺ + 1, 13), 201 (5), 183 (4), 125 (14), 111 (18), 43 (100); HRMS calcd. for C₁₃H₁₇SiO 217.1049, found 217.1046.

5-(3-methoxyphenyl)-1-(trimethylsilyl)pent-1-yn-3-one (6b).⁴ (87%, colorless oil). R_f = 0.38 (10% Et₂O/hexanes); ¹H NMR δ 7.26 (m, 1H), 6.80 (m, 3H), 3.81 (s, 3H), 2.96 (m, 4H), 0.27 (s, 9H); ¹³C NMR δ 186.3 (C), 159.6 (C), 141.6 (C), 129.4 (CH), 120.5 (CH), 114.0 (CH), 111.4

¹ Gill, T. P.; Mann, K.R. *Organometallics* **1982**, *1*, 485.

² Haack, K.-J.; Hashiguchi, S.; Fujii, A.; Ikariya, T.; Noyori, R. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 285.

³ (a) Cornwall, N. J.; Linehan, S.; Weavers, R. T.; *Tetrahedron Lett.* **1997**, *38*, 2919-2922. (b) Birkofer, L.; Ritter, A.; Uhlenbrauck, H. *Chem. Ber.* **1963**, *96*, 3280-3288.

⁴ The required propanoyl chloride **5b** was prepared from commercial available 3-(3-methoxyphenyl)propanoic acid and SOCl₂ under standard conditions. The crude residue was used without further purification.

(CH), 101.8 (C), 98.0 (C), 54.9 (CH₃), 46.5 (CH₂), 29.6 (CH₂), -0.93 (CH₃); LRMS (CI, m/z) 261 (M⁺ + 1, 53), 245 (44), 242 (30), 227 (13), 171 (23), 163 (100), 149 (20); HRMS calcd. for C₁₅H₂₁SiO₂ 261.1311, found 261.1312.

1-cyclohexenyl-4-(trimethylsilyl)but-3-yn-2-one (6c).⁵ ¹H NMR δ 5.63 (s, 1H), 3.13 (s, 2H), 2.02 (m, 4H), 1.59 (m, 4H), 0.21 (s, 9H). ¹³C-RMN δ 184.9 (C), 130.1 (C), 126.3 (CH), 101.3 (C), 99.5 (C), 43.3 (CH₂), 28.3 (CH₂), 25.2 (CH₂), 22.6 (CH₂), 21.8 (CH₂), -1.07 (CH₃). LRMS (CI, m/z) 221 (M⁺ + 7, 53), 205 (22), 176 (100); HRMS calcd. for C₁₃H₂₀OSi 220.1283, found 220.1298.

Procedure for the synthesis of ketones **6d-e** (exemplified for **6d**)

6-methyl-1-(trimethylsilyl)hept-6-en-1-yn-3-one (6d). A solution of *rac*-**1d**⁶ (920 mg, 4.69 mmol) in CH₂Cl₂ (10 mL) was added over 5 min to a suspension of PDC (2.63 g, 7.01 mmol) and MS (4 Å, 2 g) in CH₂Cl₂ (40 mL). After stirring for 45 min, the reaction mixture was filtered over celite and silica gel (1 :1 aprox.) and washed with Et₂O. The organic phases were concentrated and bulb-to-bulb distilled (18 mmHg, 75 °C) to afford **6d** [715 mg, 78%, R_f = 0.61 (5% EtOAc/hexanes)]. Ketone **6d** has been previously reported.⁷

1,6-bis(trimethylsilyl)hept-6-en-1-yn-3-one (6e). [65%, R_f = 0.73 (10% Et₂O /hexanes)]. ¹H NMR δ 5.48 (s, 1H), 5.26 (s, 1H), 2.61 (m, 2H), 2.36 (m, 2H), 0.25 (s, 9H), 0.03 (s, 9H). ¹³C NMR δ 187.2 (C), 149.8 (C), 124.1 (CH₂), 101.8 (C), 97.7 (C), 44.1 (CH₂), 29.2 (CH₂), -0.86 (CH₃), -1.80 (CH₃); LRMS (CI, m/z): 253 (M⁺ + 1, 25), 237 (21), 221 (6), 177 (10), 30 (100); HRMS calcd. for C₁₃H₂₅Si₂O 253.1444, found 253.1441.

Procedure for the asymmetric synthesis of trimethylsilylalkynols (*S*)-**1**. (exemplified for **1a**).

(2S)-4-(trimethylsilyl)-1-phenylbut-3-yn-2-ol (1a). A mixture of (*S,S*)-**7** (7.7 mg, 0.013 mmol) and ketone **6a** (100 mg, 0.52 mmol) in *i*-PrOH (5.2 mL) was stirred at rt for 20 h. The reaction mixture was then concentrated under reduced pressure. The residue was subjected to

⁵ 2-cyclohexenylacetyl chloride **5c** was prepared from 2-cyclohexenylacetic acid by treatment with SOCl₂, see: Belleau, B.; Gulini, U.; Gour-Salin, B.; Ahmed, F. R. *Can. J. Chem.* **1985**, 63, 1268.

⁶ López, F.; Castedo, L.; Mascareñas, J. L. *J. Am. Chem. Soc.* **2002**, 124, 4218-4219.

⁷ Krafft, M. E.; Boñaga, L. V. R.; Felts, A. S.; Hirose, C.; Kerrigan, S. *J. Org. Chem.* **2003**, 68, 6039-6042.

flash chromatography on silica gel to afford (*S*)-**1a** as a colorless oil. (96 mg, 97%, ee > 97%).⁸ $[\alpha]_{\text{D}}^{20} = -7.4$ (c 1.0, CHCl₃), [lit.⁹ (*R*)-**1a**, $[\alpha]_{\text{D}}^{20} = +7.2$ (c 1.0, CHCl₃)]; ¹H NMR δ 7.12 (5H, m), 4.36 (1H, m), 2.81 (2H, d, *J* = 6.3 Hz), 2.24 (1H, s), 0.05 (9H, s); ¹³C NMR δ 137.0 (C), 130.4 (CH), 128.7 (CH), 127.3 (CH), 106.5 (C), 91.0 (C), 63.9 (CH), 44.4 (CH₂), 0.25 (CH₃); LRMS *m/z* 218 (M⁺, 1), 201 (1), 169 (2), 92 (4), 58 (100); HRMS calcd. for C₁₃H₁₈SiO 218.1127, found 218.1119.

(3*S*)-5-(3-methoxyphenyl)-1-(trimethylsilyl)pent-1-yn-3-ol (1b). [96%, ee > 97%, colorless oil]. $[\alpha]_{\text{D}}^{20} = -8.0$ (c 2.2, CHCl₃); ¹H NMR δ 7.25 (1H, m), 6.82 (3H, m), 4.41 (1H, m), 3.83 (3H, s), 2.83 (3H, m), 2.09 (2H, m), 0.25 (9H, s); ¹³C NMR δ 160.0 (C), 143.4 (C), 129.8 (CH), 121.4 (CH), 114.7 (CH), 111.8 (CH), 107.3 (C), 90.0 (C), 62.4 (CH), 55.5 (CH₃), 39.5 (CH₂), 32.0 (CH₂), 0.16 (CH₃); LRMS *m/z* 262 (M⁺, 10), 261 (17), 247 (18), 231 (72), 215 (18), 135 (25), 122 (100); HRMS calcd. for C₁₅H₂₂SiO₂ 262.1389, found 262.1388.

(2*S*)-1-cyclohexenyl-4-(trimethylsilyl)but-3-yn-2-ol (1c). [84%, ee > 97%, colorless oil]. $[\alpha]_{\text{D}}^{20} = -1.54$ (c 0.6, CHCl₃); ¹H NMR δ 5.53 (1H, s), 4.41 (1H, t, *J* = 6.6 Hz), 2.32 (2H, d, *J* = 6.5 Hz), 2.22 (1H, s), 1.96 (4H, m), 1.56 (4H, m), 0.13 (9H, s); ¹³C NMR δ 133.0 (C), 125.8 (CH), 106.7 (C), 89.0 (C), 60.8 (CH), 46.2 (CH₂), 28.3 (CH₂), 25.2 (CH₂), 22.7 (CH₂), 22.1 (CH₂), -0.8 (CH₃); HRMS calcd. for C₁₃H₂₂SiO 222.1440, found 222.1432.

(3*S*)-6-methyl-1-(trimethylsilyl)hept-6-en-1-yn-3-ol (1d). [93%, ee > 97%, colorless oil]. $[\alpha]_{\text{D}}^{20} = -15.1$ (c 3.1, CHCl₃); ¹H NMR δ 4.74 (2H, s), 4.38 (1H, s), 2.17 (2H, m), 1.94 (1H, s), 1.86 (2H, m), 1.75 (3H, s), 0.18 (9H, s); ¹³C NMR δ 144.7 (C), 110.4 (CH₂), 106.7 (C), 89.2 (C), 62.2 (CH), 35.5 (CH₂), 33.2 (CH₂), 22.3 (CH₃), -0.25 (CH₃); LRMS *m/z* 196 (M⁺, 3), 195 (3), 191 (19), 155 (12), 149 (66), 127 (15), 111 (30), 57 (100); HRMS calcd. for C₁₁H₂₀SiO 196.1283, found 196.1287.

(3*S*)-1,6-bis(trimethylsilyl)hept-6-en-1-yn-3-ol (1e). [96%, ee > 97%, colorless oil]. $[\alpha]_{\text{D}}^{20} = -12.4$ (c 1.6, CHCl₃); ¹H NMR δ 5.59 (1H, s), 5.34 (1H, s), 4.37 (1H, m), 2.29 (3H, m), 1.81 (2H, m), 0.17 (9H, s), 0.09 (9H, s); ¹³C NMR δ 151.1 (C), 124.2 (CH₂), 106.8 (C), 89.1 (C), 62.2 (CH), 36.8 (CH₂), 31.0 (CH₂), -0.25 (CH₃), -1.65 (CH₃); LRMS *m/z* 254 (M⁺, 1), 253 (2), 239 (1), 221 (7), 181 (1), 149 (18), 141 (16), 73 (100); HRMS calcd. for C₁₃H₂₆Si₂O 254.1522, found 254.1520.

⁸ See following section (page S5) for enantiomeric excess determination of alkynols **1**.

⁹ Kurihara, M.; Ishii, K.; Kasahara, Y.; Miyata, N. *Tetrahedron Lett.* **40**, **1999**, 3183-3184.

Determination of the enantiomeric excess (procedure exemplified for **1d**).

[(R)-MTPA-(S)-1d]. DMAP (4 mg, 0.03 mmol) was added to a solution of (R)-trifluoromethyl-methoxyphenyl acetic acid (52 mg, 0.223 mmol), DCC (66 mg, 0.320 mmol) and (S)-**1d** (25 mg, 0.127 mmol) in Et₂O (5 mL). After stirring for 12 h at rt, the mixture was filtered and concentrated. The residue was chromatographed in silica gel (2-5% Et₂O/hexanes) affording 50 mg of [(R)-MTPA-(S)-**1d**] (95% yield, colorless oil). *R_f* = 0.55 (5% Et₂O/hexanes); ¹H NMR δ (ppm): 7.38 (m, 2H), 7.25 (m, 3H), 5.37 (t, *J* = 6.7 Hz, 1H), 4.62 (s, 1H), 4.56 (s, 1H), 3.43 (s, 1H), 2.02 (m, 2H), 1.87 (m, 2H), 1.60 (s, 3H), 0.04 (s, 9H). ¹⁹F NMR δ –72.49 (s, 1CF₃).

[(R)-MTPA-(±)-1d]. The same procedure as described above was followed but using (±)-**1d**. ¹H NMR δ 7.38 (m, 2H), 7.25 (m, 3H), 5.37 (m, 1H), 4.62 (m 1H), 4.56 (s, 0.5H), 4.46 (s, 0.5H), 3.43 (s, 1H), 2.02-1.70 (m, 4H), 1.60 (s, 3 x 0.5H), 1.56 (s, 3 x 0.5H) 0.04 (s, 9H). ¹⁹F NMR δ –72.492 (s, 0.5 CF₃), –72.093 (s, 0.5 CF₃).

[(R)-MTPA-(S)-1e]. [98%, *R_f* = 0.95 (10% Et₂O/hexanes)]; ¹H NMR δ 7.55 (m, 2H), 7.39 (m, 3H), 5.68 (s, 1H), 5.52 (t, *J* = 6.7 Hz, 1H), 5.37 (d, *J* = 1.8 Hz, 1H), 3.58 (s, 3H), 2.28 (m, 2H), 1.96 (m, 2H), 0.18 (s, 9H), 0.10 (s, 9H); ¹⁹F NMR δ –72.474 (s, 1CF₃).

[(R)-MTPA-(±)-1e]. ¹H NMR δ 7.55 (m, 2H), 7.39 (m, 3H), 5.56 (s, 1H), 5.54 (m, 1H), 5.36 (s, 0.5H), 5.34 (s, 0.5H), 3.60 (s, 3 x 0.5H), 3.58 (s, 3 x 0.5H), 2.33 - 1.90 (m, 4H), 0.20 (s, 9 x 0.5H), 0.18 (s, 9 x 0.5H), 0.10 (s, 9 x 0.5H), 0.07 (s, 9 x 0.5H); ¹⁹F NMR δ –72.474 (s, 0.5CF₃), –72.079 (s, 0.5CF₃)

[(R)-MTPA-(S)-1a]. [95%, *R_f* = 0.65 (10% Et₂O/hexanes)]; ¹H NMR δ 7.42-7.23 (m, 10H), 5.71 (t, *J* = 5.9 Hz, 1H), 3.34 (s, 3H), 3.25 (m, 2H), 0.20 (s, 9H); ¹⁹F NMR δ –72.213 (s, 1CF₃).

[(R)-MTPA-(±)-1a]. ¹H NMR δ 7.42-7.23 (m, 10H), 5.71 (m, 1H), 3.55 (s, 3 x 0.5H), 3.34 (s, 3 x 0.5H), 3.25-3.10 (m, 2H), 0.20 (s, 9H); ¹⁹F NMR δ –72.213 (s, 0.5CF₃), –72.170 (s, 0.5CF₃).

[(R)-MTPA-(S)-1b]. [86%, *R_f* = 0.52 (10% Et₂O/hexanes)]; ¹H NMR δ 7.38 (m, 2H), 7.26 (m, 3H), 7.04 (t, *J* = 7.9 Hz, 1H), 6.57 (m, 3H), 5.33 (t, *J* = 7.0 Hz, 1H), 3.62 (s, 3H), 3.40 (s, 3H), 2.57 (m, 2H), 1.99 (m, 2H), 0.00 (s, 9H). ¹⁹F NMR δ –72.453 (s, 1CF₃).

[(R)-MTPA-(±)-1b]. ¹H NMR δ 7.38 (m, 2H), 7.26 (m, 3H), 7.04 (m, 1H), 6.57 (m, 3H), 5.33 (m, 1H), 3.62 (s, 3H), 3.40 (s, 3H), 2.50 (m, 2H), 1.90 (m, 2H), 0.00 (s, 9H). ¹⁹F NMR δ –72.30 (s, 0.5CF₃), –72.453 (s, 0.5CF₃).

General procedure for the synthesis of mixed acetals (exemplified for **2a).**

((Z)-((S)-2-benzyl-dihydro-6-methoxy-2H-pyran-3(4H)-ylidene)methyl)trimethylsilane

(2a): [CpRu(CH₃CN)₃]PF₆ (44 mg, 0.102 mmol) was added to a solution of trimethylsilylalkynol **1a** (200 mg, 1.02 mmol) and allyl ethyl ether (131 mg, 1.53 mmol) in DMF (1 mL). After being stirred at rt for 2 hours the mixture was diluted in MeOH (10 mL) and HCl (10% aq., 0.1 mL) was added dropwise. The reaction was stirred at rt for 5 min, poured into a saturated solution of NaHCO₃, extracted with Et₂O, dried and concentrated. The crude was flash chromatographed (1-5% Et₂O/hexanes) to afford 165 mg of **2a** as a mixture of diastereoisomers [67% yield, 86% based on recovered **1a**, colorless oil] [α_D^{20} = 1.2 (c 0.23 CHCl₃); ¹H NMR δ 7.14 (5H, m), 5.25 (0.68H, s), 5.20 (0.32H, s), 4.73 (1H, m), 4.70-4.55 (0.68+ 0.32H, m), 3.22 (3x 0.32H, s), 3.12 (3x 0.68H, s), 3.15 (1H, dd, *J* = 9.3 and 16.1 Hz), 2.75 (1H, m), 2.49 (0.68H, m), 2.16-1.85 (0.32+ 2H, m), 1.52 (1H, m), 0.01 (9x 0.68H, s), -0.18 (9x 0.32H, s); ¹³C NMR δ 154.4 (C), 153.9 (C), 138.7 (C), 138.2 (C), 129.3 (CH), 129.2 (CH), 128.2 (CH), 128.1 (CH), 126.2 (CH), 126.1 (CH), 124.7 (CH), 124.0 (CH), 98.9 (CH), 96.9 (CH), 77.1 (CH), 75.7 (CH), 55.8 (CH₃), 55.5 (CH₃), 42.1 (CH₂), 39.6 (CH₂), 33.1 (CH₂), 32.8 (CH₂), 29.6 (CH₂), 29.1 (CH₂), 0.18 (CH₃), -0.08 (CH₃); LRMS *m/z* 290 (M⁺, 1), 259 (1), 200 (15), 199 (92), 169 (9), 141 (28), 91 (100); HRMS calcd. for C₁₇H₂₆SiO₂ 290.1702, found 290.1710.

((Z)-((S)-2-(3-methoxyphenethyl)-dihydro-6-methoxy-2H-pyran-3(4H)-ylidene)methyl)trimethylsilane (2b). [65% yield, 88% based on recovered **1b**, colorless oil]. [α_D^{20} = 3.2 (c 0.42 CHCl₃); ¹H NMR δ 7.19 (1H, m), 6.72 (3H, m), 5.20 (1H, s), 4.73 (0.65H, m), 4.64 (0.35H, m), 4.43 (0.65H, m), 4.35 (0.35H, m), 3.74 (3H, s), 3.47 (3x 0.35H, s), 3.43 (3x 0.65H, s), 3.01-2.35 (3H, m), 2.28-1.75 (3H, m), 1.57 (2H, m), -0.02 (9x 0.35H, s), -0.05 (9x 0.65H, s); ¹³C NMR δ 159.7 (C), 156.1 (C), 155.1 (C), 143.7 (C), 143.3 (C), 129.4 (CH), 129.3 (CH), 123.4 (CH), 123.1 (CH), 121.0 (CH), 120.9 (CH), 114.4 (CH), 114.3 (CH), 111.2 (CH), 111.1 (CH), 99.2 (CH), 97.0 (CH), 76.8 (CH), 74.6 (CH), 56.0 (CH₃), 55.9 (CH₃), 55.0 (CH₃), 38.6 (CH₂), 35.2 (CH₂), 33.2 (CH₂), 32.9 (CH₂), 32.6 (CH₂), 32.0 (CH₂), 31.7 (CH₂), 29.1 (CH₂), 0.12 (CH₃), 0.01 (CH₃); LRMS *m/z* 334 (M⁺, 31), 302 (5), 261 (16), 213 (14), 199 (32), 167 (43), 121 (76), 105 (43), 73 (100); HRMS calcd. for C₁₉H₃₀SiO₃ 334.1964, found 334.1964.

((Z)-((S)-2-(cyclohexenylmethyl)dihydro-6-methoxy-2H-pyran-3(4H)-ylidene)methyl)trimethylsilane (2c). [43% yield, 82% based on recovered **1c**, colorless oil]. ¹H NMR δ 5.45 (1H, s), 5.16 (1H, s), 4.62 (2H, m), 3.26 (3H, s), 2.51 (2H, m), 2.11 (2H, m), 1.91 (6H, m), 1.54 (4H, m), 0.04 (9H, s); ¹³C NMR δ 154.6 (C), 134.9 (C), 124.3 (CH), 124.0 (CH), 96.3 (CH), 74.0 (CH), 56.4 (CH₃), 42.0 (CH₂), 33.4 (CH₂), 32.4 (CH₂), 27.6 (CH₂), 25.8 (CH₂), 23.1 (CH₂), 22.9

(CH₂), 0.52 (CH₃); LRMS (CI, *m/z*): 295 (M⁺ +1, 11), 294 (5), 279 (40), 264, (20), 199 (100); HRMS calcd. for C₁₇H₃₁O₂Si 295.2093, found 295.2097.

((Z)-((S)-dihydro-6-methoxy-2-(3-methylbut-3-enyl)-2H-pyran-3(4H)-ylidene)methyl)trimethylsilane (2d). [58% yield, 77% based on recovered **1d**, colorless oil]. ¹H NMR δ 5.26 (1H, s), 4.72 (2+0.65H, m), 4.65 (0.35H, t, *J* = 3.6 Hz), 4.47 (0.65H, m), 4.33 (0.35H, m), 3.44 (3H, s), 2.61 (1H, m), 2.30-1.85 (6H, m), 1.76 (3x 0.65H, s), 1.74 (3x 0.35H, s), 1.61 (1H, m), 0.11 (9H, s); ¹³C NMR δ 156.3 (C), 155.5 (C), 145.6 (C), 145.3 (C), 123.1 (CH), 122.9 (CH), 110.2 (CH₂), 109.9 (CH₂), 99.0 (CH), 96.9 (CH), 77.4 (CH), 75.3 (CH), 55.8 (CH₃), 35.1 (CH₂), 33.9 (CH₂), 32.8 (CH₂), 32.7 (CH₂), 31.9 (CH₂), 31.8 (CH₂), 29.1 (CH₂), 22.7 (CH₃), 0.25 (CH₃), 0.17 (CH₃); LRMS *m/z* 253 (M⁺ - CH₃, 1), 237 (M⁺ - OCH₃, 4), 221 (10), 147 (37), 121 (14), 79 (100), 63 (12); HRMS calcd. for C₁₄H₂₅SiO 237.1675, found 237.1679.

((Z)-((S)-dihydro-6-methoxy-2-(3-(trimethylsilyl)but-3-enyl)-2H-pyran-3(4H)-ylidene)methyl)trimethylsilane (2e). [72% yield, 85% based on recovered **1e**, colorless oil]. ¹H NMR δ 5.60 (1H, s), 5.3 (2H, m), 4.70 (1H, m), 4.48 (0.8H, m), 4.37 (0.2H, m), 3.45 (3H, s), 2.85-2.36 (2H, m), 2.28-1.85 (4H, m), 1.77-1.43 (2H, m), 0.11 (18H, s); ¹³C NMR δ 156.0 (C), 155.0 (C), 152.0 (C), 151.9 (C), 124.1 (CH₂), 123.6 (CH₂), 123.4 (CH), 123.1 (CH), 99.0 (CH), 96.8 (CH), 77.6 (CH), 75.9 (CH), 55.9 (CH₃), 55.7 (CH₃), 35.9 (CH₂), 33.0 (CH₂), 32.9 (CH₂), 32.8 (CH₂), 31.9 (CH₂), 31.7 (CH₂), 29.0 (CH₂), 0.35 (CH₃), 0.25 (CH₃), -1.53 (CH₃), -1.57 (CH₃); LRMS *m/z* 327 (M⁺ + 1, 1), 311 (2), 295 (4), 279 (6), 221 (5), 203 (12), 133 (16), 79 (100), 73 (30); HRMS calcd. for C₁₇H₃₅Si₂O₂ 327.2176, found 327.2180.

General procedure for the Lewis-acid promoted cyclization of the mixed acetals (exemplified for **3a).**

(1S,9S)-10-[(Z)-1-(trimethylsilyl)methyliden]-13-oxatricycle[7.3.1.0^{2,7}]trideca-2(7),3,5-triene (3a**)** SnCl₄ (1.87 mmol, 1.0 M in CH₂Cl₂) was added dropwise over 20 min into a solution of **2a** (340 mg, 1.25 mmol) in CH₂Cl₂ (10 mL) cooled at -78 °C. After stirring at that temperature

for 20 min the reaction was quenched by adding a saturated solution of NaHCO₃ and the resulting mixture was warmed to room temperature. The layers were separated and the aqueous fraction was extracted with CH₂Cl₂. The combined organic phases were washed with brine, dried and concentrated. The crude residue was flash chromatographed (0.1-0.5% Et₂O/hexanes) to afford 280 mg of **3a** [92% yield, colorless oil]. [α]_D²⁰ = 6.2 (c 0.22 CHCl₃); ¹H NMR δ 7.15 (3H, m), 7.02 (1H, m), 5.26 (1H, s), 4.95 (2H, m), 3.47 (1H, dd, *J* = 17.2 and 7.8 Hz), 2.62 (1H, d, *J* = 17.2 Hz), 2.29-2.09 (3H, m), 1.76 (1H, d, *J* = 7.2 Hz), 0.19 (9H, s); ¹³C NMR δ 156.5 (C), 137.0 (C), 133.6 (C), 128.1 (CH), 126.6 (CH), 125.7 (CH), 124.8 (CH), 123.3 (CH), 71.8 (CH), 69.9 (CH), 34.2 (CH₂), 32.9 (CH₂), 29.6 (CH₂), 0.35 (CH₃); LRMS *m/z* 258 (M⁺, 75), 257 (12), 244 (12), 243 (58), 215 (12), 167 (62), 153 (42), 141 (39), 73 (100); HRMS calcd. for C₁₆H₂₂SiO 258.1440, found 258.1440. Anal. Calcd. for C₁₆H₂₂SiO : C, 74.36; H, 8.58, found C, 74.44; H, 8.68. The enantiomeric excess was determined by HPLC analysis by comparison with racemic samples (Chiralcel AS, 0-0.5% iPrOH/hexanes). Retention times: 7.38 and 9.15 min.

(1S,10S)-5-Methoxy-11-[(Z)-1-(trimethylsilyl)methylidene]-14-oxatricyclo[8.3.1.0^{2,7}]tetra deca-2(7),3,5-triene (3b). [88% yield, colorless oil]. [α]_D²⁰ = 8.2 (c 0.42 CHCl₃); ¹H NMR δ 6.95 (1H, d, *J* = 8.2 Hz), 6.72 (2H, m), 5.26 (1H, s), 5.13 (1H, t, *J* = 5.9 Hz), 4.68 (1H, t, *J* = 8.2 Hz), 3.80 (3H, s), 3.32 (1H, t, *J* = 13.0 Hz), 2.69 (1H, dd, *J* = 6.7 Hz and 15.3 Hz), 2.47-2.18 (4H, m), 1.96 (2H, m), 0.15 (9H, s); ¹³C NMR δ 158.1 (C), 156.8 (C), 142.6 (C), 135.3 (C), 128.1 (CH), 122.4 (CH), 116.1 (CH), 111.5 (CH), 75.5 (CH), 74.8 (CH), 55.6 (CH₃), 36.6 (CH₂), 33.7 (CH₂), 32.0 (CH₂), 30.2 (CH₂), 0.74 (CH₃); LRMS *m/z* 287 (M⁺-CH₃, 12), 269 (4), 211 (4), 197 (6), 191 (14), 186 (14), 169 (11), 147 (21), 127 (16), 85 (45), 57 (100); HRMS calcd. for C₁₇H₂₃SiO₂ 287.1467, found 287.1473. Anal. Calcd. for C₁₇H₂₃SiO₂ : C, 71.47; H, 8.66, found: C, 71.67; H, 8.76.

(1S,9S)-10-[(Z)-1-(trimethylsilyl)methylidene]-13-oxatricyclo[7.3.1.0^{2,7}]tridec-2(7)-ene (3c). [α]_D²⁰ = 7.3 (c 0.54 CHCl₃); [76% yield, colorless oil]. ¹H NMR δ 5.19 (1H, s), 4.72 (1H, d), 4.03 (1H, s), 2.61 (2H, m), 2.11 (1H, m), 1.84 (4H, m), 1.65 (5H, m), 1.23 (2H, m), 0.14 (9H, s); ¹³C NMR 157.2 (C), 128.4 (C), 127.1 (C), 122.0 (CH), 71.6 (CH), 69.9 (CH), 34.8 (CH₂), 29.9 (CH₂), 29.3 (CH₂), 29.1 (CH₂), 28.9 (CH₂), 26.0 (CH₂), 22.5 (CH₂), 0.10 (CH₂); LRMS *m/z* 263 (M⁺ + 1, 28), 262 (23), 261 (29), 248 (21), 247 (100), 245 (37), 171 (76); HRMS calcd. for C₁₆H₂₇SiO 263.1831, found 262.1833. The enantiomeric excess was determined by HPLC analysis by comparison with racemic samples (Chiralcel AS, 0-0.5% iPrOH/hexanes). Retention times: 14.1 and 17.5 min.

((Z)-((1S,6S)-3-chloro-3-methyl-10-oxa-bicyclo[4.3.1]decan-7-ylidene)methyl)trimethyl silane (3d). [82% yield, colorless oil]. [α]_D²⁰ = 3.4 (c 0.22 CHCl₃); ¹H NMR δ 5.24 (1H, s), 4.34 (1+0.2H, m), 4.02 (0.8H, m), 2.72 (2H, m), 2.35-1.75 (6H, m), 1.78 (3x 0.8H), 1.62 (3x 0.2H), 1.58 (1H, m), 0.08 (9H, s); ¹³C NMR δ 154.8 (C), 154.4 (C), 123.9 (CH), 123.3 (CH), 74.5 (CH), 73.4 (C), 70.5 (C), 67.9 (CH), 67.6 (CH), 46.0 (CH₂), 44.5 (CH₂), 40.2 (CH₂), 40.1 (CH₂), 36.2 (CH₃), 33.3 (CH₃), 31.0 (CH₂), 30.2 (CH₂), 30.1 (CH₂), 29.3 (CH₂), 28.6 (CH₂), 28.4 (CH₂),

0.30 (CH₃); LRMS m/z 273 ($M^+ + 1$, 1), 257 ($M^+ - \text{CH}_3$, 2), 237 (10), 221 (42), 203 (9), 183 (4), 149 (18), 147 (100), 119 (23); HRMS calcd for C₁₄H₂₆SiClO 273.1441, found 273.1441.

(Z,1S,6S)-3-chloro-3-(trimethylsilyl)-7-((trimethylsilyl)methylene)-10-oxabicyclo[4.3.1]decane (3e). [86% yield, colorless oil]. ¹H NMR δ 5.22 (1H, s), 4.39 (1H, t, J = 9.6 Hz), 4.06 (1H, t, J = 6.4 Hz), 2.72 (1H, m), 2.49-1.83 (9H, m), 0.18 (9H, s), 0.10 (9H, s); ¹³C NMR δ 155.9 (C), 122.3 (CH), 74.1 (CH), 70.1 (CH), 66.8 (C), 42.0 (CH₂), 32.9 (CH₂), 30.0 (CH₂), 29.3 (CH₂), 28.4 (CH₂), 0.30 (CH₃), -3.57 (CH₃); LRMS m/z 315 ($M^+ - \text{CH}_3$, 12), 297 (3), 295 (28), 225 (18), 207 (18), 205 (14), 133 (27), 129 (38), 73 (33), 17 (100); HRMS calcd. for C₁₅H₂₈Si₂OCl 315.1367, found 315.1361.

General procedure for the reductive opening of compounds 3a and 3b (exemplified for the synthesis of 4b).

(5S,E)-6,7,10,11-tetrahydro-2-methoxy-8-((trimethylsilyl)methyl)-5H-benzo[9]annulen-5-ol (4b). A solution of freshly prepared sodium naphthalenide (0.5M in THF) was added dropwise over a solution of **3b** (50 mg, 0.165 mmol) in THF (17 mL, 0.01M) at 23 °C until the green color persisted. After stirring at that temperature for 15 min the reaction was quenched by adding 5 mL of a saturated solution of NaHCO₃ and the resulting mixture was allowed to warm to room temperature. The layers were separated and the aqueous fraction was extracted with Et₂O. The organic phases were dried, concentrated and flash chromatographed (10% EtOAc/hexanes) to afford 42 mg of **4b** as a white solid [80%, R_f 0.15 (10% EtOAc/hexanes)]. [α]_D²⁰ = -26.6 (c 1.22 CHCl₃); ¹H NMR δ 7.50 (1H, d, J = 8.5 Hz), 6.83 (1H, dd, J = 2.5 Hz and 8.4 Hz), 6.62 (1H, s), 5.34 (1H, t, J = 8.5 Hz), 4.82 (1H, m), 3.81 (3H, s), 2.73-2.52 (2H, m), 2.47-1.91 (3H, m), 1.72 (4H, m), 1.54 (2H, ABq, J = 13.8 Hz, ΔAB = 28.4 Hz), 0.01 (9H, s); ¹³C NMR δ 158.6 (C), 143.4 (C), 138.2 (C), 135.1 (C), 126.9 (CH), 122.6 (CH), 114.5 (CH), 112.2 (CH), 68.8 (CH), 55.2 (CH₃), 38.3 (CH₂), 34.4 (CH₂), 28.7 (CH₂), 28.3 (CH₂), 25.8 (CH₂), -1.30 (CH₃); LRMS m/z 289 ($M^+ - \text{CH}_3$, 11), 287 (7), 271 (15), 243 (16), 215 (10), 213 (26), 149 (14); HRMS calcd. for C₁₈H₂₈SiO₂ - CH₃ 289.1624, found 289.1645. Anal. Calcd. for C₁₈H₂₈SiO₂: C, 71.00; H, 9.27. Found: C, 71.11; H, 9.37. The enantiomeric excess was determined by HPLC analysis by comparison with racemic samples (Chiralcel OD-H, 0-10% iPrOH/hexanes). Retention times: 29.2 and 31.8 min.

(5*S,E*)-5,6,7,10-tetrahydro-8-((trimethylsilyl)methyl)benzo[8]annulen-5-ol (4a).¹⁰ [52% yield]. $[\alpha]_D^{20} = -32.5$ (c 0.9 CHCl₃); ¹H NMR δ 7.55 (1H, m), 7.18 (3H, m), 5.22 (2H, m), 3.69 (1H, m), 3.28 (1H, m), 2.21 (2H, m), 1.95 (1H, s), 1.79 (1H, m), 1.43 (2H, ABq, $J = 13.3$ Hz, $\Delta AB = 36.2$ Hz), 1.40 (1H, m), -0.03 (9H, s); ¹³C NMR δ 143.0 (C), 139.6 (C), 139.1 (C), 128.8 (CH), 127.3 (CH), 126.9 (CH), 126.4 (CH), 123.5 (CH), 69.2 (CH), 45.4 (CH₂), 41.3 (CH₂), 30.5 (CH₂), 23.4 (CH₂), -1.30 (CH₃); LRMS m/z ($M^+ - 1$, 10), 245 ($M^+ - CH_3$, 9), 243 (15), 229 (5), 197 (14), 171 (94), 169 (70), 137 (30); HRMS calcd. for C₁₆H₂₃SiO 259.1518, found 259.1530.

Ring opening of 3a under basic conditions

(5*S,8Z,9Z*)-5,6,7,8-tetrahydro-8-((trimethylsilyl)methylene)benzo[8]annulen-5-ol $n\text{-BuLi}$ (0.13 mL, 0.30 mmol, 2.25 M en hexanes) was slowly added dropwise to a solution of **3a** (50 mg, 0.20 mmol) in THF (5 mL) cooled at 0 °C. After stirring for 5 min at this temperature, the reaction mixture was allowed to reach rt, poured over brine and extracted with Et₂O (3 x 2 mL). The organic phases were dried, filtered, concentrated and chromatographed (1-4% EtOAc/hexanes) to afford **8a** as colorless oil (49 mg, 98%). $R_f = 0.70$ (3% EtOAc/hexanes); $[\alpha]_D^{20} = -12.5$ (c 0.6 CHCl₃); ¹H NMR δ 7.60 (d, $J = 8.5$ Hz, 1H), 7.28 (m, 3H), 6.52 (d, $J = 13.2$ Hz, 1H), 6.35 (d, $J = 12.6$ Hz), 5.61 (s, 1H), 5.21 (dd, $J = 11.3$ and 4.3 Hz, 1H), 2.17 (m, 2H), 1.71 (s broad, 1H), 1.69 (m, 2H); ¹³C NMR δ 154.6 (C), 142.1 (C), 136.6 (CH), 135.6 (C), 133.5 (CH), 129.3 (CH), 127.9 (CH), 127.1 (CH), 126.6 (CH), 123.8 (CH), 69.7 (CH), 41.0 (CH₂), 34.2 (CH₂), 0.24 (CH₃); LRMS (CI, m/z): 259 ($M^+ + 1$, 4), 241 (2), 219 (6), 165 (8), 125 (25), 79 (87), 41 (100); HRMS calcd. for C₁₆H₂₃SiO 259.1518, found 259.1514.

¹⁰ In addition to **4a**, we also obtained a 43 % yield of the isomer **9a**. See: ref 6.