Supporting Information for
Practical Carbonyl-Ene Reactions of α -Methylstyrenes with Paraformaldehyde
Promoted by a Combined System of Boron Trifluoride and Molecular Sieves 4A

Takahiro Okachi, Katsuhiko Fujimoto, and Makoto Onaka*

Department of Chemistry, Graduate School of Arts and Sciences, The University of Tokyo, Komaba, Meguro, Tokyo 153-8902, Japan, and Process Development Laboratories, Sankyo Co., Ltd., Hiratsuka, Kanagawa, 254-8560, Japan.

General Experimental:

Infrared (IR) spectra were recorded on a JASCO FT/IR-550 Fourier transform infrared spectrometer. ^1H NMR spectra were recorded on a JNM-GSX 270 (270 MHz). ^{13}C NMR spectra were recorded on a JNM-GSX 270 (68 MHz). High-resolution mass spectra (HRMS) were recorded on a JEOL JMS-700V GCmate. Analytical TLC was performed on Merck DC-Platten Kieselgel 60 F254 plates. Silica gel column chromatography was carried out with Merck Kieselgel (60 mesh). MS 4A, NaX, NaY, and Na-mordenite were purchased in a powder form, and activated at 100 °C under vacuum for 16 hours. Dichloromethane was dried over MS 4A. α -Methylstyrenes (**1a-c** and **1e**) were prepared from the corresponding acetophenones and methyltriphenylphosphonium bromide by the Wittig reaction. All reactions were carried out under nitrogen atmosphere.

Typical experimental procedure for the carbonyl-ene reaction using a combined system of boron trifluoride and molecular sieves 4A:

A mixture of $\text{BF}_3 \cdot \text{OEt}_2$ (31.7 mmol) and molecular sieves 4A (5.0 g) in CH_2Cl_2 (50 ml) was stirred at room temperature for 1 hour under nitrogen, and then was allowed to cool to -10°C . A solution of 3,4-dichloro- α -methylstyrene (26.7 mmol) in CH_2Cl_2 (50 ml) and paraformaldehyde (26.7 mmol) were added successively at -10°C , and the reaction mixture was stirred at -10°C for 3-6 hours. To the reaction mixture was added saturated aq. NaHCO_3 to quench the reaction, and the mixture was filtered through a Celite pad. The organic layer was separated, washed with saturated aq. NaHCO_3 and brine, and dried over Na_2SO_4 . After filtration and evaporation under reduced pressure, the crude products were chromatographed on silica gel to give the desired adduct **2a** in 64% yield.

3-(3,4-Dichlorophenyl)-3-buten-1-ol (2a):

colorless oil; Bp. 135-140 °C (bath temperature)/ 1.5 mmHg. IR (CCl₄) 3628, 3393, 3087, 2951, 2881, 1626, 1549, 1473, 1384, 682 cm⁻¹. ¹H NMR (CDCl₃): δ 1.55 (s, 1H), 2.72 (dt, *J* = 1.1 and 6.5 Hz, 2H), 3.73 (t, *J* = 6.5 Hz, 2H), 5.21 (d, *J* = 1.1 Hz, 1H), 5.41 (d, *J* = 1.1 Hz, 1H), 7.24 (dd, *J* = 2.2 and 8.6 Hz, 1H), 7.40 (d, *J* = 8.6 Hz, 1H), 7.49 (d, *J* = 2.4 Hz, 1H). ¹³C NMR (CDCl₃): δ 38.2, 60.8, 115.9, 125.3, 128.0, 130.2, 131.4, 132.4, 140.5, 142.7. HRMS: Calcd for C₁₀H₁₀OCl₂ (M⁺) 216.0109. found 216.0103. Anal. Calcd for C₁₀H₁₀OCl₂: C, 55.33; H, 4.64; Cl, 32.66. found: C, 55.18; H, 4.62; Cl, 32.74.

3-(3,4-Difluorophenyl)-3-buten-1-ol (2b):

colorless oil; Bp. 100-110 °C (bath temperature)/ 1.5 mmHg. IR (CCl₄) 3628, 3502, 3087, 2948, 2881, 1600, 1518, 1422 cm⁻¹. ¹H NMR (CDCl₃) δ 1.55 (s, 1H), 2.74 (dt, *J* = 1.1 and 6.5 Hz, 2H), 3.73 (t, *J* = 6.5 Hz, 2H), 5.19 (s, 1H), 5.37 (s, 1H), 7.10 (m, 3H). ¹³C NMR (CDCl₃) δ 38.4, 60.8, 114.8, 115.0, 115.2, 115.3, 115.7, 122.0, 137.5, 142.9. HRMS: Calcd for C₁₀H₁₀OF₂ (M⁺) 184.0700. found 184.0701. Anal: Calcd for C₁₀H₁₀OF₂: C, 65.21; H, 5.47; F, 20.63. found: C, 65.28; H, 5.44; F, 20.67.

3-(4-Fluorophenyl)-3-buten-1-ol (2c):

colorless oil; Bp. 135-140 °C (bath temperature)/ 1.5 mmHg. IR (CCl₄) 3628, 3445, 3086, 3054, 2947, 2880, 1604, 1509, 1399 cm⁻¹. ¹H NMR (CDCl₃) δ 1.56 (s, 1H), 2.75 (dt, *J* = 1.1 and 6.5 Hz, 2H), 3.72 (t, *J* = 6.5 Hz, 2H), 5.13 (d, *J* = 1.1 Hz, 1H), 5.35 (d, *J* = 1.1 Hz, 1H), 7.02 (t, *J* = 8.9 Hz, 2H), 7.38 (dd, *J* = 5.4 and 8.9 Hz, 2H). ¹³C NMR (CDCl₃) δ 38.7, 60.9, 114.4, 115.0, 115.3, 127.4, 136.4, 143.7. HRMS: Calcd for C₁₀H₁₁OF (M⁺) 166.0794. found 166.0786. Anal: Calcd for C₁₀H₁₁OF: C, 72.27; H, 6.67; F, 11.43. found: C, 72.03; H, 6.72; F, 11.65.

3-Phenyl-3-buten-1-ol (2d):

colorless oil; Bp. 110-120 °C (bath temperature)/ 1.5 mmHg. IR (CCl₄) 3628, 3365, 3084, 3059, 3026, 2948, 2880, 1627, 1574, 1495, 1445, 705 cm⁻¹. ¹H NMR (CDCl₃) δ 1.47 (s, 1H), 2.79 (dt, *J* = 1.4 and 6.5 Hz, 2H), 3.73 (t, *J* = 6.5 Hz, 2H), 5.16 (d, *J* = 1.4 Hz, 1H), 5.41 (d, *J* = 1.4 Hz, 1H), 7.25-7.44 (m, 5H). ¹³C NMR (CDCl₃) δ 38.6, 61.0, 114.5, 126.0, 127.6, 128.3, 140.2, 144.7. HRMS: Calcd for C₁₀H₁₂O (M⁺) 148.0888. found 148.0887. Anal: Calcd for C₁₀H₁₂O: C, 81.04; H, 8.16. found: C, 80.91; H, 8.36.