Supporting Information

Practical synthesis of azobenzenophanes

*Hong-Min Kang, Hee-Yeon Kim, Jin-Woo Jung and Cheon-Gyu Cho**

Department of Chemistry, Hanyang University, Seoul 131-791, Korea

Table of Contents

General Experimental Methods	S2
Copies of $^1\text{H}/^{13}\text{C}$ NMR spectra of 12	S3 – S4
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 13	S5 – S6
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 2a	S7 – S8
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 15	S9 – S10
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 16	S11 – S12
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 17	S13 – S14
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 19	S15 – S16
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 20	S17 – S18
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 22	S19 – S20
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 2c	S21 – S22
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 23	S23 – S24
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 24	S25 – S26
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 25	S27 – S28
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 26	S29 – S30
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of bis(3-iodophenyl)methanol	S31 – S32
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 28	S33 – S34
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 29	S35 – S36
Copy of $^1\text{H}/^{13}\text{C}$ NMR spectra of 30	S37 – S38

General Methods. ^1H NMR spectra were recorded at 400 MHz and ^{13}C NMR spectra at 100 MHz, with either TMS ($\delta = 0$) or the signal for residual CHCl_3 in the CDCl_3 solvent ($\delta = 7.24$) as internal standards. J values are reported in Hz. High resolution mass spectra were measured by using FAB method. Flash column chromatography was performed with 230-400 mesh grade silica-gel. All solvents used were purified according to standard procedures.