Supporting Information

Practical Synthesis of Sultams via Sulfonamide Dianion Alkylation: Application in the synthesis of Chiral Sultams

Jaemoon Lee,* Yong-Li Zhong, Robert A. Reamer, and David Askin

Department of Process Research, Merck Research Laboratories, P.O. Box 2000, Rahway, New Jersey 07065jaemoon.lee@merck.com**General Experimental Information**

All reactions were carried out under nitrogen. Flash chromatography was carried out with EM science Silica gel 60 (neutral, 230-400 mesh). Thin-layer chromatography (TLC) was performed on EM science Silica gel (0.25mm, precoated on glass).

Visualization of compounds on TLC was accomplished by UV illumination or by staining with a solution prepared from 25g of ammonium molybdate and 1 g of ceric sulfate in 500 mL of 10% aq. sulfuric acid, followed by heating.

^1H NMR and ^{13}C NMR spectra were recorded on a Bruker 400 MHz with chemical shifts reported in ppm relative to the residual deuterated solvent. Optical rotations were measured using a Perkin-Elmer 241 Polarimeter at 23 °C using sodium lamp (D line). HPLC analysis was performed on a Hewlett Packard 1100 MSD instrument.

General procedures for the conversion of ω -halo alkylamine to sultam:

To a stirred slurry of haloalkylamine-HX salt (108 mmol) either in THF (X=Br, 300 mL) or acetonitrile (X=Cl) were added triethylamine (220mmol) and methanesulfonyl chloride (108 mmol) simultaneously via dropping funnels over 30 min (keeping the internal temp 5~10 °C). After 1 h at this temperature, the reaction was warmed to room temperature and stirred for an additional 1.5 h.

The triethylamine-HX salt was removed by filtration and the cake was washed with THF (40 mL). The THF solution of crude bromosulfonamide was used for next step without further purification.

The solution of halosulfonamide (106 mmol) in THF (340 mL) was placed in a 3-neck flask and cooled to -30°C. After addition of diisopropylamine (26.5 mmol) and 1,10-phenanthroline (30 mg), a solution *n*-BuLi in hexane (240 mmol, 1.6 M) was slowly added via a dropping funnel over period of 30 min maintaining the internal temperature range of -30 °C to -10 °C. During this time, a brown color appeared which indicated that 1.25 equiv of *n*-BuLi had been added. Then, the resulting solution was slowly warmed to 0 °C over 2 h. After 2h at 0 °C, the reaction was quenched by addition of 2N HCl (pH adjusted to ~5). After addition of 20% brine (75 mL), the phases were separated and the aq. layer was back-extracted with 100 mL of ethyl acetate. Concentration of the combined organic layers followed by crystallization gave the desired product.

1,2-thiazinane 1,1-dioxide (1)

Conditions: 3-bromopropylamine-HBr (23.6g, 108 mmol), methanesulfonyl chloride (8.4 ml, 108 mmol), triethylamine (30.8 mL, 220 mmol) in THF (300mL); diisopropylamine (3.7 mL, 26.5 mmol), *n*-BuLi in hexane (163 mL, 240 mmol, 1.6 M). reaction temperature ;~ -30 °C to -10 °C ; then 0 °C for 2h. Crystallization (ethyl acetate-heptane , 1:1) of the crude reaction mixture afforded the desired product, **1** (9.4g, 72%) as a slightly yellow solid, mp 112~113 °C (lit.¹ 113 °C). ¹H NMR (CDCl₃, 400 MHz) δ 4.62 (br s, 1H), 3.41~3.37 (m, 1H), 3.10~3.07 (m, 2H), 2.22~2.16 (m, 2H), 1.64~1.58 (m, 2H); ; ¹³C NMR (CDCl₃, 100 MHz) δ 50.2, 45.4, 24.1, 23.7 ppm.

6-methyl-1,2-thiazinane 1,1-dioxide (6)

Conditions: 3-Chloropropylamine-HCl (6.3g, 49 mmol), ethanesulfonyl chloride (5.12 ml, 53.9 mmol), triethylamine (15.0 mL, 107 mmol) in acetonitrile (80 mL); diisopropylamine (1.55 mL, 11 mmol), *n*-BuLi in hexane (63 mL, 100.8 mmol, 1.6 M). THF (130 mL) reaction temperature ;~ -30 °C to -10 °C ; then 0 °C for 2h. Filtration of the crude reaction mixture over small plug of silica gel (ethyl acetate-hexane-methylene chloride , 3:6:1) afforded the desired product, **6** (5.4g, 72%) as a slightly yellow oil. ¹H NMR (CDCl₃, 400 MHz) δ 4.30 (br s, 1H), 3.40~3.33 (m, 2H), 3.08~3.02 (m, 2H), 2.08~2.14 (m, 1H), 1.88~1.98 (m, 1H), 1.72~1.77 (m, 1H), 1.57~1.60 (m, 1H), 1.39 (d, *J* = 6.75Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 56.1, 45.4, 31.6, 25.2, 14.0 ppm.

6-phenyl-1,2-thiazinane 1,1-dioxide (8)

Conditions: 3-Bromopropylamine-HBr (6.1g, 27.8mmol), α-toluenesulfonyl chloride (5.5 g, 29 mmol), triethylamine (8.2 mL, 58.38 mmol) in THF (100 mL); diisopropylamine (0.97 mL, 6.9 mmol), *n*-BuLi in hexane (63 mL, 39.2 mmol, 1.6 M). THF (130 mL) reaction temperature ;~ -40 °C to -10 °C ; then 0 °C for 2h. Filtration of the crude reaction mixture over small plug of silica gel (ethyl acetate-hexane-methylene chloride, 3:6:1) afforded the desired product **8** as a yellow solid (4.4g, 75%). ¹H NMR (CDCl₃, 400 MHz) δ 7.45-7.35 (m, 5H), 4.36 (br s, 1H), 4.11 (dd, 1H, *J* = 12.8, 3.4Hz), 3.65~3.55 (m, 1H), 3.45~3.38 (m, 1H), 2.68~2.57 (m, 1H), 2.33~2.27 (m, 1H), 1.98~1.91 (m, 1H), 1.82~1.70; ¹³C NMR (CDCl₃, 100 MHz) δ 129.3, 128.9, 128.1, 67.1, 45.4, 30.5, 26.8 ppm

isothiazolidine 1,1-dioxide (11)

Conditions: 2-Chloroethylamine-HCl (12.2g, 106 mmol), methanesulfonyl chloride (8.6, 111 mmol), triethylamine (31.2 mL, 223 mmol) in acetonitrile (150 mL); diisopropylamine (3.5 mL, 25 mmol), *n*-BuLi in hexane (135 mL, 216 mmol, 1.6 M). THF (330 mL), reaction temperature; ~ -60 °C to -50 °C; then warmed to 0 °C over a period of 2h. Filtration of the crude reaction mixture over small plug of silica gel (ethyl acetate-hexane-methylene chloride, 3:6:1) afforded the desired product **11** (7.6g, 60%) as an oil.²

¹H NMR (CDCl₃, 400 MHz) δ 4.27 (br s, 1H), 3.43 (m, 2H), 3.09 (t, 2H, *J* = 7.54 Hz), 2.46 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 46.5, 42.2, 23.9 ppm.

General procedures for the conversion of β-amino alcohol to 1,3-sultam:

Typical procedure; To a stirred solution of the amino alcohol (36.6 mmol) and triethylamine (75 mmol) in THF were added methanesulfonyl chloride (75 mmol) at 0°C. After aging for 4h at room temperature, 30 ml of MTBE was added. After filtration of the triethylamine-HCl salt, the filtrate was concentrated to afford bis-mesylate.

The resulting presumed bis-mesylate was treated with NaCl (55 mmol) in DMF (45 mL) and warmed to 80 °C. After aging for 16h at this temperature, the reaction was cooled to room temperature and diluted with water (30 mL) and EtOAc-hexanes (50 mL, 2:1). After phase separation, the organic layer was washed sequentially with 0.5N HCl and brine. The solution was dried over Na₂SO₄, filtered and concentrates under reduced pressure. The crude material was eluted through a small plug of silica gel (hex/EtOAc, 2:1) to afford the chloromethanesulfonate.

The solution of chlorosulfonamide (34 mmol) in THF (100 mL) was placed in 500 mL RBF and cooled to -60°C under nitrogen atmosphere. After addition of diisopropylamine (8.3 mmol) and 1,10-phenanthroline (15 mg), a solution *n*-BuLi in hexane (73 mmol, 1.6 M) was slowly added via a dropping funnel maintaining the internal temperature below -50 °C over a period of 30 min. Then, the resulting solution was slowly warmed to 0 °C over 4 h. After 2h at 0 °C, the reaction was quenched by addition of 2N HCl (pH adjusted to ~5). After addition of 20% brine (30 mL), the phases were separated. The aq. layer was back-extracted with ethyl acetate (40 mL). Concentration of the combined organic layers followed by flash or filtering silica gel chromatography gave the desired product.

***N*-[(1*R*)-1-(chloromethyl)propyl]methanesulfonamide (16)**

$[\alpha]_D^{23} +16.8^\circ$ (c 0.037, EtOH); NMR (CDCl₃, 400 MHz) δ 4.50 (br s, 1H), 3.74 (dd, 1H, $J = 11.20, 4.04$), 3.66~3.58 (m, 2H), 3.04 (s, 3H), 1.68 (m, 2H), 1.01 (t, 3H, $J = 7.40$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 56.1, 48.1, 42.1, 26.1, 10.3 ppm.

(3*R*)-3-ethylisothiazolidine 1,1-dioxide (17)

$[\alpha]_D^{23} +13.3^\circ$ (c 0.037, EtOH); ¹H NMR (CDCl₃, 400 MHz) δ 4.30 (br s, 1H), 3.54~3.50 (m, 1H), 3.24~3.14 (m, 1H), 3.04~3.04 (m, 1H), 2.54~2.46 (m, 1H), 2.12~2.02 (m, 1H), 1.70~1.57 (m, 2H), 0.99 (t, 3H, $J = 7.4$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 56.6, 48.1, 29.4, 28.9, 10.4 ppm.

***N*-[(1*R*)-1-benzyl-2-chloroethyl]methanesulfonamide (19)**

$[\alpha]_D^{23} -33.9^\circ$ (c 0.038, EtOH); ¹H NMR (CDCl₃, 400 MHz) δ 7.37-7.25 (m, 5H), 4.84 (br d, $J = 8.8$ Hz, 1H), 3.88 (m, 1H), 3.70 (dd, $J = 11.8, 4.8$ Hz, 1H), 3.63 (dd, $J = 11.8, 4.8$ Hz, 1H), 3.02 (dd, $J = 13.8, 6.4$ Hz, 1H), 2.86 (dd, $J = 13.8, 6.4$ Hz, 1H), 2.58 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 136.8, 129.6, 129.0, 127.4, 56.4, 48.1, 42.4, 39.2 ppm.

(3*R*)-3-benzylisothiazolidine 1,1-dioxide (20)

$[\alpha]_D^{23} -22.3^\circ$ (c 0.031, EtOH)³; ¹H NMR (CDCl₃, 400 MHz) δ 7.40-7.15 (m, 5H), 4.62 (br s, 1H), 3.90~3.80 (m, 1H), 3.26~3.20 (m, 1H), 3.16~3.06 (m, 1H), 3.00~2.84 (m, 2H), 2.53~2.45 (m, 1H), 2.21~2.15 (m, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 129.1, 128.9, 128.7, 127.2, 56.11, 47.9, 42.2, 28.9 ppm.

***N*-[(1*R*)-2-chloro-1-phenylethyl]methanesulfonamide (22)**

$[\alpha]_D^{23}$ -49.2° (c 0.031, EtOH); ^1H NMR (CDCl_3 , 400 MHz) δ 7.42-7.34 (m, 5H), 5.30 (br d, J = 6.8 Hz, 1H), 4.81 (m, 1H), 3.88 (dd, J = 11.4, 7.2 Hz, 1H), 3.77 (m, J = 11.4, 7.2 Hz, 1H), 2.79 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 137.8, 129.1, 128.8, 126.9, 58.7, 48.3, 42.0 ppm.

(3*S*)-3-phenylisothiazolidine 1,1-dioxide (23)

$[\alpha]_D^{23}$ -50.5° (c 0.033, EtOH)⁴; ^1H NMR (CDCl_3 , 400 MHz) δ 7.42-7.30 (m, 5H), 4.76-4.71 (m, 1H), 4.62 (br s, 1H), 3.40-3.33 (m, 1H), 3.26-3.18 (m, 1H), 2.82-2.79 (m, 1H), 2.46-2.36 (m, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 140.2, 129.0, 128.5, 126.0, 58.2, 48.2, 32.2 ppm.

***N*-[(1*S*, 2*S*)-2-chloro-2,3-dihydro-1*H*-inden-1-yl]methanesulfonamide (25)**

$[\alpha]_D^{23}$ +91.3° (c 0.016, EtOH); ^1H NMR (CDCl_3 , 400 MHz) δ 7.43 (m, 1H), 7.32 (m, 2H), 7.31 (m, 1H), 5.03 (t, J = 8.4 Hz, 1H), 4.80 (br d, 1H), 4.29 (dd, J = 15.5, 7.6 Hz, 1H), 3.49 (dd, J = 15.9, 7.6 Hz, 1H), 3.20 (s, 3H), 3.14 (m, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 139.5, 138.7, 129.3, 128.1, 124.7, 124.3, 67.0, 63.2, 42.67, 40.1 ppm.

(3*aR*, 8*bR*)-3,3*a*,4,8*b*-tetrahydro-1*H*-indeno[1,2-*c*]isothiazole 2,2-dioxide (26)

$[\alpha]_D^{23}$ -101.2° (c 0.023, EtOH); ^1H NMR (CDCl_3 , 400 MHz) δ 7.36-7.27 (m, 4H), 5.05 (d, J = 7.3 Hz, 1H), 3.63-3.56 (m, 1H), 3.40-3.50 (m, 2H), 3.03 (dd, J = 16.6, 4.0 Hz, 1H), 2.89 (dd, 1H, J = 12.7, 7.9 Hz); ^{13}C NMR (CDCl_3 , 100 MHz) δ 141.3, 139.9, 129.5, 127.9, 125.4, 125.1, 63.0, 52.9, 39.8, 37.5 ppm.

References

1. White, E. H.; Lim, H. M. *J. Org. Chem.* **1987**, *52*, 2162.
2. Bliss, A. D.; Cline, W. K.; Hamilton, C. E.; Sweeting, O. *J. Org. Chem.* **1963**, *28*, 3537.
3. The antipode of sultam **20** has been reported in literature. However, no optical rotation value was given, see; Spaltenstein, A., Almond, M. R., Bock, W. J., Cleary, D. G., Furfine, E. S., Hazen, R. J., Kazmierski, W. M., Salituro, F. G., Tung, R. D., Wright, L. L. *Bioorganic. Med. Chem. Lett.* **2000**, *10*, 1159.
4. For the racemic *N*-methyl sultam **23** synthesis, see; Togo, H.; Harada, Y.; Yokoyama, M. *J. Org. Chem.* **2000**, *65*, 926.

nmr400a h-1

Current Data Parameters
NAME 66684-288
EXPNO 13
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030724
Time 10.06
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 17
DS 4
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4904180 sec
RG 128
DM 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.50 usec
PL1 -3.00 dB
SFO1 400.2522014 MHz

F2 - Processing parameters
SI 16384
SF 400.2500553 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 40.00 cm
F1P 10.000 ppm
F1 4002.50 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCX 0.25000 ppm/cm
RECN 100.06252 Hz/cm

nmr400a h-1 decoupled c-13

Current Data Parameters
 NAME 6684-288
 EXPNO 12
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030724
 Time 10.02
 INSTRUM spect
 PROBRD 5 mm QNP 1H
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 106
 DS 0
 SWH 26315.789 Hz
 FIDRES 0.401547 Hz
 AQ 1.2452340 sec
 RG 8152
 DW 19.000 usec
 DE 6.00 usec
 TE 300.0 K
 D1 0.10000000 sec
 d11 0.03000000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 2.00 usec
 PL1 0.00 dB
 SF01 100.613737 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 P2 9.00 usec
 PL2 120.00 dB
 PL12 18.00 dB
 SF02 400.2530012 MHz

F2 - Processing parameters
 SI 32768
 SF 100.612558 MHz
 WDW EM
 SSB EM
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR Plot Parameters
 CX 40.00 cm
 FIP 200.000 ppm
 F1 20328.59 Hz
 F2 0.000 ppm
 F2 0.00 Hz
 FWHM 5.00000 ppm/cm
 BIZN 503.21478 Hz/cm

76.74
77.06
77.38

45.39

50.19

24.12
23.67

nmr400b h-1

Current Data Parameters
NAME 66684-272
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030721
Time 10.34
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 32
DS 2
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4504180 sec
RG 256
DW 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

***** CHANNEL f1 *****

NUC1 1H
P1 10.50 usec
PL1 6.00 dB
SFO1 399.872692 MHz

F2 - Processing parameters

SI 16384
SF 399.870207 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.40

1D NMR plot parameters

CX 40.00 cm
FIP 10.000 ppm
F1 3998.70 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCH 0.25000 ppm/cm
HZCM 99.96751 Hz/cm

nmr400b h-1 decoupled c-13

Current Data Parameters
NAME 66684-272
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030721
Time 10.40
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zgpg30
TD 65536
FIDRES 0.000000
SOLVENT CDCl3
NS 301
DS 4

SWH 26315.789 Hz
FIDRES 0.401547 Hz
AQ 1.2457340 sec
RG 8192
DM 19.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec
D11 0.03000000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 3.00 usec
PL1 5.00 dB
SFO1 100.5584112 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.70 dB
SFO2 399.8719994 MHz

F2 - Processing parameters
SI 32768
SF 100.5473900 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 40.00 cm
F1P 200.000 ppm
F1 20109.48 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCH 5.00000 ppm/cm
HZCH 502.73697 Hz/cm

77.391
77.074
76.755
56.183
45.406
31.671
25.219
14.044

ppm

Current Data Parameters
NAME 66684-274
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030718
Time 9.09
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 32
DS 4
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4904180 sec
RG 912.3
DN 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 10.50 usec
PL1 -1.00 dB
SFO1 400.2522014 MHz

F2 - Processing parameters
SI 16384
SF 400.2500593 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 40.00 cm
F1P 10.000 ppm
F1 4002.50 Hz
F2P 0.000 ppm
F2 0.00 Hz
FREQ 0.25000 ppm/cm
HZCH 100.06252 Hz/cm

nmr400a h-1 decoupled c-13

129.3
128.9
128.7

77.3
77.0
76.7

67.0

45.4

30.5

26.8

Current Data Parameters
NAME 6664-274
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030718
Time 9.17
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zgpgc
TD 65536
SOLVENT CDCl3
NS 380
DS 0
SWH 26315.789 Hz
FIDRES 0.401547 Hz
AQ 1.2452340 sec
RG 5180.6
DW 13.000 usec
DE 300.0 K
TE 300.0 K
D1 0.10000000 sec
D11 0.03000000 sec
D12 0.03000000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 2.00 usec
PL1 0.00 dB
SFO1 100.630737 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 93.00 usec
PL2 120.00 dB
PL12 120.00 dB
SFO2 400.2510012 MHz

F2 - Processing parameters
SI 32768
SF 100.642588 MHz
WDW EM
SS 0
SB 1.00 Hz
GB 0
PC 1.47

1D NMR plot parameters
AQ 40.00 cm
SI 200.000 ppm
F2 2028.55 Hz
WDW EM
SS 0.000 ppm
GB 0.000 ppm
PC 5.00000 ppm/cm
F2 503.21478 Hz/cm

nmr400a h-1

Current Data Parameters
NAME 66684-281
EXPNO 5
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030626
Time 15.01
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 32
DS 4
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.490480 sec
RG 406.4
EQ 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

***** CHANNEL f1 *****

NUC1 1H
P1 10.50 usec
PL1 -3.00 dB
SFO1 400.252014 MHz

F2 - Processing parameters

SI 16384
SF 400.2500591 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.40

1D NMR plot parameters

CX 40.00 cm
F1P 10.000 ppm
F1 4002.50 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCH 0.25000 ppm/cm
HZCH 100.06252 Hz/cm

nmr400a h-1 decoupled c-13

nmr400a h-1

nmr400a h-1 decoupled c-13

Current Data Parameters
 NAME 66684-279
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030718
 Time 11.15
 INSTRUM spect
 PROBED 5 mm QNP 1H
 PULPROG zg30
 TD 32768
 SOLVENT CDCl3
 NS 32
 DS 2
 SWH 6578.947 Hz
 FIDRES 0.200774 Hz
 AQ 2.4904180 sec
 RG 256
 DW 76.000 usec
 DE 6.00 usec
 TE 300.0 K
 D1 0.1000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 10.50 usec
 PL1 6.00 dB
 SF01 399.8724592 MHz

F2 - Processing parameters
 SI 16384
 SF 399.8700207 MHz
 MDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 40.00 cm
 P1P 10.000 ppm
 F1 3998.70 Hz
 F2P 0.000 ppm
 F2 0.00 Hz
 PPMCN 0.25000 ppm/cm
 HZCN 99.86751 Hz/cm

nmr400b h-1 decoupled c-13

nmr400b h-1

Current Data Parameters
NAME 66684-288
EXPNO 5
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030916
Time 8.33
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl₃
NS 32
DS 2
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4904180 sec
RG 101.6
DW 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.50 usec
PL1 6.00 dB
SFO1 399.872452 MHz

F2 - Processing parameters
SI 16384
SF 399.870207 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CK 40.00 cm
F1P 10.000 ppm
F1 3998.70 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCH 0.25000 ppm/cm
HZCM 99.96751 Hz/cm

nmr400b h-1 decoupled c-13

nmr400b h-1

Current Data Parameters
NAME 66684-288
EXPNO 4
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030801
Time 15.56
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 24
DS 2
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4904180 sec
RG 181
RG 181
DW 76.000 usec
DE 6.00 usec
TE 300.0 K
D1 0.10000000 sec

***** CHANNEL f1 *****

NUC1 1H
P1 10.50 usec
PL1 6.00 dB
SFO1 399.872452 MHz

F2 - Processing parameters

SI 16384
SF 399.8700207 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 40.00 cm
FIP -10.000 ppm
F1 3998.70 Hz
F2 0.000 ppm
FZ 0.00 Hz
PPMCH 0.25000 ppm/cm
HZCH 99.96751 Hz/cm

nmr400a h-1 decoupled c-13

Current Data Parameters
 NAME 6684-289
 EXPNO 12
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20030918
 Time 11:20
 INSTRUM spect
 PROBRD 5 mm QNP 1H
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl₃
 NS 321
 DS 0
 SWH 26315.789 Hz
 FIDRES 0.401547 Hz
 AQ 1.2453340 sec
 RG 4597.9
 DW 15.000 usec
 DE 6.00 usec
 TE 300.0 K
 D1 0.10000000 sec
 DI 11
 0.03000000 sec

***** CHANNEL f1 *****

NUC1 13C
 P1 2.00 usec
 PL1 0.00 dB
 SF01 100.6339737 MHz

***** CHANNEL f2 *****

CPDPRG2 waltz16
 NUC3 1H
 PCPD2 93.00 usec
 PL2 120.00 dB
 PL12 18.00 dB
 SF02 400.2520012 MHz

F2 - Processing parameters

SI 32768
 SF 100.6229588 MHz
 NQW no
 SSB no
 LB 0.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters

CX 40.00 cm
 FIP 200.000 ppm
 F1 20128.59 Hz
 F2 0.000 ppm
 F2 0.00 Hz
 PPMCN 5.000000 ppm/cm
 HZCN 503.21478 Hz/cm

Current Data Parameters
NAME 66684-299
EXPNO 11
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030918
Time 11.15
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 23
DS 4
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4994180 sec
RG 287.4
DWD 76.000 usec
DE 6.00 usec
TE 300.0 K
TD 0.1000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 -3.00 dB
SFO1 400.2522014 MHz

F2 - Processing parameters

SI 16384
SF 400.2500593 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CK 40.00 cm
F1P 10.000 ppm
F1 4002.50 Hz
F2P 0.000 ppm
F2 0.00 Hz
PPMCK 0.25000 ppm/cm
HZCK 100.06252 Hz/cm

NH₂SO₂Me

nmr400b h-1

Current Data Parameters
NAME 66684-277
EXNO 7
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030916
Time 8.26
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 32768
SOLVENT CDCl₃
NS 21
DS 2
SWH 6578.947 Hz
FIDRES 0.200774 Hz
AQ 2.4904180 sec
RG 228.1
DM 76.000 usec
DE 6.00 usec
TE 300.0 K
TS 0.10000000 sec
D1

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 6.00 dB
SFO1 399.8724532 MHz

F2 - Processing parameters

SI 16384
SF 399.8700207 MHz
WDW no
SSB 0
LA 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 40.00 cm
P1P 10.000 ppm
P1 3998.70 Hz
P2P 0.000 ppm
P2 0.00 Hz
PPMCH 0.25000 ppm/cm
HZCH 99.96751 Hz/cm

nmr400b h-1 decoupled c-13

Current Data Parameters
 NAME 66684-277
 EXNO 2
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20030612
 Time_ 8.50
 INSTRUM spect
 PROBED 5 mm QNP 1H
 PULPROG zg30
 TD 32768
 SOLVENT CDCl3
 NS 23
 DS 4
 SWH 6578.947 Hz
 FIDRES 0.200774 Hz
 AQ 2.4904180 sec
 RG 574.7
 DW 76.000 usec
 DE 6.90 usec
 TE 300.0 K
 D1 0.1000000 sec

===== CHANNEL f1 =====

MIX1 1H
 P1 10.50 usec
 PL1 -3.00 dB
 SFO1 400.2522014 MHz

F2 - Processing parameters

SI 16384
 SF 400.2500553 MHz
 WDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters

CX 40.00 cm
 P1P 10.000 ppm
 F1 4002.50 Hz
 P2P 0.000 ppm
 F2 0.00 Hz
 PRCN 0.25000 ppm/cm
 HZCN 100.06252 Hz/cm

nmr400a h-1 decoupled c-13