Supporting Information

Practical and Efficient Suzuki-Miyaura Cross-Coupling of 2-Iodocycloenones with Arylboronic Acids Catalyzed by Recyclable Pd/C

François-Xavier Felpin

Université Bordeaux-I, Laboratoire de Chimie Organique et Organométallique, 351 Cours de la Libération, 33405 Talence Cedex, France

fx.felpin@lcoo.u-bordeaux1.fr

Contents

General Methods.....	S2
Preparation of compounds.....	S2-S12
NMR Spectra.....	S13-S52

General Methods: Chemical shifts from proton and carbon NMR spectra are reported in ppm relative to the CDCl₃ peak at 7.26 ppm (¹H) or 77.0 ppm (¹³C). Infrared (IR) spectra were recorded as neat samples on NaCl plates or with KBr pellets. Yields refer to isolated material determined to be pure by NMR spectroscopy and thin layer chromatography (TLC), unless specified otherwise in the text. Compounds **1**,¹ **2**,² **12**,³ **14**,⁴ **15**,¹ **16**,³ **17**,⁵ **18**⁶ were prepared according literature procedures.

General procedure for Suzuki cross-coupling reaction. To a solution of iodoenone (1.5 mmol) in DME (4.5 mL) and H₂O (4.5 mL) was added Na₂CO₃ (318 mg, 3 mmol), Ar-B(OH) (3 mmol) and Pd/C (79.5 mg, 5 mol%). The mixture was stirred for the specified time at 25, 50 or 80 °C as indicated in tables 2 and 3. Pd/C was filtered, and the filtrate was diluted with H₂O (4.5 mL) and extracted with Et₂O (3 x). The collected organic extracts were dried (MgSO₄) and concentrated under reduced pressure.

Eventually, the catalyst was washed twice with water and twice with Et₂O, dried under vacuum (4 h) and reused in a new run.

2-Phenylcyclohex-2-enone (3). Purification by flash chromatography (10%

EtOAc-Petroleum ether) gave **3** as a white solid (222 mg, 86%). mp 93-94 °C [lit.⁷ 92-93 °C]. IR (KBr) ν 1662, 2933, 3016 cm⁻¹. ¹H NMR (CDCl₃, 300 MHz) δ 2.09-2.17 (m, 2H), 2.53-2.64 (m, 4H), 7.06 (app t, 1H, J = 4.1 Hz), 7.28-7.40 (m, 5H). ¹³C NMR (CDCl₃, 75 MHz) δ 22.9, 26.5, 39.0, 127.5, 127.9, 128.5, 136.5, 140.3, 147.9, 197.9. HRMS (EI) calcd for C₁₂H₁₂O (M⁺) 172.0888, found 172.0886.

2-*o*-Tolylcyclohex-2-enone (4). Purification by flash chromatography (10%

EtOAc-Petroleum ether) gave **4** as a colorless oil (223 mg, 80%). IR (neat) ν 1682, 2949, 3059 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.14 (s, 3H), 2.14-2.18 (m, 2H), 2.54-2.61 (m, 4H), 6.89-6.91 (m, 2H), 7.00-7.03 (m, 1H), 7.17-7.26 (m, 3H). ^{13}C NMR (CDCl_3 , 75 MHz) δ 20.0, 23.1, 26.3, 38.7, 125.5, 127.8, 129.5, 129.7, 136.3, 136.9, 141.9, 148.6, 198.0. HRMS (EI) calcd for $\text{C}_{13}\text{H}_{14}\text{O}$ (M^+) 186.1045, found 186.1042.

2-(4-Methoxyphenyl)cyclohex-2-enone (5). Purification by flash

chromatography (20% EtOAc-Petroleum ether) gave **5** as a white solid (264 mg, 87%). mp 58-59 $^{\circ}\text{C}$ [lit.⁸ 59 $^{\circ}\text{C}$]. IR (KBr) ν 1573, 1604, 1669, 2931 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.07-2.15 (m, 2H), 2.51-2.62 (m, 4H), 3.82 (s, 3H), 6.90 (dm, 2H, $J = 9.0$ Hz), 7.00 (app t, 1H, $J = 4.5$ Hz), 7.27 (dm, 2H, $J = 9.0$ Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.9, 26.5, 39.0, 55.2, 113.4, 128.9, 129.7, 139.7, 146.9, 159.0, 198.2. HRMS (EI) calcd for $\text{C}_{13}\text{H}_{14}\text{O}_2$ (M^+) 202.0994, found 202.0995.

2-(2,3-Dimethoxyphenyl)cyclohex-2-enone (6). After the usual work-up,

the organic extracts were washed two times with 10% aqueous NaOH solution. Purification by flash chromatography (20% EtOAc-Petroleum ether) gave **6** as a white solid (313 mg, 90%). mp

94-95 °C [lit.⁹ 95-96 °C]. IR (KBr) ν 1578, 1594, 1675, 2926 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.09-2.17 (m, 2H), 2.48-2.54 (m, 2H), 2.58-2.62 (m, 2H), 3.72 (s, 3H), 3.85 (s, 3H), 6.67 (dd, 1H, J = 1.1, 7.5 Hz), 6.87-6.91 (m, 2H), 7.01 (t, 1H, J = 8.3 Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ 23.1, 26.3, 38.6, 55.8, 60.4, 112.4, 122.5, 123.5, 131.5, 138.6, 146.8, 148.2, 152.5, 197.8. HRMS (LSIMS) calcd for $\text{C}_{14}\text{H}_{17}\text{O}_3$ ($\text{M}+\text{H}^+$) 233.1178, found 233.1172.

2-(3-Nitrophenyl)cyclohex-2-enone (7). Purification by flash

chromatography (20% EtOAc-Petroleum ether) gave **7** as a colorless oil (260 mg, 80%) which solidified on standing. mp 45 °C [lit.⁶ 46 °C]. IR (KBr) ν 1574, 1595, 1613, 1682, 2948, 3085 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.10-2.18 (m, 2H), 2.57-2.64 (m, 4H), 7.14-7.18 (m, 1H), 7.50 (t, 1H, J = 8.3 Hz), 7.65-7.68 (dd, 1H, J = 1.1, 7.9 Hz), 8.12-8.17 (m, 2H). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.7, 26.6, 38.7, 122.4, 123.6, 128.8, 134.9, 138.0, 138.5, 148.0, 149.7, 197.1. HRMS (LSIMS) calcd for $\text{C}_{12}\text{H}_{12}\text{NO}_3$ ($\text{M}+\text{H}^+$) 218.0817, found 218.1108.

2-(2-Fluorophenyl)cyclohex-2-enone (8). Purification by flash chromatography

(10% EtOAc-Petroleum ether) gave **8** as a white solid (228 mg, 80%). mp 64-65 °C. IR (KBr) ν 1577, 1607, 1673, 2935, 3075 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.10-2.19 (m, 2H), 2.53-2.64 (m, 4H), 7.03-7.34 (m, 5H). ^{13}C NMR (CDCl_3 , 75 MHz), (2 rotamers) δ 22.8, 26.4, 38.5, 115.2, 115.5, 123.7, 123.7, 124.4, 124.6, 129.3, 129.4, 131.1, 131.2, 136.0, 149.8, 158.3, 161.5, 196.7. HRMS (EI) calcd for $\text{C}_{12}\text{H}_{11}\text{OF}$ (M^+) 190.0794, found 190.0787.

2-(4-Acetylphenyl)cyclohex-2-enone (9). Purification by flash chromatography (30% EtOAc-Petroleum ether) gave **9** as a white solid (314 mg, 98%). mp 81-82 °C. IR (neat) ν 1600, 1675, 2926, 3058 cm^{-1} . ^1H NMR (CDCl_3 , 250 MHz) δ 2.08-2.18 (m; 2H), 2.54-2.64 (m, 7H), 7.11 (t, 1H, $J = 4.3$ Hz), 7.41 (dd, 2H, $J = 1.8, 6.7$ Hz), 7.93 (dd, 2H, $J = 1.5, 6.4$ Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.8, 26.6, 38.9, 128.0, 128.8, 136.1, 139.6, 141.3, 149.1, 197.4, 197.8. HRMS (LSIMS) calcd for $\text{C}_{14}\text{H}_{15}\text{O}_2$ ($\text{M}+\text{H}^+$) 215.1072, found 215.1070.

2-(Thiophen-2-yl)cyclohex-2-enone (10). Purification by flash chromatography (7% EtOAc-Petroleum ether) gave **10** as a colorless oil which turns rapidly blue under air atmosphere (216 mg, 81%). IR (neat) ν 1517, 1677, 2947, 3070 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.04-2.12 (m, 2H), 2.53-2.62 (m, 4H), 7.02 (dd, 1H, $J = 3.8, 5.3$ Hz), 7.27-7.36 (m, 3H). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.5, 26.5, 38.7, 125.0, 125.9, 126.5, 133.0, 136.9, 145.3, 196.8. HRMS (EI) calcd for $\text{C}_{10}\text{H}_{10}\text{OS}$ (M^+) 178.0452, found 178.0457.

2-(Benzofuran-2-yl)cyclohex-2-enone (11). Purification by flash chromatography (10% EtOAc-Petroleum ether) gave **11** as a white solid (300 mg, 94%). mp 103-104 °C. IR (KBr) ν 1549, 1677, 2947, 3052 cm^{-1} . ^1H NMR (CDCl_3 , 250 MHz) δ 2.06-2.17 (m,

2H), 2.58-2.66 (m, 4H), 7.16-7.29 (m, 2H), 7.41-7.44 (m, 2H), 7.56-7.59 (m, 1H), 7.75 (app t, 1H, $J = 5.5$ Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.4, 26.1, 39.1, 106.4, 110.6, 121.5, 122.7, 124.5, 129.2, 129.3, 146.4, 150.1, 153.7, 196.0. HRMS (EI) calcd for $\text{C}_{14}\text{H}_{12}\text{O}_2$ (M^+) 212.0837, found 212.0844.

2-Phenylcyclopent-2-enone (20). Purification by flash chromatography (10%

EtOAc-Petroleum ether) gave **20** as a white solid (191 mg, 81%). mp 71-72 °C [lit.¹⁰ 68-69 °C]. IR (KBr) ν 1596, 1618, 1692, 2951, 3046 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 2.60-2.63 (m, 2H), 2.71-2.75 (m, 2H), 7.32-7.43 (m, 3H), 7.68-7.72 (m, 2H), 7.84 (app t, 1H, $J = 2.6$ Hz), ^{13}C NMR (CDCl_3 , 75 MHz) δ 26.2, 35.7, 127.0, 128.3, 128.4, 131.6, 143.5, 158.9, 207.5. HRMS (LSIMS) calcd for $\text{C}_{11}\text{H}_9\text{O}$ ($\text{M}-\text{H}^+$) 157.0657, found 157.0653.

Diethyl 3,3a,4,5-tetrahydro-5-oxo-6-phenylpentalene-2,2(1H)-

dicarboxylate (21). Purification by flash chromatography (15% EtOAc-Petroleum ether) gave **21** as a colorless oil (236 mg, 46%). IR (KBr) ν 1650, 1706, 1730, 2932, 3057 cm^{-1} . ^1H NMR (CDCl_3 , 250 MHz) δ 1.22 (t, 3H, $J = 7.0$ Hz), 1.31 (t, 3H, $J = 7.0$ Hz), 1.76 (t, 1H, $J = 12.8$ Hz), 2.31 (dd, 1H, $J = 3.3, 18.0$ Hz), 2.78-2.88 (m, 2H), 3.12-3.16 (m, 1H), 3.29 (d, 1H, $J = 19.5$ Hz), 3.65 (d, 1H, $J = 19.2$ Hz), 4.12-4.21 (m, 2H), 4.28 (q, 2H, $J = 7$ Hz), 7.29-7.44 (m, 3H), 7.53-7.58 (m, 2H). ^{13}C NMR (CDCl_3 , 62.5 MHz) δ 14.0, 14.0, 35.9, 38.7, 42.7, 42.9, 61.3, 62.0, 62.2,

128.2, 128.4, 130.9, 135.5, 170.7, 171.6, 179.0, 207.2. HRMS (LSIMS) calcd for $C_{20}H_{23}O_5$ ($M+H^+$) 343.1545, found 343.1546.

2-Phenylcyclohept-2-enone (22). Purification by flash chromatography (10%

EtOAc-Petroleum ether) gave **22** as a colorless oil (198 mg, 71%). IR (neat) ν 1675, 2935 cm^{-1} .

1H NMR ($CDCl_3$, 300 MHz) δ 1.80-1.98 (m, 4H), 2.55 (app q, 2H, $J = 6.0$ Hz), 2.77 (app t, 2H, $J = 6.0$ Hz), 6.77 (t, 1H, $J = 6.4$ Hz), 7.28-7.37 (m, 5H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ 22.1, 24.8, 28.0, 43.2, 127.4, 128.0, 128.1, 138.9, 141.6, 144.7, 205.6. HRMS (EI) calcd for $C_{13}H_{14}O$ (M^+) 186.1045, found 186.1047.

2-Phenyl-4-tert-butylcyclohex-2-enone (23). Purification by flash

chromatography (10% EtOAc-Petroleum ether) gave **23** as a yellow oil (291 mg, 85%). IR (neat)

ν 1599, 1684, 2959, 3056 cm^{-1} . 1H NMR ($CDCl_3$, 250 MHz) δ 1.03 (s, 9H), 1.77-1.91 (m, 1H), 2.14-2.19 (m, 1H), 2.35-2.57 (m, 2H), 2.67-2.73 (m, 1H), 7.06 (s broad, 1H), 7.33-7.36 (m, 5H). ^{13}C NMR ($CDCl_3$, 62.5 MHz) δ 24.3, 27.3, 33.2, 38.7, 47.5, 127.5, 127.9, 128.6, 136.7, 140.3, 150.7, 198.3. HRMS (EI) calcd for $C_{12}H_{12}O$ ($M^+ - C_4H_8$) 172.0888, found 172.0884.

3-Methyl-2-phenylcyclohex-2-enone (24). Purification by flash chromatography

(15% EtOAc-Petroleum ether) gave **24** as a colorless oil (218 mg, 78%). IR (neat) ν 1598, 1621, 1666, 2936, 3054 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.83 (s, 3H), 2.09 (s br, 2H), 2.50-2.53 (m, 4H), 7.04-7.07 (m, 1H), 7.26-7.38 (m, 3H). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.1, 22.7, 32.5, 38.0, 126.9, 127.9, 129.9, 135.8, 138.1, 157.4, 198.1. HRMS (EI) calcd for $\text{C}_{13}\text{H}_{14}\text{O}$ (M^+) 186.1045, found 186.1037.

3-phenyl-4H-chromen-4-one (25). Purification by flash chromatography

(10% EtOAc-Petroleum ether) gave **25** as a white solid (316 mg, 95%). mp 128-129 $^{\circ}\text{C}$ [lit.¹¹ 131.5-132 $^{\circ}\text{C}$]. IR (KBr) ν 1570, 1618, 1640, 3052 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 7.36-7.50 (m, 5H), 7.56-7.59 (m, 2H), 7.66-7.72 (m, 1H), 8.03 (s, 1H), 8.33 (dd, 1H, $J = 1.5, 7.9$ Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ 118.0, 124.6, 125.2, 125.4, 126.4, 128.2, 128.5, 128.9, 131.8, 133.6, 153.0, 156.2, 176.2. HRMS (EI) calcd for $\text{C}_{15}\text{H}_{10}\text{O}_2$ (M^+) 222.0681, found 222.0674.

1-Benzoyl-2,3-dihydro-5-phenylpyridin-4(1H)-one (26). After the usual

work-up, the organic extracts were washed two times with 10% aqueous NaOH solution. Purification by flash chromatography (30% EtOAc-Petroleum ether) gave **26** as a white solid

(399 mg, 96%). mp 133-134 °C. IR (KBr) ν 1494, 1602, 1661, 2921, 3058 cm^{-1} . ^1H NMR (CDCl_3 , 250 MHz) δ 2.84 (t, 2H, $J = 7.3$ Hz), 4.25 (t, 2H, $J = 7.3$ Hz), 7.25 (m, 5H), 7.46-7.64 (m, 5H), 7.79 (s, 1H). ^{13}C NMR (CDCl_3 , 62.5 MHz) δ 36.6, 43.2, 119.9, 127.5, 128.2, 128.6, 128.6, 128.8, 131.9, 132.7, 133.8, 142.5, 170.2, 191.6. HRMS (LSIMS) calcd for $\text{C}_{18}\text{H}_{16}\text{NO}_2$ ($\text{M}+\text{H}^+$) 278.1181, found 278.1186.

2-Phenyl-5 α -cholest-1-en-3-one (27). Purification by flash

chromatography (5% EtOAc-Petroleum ether) gave **27** as a white solid (628 mg, 91%). mp 128-129 °C. IR (KBr) ν 1675, 2937 cm^{-1} . ^1H NMR (CDCl_3 , 250 MHz) δ 0.71 (s, 3H), 0.87 (d, 6H, $J = 6.7$ Hz), 0.91 (d, 3H, $J = 6.4$ Hz), 1.10 (s, 3H), 0.85-1.59 (m, 20H), 1.71-1.86 (m, 3H), 1.98-2.12 (m, 2H), 2.38 (dd, 1H, $J = 4.3, 17.5$ Hz), 2.54 (dd, 1H, $J = 13.7, 17.5$ Hz), 7.17 (s, 1H), 7.26-7.36 (m, 5H). ^{13}C NMR (CDCl_3 , 62.5 MHz) δ 12.2, 13.0, 18.6, 21.3, 22.5, 22.8, 23.8, 24.1, 27.4, 28.0, 28.2, 31.3, 35.7, 35.8, 36.1, 39.5, 39.8, 41.6, 42.7, 44.1, 50.3, 56.2, 56.4, 127.5, 128.0, 128.8, 136.7, 138.0, 156.4, 198.2. HRMS (LSIMS) calcd for $\text{C}_{33}\text{H}_{49}\text{O}$ ($\text{M}+\text{H}^+$) 461.3783, found 461.3787.

Diethyl 3,3a,4,5-tetrahydro-6-(trimethylsilyl)-5-oxopentalene-2,2(1H)-

dicarboxylate (28). To a solution of diethyl 1- (trimethylsilyl)- 6-hepten- 1-yne- 4, 4-dicarboxylate¹² (1 g, 3.23 mmol) in Et₂O (32 mL) at room temperature was added Co₂(CO)₈ (1.32 g, 3.88 mmol). After 1 h of stirring, the volatiles were evaporated under reduced pressure. The brown crude was dissolved in THF (32 mL) and treated with TMANO.2H₂O (2.36 g, 21.3 mmol). After 3 h of stirring, the resulting mixture was filtered and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (15% EtOAc-Petroleum ether) gave the desired enone **28** as a white solid (872 mg, 80%). mp 65-66 °C [lit.¹³ 66-68 °C]. IR (KBr) ν 1575, 1689, 2932 cm⁻¹. ¹H NMR (CDCl₃, 300 MHz) δ 0.19 (s, 9H), 1.26 (q, 6H, J = 7.1 Hz), 1.66 (t, 1H, J = 12.8 Hz), 2.06 (dd, 1H, J = 4.1, 17.7 Hz), 2.57 (dd, 1H, J = 6.4, 17.3 Hz), 2.77 (dd, 1H, J = 7.9, 12.8 Hz), 3.01 (m, 1H), 3.19-3.35 (m, 1H), 4.16-4.27 (m, 4H). ¹³C NMR (CDCl₃, 75 MHz) δ -1.3, 14.0, 36.2, 38.5, 42.9, 46.5, 60.8, 61.9, 62.0, 136.6, 171.0, 171.6, 192.4, 213.1. HRMS (LSIMS) calcd for C₁₇H₂₇O₅Si (M+H⁺) 339.1628, found 339.1629.

Diethyl 3,3a,4,5-tetrahydro-6-iodo-5-oxopentalene-2,2(1H)-dicarboxylate

(13). A solution of **28** (800 mg, 2.37 mmol) in CH₂Cl₂ (4 mL) was added ICl (1M in CH₂Cl₂, 2.2 mL) at 0 °C. The resulting mixture was allowed to warm to room temperature and stirred for 2 h. The reaction was quenched with saturated Na₂S₂O₃ aqueous solution, extracted with ether (3 x),

washed with brine, dried (MgSO₄) and concentrated under reduced pressure. Purification by flash chromatography (17% EtOAc-Petroleum ether) gave the iodoenone **13** as a colorless oil (835 mg, 90%). IR (KBr) ν 1634, 1729, 2982 cm⁻¹. ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, 3H, J = 6.8 Hz), 1.28 (t, 3H, J = 7.2 Hz), 1.79 (t, 1H, J = 12.8 Hz), 2.25 (dd, 1H, J = 3.0, 17.7 Hz), 2.81-2.90 (m, 2H), 3.12 (d, 1H, J = 20.0 Hz), 3.15-3.24 (m, 1H), 3.37 (d, 1H, J = 20.0 Hz), 4.24 (q, 2H, J = 7.2 Hz), 4.27 (q, 2H, J = 7.2 Hz). ¹³C NMR (CDCl₃, 75 MHz) δ 14.0, 37.2, 39.1, 40.0, 46.4, 59.8, 62.1, 62.3, 94.6, 170.3, 171.1, 188.4, 203.2. HRMS (LSIMS) calcd for C₁₄H₁₈IO₅ (M+H⁺) 393.0199, found 393.0198.

2-Iodo-5 α -cholest-1-en-3-one (19). A solution of 5 α -cholest-1-en-3-one¹⁴ (1.02 g, 2.66 mmol) in CCl₄ (10 mL) and pyridine (10 mL) was treated with I₂ (2.02 g, 7.97 mmol) and stirred at room temperature for 14 h. The mixture was washed with saturated Na₂S₂O₃ aqueous solution followed by brine. The organic extract was dried (MgSO₄) and concentrated under reduced pressure. Purification by flash chromatography (5% EtOAc-Petroleum ether) gave **19** as a white solid (1.33 mg, 98%). mp 128-129 °C. IR (KBr) ν 1575, 1689, 2932 cm⁻¹. ¹H NMR (CDCl₃, 250 MHz) δ 0.69 (s, 3H), 0.86 (d, 6H, J = 6.7 Hz), 0.91 (d, 3H, J = 6.4 Hz), 1.03 (s, 3H), 0.85-2.07 (m, 25H), 2.50-2.54 (m, 2H), 7.90 (s, 1H). ¹³C NMR (CDCl₃, 75 MHz) δ 12.2, 12.7, 18.6, 21.3, 22.5, 22.8, 23.8, 24.1, 27.3, 28.0, 28.2, 31.0, 35.5, 35.7, 36.1, 39.5, 39.6, 39.8, 42.6,

44.1, 44.3, 49.8, 56.1, 56.3, 103.3, 167.1, 192.4. HRMS (EI) calcd for C₂₇H₄₄IO (M+H⁺) 511.2437, found 511.2451.

-
- (1) William, A. D.; Kobayashi, Y. *J. Org. Chem.* **2002**, *67*, 8771-8782.
- (2) Kim, J. G.; Waltz, K. M.; Garcia, I. F.; Kwiatkowski, D.; Walsh, P. J. *J. Am. Chem. Soc.* **2004**, *126*, 12580-12585.
- (3) Rossi, R.; Bellina, F.; Ciucci, D. *J. Organomet. Chem.* **1997**, *542*, 113-120.
- (4) Mayasundari, A.; Young, D. G. *J. Tetrahedron Lett.* **2001**, *42*, 203-206.
- (5) Lin, G.-Q.; Hong, R. *J. Org. Chem.* **2001**, *66*, 2877-2880.
- (6) Prasad, A. S. B.; Knochel, P. *Tetrahedron* **1997**, *53*, 16711-16720.
- (7) Kraus, G. A.; Frazier, K. *J. Org. Chem.* **1980**, *45*, 2579-2581.
- (8) Arumugam, N.; Shenbagamurthi, P. *Indian J. Chem. Sect. B* **1976**, *14*, 831-833.
- (9) Barnes, R. A.; Reinhold, D. F. *J. Am. Chem. Soc.* **1952**, *74*, 1327-1329.
- (10) Donkervoort, J. G.; Gordon, A. R.; Johnstone, C.; Kerr, W. J.; Lange, U. *Tetrahedron* **1996**, *52*, 7391-7420.
- (11) Hoshino, Y.; Miyaura, N.; Suzuki, A. *Bull. Chem. Soc. Jpn.* **1988**, *61*, 3008-3010.
- (12) Kobayashi, T.; Koga, Y.; Narasaka, K. *J. Organomet. Chem.* **2001**, *624*, 73-87.
- (13) Morimoto, T.; Chatani, N.; Fukumoto, Y.; Murai, S. *J. Org. Chem.* **1997**, *62*, 3762-3765.
- (14) Win, W. W.; Franck, R. W. *J. Org. Chem.* **1997**, *62*, 4510-4512.