This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Preferential transport of microplastics by wind

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1016/j.atmosenv.2020.118038>

PUBLISHER

Elsevier

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This paper was accepted for publication in the journal Atmospheric Environment and the definitive published version is available at <https://doi.org/10.1016/j.atmosenv.2020.118038>.

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Bullard, Joanna, Annie Ockelford, Patrick O'Brien, and Cheryl McKenna Neuman. 2020. "Preferential Transport of Microplastics by Wind". Loughborough University. <https://hdl.handle.net/2134/13148378.v1>.

Title: Preferential transport of microplastics by wind

Authors: *Joanna E. Bullard¹, Annie Ockelford², Patrick O'Brien³, Cheryl McKenna
Neuman³

Affiliations:

¹ Geography and Environment, Loughborough University, Leicestershire LE11 3TU UK.

² School of Environment and Technology, University of Brighton, Brighton, BN2 4GJ
UK.

³ Trent Environmental Wind Tunnel Laboratory, School of the Environment, 1600 West
Bank Drive, Trent University, Peterborough, Ontario, Canada K9J 0G2.

*Correspondence to: j.e.bullard@lboro.ac.uk.

Abstract

Contamination of terrestrial and marine environments by plastic waste has been widely documented. Most research into the distribution of microplastics has focused on water but here we show that wind transport can be very effective in mobilising microplastic particles. A series of wind tunnel experiments using two different substrates (sand and soil), two different microplastics (microbeads and fibres) and 5 different concentrations of microplastics (ranging from 0 mg kg⁻¹_{dw} to 1040 mg kg⁻¹_{dw}) is used to demonstrate that microplastics are preferentially transported by wind compared to sand and soil. When compared to either of the untreated substrate beds (0 mg kg⁻¹_{dw}), the inclusion of microplastics was not found to significantly affect the wind erosion threshold for any of the concentrations or geometric forms (fibres or beads) tested. Averaged over all concentrations of microplastics and both substrate types, microplastic enrichment was lower for microbeads than fibres. The enrichment of microplastic fibres within the entrained particulate matter was one to two orders of magnitude higher for both test bed substrates, ranging from 98 to 498 for the sand and 278 to 726 for the soil. This suggests microplastic shape needs to be carefully parameterized in models of atmospheric microplastic transport. We suggest that microplastic research could benefit from previous investigations into the wind erosion of soil organic carbon.

37 Keywords

38 Wind erosion

39 Microplastic entrainment

40 Particle shape

41 Microbead

42 Fibre

43 Plastic cycle

44

45 Highlights

46 • We ran experiments to determine the susceptibility of microplastics to wind
47 erosion

48 • Fibrous microplastics are preferentially transported compared to microbeads

49 • Particle shape may affect the atmospheric transport distance for microplastics

50

51

1. Introduction

Plastics are highly versatile, low-cost, lightweight, materials and hence in high societal demand for innovation-driven growth and development (North and Halden, 2013; PlasticsEurope, 2019). The properties that make plastics useful - strength, flexibility, durability - also make them difficult to dispose of, and their resistance to degradation has introduced persistent, complex materials to the environment that may have serious consequences for environmental pollution (e.g. Andrady, 2011; Barnes et al., 2009; Free et al., 2014; Jambeck et al., 2015), environmental health (e.g. Gasperi et al., 2018; He, D. et al., 2018; Wright and Kelly, 2017) and ecosystem functionality (e.g. Hu et al., 2019; Lwanga et al., 2016; Arias-Andres et al., 2018). This paper focuses on microplastics, defined as solid synthetic-polymer-containing particles less than 5 mm in size (NOAA/European Marine Strategy Directive). Microplastics can be purposefully produced to be small in size (primary microplastics) or derived from the breakdown of macroplastics (to form secondary microplastics) by chemical (Mailhot et al., 2000; Song et al., 2017), microbial (Yuan et al., 2020) and mechanical processes (Song et al., 2017). Breakdown of plastics since the onset of mass production 70 years ago has resulted in a decrease in the average size of plastic particles and an increase in the abundance and distribution of secondary microplastics, which are pervasive and difficult to remove from the environment (Barnes et al., 2009).

Microplastics have potentially deleterious effects on terrestrial, freshwater and marine environments (e.g. recent reviews by Beaumont et al. 2019; Wang et al., 2020; Li et al., 2020). Research into their long-term effects is in its infancy, but the risks may include damage to organisms' digestive systems (von Moos et al., 2012; Tanaka & Takada, 2016; Wright et al., 2013), facilitating the transfer of harmful chemicals, including persistent organic pollutants (e.g. Hidalgo-Ruz et al., 2012; McCormick et al., 2014; Carbery et al.,

2018), and alterations to soil stability (Boots et al., 2019; Barnes et al., 2009; Lehmann et al., 2019; Zhang et al., 2019; Machado et al., 2018).

Microplastics derive from land-based anthropogenic activity with >50% of all microplastics estimated to be retained in the terrestrial environment (Boucher and Friot, 2017; Zalasiewicz et al., 2016). Despite this, most scientific and media attention has focused on microplastics in the marine depositional sink and presence/absence in the terrestrial environment. The main focus for research in to microplastic transport pathways has been via rivers to the oceans, however recent studies of soils (Scheurer and Bigalke 2018; Allen et al., 2019; Rezaei et al., 2019), snow (Bergmann et al., 2019), supraglacial debris (Ambrosini et al., 2019) and total atmospheric fallout (Dris et al., 2016) indicate airborne transport may be important. The wind erosion and transport of microplastic can not only redistribute particles within the terrestrial environment but atmospheric fallout is also likely to contribute to marine pollution (Evangelidou et al., 2020).

A recent review of atmospheric microplastics (Zhang et al., 2020) highlights that most research has focused on atmospheric deposition of plastic particles. In particular, studies of atmospheric deposition have focused on cities in China (e.g. Cai et al., 2017; Liu et al., 2019; Zhou et al., 2017), France (Dris et al., 2015, 2016, 2017), Germany (Klein and Fischer et al., 2019), Iran (Dehghani et al., 2017; Abbasi et al., 2019) and the UK (Stanton et al., 2019; Wright et al., 2020), where outdoor microplastic abundance ranges from <5 to >1000 particles m⁻² d⁻¹, or 2 - >9000 mg kg⁻¹. Studies of rural and remote environments have identified microplastics in snow in the Pyrenees mountains (Allen et al., 2019), Alps (Ambrosini et al., 2019) and Arctic (Bergmann et al., 2010) and in remote lake catchments in Ireland (Roblin and Aherne, 2020). Microplastics have also been found in floodplain

soils in nature reserves in Switzerland (Scheurer and Bigalke, 2018). The distance of these rural or remote locations from urban areas suggest that airborne, rather than fluvial, transport is a more likely source of at least a proportion of the microplastics found there.

Microplastics take a variety of forms including fragments (flattened, angular), fibres (even-thickness, linear, flexible), films (thin and typically transparent), foams (compressible, sponge-like texture) and beads (granular, rounded). The shape depends both on the specific type of plastic, the original form of the primary plastic, the processes operating to degrade macroplastics to microplastics, age and environmental residence time. For example, fibres are often associated with shedding from textiles, beads with personal care products, foams with insulation or food packaging, and fragments as by-products of manufacturing or the breakdown of macroplastics (Rochman et al., 2019). The most common shapes identified for atmospheric microplastics are fibres e.g. >90% in Paris (Dris et al., 2015); >80% in Dongguan (Cai et al., 2017); 95% in Yantai (Zhou et al., 2017), China; and 92% in London (Wright et al., 2020): or fragments e.g. >90% in Hamburg (Klein and Fischer, 2019) and 68% in the Pyrenees (Allen et al., 2019). This is likely to reflect a combination of the prevalence of shapes at nearby sources (e.g. urban areas that are production centres for synthetic fibres) and the influence of shape on atmospheric transport and residence time.

The small size and often low density of microplastics means that when they are exposed to wind erosion they are likely to be very susceptible to entrainment and transport. This may mean microplastics form a potential risk to human (Gasperi et al., 2018) and ecosystem health in locations far-removed from their origin (Rezaei et al., 2019). To date

there have been very few studies of the entrainment potential of microplastics by wind nor of the impact of particle properties on the mode and distance of airborne transport (Waldschläger et al., 2020). Rezaei et al., (2019) used a field wind tunnel study to demonstrate that wind-eroded sediments from both agricultural and natural areas were enriched with microplastics compared to the original soils but there is no indication of whether particular forms (shapes) of microplastics are preferentially eroded.

The aim of this paper is to explore the extent to which microplastic particles are preferentially-transported compared to natural sediments using a series of controlled wind tunnel experiments. Two different substrates (sand and soil) and two different shapes of microplastic (bead and fibre) were used to determine the impact of substrate and shape on microplastic entrainment.

2. Materials and Methods

2.1 Materials

Experiments were performed in the Trent University Environmental Wind Tunnel (TEWT), a boundary layer simulation tunnel with an open-loop suction design. The tunnel working section is 12.5 m long, with a cross section 0.7 m wide by 0.76 high, and the whole tunnel is housed in an environmental chamber which, for this study, was held constant at 20 °C and 20% relative humidity to control electrostatics. Further details of the TEWT facility are provided in previous publications (Nickling and McKenna Neuman 1997; McKenna Neuman and Nickling, 2000; Li and McKenna Neuman, 2012).

Experiments were conducted using two substrates and two types of microplastic. With a particle density of 2.65 g cm³, substrate one was well-sorted quartz sand (hereafter

'sand') similar to that found on beaches (Constant et al., 2019) and lakeshores (Zhang et al., 2016) from which high concentrations of macro- and microplastics have been reported (Fig. 1a; Table 1). Substrate two was a poorly-sorted soil containing 13.2% organic matter (calculated by loss on ignition at 850 °C) (hereafter 'soil') (Fig.1b; Table 2). Both substrates were air-dried and stored in the same environmental chamber that houses the wind tunnel. The sand is likely to have similar erodibility to dry, loose *in situ* sand e.g. on dry beaches or dunes, because it is well-sorted and, in the absence of organic crusting, not subject to aggregation in the field. The soil may be less-representative of *in situ* field conditions as the degree of aggregation and sorting may have been changed during the preparation process, however the soil experiments are comparable with each other in relative terms. The microplastics were (i) fluorescent, red, polyethylene microspheres in the size range 212-250 µm with a density of 1.2 g cm³ (hereafter 'microbeads'), and chosen to be similar in size/shape, but lower in density to the sand bed; and (ii) 5 mm long polyester fibres (hereafter 'fibres') having a width of 0.5-1 mm and a density of 1.38 g cm³. Plastics were mixed with the substrate at concentrations of 40, 240, 640 and 1040 mg kg⁻¹_{dw}. For the microbeads, the associated range in particle count per kg is from 6200 to 160,000 (particle count for fibres could not be calculated). The lower concentrations are comparable with field measurements (Scheurer and Bigalke, 2018; Zhang et al., 2016) and predicted (Amec Foster Wheeler Environment & Infrastructure, 2017) environmental concentrations for agricultural soils, whilst the highest is at the lower end of that reported for industrial areas (Fuller and Gautam, 2016).

<< **Figure 1:** (A) Particle size distribution of substrate one (sand) and (B) substrate 2 (soil). >>

2.2 Methods

For each experiment, test surfaces were prepared by filling a 0.025 m by 0.35 m by 1.0 m metal tray with the given mixture of test material (substrate, or substrate + microplastic), which was prepared by weighing and manually mixing the appropriate ratios of substrate to plastic (Table 3). The test material was levelled such that the substrate surface was level with the tunnel floor and finally an airtight seal was secured around the tray perimeter. The upwind fetch was 8 m, wherein the flow travelled over a loose bed of randomly distributed gravel to adjust and achieve the desired aerodynamic roughness upon reaching the test bed.

For each threshold experiment, the freestream wind speed was increased from well below threshold up to 14 m s^{-1} in 0.25 m s^{-1} increments to determine wind speeds for the inception of intermittent and continuous saltation (Davidson-Arnott and Bauer, 2009); i.e. particles moving across the bed surface in low ballistic trajectories. The freestream wind velocity was sampled throughout the experiments at an elevation of 0.35 m using a micro pitot tube positioned 1.42 m upwind of the test bed. Freestream wind speed was controlled by a stepper motor with each “step” lasting 30 seconds with this stepping wind speed pattern being typical of emissions threshold testing for dust and soils (Ogungbemide, 2017). Experiments were performed in triplicate for each test material (ratio of substrate to plastic). Vertical velocity profiles were collected at each wind speed step and used to calculate the aerodynamic roughness and friction velocity by fitting the velocity data to the Law of the Wall equation.

Particle transport was approximated by particle counts which were measured using two fork-type Wenglor™ gate sensors secured to the bed 2 cm downwind of the test surface. These sensors were attached equidistant from the walls of the tunnel, and particle counts were measured approximately 0.5 cm above the surface. Due to natural variability in the streamwise distribution of particle transport, data from both Wenglors were summed during analysis.

Trapped samples of mobilised material were retrieved after each run at distances of 5, 30, 60, 100, 150 and 200 cm downwind of the test bed using glycerol-coated microscope slides attached to the wind tunnel base. The trapped samples were used to determine microplastic enrichment ratios (Sharpley, 1985) – the ratio of microplastic content in the particulate matter transported by creep and saltation to that in the original substrate.

Published relationships between saltation and suspension are expected to provide insight into long distance microplastic transport potential.

2.3 Quantification of Microplastic Entrainment

Although each wind tunnel experiment was repeated three times, quantities of entrained sediment and microplastics were small and therefore the material captured on the glycerol-coated slides was bulked together for replicate tests. Particles trapped on the slides were transferred onto cross hatched filter papers and distributed to ensure that samples were only one grain thick. Where large amounts of sediment were transported, multiple filter papers were used to ensure a single thickness of grains. Averages were taken of the derived data if more than one filter paper was needed per sample. Images of the filter papers were captured using a Leica S9D stereomicroscope microscope. The images were imported into ImageJ, a free image processing and analysis software package. The version used in this work was ImageJ 1.8.0_172 with the extension plug-in *Threshold Colour*.

For each image a central sub region was cropped and the scale applied such that images were scaled to their actual size. The total area of the image covered by sediment was calculated using the Threshold Colour plug-in. This plugin can threshold a specific colour from an image by selecting a range of values within the HSB (Hue, Saturation, Brightness) range. Given photographs of the samples were taken under constant light conditions, once the HSB value was derived for the first image it was kept constant for all subsequent analysis. Following this thresholding procedure, a binary transformation was applied to separate the selected colour from all other pixels. The surface area of the thresholded pixels was then automatically calculated using ImageJ.

In addition to the total area covered by sediment in each image, the total number of microplastic particles was manually counted. The microplastic spheres were of a known calibrated diameter of 212 μm so once the total number per image was known the total area they occupied could be calculated. A greater range in sizes was observed for the fibres so for each image alongside the total number of fibres, the length, width and total area of individual fibres was also measured. For all fibres analysed the median area was 3.8 mm^2 , mean area was 4.35 mm^2 and the standard deviation of the area was 4.36 mm^2 . In the results presented herein the median area was multiplied by the number of fibres to give the total area per image.

3. Results

3.1 Velocity-flux profiles

For the test beds without microplastics, the threshold for intermittent saltation was 7.25-7.5 m s^{-1} for sand and 11-12 m s^{-1} for soil (Table 4). The threshold for continuous saltation was higher in each case at 7.5-8 m s^{-1} for sand and 12.5-13.5 m s^{-1} for soil. Mineral particles and microplastics were entrained and transported during all the experiments. For the sand and soil test beds including microplastics intermittent saltation occurred in the range 6.75 to 7.5 m s^{-1} and 10.0 to 13.5 m s^{-1} , respectively, as compared to the higher wind speeds required for continuous saltation which were 7.5 to 8 m s^{-1} (sand) and 11.5 to 14 m s^{-1} (soil)(Table 4). When compared to either of the untreated test beds, the inclusion of microplastics was not found to significantly affect the wind erosion threshold (continuous or intermittent) for any of the concentrations or geometric forms (fibres or beads) tested (Table 4).

The velocity-flux profiles for Run 1 of each experiment are shown in Figure 2. For the sand beds the similarity of threshold is evident but there are differences in the relationship between total particle count (mineral particle + microplastic) and wind velocity. Above threshold for a given wind speed the flux over the sand bed is higher for experiments including microplastics than for sand without microplastics, and higher where the microplastics are fibres than for experiments including beads. For example, for a wind speed of 7.75 m s^{-1} the flux is $7.6 \text{ counts cm}^{-1}\text{s}^{-1}$ for the pure sand bed, $18\text{-}16.5 \text{ counts cm}^{-1}\text{s}^{-1}$ for the sand bed including microplastics, and $11\text{-}31.5 \text{ counts cm}^{-1}\text{s}^{-1}$ for the sand bed including fibres. Experiments using the soil bed are less systematically variable. At low wind speeds the flux fluctuates but following the onset of continuous saltation the soil beds including fibres generally have a higher flux than those containing beads for the same wind speed. One exception is the run with the lowest concentration of fibres ($40 \text{ mg kg}^{-1}_{\text{dw}}$), where flux is lower for beds including microplastics than for the soil bed with no microplastics added.

<< **Figure 2:** Relationship between wind speed and flux (mineral particle + microplastics) for A) sand bed, and B) soil bed. >>

3.2 Microplastic Enrichment Ratios

The susceptibility of microplastic particles to wind entrainment was determined by calculating the ratio of microplastic content in the transported material to that in the original substrate. The mean number of microplastic particles transported in each experiment ranged from 11 to 78 for the microbeads and 3 to 191 for fibres. Averaged over replicates of all concentrations of microplastics, distances and both substrate types, microplastic enrichment was lower for microbeads than fibres (Figure 3; Table 3). For the soil tests, the enrichment ratio for microbead entrainment was lowest and close to 1

for most concentrations. This suggests the relative quantity of microbeads transported is similar to that for natural particles in the soil population. Enrichment ratios for microbeads entrained from sands ranged from 5.03 for the lowest concentration to 3.12 for the highest concentration. Notably, the enrichment of microplastic fibres within the entrained particulate matter was one to two orders of magnitude higher for both test bed substrates, ranging from 98.15 to 498.25 for the sand and 278.14 to 726 for the soil (Table 3).

Figure 3. Summary of the ratio of microplastic content in eroded material compared to that in the original substrate. Box indicates median (centre line), 25th and 75th percentiles with outliers (>1.5 times inter-quartile range) indicated by '+'

In order to determine the likelihood of microplastic particles being transported long distances entrained material was trapped at differences downwind, up to 2 metres from the downwind edge of each test bed. The enrichment ratios for each distance downwind do not demonstrate any significant or systematic variability which suggests once entrained into the airflow particles may be transported away from the source (Figure 4; Table 5).

Figure 4: Change in microplastic enrichment ratio with distance downwind of the test bed for (a) sand, and (b) soil. Lines indicate best-fit regression. R^2 values in Table 4.

4. Discussion

Microplastics have recently been detected in the atmosphere, and in remote sites where the most likely source of contamination is via atmospheric deposition (Zhang et al., 2020). There has been very little research to date concerning the entrainment of microplastics by wind at source, despite the fact that microplastic occurrence can be high in soils susceptible to wind erosion (Rezaei et al., 2019). Similarly, very little is known about the processes governing airborne microplastic transport (Horton & Dixon, 2017). The data presented in this paper provide insights for a better understanding of the entrainment, transport and deposition of microplastics by wind when considered alongside previous research relating to the effects of particle density and shape on wind erosion and dust emission.

The low density of plastics is expected to affect both their entrainment potential at source and settling velocity, which will determine long-distance transport potential and deposition rates. In this experiment, the density of microplastic particles ($1.2\text{--}1.38\text{ g cm}^3$) is similar to that of organic soil components, which range from $1.0\text{--}1.5\text{ g cm}^3$ depending on the degree of decomposition (Zenchelsky et al., 1976; Skopp, 2000). Wind-eroded material from soils is often enriched in organic matter ($\text{ER} = 1$ to 10) as compared with the substrate from which it is entrained (Rühlmann et al., 2006; Webb et al., 2013; Iturri et al., 2017). Whilst this enrichment varies on an event-by-event basis (Panebianco et al., 2016), a common observation is that soil organic carbon enrichment is greater for soils with a high sand content when compared to clay-rich soil (Webb et al., 2013; Panebianco et al., 2016). This outcome is consistent with the experimental results presented here for microbead entrainment where the ER is slightly higher for the sand as compared to the soil. Microbead ERs observed here are similar in magnitude to ERs measured in the only published field study of microplastic wind entrainment at source (Rezaei et al., 2019) (ER

= 0 to 7.63), which also identifies a positive relationship between microplastic enrichment and sand content, but does not provide information on particle shape. In contrast, the ER outcomes for fibre entrainment (Table 3) have a less clear association with substrate type where enrichment is generally, but not always, higher for soil rather than sand. Building on existing wind erosion studies of organic matter enrichment as an analogue, future research carried out in the laboratory and field should aim to quantify the relationship between microplastic enrichment and increasing wind speed, given the mobilisation of larger soil particle diameters with increased fluid drag.

A substantial unknown is the effect of low particle density on the vertical displacement of particles at the soil surface via saltation impact. Saltation is well-known as a process for driving not only the mobilization of surface particles by creep and dynamic saltation, but also for the ejection of small and/or light particles higher in to the airstream where it is transported via suspension and may amplify long range transport (Iturri et al., 2017). The impact of particle density on the rate and nature of aeolian particle transport is poorly understood particularly for mixed substrates (Sherman, 2020). In a study to consider the behaviour of snow particles, Gordon and McKenna Neuman (2011) compared the particle splash dynamics of quartz sand (density 2.63 g cm^{-3}) and acrylic particles with a density of 1.21 g cm^{-3} and mean diameter of 192 μm . They found the acrylic particles impacted the bed surface at lower velocities and angles but caused the ejection of a greater number of particles than the sand particles. The shape of the particles was argued to be important due to its impact on particle packing and bed porosity. Owing to their low density, we suggest that microplastic particles are unlikely to have sufficient kinetic energy to eject mineral particles from the bed surface and thereby contribute to the development and maintenance of the saltation cloud. This remains to be confirmed, however, as no statistics exist on the number, launch angle and velocity components for microplastic

particles splashed from a mixed bed surface as a consequence of particle impact. Analogous to dust emission modelling, such information is required for estimating the sandblasting efficiency associated with release of microplastic particles to the atmosphere. Microplastic shape is also likely to influence the transfer of momentum from dynamic to static particles where compact (spherical) particles may be more effective than pliant (fibrous) particles. The relative proportion of mineral and microplastic particles would there be expected to affect overall aeolian transport.

With regards to the effect of particle shape, existing studies of microplastic deposition in both cities (Dris et al., 2016) and at remote mountain sites (Free et al., 2014; Allen et al., 2019) have found that fragments, films and fibres are more common than beads and pellets. Results presented in this paper indicate that the entrainment of microbeads is similar to, or slightly higher than, the entrainment of sand or soil, but that fibres are considerably more likely to be entrained than either substrate. Although it was not possible explicitly to differentiate whether microplastic particles were entrained at lower wind speeds than mineral particles in the experiments reported here, Figure 2 suggests that in the case of the sand bed this is likely to have been the case. Particle counts and higher flux rates occurred at lower wind speeds for sand beds containing microplastics than those without. This may be because the bed was loose and well-sorted and the low density microplastic particles were more readily entrained than the mineral particles at low wind speeds between intermittent and continuous saltation. Once the wind speed is above the continuous saltation threshold the whole bed, including both sand and microplastics, is mobilised. The dynamics of the soil beds are somewhat different to those of the sand beds because the poorly-sorted soil substrate is more susceptible to armouring as winnowing of fine particles takes place. Spherical microbeads are more likely to be sieved down through the soil during the armouring process which would

protect them from erosion whereas microplastic fibres may be less susceptible to sieving due to their shape and pliability. This may help to explain why fibres are more likely to be entrained but more research will be required to establish this. It is likely that fibres lying horizontally on the soil surface can be easily entrained but those caught in sediment interspaces and partially buried are likely to require higher shear stresses to be mobilized (Waldschläger and Schüttrumpf, 2019). Waldschläger and Schüttrumpf, (2019) suggest that within water the drag force is probably more important for the erosion of beads and the lift force more important for fibres and this may be the same for transport in air. The preferential entrainment of fibrous microplastics may also be due to their impact on soil porosity (de Souza Machado et al., 2018; Zhang et al., 2019).

Particle shape can be summarised by aspect ratio (λ – the ratio of particle length to width) and for natural mineral particles λ is usually around 1.5 with a maximum of 5, although for some mineralogies λ of up to 10 can occur (Ginoux, 2003). For microplastics, λ varies substantially depending on type and origin. For the present study, $\lambda = 1$ for the microbeads, and for the fibres the modal $\lambda = 5$ and mean $\lambda = 8$ (range 1-40). Particles with high λ generally have an increased surface area to volume ratio, which for microplastics descending in air, increases the opposing fluid drag relative to the gravitational force. Neglecting particle interactions and Magnus effects (spin), the higher drag associated with aspherical particles contributes to lower settling velocities, meaning that once entrained the particles stay in the atmosphere for longer periods of time than do spherical particles (Yang et al., 2003). Particle size modifies this effect to some extent such that for a given shape, settling velocity is decreased less for small particles ($<5 \mu\text{m}$) than larger ones. Ginoux (2003) determined that for $100 \mu\text{m}$ particles with $\lambda = 10$, the settling velocity was close to an order of magnitude slower than for a sphere of the same size. This has implications for the residence times of particles in the atmosphere, where

$\lambda = 10$ may result in a doubling of particle lifetime due to gravitational settling and suggests that once microplastic fibres have been entrained they are more likely to travel longer distances than plastic microbeads.

Allen et al. (2019) estimated the distance over which microplastics deposited in a remote mountain catchment might have travelled (by assuming a settling velocity equivalent to that for a 25 μm dust particle) to be up to 95 km. Using the same approach, and accounting for the size and density of the microbeads used here (particle diameter 250 μm , particle density 1.2 g cm^3), we estimate that dispersion up to 126 km from the source could occur for wind speeds averaging 7 m s^{-1} . It is not yet possible to do the same calculation for fibrous microplastics and given the likely influence of particle shape on suspension and settling, we suggest that microplastic shape needs to be carefully parameterized to predict the transport distance for microplastic fibres and fragments.

5. Conclusions

Novel laboratory experiments have quantified, for the first time, the extent to which microplastic particles are preferentially-transported compared to natural sediments. Specifically, using sand and soil substrates containing either microplastic beads or fibres, wind tunnel experiments were used to determine the impact of substrate and shape on microplastic entrainment. At the concentrations tested here, results show that although the inclusion of microplastics within a substrate was not found significantly to affect wind erosion thresholds, overall particle flux for beds containing plastic particles was higher in conditions above threshold. There is an enrichment of both types of microplastic particles within the entrained particulate matter. This enrichment is of one to two orders of magnitude for the microplastic fibres from both test bed substrates. The enrichment ratios

452 of plastic particles do not demonstrate any significant or systematic variability with
453 distance down wind (over 2 m) which suggests once entrained in to the air flow, particles
454 may be transported away from source.

455
456 The results reported here have provided first insights into the entrainment and transport
457 of microplastics from source substrates but questions remain as to the detailed particle-
458 scale processes by which microplastics are entrained in to the airflow. Considerably more
459 research into the roles of microplastic shape and density on their susceptibility to wind
460 erosion and long distance transport needs to be undertaken to effectively parameterize
461 models of atmospheric microplastic transport. Previous investigations into the wind
462 erosion and modelling of soil organic carbon enrichment and dispersal may provide a
463 useful starting point for future research in to airborne microplastics.

References

- Abbasi, S., Kechavarzi, B., Moore, F., Turner, A., Kelly, F.J., Dominguez, A.O., Jaafarzadeh, N. 2019. Distribution and potential health impacts of microplastics and microrubbers in air and street dusts from Asaluyeh County, Iran. *Environmental Pollution*, 244, 153-164.
- Allen, S., Allen, D., Phoenix, V.R., Le Roux, G., Durántez Jiménez, P., Simmoneau, A., Binet, S., Galop, D. 2019. Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nature Geoscience*, doi: 10.1038/s41561-019-0335-5
- Ambrosini, R., Azzoni, R.S., Pillino, F., Diolaiuti, G., Franzetti, A., Parolini, M. 2019. First evidence of microplastic contamination in the supraglacial debris of an alpine glacier. *Environmental Pollution* 253, 197-301
- Amec Foster Wheeler Environment & Infrastructure UK Ltd. Intentionally added microplastics in products – final report for European Commission (DG Environment). Amec Foster Wheeler Environment & Infrastructure UK Ltd. 2017.
- Andrady, A.L. 2011. Microplastics in the marine environment. *Marine Pollution Bulletin*, 62, 1596-1605.
- Arias-Andres, M., Kettner, M.T., Miki, T., Grossar, H.P. 2018. Microplastics: new substrates for heterotrophic activity contribute to altering organic matter cycles in aquatic ecosystems. *Science of the Total Environment*, 635, 1152-1159.
- Barnes, D.K.A., Galgani, F., Thompson, R.C., Marlaz, M. 2009. Accumulation and fragmentation of plastic debris in global environments. *Philosophical Transactions of the Royal Society B* 364, 1985-1990
- Beaumont, N.J., Aanesen, M., Austen, M.C., Borger, T., Clarke, J.R., Cole, M., Hooper, T., Lindeque, P.K., Pasco, C., Wyles, K.J. 2019. Global ecological, social and economic impacts of marine plastic. *Marine Pollution Bulletin*, 142, 189-195.
- Bergmann, M., Mützel, M.S., Primpke, S., Tekman, M.B., Trachsel, J., Gerdts, G. 2019. White and wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Science Advances* 5, eaax1157

493 Boots, B., Russell, C.W. and Green, D.S. 2019. Effects of microplastics in soil
 494 ecosystems: above and below ground. *Environmental Science and Technology*,
 495 53, 19, 11496-11506.

496 Boucher, J., Friot, D. 2017. *Primary microplastics in the oceans: a global evaluation of*
 497 *sources*, IUCN, Switzerland.

498 Cai, L., Wang, J., Peng, J., Tan, Z., Zhan, Z., Tan, X., Chen, Q. 2017. Characteristic of
 499 microplastics in the atmospheric fallout from Dongguan city, China: preliminary
 500 research and first evidence. *Environmental Science Pollution Research* 24, 24928-
 501 24935

502 Carbery, M., O'Connor, W., Thavamani, P. 2018. Trophic transfer of microplastics and
 503 mixed contaminants in the marine food web and implications for human health.
 504 *Environment International*, 115, 400-409.

505 Constant, M., Kerhervé, P., Mino-Vercellio-Verollet, M., Dumontier, M., Sánchez Vidal,
 506 A., Canals, M., Heissner, S. 2019. Beached microplastics in the Northwestern
 507 Mediterranean Sea, *Marine Pollution Bulletin* 142, 264-273

508 Davidson-Arnott, R.G.D., Bauer, O. 2009. Aeolian sediment transport on a beach:
 509 thresholds, intermittency, and high frequency variability. *Geomorphology* 105, 117-
 510 126

511 Dehghani, S., Moore, F., Akhbarizadeh, R. 2017. Microplastic pollution in deposited urban
 512 dust, Tehran metropolis, Iran. *Environmental Science and Pollution Research*, 24,
 513 20360-20371.

514 de Souza Machado, A.A., Lau, C.W., Till, J., Kloas, W., Lehmann, A., Becker, R., Rillig,
 515 M.C. 2018. Impacts of microplastics on the soil biophysical environment.
 516 *Environmental Science and Technology*, 52, 9656-9665.

517 Dris, R., Gasperi, C.J., Rocher, A.V., Saadi, B.M., Renault, N., Tassin, B. 2015.
 518 Microplastic contamination in an urban area: a case study in Greater Paris.
 519 *Environmental Chemistry*, 12, 592-599.

520 Dris, R., Gasperi, J., Saad, M., Mirande, C., Tassin, B. 2016. Synthetic fibers in
 521 atmospheric fallout: a source of microplastics in the environment? *Marine Pollution*
 522 *Bulletin* 104, 290-293

523 Dris, R., Gasperi, J., Mirande, C., Mandin, C., Guerrouache, M., Langlois, V., Tassin, B.
524 2017. A first overview of textile fibers, including microplastics, in indoor and outdoor
525 environments. *Environmental Pollution*, 221, 453-458.

526 Evangeliou, N., Grythe, H., Klimont, Z., Heyes, C., Eckhardt, C., Lopez-Aparicio, S.,
527 Stohl, A. 2020. Atmospheric transport is a major pathway of microplastics to remote
528 regions. *Nature Communications*, 11, 3381.

529 Free, C.M., Jensen, O.P., Mason, S.A., Eriksen, M., Williamson, N.J., Boldgiv, B. 2014.
530 High-levels of microplastic pollution in a large, remote, mountain lake. *Marine*
531 *Pollution Bulletin* 85, 156-163

532 Folk, R.L., Ward, W.C. 1957. Brazos River bar: a study in the significance of grain size
533 parameters. *Journal of Sedimentary Petrology*, 27, 3-26.

534 Fuller, S., Gautam, A. 2016. procedure for measuring microplastics using pressurized
535 fluid extraction. *Environmental Science and Technology* 50, 5774-5780

536 Gasperi, J., Wright, S.L., Dris, R., Collard, F., Mandin, C., Guerrouache, M., Langlois, V.,
537 Kelly, F.J., Tassin, B. 2018. Microplastics in air: are we breathing it in? *Current*
538 *Opinion in Environmental Science and Health* 1, 1-15

539 Ginoux, P. 2003. Effects of nonsphericity on mineral dust modelling. *Journal of*
540 *Geophysical Research* 108

541 He, D., Luo, Y., Lu, S., Liu, M., Song, Y., Lei, L. 2018. Microplastics in soils: analytical
542 methods, pollution characteristics and ecological risks. *Trends in Analytical*
543 *Chemistry*, 109, 163-172.

544 Hidalgo-Ruz, V., Gutow, L., Thompson, R.C. Thiel, M. 2012. Microplastics in the marine
545 environment: a review of the methods used for identification and quantification.
546 *Environmental Science and Technology*, 46, 6, 3060-3075.

547 Horton, A., Dixon, S.J. 2017. Microplastics: an introduction to environmental transport
548 processes. *WIREs Water*, 2018, 5, e1268.

549 Hu, D., Shen, M., Zhang, Y., Li, H., Zeng, G. 2019. Microplastics and nanoplastics: would
550 they affect global biodiversity change? *Environmental Science and Pollution*
551 *Research*, 26, 19997-20002.

552 Iturri, L.A., Funk, R., Leue, M., Sommer, M., Buschiazzi, D.E. 2017. Wind sorting affects
553 differently the organo-mineral composition of saltating and particulate materials in
554 contrasting texture agricultural soils. *Aeolian Research* 28, 39-49

555 Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan,
556 R., Law, K.L. 2015. Plastic waste inputs from land into the ocean. *Science*, 347,
557 6223, 768-771.

558 Klein, M., Fischer, E.K. 2019. Microplastic abundance in atmospheric deposition within the
559 Metropolitan area of Hamburg, Germany. *Science of the Total Environment* 685, 96-
560 103

561 Lehmann, A., Fritschen, K., Rillig, M. 2019. Abiotic and biotic factors influencing the effect
562 of microplastic on soil aggregation. *Soil Systems*, 3, 21.

563 Li, B., McKenna Neuman, C., 2012. Boundary-layer turbulence characteristics during
564 aeolian saltation. *Geophysical Research Letters* 39. DOI: 10.1029/2012GL052234

565 Li, C.R., Busquets, R., Campos, L.C. 2020. Assessment of microplastics in freshwater
566 systems : a review. *Science of the Total Environment*, 707, art. no. 135578, doi :
567 10.1016/j.scitotenv.2019.135578.

568 Liu K., Wang, X., Fang, T., Xu, P., Zhu, L., Li, D. 2019. Source and potential risk
569 assessment of suspended atmospheric microplastics in Shanghai. *Science of the*
570 *Total Environment*, 675, 462-471.

571 Lwanga, E.H., Gertsen, H., Gooren, H., Peters, P., Salánki, T., van der Ploeg, M.,
572 Besseling, E., Koelmans, A.A., Geissen, V. 2016. *Environmental Science and*
573 *Technology*, 50, 2685-2691.

574 Machado, A., Lau, C.W., Till, J., Kloas, W., Lehmann, A., Becker, R., Rillig, M.C. 2018.
575 Impacts of microplastics on the soil biophysical environment. *Environmental*
576 *Science and Technology*, 52, 9656-9665.

577 Mailhot, B., Morlat, S., Gardette, J.-L. 2000. Photooxidation of blends of polystyrene and
578 poly_vinyl methyl ether): FTIR and AFM studies. *Polymer*, 41, 1981-1988.

579 McCormick, A., Hoellein, T.J., Mason, S.A., Schluep, J., Kelly, J.J. 2014. Microplastic is
580 an abundant and distinct microbial habitat in an urban river. *Environmental Science*
581 *and Technology* 48, 11863-11871

582 McKenna Neuman, C., Nickling, W. G. 2000. Aeolian transport measurements:
583 instruments and techniques. *Proceedings from the CCRG Workshop on the*
584 *Measurement and Simulation of Sediment Transport by Wind* Trent University,
585 Ontario

586 Nickling, W., McKenna Neuman, C., 1997. Wind tunnel evaluation of a wedge-shaped
587 aeolian. *Geomorphology* 18, 333-345

588 North, E.J., Halden, R.U. 2013. Plastics and environmental health: the road ahead.
589 *Reviews on Environmental Health*, 28, 1,

590 Ogungbemde, D., 2017. Wind tunnel simulation and modelling of dust emission from
591 mine tailings (nepheline syenite), PhD Dissertation, Trent University.

592 PlasticsEurope, *Plastics–The Facts 2019, An Analysis of European Plastics Production,*
593 *Demand and Waste Data*, PlasticsEurope, Brussels, Belgium 2019.

594 Panebianco, J.E., Mendez, M.J., Buschiazzo, D.E. 2016. PM10 emission, sandblasting
595 efficiency and vertical entrainment during successive wind-erosion events: a wind-
596 tunnel approach. *Boundary-layer Meteorology* 161, 335-353

597 Rezaei, M., Riksen, M.J.P.N., Sirjani, E., Sameni, A., Geissen, V. 2019. Wind erosion as
598 a driver for transport of light density microplastics. *Science of the Total Environment*
599 669, 273-281

600 Roblin, B., Aherne, J. 2020. Moss as a biomonitor for the atmospheric deposition of
601 anthropogenic microfibres. *Science of the Total Environment*, 715, 136973.

602 Rochman, C.M., Brookson, C., Bikker, J., Djuric, N., Earn, A., Bucci, K., Athey, S.,
603 Huntington, A., McIlwraith, H., Munno, K., De Frond, H., Kolomijeca, A., Erdle, L.,
604 Grbic, J., Bayoumi, M., Borrelle, S.B., Wu, T., Santoro, S., Werbowski, L.M., Zhu,
605 X., Giles, R.K., Hamilton, B.M., Thaysen, C., Kaura, A., Klasios, N., Ead, L., Kim, J.,
606 Sherlock, C., Ho, A., Hung, C. 2019. Rethinking microplastics as a diverse
607 contaminant suite. *Environmental Toxicology and Chemistry*, 38, 4, 703-711.

608 Rühlmann, J., Körschens, M., Graefe, J. 2006. A new approach to calculate the particle
609 density of soils considering properties of the soil organic matter and the mineral
610 matrix. *Geoderma* 130, 272-283

611 Scheurer, M., Bigalke, M. 2018. Microplastics in Swiss floodplain soils. *Environmental*
612 *Science and Technology* 52, 3591-3598

613 Sharpley, A.N. 1985. The selective erosion of plant nutrients in runoff. *Soil Science*
614 *Society of America Journal* 49, 1527-1534

615 Sherman, D.J. 2020. Understanding wind-blown sand: six vexations. *Geomorphology*,
616 366, 107193.

617 Skopp, J.M. 2000. Physical properties of primary particles. *Handbook of Soil Science* H.
618 E. Summer, Ed. (CRC Press, Boca Raton, Florida, PP. A3-A17)

619 Song, Y.K., Hong, S.H., Jang, M., Han, G.M., Jung, S.W., Shim, W.J., 2017. Combined
620 effects of UV exposure duration and mechanical abrasion on microplastic
621 fragmentation by polymer type. *Environmental Science and Technology* 8, 4368-
622 4376

623 Stanton, T., Matthew, J., Nathanail, P., MacNaughtan, W., Gomes, R.L. 2019. Freshwater
624 and airborne textile fibre populations are dominated by 'natural', not microplastic
625 fibres. *Science of the Total Environment*, 666, 377-389.

626 Tanaka, K., Takada, H. 2016. Microplastic fragments and microbeads in digestive tracts
627 of planktivorous fish from urban coastal waters. *Scientific Reports*, 6, 34351.

628 von Moos, N., Burkhardt-Holm, P., Köhler, A. 2012. Uptake and effects of microplastics
629 on cells and tissue of the Blue Mussel *Mytilus edulis* L. after an experimental
630 exposure. *Environmental Science and Technology*, 46, 11327-11335.

631 Waldschläger, K., Lechthaler, S., Stauch, G., Schtüttrumpf, H. 2020. The way of
632 microplastic through the environment – application of the source-pathway-receptor
633 model (review). *Science of the Total Environment*, 713, 136584.

634 Waldschläger, K., Schtüttrumpf, H. 2020. Erosion behavior of different microplastic
635 particles in comparison to natural sediments. *Environmental Science and*
636 *Technology*, 53, 22, 13219-13227.

637 Wang, W.F., Ge, J., Yu, X.Y., Li, H. 2020. Environmental fate and impacts of
638 microplastics in soil ecosystems: progress and perspective. *Science of the Total*
639 *Environment*, 708, art. No. 134841, doiL 10.1016/j.scitotenv.2019.134841.

640 Webb, N.P., Strong, C.L., Chappell, A., Marx, S.K., McTainsh, G.H. 2013. Soil organic
641 carbon enrichment of dust emissions: magnitude, mechanisms and its implications
642 for the carbon cycle. *Earth Surface Processes and Landforms* 38, 1662-1671

643 Wright, S.L, Thompson, R.C., Galloway, T.S. 2013. The physical impacts of microplastics
644 on marine organisms: a review. *Environmental Pollution* 178, 483-492

645 Wright, S.L., Ulke, J., Font, A., Chan, K.L.A., Kelly, F.J. 2020. Atmospheric microplastic
646 deposition in an urban environment and an evaluation of transport. *Environment*
647 *International*, 136, 105411.

648 Wright, S.L., Kelly, F.J. 2017. Plastic and human health: a micro issue? *Environmental*
649 *Science and Technology*, 51, 6634-6647.

650 Yang, W., Marshak, A., Kostinski, A.B., Várnai, T. 2003. Shape-induced gravitational
651 sorting of Saharan dust during transatlantic voyage: evidence from CALIOP lidar
652 depolarization measurements. *Geophysical Research Letters* 40, 3281-3286

653 Zenchelsky, S.T., Delany, A.C., Pickett, R.A. 1976. The organic component of wind-blown
654 soil aerosol as a function of wind velocity. *Soil Science* 122, 129-132.

655 Yuan, J., Ma, J., Sun, Y., Zhou, T., Zhao, Y., Yu, F. 2020. Microbial degradation and other
656 environmental aspects of microplastics/plastics. *Science of the Total Environment*,
657 715, 136968.

658 Zalasiewicz, J., Waters, C.N., Ivar do Sul, J.A., Corcoran, P.L., Barnosky, A.D., Cearreta,
659 A., Edgeworth, M., Galuszka, A., Jeandel, C., Leinfelder, R., McNeill, J.R. Steffen,
660 W., Summerhayes, C., Wapreigh, M., Williams, M., Wolfe, A.P., Yonan, Y. 2016.
661 The geological cycle of plastics and their use as a stratigraphic indicator of the
662 Anthropocene. *Anthropocene*, 13, 4-17.

663 Zhang, K., Su, J., Ziong, X., Wu, X., Wu, C., Liu, J. 2016. Microplastic pollution of
664 lakeshore sediments from remote lakes in Tibet plateau, China. *Environnemental*
665 *Pollution* 219, 450-455

666 Zhang, G.S., Zhang, F.X., Li, X.T. 2019. Effects of polyester microfibers on soil physical
667 properties: perception from a field and pot experiment. *Science of the Total*
668 *Environment*, 670, 107.

669 Zhang, Y., Kang, S., Allen, S., Allen, D., Tanguang, G., Sillanpää, M. 2020. Atmospheric
670 microplastics : a review on current status and perspectives. *Earth-Science*
671 *Reviews*, 203, 103118.

672 Zhou, Q., Zhang, H., Fu, C., Zhou, Y., Dai, Z., Li, Y., Tu, C., Luo, Y. 2018. The distribution
673 and morphology of microplastics in coastal soils adjacent to the Bohai Sea and the
674 Yellow Sea. *Geoderma*, 322, 201-208.

675

676 **Acknowledgments:**

677 The authors would like to thank Stacey Johnston and Taylor Brooks for their assistance
678 with the wind tunnel experiments. **Funding:** This research did not receive any specific
679 grant from external funding agencies in the public, commercial or not-for-profit sector.

680 **Author contributions:** All authors were involved in conceptualization and methodology.

681 PO'B and CMN analysed wind tunnel airflow and sediment transport data. JEB and AO
682 analysed microplastic and sediment samples. All authors interpreted data. All authors

683 wrote the manuscript. **Competing interests:** Authors declare no competing interests.

684 **Data and materials availability:** All data are available in the main text.

685

Table 1: Folk and Ward (1957) statistics for substrate one (sand)

	Geometric (μm)	Logarithmic (ϕ)	Description
Mean	254.2	1.976	Medium sand
Sorting	1.412	0.497	Well sorted
Skewness	-0.129	0.129	Fine skewed
Kurtosis	1.099	1.099	Mesokurtic

Table 2: Folk and Ward (1957) statistics for substrate two (soil)

	Geometric (μm)	Logarithmic (ϕ)	Description
Mean	671.5	0.574	Coarse sand
Sorting	2.789	1.480	Poorly sorted
Skewness	-0.025	0.025	Symmetrical
Kurtosis	0.832	0.832	Platykurtic

Table 3: Initial microplastic concentrations and average enrichment ratio (all distances) of microplastics in wind-eroded material.

Substrate	Concentration (mg kg ⁻¹ _{dw})	Ratio of microplastic:substrate	Average enrichment ratio (all distances)	
			Microbead	Fibre
Sand	40	1:25,000	5.03	129.05
	240	1:4167	5.38	498.25
	640	1:1563	3.84	98.15
	1040	1:962	3.12	193.64
Soil	40	1:25,000	3.49	726.40
	240	1:4167	1.59	278.14
	640	1:1563	0.89	606.67
	1040	1:962	1.06	305.40

Table 4: Thresholds for intermittent and continuous saltation for each substrate/microplastic combination and r^2 regression coefficients for best-fit relationships between enrichment ratio and distance downwind for different concentrations of microplastic (plotted in Figure 3 main text).

Substrate_microplastic type	Microplastic concentration (mg kg ⁻¹ _{dw})	Intermittent saltation threshold (m s ⁻¹)			Continuous saltation threshold (m s ⁻¹)			r^2 relationship between enrichment ratio and distance downwind
		Run 1	Run 2	Run 3	Run 1	Run 2	Run 3	
Sand (no microplastic)	0	7.5	7.25	7.5	8	7.5	8	-
Sand_microbead	1040	7.25	7.5	7.25	7.75	7.75	7.75	0.305
Sand_microbead	640	7.5	7.25	7.25	8	8	7.75	0.038
Sand_microbead	240	7.25	7.25	7.25	8	8	8	0.021
Sand_microbead	40	7.5	7.5	7	8	8	8	0.754
Sand_fibre	1040	7	7.25	7	7.75	8	7.75	0.556
Sand_fibre	640	6.75	7	7	7.75	7.75	7.5	0.086
Sand_fibre	240	7.5	7	7.25	8	7.75	7.75	0.438
Sand_fibre	40	7.5	7.5	7.25	7.75	8	8	0.017
Soil (no microplastic)	0	12	11	12	13	12.5	13.5	-
Soil_microbead	1040	12.5	11	11.5	13	12.5	13	0.011
Soil_microbead	640	13.5	13	11	14	13.5	12.5	0.197
Soil_microbead	240	12	10.5	11	13.5	12	12	0.138
Soil_microbead	40	12	11.5	11.5	13	12.5	13	0.255
Soil_fibre	1040	12	10.5	10	13	12	13	0.110
Soil_fibre	640	10	12	12	11.5	13.5	13.5	0.636
Soil_fibre	240	12	12	11	13.5	13	12	0.195
Soil_fibre	40	11	12	12	14	13	13	0.163

Table 5: Microplastic enrichment for each substrate/microplastic combination, concentration and distance.

Microplastic concentration (mg kg ⁻¹ _{dw})	Distance downwind (cm)	Enrichment ratio			
		Sand_microbead	Sand_fibre	Soil_microbead	Soil_fibre
1040	200	4.73	59.36	0.53	181.95
1040	150	2.14	143.83	1.39	221.07
1040	100	3.31	119.50	0.67	333.57
1040	60	3.44	348.13	2.30	332.26
1040	30	3.22	281.66	0.91	652.55
1040	5	1.86	209.38	0.55	111.00
640	200	4.10	76.98	0.37	241.75
640	150	3.73	93.12	1.23	422.07
640	100	3.34	92.51	0.95	431.65
640	60	5.34	109.16	0.55	621.91
640	30	5.02	153.46	0.77	574.40
640	5	1.51	63.64	1.47	1348.23
240	200	4.75	76.99	0.67	118.73
240	150	5.79	435.57	1.67	254.31
240	100	4.67	213.66	1.89	308.78
240	60	10.71	1049.70	0.50	302.81
240	30	4.06	393.94	3.52	531.05
240	5	2.16	819.63	1.28	153.17
40	200	8.94	150.34	0	778.83
40	150	7.47	0	2.03	0
40	100	3.08	0	5.02	598.38
40	60	6.17	204.26	7.19	987.48
40	30	30.08	0	5.09	0
40	5	1.46	419.72	1.63	1993.68