SUPPORTING INFORMATION

Preparation, isolation and characterization of N^α -Fmoc-peptide isocyanates: Solution synthesis of oligo- α -peptidyl ureas

Vommina V. Suresh Babu*, Basanagoud S. Patil, and Rao Venkataramanarao

*Department of studies in Chemistry, Central College Campus, Bangalore University,
Dr. B. R. Ambedkar Veedhi, Bangalore 560 001, INDIA*

e-mail: hariccb@rediffmail.com

Table of contents in supporting information

Page No.	Description
S3	Experimental-General
S4 & S5	General procedure: N^α -Fmoc-peptide acid azides
S6 – S11	Characteristic data for the N^α -Fmoc-peptide isocyanates
S11 – S13	General procedure : N^α -Fmoc-peptidyl urea acids 7a-c and characteristic data for 7a - 7b , 8a-c and 10a .
S14 – S15	General procedure: N^α -Fmoc-peptidyl urea esters 7d-f and characteristic data for 7d , 7f , and 10b .
S15 – S16	General procedure: Deprotection of Fmoc group from the N^α -Fmoc-peptidyl urea acids 12 and characteristic data for 12a , 12b , and 12d .
S16 – S19	Synthetic procedure for [Leu ⁵]enkephalin analogs and characteristic data for 13 , 14 , 15 and 18 .
S20 - S21	Synthetic procedure for Bioelastic polymer analogs and characteristic data for 22 and 23 .
S22 – S23	Table: N^α -Fmoc-peptide isocyanates
S24	IR spectra of Fmoc-Ser(Bu ^t)-Phe-CON ₃ 2o
S25	IR spectra of Fmoc-Val-Pro-NCO 3g

- S26** RP-HPLC of Fmoc-Val-Phe-CON₃ **2e** and
RP-HPLC of Fmoc-Val-Phe-NCO **3e**
- S27** Monitoring of Curtius rearrangement of **2g** to **3g** using IR spectrum
- S28** ¹H NMR spectra of Fmoc-Val-Ala-NCO **3b**
- S29** ¹³C NMR spectra of Fmoc-Val-Ala-NCO **3b**
- S30** MALDI-mass spectra of Fmoc-Val-Pro-Gly-NCO **4c**
- S31** ¹H NMR spectra of Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH **7a**
- S32** ¹³C NMR spectra of Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH **7a**
- S33** ES-MS of Fmoc- Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH **7a**
- S34** ¹H NMR spectra of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**
- S35** ¹³C spectra NMR of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**
- S36** ES-MS of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**
- S37** ¹³C NMR spectra of Tyr-Gly-Gly-ψ(NH-CO-NH)-Phe-Leu-OH **14**
- S38** MALDI-mass of Tyr-Gly-ψ(NH-CO-NH)-Gly-Phe-Leu-OH **13**
- S39** ES-MS of Tyr-Gly-ψ(NH-CO-NH)-Gly-ψ(NH-CO-NH)-Gly- ψ(NH-CO-NH)-Phe-ψ(NH-CO-NH)-Leu-OH **21**
- S40** ¹H NMR spectra of Val-Pro-Gly-ψ(NH-CO-NH)-Val-Gly-OH **22**
- S41** MALDI-mass spectra of Pro-Gly-Val-Gly-ψ(NH-CO-NH)-Val-OH **23a**
- S42** ¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-R-(+)-1-phenylethylamine **25a**
- S43** ¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-R-(+)-1-phenylethylamine **25a**
- S44** ¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-S-(-)-1-phenylethylamine **25b**

- S45** ^1H NMR spectra of Fmoc-Leu-Val- $\psi(\text{NH-CO-NH})$ -S-(-)-1-phenylethylamine **25b**
- S46** ^1H NMR spectra of Fmoc-Leu-Val- $\psi(\text{NH-CO-NH})$ -(R&S)-1-phenylethylamine **25c**
- S47** ^1H NMR spectra of Fmoc-Leu-Val- $\psi(\text{NH-CO-NH})$ -(R&S)-1-phenylethylamine **25c**
- S48** HR MS of Fmoc-Val-Ala-NCO **3b**
- S49** HR MS of Fmoc-Ala-Leu-NCO **3d**
- S50** HR MS of Fmoc-Ser(Bu^t)-Phe- $\psi(\text{NH-CO-NH})$ -Leu-OMe **7e**
- S51** HR MS of Fmoc-Leu-Val- $\psi(\text{NH-CO-NH})$ -S-(-)-1-phenylethylamine **25b**
- S52** HR MS of Fmoc-Val-Pro-Gly- $\psi(\text{NH-CO-NH})$ -Val-Gly-OH **22a**

Experimental Section

Melting points were determined using capillary method and are uncorrected. LG domestic microwave oven operating at 2450 MHz was used for the preparation of isocyanates. The ultrasonication reactions were carried out using a sonic bath (35 kHz, Elma, T 310 / H German make) at ambient temperature. The TLC was effected with silica gel GF254 on precoated glass plates using as mobile phases the following solvent systems : (a) Chloroform : Methanol : Acetic acid (40:2:1) and (b) Ethyl acetate : Hexane (7:3). All solvents were freshly distilled prior to use. The *N*^α-Fmoc-di, tri and tetrapeptide acids have been prepared via mixed anhydride method by employing *O,N*-bis-trimethylsilylamino acids using the reported procedures^{31,32} which were recrystallized and characterized prior to use. *N*^α-Fmoc-peptide acid azides were synthesized and characterized.³⁴

General procedure for the synthesis of *N*^α-Fmoc-peptide acid azides:

Method A: Employing mixed anhydride method

N^α-Fmoc-peptide acid (1 mmole) was dissolved in dry THF (5 mL) and cooled in an ice-salt bath. EtOCOC₂H₅ (0.1 mL, 1 mmole) and NMM (0.11 mL, 1 mmole) were added to the above cooled solution and stirred for 10 min. while maintaining the temperature at -10 °C. The resulting reaction mixture was treated with aqueous NaN₃ (0.1 g, 1.5 mmole in 1 mL of water) at the same temperature. The reaction, as monitored by TLC as well as IR analysis, was complete in about 15 min. The THF solution was evaporated under reduced pressure at r.t. and the residue was taken in CH₂Cl₂ (15 mL). The CH₂Cl₂ layer was washed with cold water (2 × 10 mL), dried over anhydrous Na₂SO₄ and concentrated

the half of the volume and *n*-hexane (10 mL) was added to obtain pure *N*^α-Fmoc-peptide acid azide as a crystalline solid.

Method B: Employing acid chloride method

To a cold solution of *N*^α-Fmoc-peptide acid chloride (1 mmole) in acetone (5 mL) was added aqueous NaN₃ (0.1 g, 1.5 mmole in 1 mL of water). The resulting reaction mixture was stirred for 15 min. at -10 °C. The separated solid was filtered and washed with cold water. The resulting solid was dissolved in CH₂Cl₂ (5 mL), dried over anhydrous Na₂SO₄ and crystallized using *n*-hexane.

Data for few selected *N*^α-Fmoc-peptide acid azides: Fmoc-Gly-Phe-N₃ 2a: Yield 92 % ; m.p. 205-06 °C ; IR 2142 cm⁻¹ ; Fmoc-Ala-Leu-N₃ 2d: Yield 89 % ; m.p. 176 °C ; IR 2139 cm⁻¹ ; Fmoc-Val-Phe-N₃ 2e: Yield 91 % ; m.p. 188 °C ; IR 2137 cm⁻¹ ; Fmoc-Gln(Trt)-Ile-N₃ 2j: Yield 84 % ; m.p. 160-62 °C ; IR 2138 cm⁻¹ ; Fmoc-Ser(Bu^t)-Phe-N₃ 2o: Yield 79 % ; m.p. 164-65 °C ; IR 2138 cm⁻¹. The HPLC analysis was carried out for Fmoc-Val-Phe-CON₃/NCO 2e/3e using isocratic system and the mobile phase was MeCN : water (60 : 40). It was confirmed that during the HPLC analysis the Fmoc-peptide acid azide and isocyanate were stable and it was confirmed by the presence of sharp peak for CON₃ and NCO groups in IR spectra (see the IR spectra and HPLC data in the supporting information).

Fmoc-Gly-Phe-NCO 3a: IR 2256 cm^{-1} ; ^1H NMR (δ , CDCl_3) 2.91 (2H, d), 3.85 (2H, m), 4.18-4.45 (4H, m), 5.34 (1H, d), 6.57 (1H, d), 7.2-7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 37.8, 43.2, 47.2, 51.9, 66.6, 120.0, 124.9, 126.6, 126.7, 126.9, 127.5, 128.8, 129.4, 137.6, 141.3, 143.6, 156.1, 175.4 ; ES MS m/z observed 442.5.

Fmoc-Val-Ala-NCO 3b: IR 2247 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.95 (6H, d), 1.38 (3H, d), 2.1 (1H, m), 3.9 (1H, m), 4.1-4.48 (4H, m), 5.35 (1H, br s), 6.5 (1H, s), 7.2-7.8 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 17.9, 19.1, 22.8, 30.1, 47.2, 57.1, 60.3, 67.2, 120.0, 124.9, 126.2, 127.1, 127.8, 141.3, 143.7, 156.5, 171.1 ; ES MS m/z observed 406.5.

Fmoc-Leu-Val-NCO 3c: IR 2257 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.86-0.98 (12H, m), 1.36 (2H, m), 1.57 (1H, m), 1.89 (1H, m), 3.68 (1H, m), 3.86 (1H, m), 4.2 (1H, t), 4.4 (2H, d), 5.36 (1H, d), 6.56 (1H, d), 7.3-7.8 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 18.4, 19.3, 21.9, 22.8, 24.6, 29.6, 41.2, 49.0, 56.2, 66.6, 112.0, 125.1, 126.4, 127.1, 127.7, 141.4, 143.9, 156.6, 170.8 ; ES MS m/z observed 450.3.

Fmoc-Ala-Leu-NCO 3d: IR 2251 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.92 (6H, d), 1.15 (3H, d), 1.42 (2H, m), 1.65 (1H, m), 3.89-4.45 (5H, m), 5.33 (1H, d), 6.59 (1H, d), 7.3-7.85 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 17.2, 22.0, 23.0, 24.6, 40.3, 47.2, 48.9, 51.2, 66.6, 119.9, 125.0, 126.2, 127.0, 127.6, 141.3, 143.8, 156.2, 171.3 ; ES MS m/z observed 422.4.

Fmoc-Val-Phe-NCO 3e: IR 2254 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.93 (6H, d), 1.84 (1H, m), 2.91 (2H, s), 3.8-4.2 (3H, m), 4.45 (2H, d), 5.35 (1H, d), 6.55 (1H, d), 7.2-7.8 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 17.2, 18.6, 19.5, 30.8, 47.3, 48.9, 58.5, 66.5, 120.0, 125.0, 126.1, 126.7, 127.1, 127.6, 128.6, 129.2, 137.5, 141.3, 143.8, 157.0, 171.3 ; ES MS m/z observed 484.5.

Fmoc-Val-Phg-NCO 3f: IR 2252 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.93 (6H, d), 1.84 (1H, m), 3.78-4.45 (5H, m), 5.32 (1H, d), 6.52 (1H, d), 7.2-7.8 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 18.4, 19.3, 29.6, 47.1, 51.9, 56.2, 66.6, 119.9, 125.0, 126.6, 126.9, 127.1, 127.6, 128.8, 129.3, 137.0, 141.2, 143.7, 156.0, 170.2 ; ES MS m/z observed 470.5.

Fmoc-Val-Pro-NCO 3g: IR 2251 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.95 (6H, d), 1.38–2.3 (5H, m), 3.61 (1H, m), 4.2 (1H, t), 4.42 (2H, m), 5.33 (1H, d), 7.3-7.8 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 18.4, 19.3, 24.2, 28.7, 29.6, 47.1, 47.2, 56.2, 60.6, 66.6, 119.9, 125.0, 126.3, 126.9, 127.6, 141.2, 143.7, 156.0, 170.2 ; ES MS m/z observed 434.5.

Fmoc-Ile-Ser(Bzl)-NCO 3h: IR 2245 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.92 (6H, m), 1.2 (1H, m), 1.82 (2H, m), 2.88 (2H, d), 3.58-4.44 (7H, m), 5.31 (1H, d), 6.52 (1H, d), 7.3-7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 11.3, 15.5, 25.4, 35.8, 37.3, 47.3, 51.8, 57.3, 62.1, 66.5, 119.9, 125.0, 126.4, 126.7, 127.0, 127.6, 128.6, 129.2, 137.5, 141.3, 143.8, 157.0, 170.9 ; ES MS m/z observed 528.5.

Fmoc-Asp(OBu^t)-Ser(Bzl)-NCO 3i: IR 2248 cm^{-1} ; ^1H NMR (δ , CDCl_3) 1.44 (9H, s), 2.52 (2H, d), 2.87 (2H, d), 3.35–3.84 (3H, m), 4.04 (1H, br s), 4.2 (1H, t), 4.4 (2H, m), 5.58 (1H, d), 6.65 (1H, m), 7.2–7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 27.9, 37.2, 37.3, 47.0, 49.9, 51.8, 62.2, 66.7, 81.3, 119.9, 124.9, 126.6, 126.7, 126.9, 127.6, 128.6, 129.2, 137.5, 141.1, 143.7, 156.2, 170.9, 170.1 ; ES MS m/z observed 586.5.

Fmoc-Cys(Bzl)-Val-NCO 3m: IR 2248 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.97 (6H, d), 1.92 (1H, m), 3.6-4.6 (9H, m), 5.45 (1H, d), 6.6 (1H, m), 7.2–7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 18.6, 19.5, 29.1, 47.3, 58.5, 66.6, 120.0, 125.0, 126.5, 126.9, 127.6, 141.2, 143.9, 156.4 ; ES MS m/z observed 530.4.

Fmoc-Ser(Bzl)-Phe-NCO 3n: IR 2258 cm^{-1} ; ^1H NMR (δ , CDCl_3) 2.85-2.94 (4H, m), 3.48 (2H, d), 3.95 (1H, m), 4.18-4.45 (4H, m), 5.35 (1H, d), 6.58 (1H, d), 7.12-7.9 (18H, m) ; ^{13}C NMR (δ , CDCl_3) 37.9, 38.1, 47.2, 47.2, 50.6, 51.9, 66.7, 119.7, 124.9, 126.2, 126.5, 126.6, 127.1, 127.3, 128.6, 128.8, 129.3, 129.3, 137.4, 137.6, 140.9, 143.5, 156.1, 174.1 ; ES MS m/z observed 562.6.

Fmoc-Ser(Bu^t)-Phe-NCO 3o: IR 2255 cm^{-1} ; ^1H NMR (δ , CDCl_3) 1.19 (9H, s), 2.9 (2H, d), 3.5-3.85 (3H, m), 4.18-4.45 (4H, d), 5.35 (1H, d), 6.58 (1H, d), 7.2-7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 27.0, 37.3, 46.7, 51.8, 54.1, 62.1, 66.5, 73.3, 119.9, 125.0, 126.4, 126.7, 127.0, 127.6, 128.7, 129.3, 137.4, 141.3, 143.8, 156.0, 170.6 ; ES MS m/z observed 528.5.

Fmoc-Thr(Bu^t)-Glu(OBzl)-NCO 3p: IR 2246 cm^{-1} ; ^1H NMR 1.19 (9H, s), 1.2 (3H, d), 1.6-2.1 (4H, m), 3.4 (2H, m), 3.92-4.45 (5H, m), 5.4 (1H, d), 6.35 (1H, m), 7.25-7.76 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 22.0, 27.0, 30.1, 37.4, 46.7, 50.2, 51.8, 63.2, 66.5, 73.3, 119.7, 124.8, 126.6, 126.7, 127.3, 140.9, 143.6, 156.2, 169.6 ; ES MS m/z observed 524.6.

Fmoc-Tyr(Bu^t)-Gly-NCO 3q: IR 2248 cm^{-1} ; ^1H NMR (δ , CDCl_3) 2.87 (2H, d), 3.88 – 4.46 (6H, m), 5.35 (1H, d), 6.6 (1H, m), 7.15-7.76 (12H, m) ; ^{13}C NMR (δ , CDCl_3) 28.8, 37.6, 43.3, 47.1, 56.1, 67.1, 78.5, 120.0, 124.3, 125.0, 127.1, 127.7, 129.7, 141.3, 143.6, 154.5, 171.3, 176.1 ; ES MS m/z observed 514.5.

N^α -Fmoc-tripeptide isocyanates 4

Fmoc-Val-Ala-Leu-NCO 4a: Yield 0.478 g (92 %) ; m.p. 166-67 $^\circ\text{C}$; IR 2253 cm^{-1} ; ^1H NMR (δ , CDCl_3) 0.86-0.95 (12H, m), 1.16 (3H, d), 1.39 (2H, m), 1.68 (1H, m), 1.78-1.84 (1H, m), 3.81-4.4 (6H, m), 5.25 (1H, d), 6.35-6.67 (2H, m), 7.25-7.75 (8H, m) ; ^{13}C NMR (δ , CDCl_3) 17.2, 18.6, 19.5, 22.0, 23.0, 24.6, 29.1, 40.3, 47.3, 48.9, 51.2, 58.5,

66.5, 119.9, 124.9, 126.4, 127.0, 127.6, 141.3, 143.8, 157.0, 170.1, 170.6 ; ES MS m/z observed 521.5.

Fmoc-Tyr(Bu^t)-Gly-Gly-NCO 4b: Yield 0.536 g (94 %) ; m.p. 189-91 °C ; IR 2255 cm⁻¹ ; ¹H NMR (δ , CDCl₃) 2.86 (2H, d), 3.82–4.44 (8H, m), 5.35 (1H, d), 6.56–6.85 (2H, m), 7.2–7.78 (12H, m) ; ¹³C NMR (δ , CDCl₃) 36.5, 43.3, 43.5, 47.2, 54.4, 66.7, 120.0, 124.4, 125.1, 126.6, 127.2, 127.9, 129.7, 132.4, 141.3, 144.2, 154.4, 156.6 ; ES MS m/z observed 571.6.

Fmoc-Val-Pro-Gly-NCO 4c: Yield 0.461 g (94 %) ; m.p. 148-50 °C ; IR 2247 cm⁻¹ ; ¹H NMR (δ , CDCl₃) 0.95 (6H, d), 1.36–2.11 (5H, m), 3.69–4.21 (4H, m), 4.42 (2H, m), 5.23 (1H, d), 6.42–6.48 (1H, m), 7.3–7.8 (8H, m) ; ¹³C NMR (δ , CDCl₃) 18.4, 19.3, 24.2, 28.7, 29.6, 47.1, 47.2, 56.2, 60.6, 66.6, 119.9, 125.0, 126.3, 126.7, 126.9, 127.6, 141.2, 143.7, 156.0 ; MS (MALDI-TOF) m/z observed 491.2.

N^α-Fmoc-tetrapeptide isocyanates 5

Fmoc-Gly-Phe-Leu-Val-NCO 5a: Yield 0.595 g (91 %) ; m.p. 176-78 °C ; IR 2248 cm⁻¹ ; ¹H NMR (δ , CDCl₃) 0.86–0.98 (12H, m), 1.33 (2H, m), 1.63 (1H, m), 1.89 (1H, m), 2.89 (2H, d), 3.84–4.44 (8H, m), 5.28 (1H, m), 6.3–6.78 (3H, m), 7.17–7.85 (13H, m) ; ¹³C NMR (δ , CDCl₃) 18.6, 19.5, 22.0, 23.0, 24.6, 29.1, 37.3, 40.3, 43.6, 47.0, 51.2, 54.1, 58.5, 66.6, 119.9, 124.9, 126.6, 126.7, 127.0, 127.6, 128.6, 129.2, 137.5, 141.2, 143.7, 157.6, 169.1, 170.1, 170.3 ; ES MS m/z observed 657.6.

Fmoc-Tyr(Bu^t)-Gly-Gly-Phe-NCO 5b: Yield 0.638 g (89 %) ; m.p. 149-51 °C ; IR 2256 cm⁻¹ ; ¹H NMR (δ , CDCl₃) 2.82–2.92 (4H, m), 3.83–4.45 (8H, m), 5.36 (1H, d), 6.48–6.96 (3H, m), 7.1–7.85 (17H, m) ; ¹³C NMR (δ , CDCl₃) 36.8, 37.2, 43.3, 43.4, 47.3, 54.2, 54.4, 66.7, 119.9, 124.4, 125.1, 126.6, 126.7, 127.2, 128.6, 127.9, 129.2, 129.7, 132.4, 137.4, 141.3, 144.2, 154.4, 156.6 ; ES MS m/z observed 718.8.

Fmoc-Pro-Gly-Val-Gly-NCO 5c: Yield 0.509 g (93 %) ; m.p. 158-59 °C ; IR 2252 cm⁻¹ ; ¹H NMR (δ, CDCl₃) 0.95 (6H, d), 1.34–2.12 (5H, m), 3.68-4.44 (9H, m), 5.29 (1H, d), 6.43–6.8 (2h, m), 7.28-7.76 (8H, m) ; ¹³C NMR (δ, CDCl₃) 18.4, 19.3, 24.2, 28.7, 29.6, 47.1, 47.2, 56.2, 60.6, 66.6, 119.9, 125.0, 126.3, 126.7, 126.9, 127.6, 141.2, 143.7, 156.0 ; ES MS *m/z* observed 548.6.

Fmoc-Ala-Leu-ψ(NH-CO-NH)-Val-NCO 9a: A 0.523 g (1 mmol) of Fmoc-Ala-Leu-ψ(NH-CO-NH)-Val-OH, following the general procedure for **3**, was converted to its isocyanate. Yield 0.478 g (92 %) ; m.p. 156-57 °C ; ¹H NMR (δ, CDCl₃) 0.88–0.98 (12H, d), 1.16 (3H, d), 1.33 (2H, m), 1.62 (1H, m), 1.89 (1H, m), 3.68–4.23 (4H, m), 4.42 (2H, d), 5.13 (1H, d), 5.9–6.75 (4H, m), 7.25–7.75 (8H, m) ; ¹³C NMR (δ, CDCl₃) 17.2, 18.6, 19.5, 22.0, 23.0, 24.6, 29.1, 40.2, 47.3, 48.8, 51.3, 58.5, 66.6, 119.9, 125.0, 126.4, 127.0, 127.6, 141.3, 143.7, 156.5, 156.9, 165.6, ; ES MS *m/z* observed 536.4.

Fmoc-Pro-Val-ψ(NH-CO-NH)-Gly-NCO 9b: A 0.493 g (1 mmol) of Fmoc-Pro-Val-ψ(NH-CO-NH)-Gly-OH, following the general procedure for **3**, was converted to its isocyanate. Yield 0.436 g (89 %) ; m.p. 161-63 °C ; ¹H NMR (δ, CDCl₃) 0.93 (6H, d), 1.21–2.05 (4H, m), 1.84 (1H, m), 3.26–3.72 (5H, m), 3.81 (1H, m), 4.0 (1H, m), 4.21 (1H, t), 4.43 (2H, m), 5.03 (1H, d), 6.46–6.72 (2H, m), 7.2-7.75 (8H, m) ; ¹³C NMR (δ, CDCl₃) 18.6, 19.5, 24.2, 28.6, 29.1, 43.3, 47.3, 47.6, 58.5, 60.8, 66.5, 120.0, 125.1, 126.7, 127.1, 127.6, 141.3, 143.7, 155.3, 156.3, 159.2 ; ES MS *m/z* observed 506.6.

Fmoc-Leu-Phe-ψ(NH-CO-NH)-Ala-NCO 9c: A 0.586 g (1 mmol) of Fmoc-Leu-Phe-ψ(NH-CO-NH)-Ala-OH, following the general procedure for **3**, was converted to its isocyanate **9c**. Yield 0.543 g (93 %) ; m.p. 181-83 °C ; IR 2247 cm⁻¹ ; ¹H NMR (δ, CDCl₃) 0.88 (6H, m), 1.21 (3H, d), 1.32 (2H, m), 1.61 (1H, m), 2.86 (2H, m), 3.82-4.46

(6H, m), 5.36 (1H, d), 6.25–6.85 (3H, m), 7.12–7.85 (13H, m) ; ^{13}C NMR (δ , CDCl_3) 17.2, 22.0, 23.0, 24.6, 37.3, 40.3, 47.2, 48.9, 51.2, 54.1, 66.6, 120.0, 125.0, 126.7, 126.9, 127.0, 127.6, 128.5, 129.2, 137.6, 141.3, 143.9, 154.2, 155.2, 169.1 ; ES MS m/z observed 514.5.

***N*^α-Fmoc-peptidyl urea acids **7a-c**: General procedure**

To a solution of *N*^α-Fmoc-peptide isocyanate **3** (1 mmol) in DCM (5 mL) at 0 °C was added freshly prepared Bis-TMS or Tris-TMS amino acid or peptide (from 1.2 mmol of amino acid or peptide refluxed with 2.5 mmol each of TEA and TMS-Cl in 10 mL of DCM or 4 mmol each of TEA and TMS-Cl in the case of Tris-TMS derivatives) and stirring was continued for 15 - 30 min. After completion of the reaction, the solvent was evaporated under reduced pressure below 40 °C. The residue was dissolved in 10 % Na_2CO_3 (20 mL) and washed with diethyl ether (20 mL x 2). The aqueous layer was acidified with dilute HCl until pH 2. The separated product was extracted with EtOAc (20 mL x 3). The water wash (20 mL x 3) was given to the combined organic layer and dried over anhydrous Na_2SO_4 . The organic layer was concentrated to obtain the title compound. The peptidyl ureas were purified either by crystallization using DMF and water or by column chromatography using the solvent system MeOH and CHCl_3 (5 % to 10 % of MeOH in CHCl_3).

Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH **7a:** A 0.483 g (1 mmol) of **3e** was coupled with freshly prepared Bis-TMS-Ala (from 0.106 g, 1.2 mmol of Ala-OH in 10 mL of DCM) to obtain **7a**. Yield 0.479 g (89 %) ; m.p. 188 - 89 °C ; ^1H NMR (δ , DMSO) 0.8 (6H, d), 1.16 (3H, d), 1.86 (1H, m), 2.9 (2H, d), 3.35 (1H, t), 3.77 (1H, t), 4.0-4.38 (4H, m), 5.32 (1H, m), 6.4-6.75 (2H, m), 7.15–7.9 (13H, m), 8.29 (1H, m) ; ^{13}C NMR (δ ,

DMSO) 14.1, 18.6, 19.0, 19.6, 30.8, 31.1, 41.0, 41.3, 47.1, 48.5, 58.5, 58.5, 60.7, 66.1, 120.5, 125.8, 126.6, 127.5, 128.1, 128.5, 129.7, 138.0, 141.1, 144.2, 144.4, 144.4, 156.4, 156.5, 171.0, 175.5 ; ES MS m/z observed 573.4.

Fmoc-Val-Pro- ψ (NH-CO-NH)-Gly-OH 7b: A 0.433 g (1 mmol) of **3g** was coupled with freshly prepared Bis-TMS-Gly (from 0.09 g, 1.2 mmol of Gly-OH in 10 mL of DCM) to obtain **7b**. Yield 0.383 g (82 %) ; m.p. 168 - 69 °C ; ^1H NMR (δ , DMSO) 0.93 (6H, d), 1.21–2.05 (4H, m), 1.84 (1H, m), 3.25–3.65 (5H, m), 3.81 (1H, m), 4.0 (1H, m), 4.21 (1H, t), 4.43 (2H, m), 5.28 (1H, d), 6.46–6.72 (2H, m), 7.2–7.75 (8H, m) ; ^{13}C NMR (δ , DMSO) 18.6, 19.5, 24.1, 28.6, 29.1, 43.2, 47.3, 47.5, 58.5, 60.8, 66.5, 119.9, 124.9, 127.0, 127.6, 141.3, 143.8, 155.2, 156.3, 170.1, 173.6 ; ES MS m/z observed 509.6.

N^α -Fmoc-tetrapeptidyl urea acids 8a-c

Fmoc-Val-Phe- ψ (NH-CO-NH)-Ala-Leu-OH 8a: A 0.483 g (1 mmol) of **3e** was coupled with freshly prepared Bis-TMS-Ala-Leu (from 0.242 g, 1.2 mmol of Ala-Leu-OH in 10 mL of DCM) to obtain **8a**. Yield 0.567 g (89 %) ; m.p. 201-02 °C ; ^1H NMR (δ , DMSO) 0.88–0.96 (12H, m), 1.16 (3H, d), 1.33 (2H, m), 1.63 (1H, m), 1.84 (1H, m), 2.87 (2H, d), 3.75–4.42 (7H, m), 5.26 (1H, d), 6.45–6.8 (2H, m), 7.12–7.85 (13H, m), 8.25–8.45 (2H, m) ; ^{13}C NMR (δ , DMSO) 17.2, 18.6, 19.5, 22.0, 23.0, 24.6, 29.1, 37.3, 40.3, 47.3, 48.9, 51.2, 54.1, 58.5, 66.6, 120.0, 125.0, 126.7, 127.0, 127.6, 128.6, 129.2, 137.5, 141.3, 143.8, 155.6, 156.8, 168.9, 170.3, 174.8 ; ES MS m/z observed 686.8.

Fmoc-Val-Phg- ψ (NH-CO-NH)-Val-Gly-OH 8b: A 0.469 g (1 mmol) of **3f** was coupled with freshly prepared Bis-TMS-Val-Gly (from 0.2 g 1.2 mmol of Val-Gly-OH in 10 mL of DCM) to obtain **8b** as a off white solid. Yield 0.486 g (88 %) ; m.p. 214-215 °C ; ^1H NMR (δ , DMSO) 0.93 (12H, m), 1.84 (2H, m), 3.76–4.52 (8H, m), 5.28 (1H, d), 6.4–6.7

(2H, m), 7.13–7.85 (13H, m), 8.15–8.35 (2H, m) ; ^{13}C NMR (δ , DMSO) 18.6, 18.6, 19.5, 19.5, 29.1, 29.1, 43.4, 47.3, 54.1, 58.5, 58.5, 66.6, 120.0, 125.0, 126.7, 127.0, 127.6, 128.6, 129.2, 137.5, 141.3, 143.8, 155.2, 156.3, 169.2, 170.2, 174.5 ; ES MS m/z observed 644.7.

Fmoc-Val-Ala- ψ (NH-CO-NH)-Val-Gly-OH 8c: A 0.07 g (1 mmol) of **3b** was coupled with freshly prepared Bis-TMS-Val-Gly (from 0.209 g, 1.2 mmol of Val-Gly-OH in 10 mL of DCM) to obtain **8c**. Yield 0.429 g (89 %) ; m.p. 194-96 °C ; ^1H NMR (δ , DMSO) 0.96 (12H, d), 1.18 (3H, d), 1.82–1.86 (2H, m), 3.6–3.95 (5H, m), 4.2 (1H, t), 4.42 (2H, m), 5.30 (1H, d), 6.5–6.85 (2H, m), 7.15–7.80 (8H, m), 8.2–8.45 (2H, m) ; ^{13}C NMR (δ , DMSO) 17.2, 18.6, 18.6, 19.5, 19.5, 29.1, 29.1, 43.6, 47.3, 48.9, 58.5, 58.5, 66.6, 120.0, 125.0, 127.0, 127.6, 141.3, 143.8, 156.1, 156.6, 168.8, 170.2, 173.2 ; ES MS m/z observed 582.7.

N^a -Fmoc-tetrapeptidyl ureas containing two ureas moieties

Fmoc-Leu-Phe- ψ (NH-CO-NH)-Ala- ψ (NH-CO-NH)-Ser-OH 10a: A 0.583 g (1 mmol) of Fmoc-Leu-Phe- ψ (NH-CO-NH)-Ala-NCO was coupled with freshly prepared Tris-TMS-Ser (from 0.126 g, 1.2 mmol of Ser-OH in 10 mL of DCM) to obtain **10a**. It was further purified by column chromatography using MeOH and CHCl_3 (5 % to 20 % of MeOH in CHCl_3). Yield 0.576 g (90 %) ; m.p. 156-57 °C ; ^1H NMR (δ , DMSO) 0.93 (6H, d), 1.16 (3H, d), 1.33 (2H, m), 1.63 (1H, m), 2.81 (2H, m), 3.4 (2H, m), 3.82-4.46 (7H, m), 5.96 (1H, s), 6.3–6.85 (4H, m), 7.12–7.85 (13H, m), 8.21 (1H, m) ; ^{13}C NMR (δ , DMSO) 17.2, 22.0, 23.0, 24.6, 37.3, 40.3, 47.2, 48.9, 51.2, 51.8, 54.1, 62.1, 66.6, 120.0, 125.0, 126.7, 127.0, 127.6, 128.5, 129.2, 137.6, 141.3, 143.9, 154.2, 155.2, 156.4, 169.1, 172.3 ; ES MS m/z observed 689.7.

General procedure: *N*^α-Fmoc-peptidyl urea esters 7d-f:

To an ice-cold mixture of amino acid or peptide ester hydrochloride or *p*-toluene sulfonate salt (1.1 mmol) in THF (5 mL) and NMM (1.21 mL, 1.1 mmol) was added a solution of *N*^α-Fmoc-peptide isocyanate (1 mmol) in DCM (5 mL) and stirred for about 15–20 min. After the completion of the reaction, the solvent was removed under reduced pressure and the residue was dissolved in EtOAc. The organic layer was washed with 10 % Na₂CO₃ (10 mL) and water (10 mL x 3) and finally dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure to obtain the *N*^α-Fmoc-peptidyl urea esters. The peptidyl ureas were purified either by the crystallization using DMF-water or by column chromatography using the solvent system MeOH and CHCl₃ (2 % to 10 % of MeOH in CHCl₃).

Fmoc-Gly-Phe-ψ(NH-CO-NH)-Gly-OMe 7d: A 0.441 g (1 mmol) of **3a** was coupled with Gly-OMe.HCl (0.137 g, 1.1 mmol) to obtain **7d**. Yield 0.478 g (94 %) ; m.p. 173-75 °C ; ¹H NMR (δ, DMSO) 2.87 (2H, d), 3.65 (3H, s), 3.92-4.48 (8H, m), 5.13 (1H, d), 6.45–6.7 (2H, m), 7.15–7.8 (13H, m), 8.26 (1H, m) ; ¹³C NMR (δ, DMSO) 37.3, 43.3, 43.6, 47.2, 54.1, 56.0, 66.6, 119.9, 124.9, 127.0, 126.6, 126.7, 128.6, 129.2, 137.5, 141.3, 143.8, 154.6, 156.7, 168.8, 173.2 ; ES MS *m/z* observed 531.7.

Fmoc-Thr(Bu^t)-Gly-ψ(NH-CO-NH)-Leu-OBzl 7f: A 0.541 g (1 mmol) of **3p** was coupled with Leu-OBzl.PTSA (0.432 g, 1.1 mmol) to obtain **7f**. Yield 0.551 g (89 %) ; m.p. 156-57 °C ; ¹H NMR (δ, DMSO) 0.88 (6H, m), 1.11–1.25 (12H, m), 1.33 (2H, m), 1.63-2.87 (7H, m), 3.36 (2H, m), 3.85-4.43 (6H, m), 5.65 (1H, d), 6.5–6.75 (2H, m), 7.1–7.85 (13H, m), 8.2 (1H, m) ; ¹³C NMR (δ, DMSO) 22.0, 22.2, 23.0, 24.6, 27.0, 30.2, 37.3, 37.4, 40.3, 46.7, 50.3, 51.2, 51.8, 62.1, 66.6, 73.3, 119.7, 124.8, 126.7, 126.7, 127.3,

128.6, 129.2, 137.5, 140.9, 143.6, 156.3, 156.8, 170.1, 176.9 ; ES MS m/z observed 745.7.

***N*^α-Fmoc-tetrapeptidyl ureas containing two ureas moieties**

Fmoc-Leu-Phe-ψ(NH-CO-NH)-Ala-ψ(NH-CO-NH)-Gly-OCH₃ 10b: A 0.583 g (1 mmol) of Fmoc-Leu-Phe-ψ(NH-CO-NH)-Ala-NCO was coupled with Gly-OMe.HCl (0.137 g, 1.1 mmol) to obtain **10b**. It was further purified by column chromatography using the solvent system CHCl₃ : MeOH : AcOH (5 % to 20 % of MeOH in CHCl₃ containing 1 % AcOH) to get analytically pure compound. Yield 0.632 g (94 %) ; m.p. 204 - 06 °C ; ¹H NMR (δ, DMSO) 0.89 (6H, d), 1.16 (3H, d), 1.33 (2H, m), 1.63 (1H, m), 2.88 (2H, m), 3.65 (3H, s), 3.86-4.48 (8H, m), 5.46 (1H, s), 6.35–6.8 (4H, m), 7.12–7.85 (13H, m), 8.26 (1H, m) ; ¹³C NMR (δ, DMSO) 17.2, 22.0, 23.0, 24.6, 37.3, 40.3, 43.4, 47.2, 48.9, 51.2, 54.1, 56.0, 66.6, 120.0, 125.0, 126.7, 127.0, 127.6, 128.4, 129.2, 137.5, 141.3, 143.9, 154.2, 155.2, 156.4, 169.6, 174.3 ; ES MS m/z observed 673.9.

Deprotection of Fmoc group from the *N*^α-Fmoc-peptidyl urea acids **12: General procedure**

A mixture of *N*^α-Fmoc-peptidyl urea (1 mmol) in DCM (5mL) and DEA (5 mL) was stirred until the deprotection was complete. After the completion of the reaction, the solvent was removed under reduced pressure below 40 °C and the residue was recrystallized using EtOAc and *n*-hexane to get the solid crystalline compounds in most of the cases. Free peptidyl ureas were purified by column chromatography using the solvent system CHCl₃ : MeOH (5 % to 15 % of MeOH in CHCl₃).

Val-Pro-ψ(NH-CO-NH)-Gly-OH 12a: A suspension of 0.493 g (1 mmol) of **9c** in 5mL of DCM was treated with 5 mL of DEA to obtain **12a**. Yield 0.235 g (87 %) ; m.p. 146-48 °C ; ¹H NMR (δ, DMSO) 0.95 (6H, d), 1.2-2.1 (5H, m), 3.2–3.85 (6H, m), 4.0 (1H, m), 6.25–6.75 (2H, m), 7.8 (1H, d), 8.34 (1H, m) 8.6 (1H, br s) ; ¹³C NMR (δ, DMSO)

18.6, 19.5, 24.1, 28.6, 29.1, 47.3, 58.5, 60.8, 154.9, 158.3, 169.8 ; ES MS m/z observed : 286.4.

Ser(Bzl)-Phe- ψ (NH-CO-NH)-Leu-OH 12b: A suspension of 0.677 g (1 mmol) of Fmoc-Ser(Bzl)-Phe- ψ (NH-CO-NH)-Leu-OH in 5mL of DCM was treated with 5 mL of DEA to obtain **12b**. Yield 0.381 g (84 %) ; m.p. 168-69 °C ; ^1H NMR (δ , DMSO) 0.92 (6H, m), 1.13 (1H, m), 1.53 (2H, m), 2.8 2.85 (4H, m), 3.3–3.85 (4H, m), 6.45-6.82 (2H, m), 7.1–7.5 (10H, m), 8.24–8.65 (3H, m) ; ^{13}C NMR (δ , DMSO) 11.3, 15.5, 25.4, 35.8, 37.3, 51.8, 54.1, 57.3, 62.1, 126.7, 126.8, 128.6, 128.7, 129.2, 129.2, 137.5, 137.5, 155.2, 158.9, 170.1 ; ES MS m/z observed 470.4.

Val-Phe- ψ (NH-CO-NH)-Ala-Leu-OH 12d: A suspension of 0.670 g (1 mmol) of **8a** in 5mL of DCM was treated with 5 mL of DEA to obtain **12d**. Yield 0.380 g (85 %) ; m.p. 172-73 °C ; ^1H NMR (δ , DMSO) 0.91–0.95 (12H, m), 1.16 (3H, m), 1.33 (2H, m), 1.59 (1H, m), 1.84 (1H, m), 2.87 (2H, d), 3.8–3.85 (3H, m), 6.81 (1H, m), 7.1–7.45 (5H, m), 8.25–8.62 (2H, m) ; ^{13}C NMR (δ , DMSO) 17.2, 18.6, 19.5, 22.0, 23.0, 24.6, 29.1, 37.3, 40.0, 48.9, 51.2, 58.5, 126.7, 128.6, 129.2, 137.5, 156.1, 158.3, 158.8, 170.6 ; ES MS m/z observed 463.5.

[Leu⁵]enkephalin analogs

Tyr-Gly- ψ (NH-CO-NH)-Gly-Phe-Leu-OH 13: A solution of 0.513 g (1 mmol) of Fmoc-Tyr(Bu^t)-Gly-NCO **3q** in DCM (5 mL) was cooled to 0 °C and freshly prepared Bis-TMS-Gly-Phe-Leu (from 0.402 g, 1.2 mmol of Gly-Phe-Leu-OH in 10 mL of DCM) was added. The resulting mixture was stirred for 30 min. and the reaction was monitored by TLC. After the completion of the reaction, the product was obtained by routine work-up. The crude Fmoc-Tyr(Bu^t)-Gly- ψ (NH-CO-NH)-Gly-Phe-Leu-OH was crystallized

using methanol and water to obtain the pure product. Yield 0.693 g (86 %) ; m.p. 144 - 46 °C ; MS (MALDI-TOF) m/z observed : 871.0 [M+Na]⁺, 887.2 [M+K]⁺. The suspension of Fmoc-pentapeptidyl urea (0.65 g, 0.779 mmol) in 2 mL of DCM was treated with TFA (4 mL in 4 mL of DCM) for 30 min. and the resulting solution was concentrated and residual TFA was removed by ether washes (2 x 10 mL). The resulting residue was dissolved in 2 mL of DCM taken in DEA (4 mL in 4 mL of DCM) and stirred for 45 min. After the completion of the deprotection (by TLC analysis), the solvent was concentrated under reduced pressure. The residue was recrystallized by adding ether. The crude peptidyl urea **13** was further purified by column chromatography using the solvent system CHCl₃ : MeOH (5 % to 15 % of MeOH in CHCl₃) to obtain the analytically pure compound. Yield 0.367 g (79 %) ; m.p. 156–158 °C ; ¹H NMR (δ, DMSO) 0.96 (6H, m), 1.39 (2H, m), 1.55 (1H, m), 2.75–4.6 (11H, m), 6.6–8.7 (16H, m) ; ¹³C NMR (δ, DMSO) 10.9, 13.7, 21.4, 22.7, 24.2, 36.1, 39.0, 41.3, 50.4, 53.6, 115.3, 115.7, 116.2, 118.7, 124.8, 126.5, 127.9, 129.2, 130.4, 131.2, 154.9, 156.6, 158.0, 158.3, 169.8 ; MS (MALDI-TOF) m/z observed 571.5.

Tyr-Gly-Gly-ψ(NH-CO-NH)-Phe-Leu-OH 14: A 0.570 g (1 mmol) of Fmoc-Tyr(Bu^t)-Gly-Gly-NCO **4b** was coupled with Bis-TMS-Phe-Leu-OH (from 0.333 g, 1.2 mmol of Phe-Leu-OH in 10 mL of DCM) to obtain Fmoc-Tyr(Bu^t)-Gly-Gly-ψ(NH-CO-NH)-Phe-Leu-OH as described above for **13**. Yield 0.67 g (84 %) ; m.p. 162 - 63 °C, MS (MALDI-TOF) m/z observed : 879.9 [M+Na]⁺, 887.2 [M+K]⁺. A 0.67 g (0.789 mmol) of *N*^α-Fmoc-pentapeptidyl urea was treated with TFA (4 mL in 4 mL of DCM) to cleave Bu^t group and finally with DEA (4 mL in 4 mL of DCM) as described above for **13** to obtain the title compound **14**. It was purified by column chromatography using the solvent

system CHCl₃ : MeOH (5 % to 15 % of MeOH in CHCl₃) to obtain the analytically pure compounds. Yield 0.351 g (78 %) ; m.p. 176-77 °C ; ¹H NMR (δ, DMSO) 1.0 (6H, m), 1.38 (2H, m), 1.56 (1H, m), 2.78–4.65 (11H, m), 6.65–8.75 (16H, m) ; ¹³C NMR (δ, DMSO) 10.8, 13.7, 21.4, 22.7, 24.2, 36.1, 39.0, 41.3, 50.4, 53.6, 115.3, 115.7, 116.2, 118.7, 124.8, 126.5, 128.0, 129.2, 130.4, 131.2, 154.9, 156.6, 158.0, 158.3, 169.7 ; MS (MALDI-TOF) *m/z* observed : 571.5.

Tyr-Gly-Gly-Phe-ψ(NH-CO-NH)-Leu-OH 15: A 0.717 g (1 mmol) of Fmoc-Tyr(Bu^t)-Gly-Gly-Phe-NCO **5b** was coupled with Bis-TMS-Leu-OH (from 0.157 g, 1.2 mmol of Leu-OH in 10 mL of DCM) to obtain the protected pentapeptidyl urea Fmoc-Tyr(Bu^t)-Gly-Gly-Phe-ψ(NH-CO-NH)-Leu-OH as described above for **13**. Yield 0.633 g (84 %) ; m.p. 163-64 °C ; MS (MALDI-TOF) *m/z* observed 871.0 [M+Na]⁺, 887.2 [M+K]⁺. A 0.633 g (0.745 mmol) of *N*^α-Fmoc-pentapeptidyl urea was treated with TFA (4 mL in 4 mL of DCM) to cleave Bu^t group and finally with DEA (4 mL in 4 mL of DCM) as described above for **13** to obtain the title compound **15**. It was purified by column chromatography using the solvent system CHCl₃ : MeOH (5 % to 15 % of MeOH in CHCl₃) to obtain the analytically pure compound. Yield 0.361 g (85 %) ; m.p. 186–188 °C ; ¹H NMR (δ, DMSO) 0.98 (6H, m), 1.41 (2H, m), 1.53 (1H, m), 2.68–4.6 (11H, m), 6.58–8.72 (16H, m) ; ¹³C NMR (δ, DMSO) 10.8, 13.6, 21.4, 22.7, 24.2, 36.1, 39.0, 41.2, 50.4, 53.6, 115.3, 115.7, 116.2, 118.7, 124.8, 126.5, 127.9, 129.2, 130.4, 131.2, 154.8, 156.5, 158.0, 158.3, 169.7 ; MS (MALDI-TOF) *m/z* observed 571.4.

Tyr-Gly-ψ(NH-CO-NH)-Gly-Phe-ψ(NH-CO-NH)-Leu-OH 18: A 0.513 g (1 mmol) of Fmoc-Tyr(Bu^t)-Gly-NCO **3q** was coupled with Bis-TMS-Gly-Phe (1.2 mmol in 10 mL of DCM) to obtain Fmoc-Tyr(Bu^t)-Gly-ψ(NH-CO-NH)-Gly-Phe-OH **16**. Yield 0.569 g (79

%) ; m.p. 178-79 °C ; MS (MALDI-TOF) m/z observed : 757.9 $[M+Na]^+$, 773.9 $[M+K]^+$.

A 0.569 g (0.79 mmol) of Fmoc-Tyr(Bu^t)-Gly- ψ (NH-CO-NH)-Gly-Phe-OH was converted to the corresponding isocyanate **17**. Yield 0.527 g (93 %) ; m.p. 184-187 °C ; IR 2257 cm^{-1} ; 1H NMR (δ , $CDCl_3$) 2.84–2.93 (4H, m), 3.85–4.46 (8H, m), 5.32 (1H, d), 6.26–6.96 (4H, m), 7.1–7.92 (17H, m) ; ^{13}C NMR (δ , $CDCl_3$) 36.9, 37.3, 43.3, 43.5, 47.2, 54.1, 54.5, 66.6, 120.0, 124.4, 125.0, 126.6, 126.7, 127.3, 128.6, 127.8, 129.1, 129.7, 132.3, 137.5, 141.3, 144.2, 154.6, 156.6 ; ES MS m/z observed 733.8. A solution of 0.527 g (0.73 mmol) of Fmoc-Tyr(Bu^t)-Gly- ψ (NH-CO-NH)-Gly-Phe-NCO **17** in DCM (5 mL) was coupled with Bis-TMS-Leu (0.88 mmol) and after the routine work-up the precursor of the title compound Fmoc-Tyr(Bu^t)-Gly- ψ (NH-CO-NH)-Gly-Phe- ψ (NH-CO-NH)-Leu-OH was obtained. The purity of the crude peptidyl urea obtained was satisfactory. It was further purified by flash column chromatography using the solvent system $CHCl_3$: MeOH (5 % to 15 % of MeOH in $CHCl_3$) to obtain the analytically pure compound. Yield 0.475 g (75 %) ; m.p. 186 - 88 °C ; MS (MALDI-TOF) m/z observed : 886.5 $[M+Na]^+$, 901.7 $[M+K]^+$. After the deprotection of Fmoc and Bu^t groups as above for **13**, the free peptidyl urea Tyr-Gly- ψ (NH-CO-NH)-Gly-Phe- ψ (NH-CO-NH)-Leu-OH **18** was obtained in good yield as well as purity. Yield 0.254 g (79 %) ; m.p. 128-29 °C ; 1H NMR (δ , DMSO) 0.98 (6H, m), 1.44 (2H, m), 1.54 (1H, m), 2.68–4.6 (11H, m), 6.48–8.72 (17H, m) ; ^{13}C NMR(δ , DMSO) 10.9, 13.7, 21.4, 22.8, 24.3, 36.0, 39.0, 41.3, 50.4, 53.7, 115.3, 115.7, 116.3, 118.7, 124.8, 126.5, 128.0, 129.2, 130.5, 131.3, 155.1, 156.6, 158.1, 158.4, 169.8 ; MS (MALDI-TOF) m/z observed 585.6.

Bioelastic polymer analogs

Val-Pro-Gly- ψ (NH-CO-NH)-Val-Gly-OH 22: A solution of 0.52 g (1 mmol) of Fmoc-Val-Pro-Gly-CON₃ in 10 mL of toluene was heated with Bis-TMS-Val-Gly-OH (1.15 mmol in 10 mL of DCM) at 80 °C until completion of the reaction by TLC using the eluent system MeOH : CHCl₃ (3:7). After completion of the reaction, the Fmoc-protected pentapeptidyl urea **22a** was obtained by routine work-up. The crude peptidyl urea was crystallized using DMF and water to obtain the pure product. Yield 0.642 g (85 %); m.p. 169-71 °C; MS (MALDI-TOF) *m/z* observed: 687.0 [M+Na]⁺, 703.1 [M+K]⁺. A 0.642 g (0.96 mmol) of Fmoc-Val-Pro-Gly- ψ (NH-CO-NH)-Val-Gly-OH was taken in 5 mL of DCM and treated with DEA (4 mL) to cleave Fmoc group as described in case of **21** to obtain the title compound **22**. The product was purified by column chromatography using the solvent system CHCl₃ : MeOH (5 % to 10 % of MeOH in CHCl₃) to obtain the analytically pure compound. Yield 0.327 g (79 %) ; m.p. 152-53 °C ; ¹H NMR (δ , DMSO) 0.81–0.9 (12H, m), 1.86–2.0 (6H, m), 2.32 (1H, m), 3.5–4.42 (12H, m), 6.32 (1H, d), 6.74 (1H, t), 8.05–8.73 (3H, m), 9.4 (1H, br s) ; ¹³C NMR (δ , DMSO) 11.2, 17.7, 17.9, 18.2, 19.0, 19.3, 25.4, 30.1, 30.6, 30.8, 41.3, 42.0, 44.6, 46.4, 57.2, 58.2, 60.0, 157.5, 168.0, 168.5, 171.5, 173.8 ; ES MS *m/z* observed 442.5.

Pro-Gly-Val-Gly- ψ (NH-CO-NH)-Val-OH 23: A 0.57 g (1 mmol) of Fmoc-Pro-Gly-Val-Gly-CON₃ was coupled with Bis-TMS-Val-OH (1.15 mmol in 10 mL of DCM) to obtain the precursor of the title compound Fmoc-Pro-Gly-Val-Gly- ψ (NH-CO-NH)-Val-OH **23a** as described above for **22a**. The crude peptidyl urea was crystallized using DMF and water to obtain the pure product. Yield 0.482 g (78 %); m.p. 176-177 °C; MS (MALDI-TOF) *m/z* observed : 686.7 [M+Na]⁺, 702.8 [M+K]⁺. A 0.482 g (0.78 mmol) of

Fmoc-Pro-Gly-Val-Gly- ψ (NH-CO-NH)-Val-OH **23a** was taken in 5 mL of DCM and treated with DEA (4 mL) and stirred until the deprotection was complete. The solution was concentrated under reduced pressure and residual DEA and side products were removed by dry ether washes (4 x 10 mL). The obtained peptidyl urea **23** was foamy solid. It was further purified by column chromatography using the solvent system CHCl₃ : MeOH (5 % to 10 % of MeOH in CHCl₃) to obtain the analytically pure compound. Yield 0.243 g (76 %) ; ¹H NMR (δ , DMSO) 0.8–0.87 (12H, m), 1.88–1.99 (6H, m), 2.3 (1H, m), 3.5–4.4 (12H, m), 6.3 (1H, d), 6.7 (1H, t), 8.0 (1H, d), 8.6–8.8 (2H, m), 9.4 (1H, br s) ; ¹³C NMR (δ , DMSO) 11.2, 17.7, 17.89, 18.1, 19.0, 19.3, 25.4, 30.1, 30.6, 30.76, 41.3, 41.96, 44.6, 46.4, 57.2, 58.3, 60.0, 157.4, 168.0, 168.5, 171.5, 173.9 ; ES MS m/z observed 442.4.

Table 1. *N*^α-Fmoc-peptide isocyanates Fmoc-NH-CHR₁-CONH-CHR₂-NCO

Product	R ₁	R ₂	Method*	Time	Yield (%)	m.p. (°C)
3a	H	CH ₂ C ₆ H ₅	A	30 min.	83	208-09
			C	40 sec.	87	
			D	20 min.	86	
3b	CH(CH ₃) ₂	CH ₃	A	30 min.	84	178-80
			C	40 sec.	92	
			D	25 min.	91	
3c	CH ₂ CH(CH ₃) ₂	CH(CH ₃) ₂	A	30 min.	85	193-94
			C	40 sec.	91	
			D	30 min.	88	
3d	CH ₃	CH ₂ CH(CH ₃) ₂	A	30 min.	84	178-80
			C	40 sec.	88	
			D	30 min.	91	
3e	CH(CH ₃) ₂	CH ₂ C ₆ H ₅	A	30 min.	81	198-99
			C	35 sec.	87	
			D	30 min.	82	
3f	CH(CH ₃) ₂	C ₆ H ₅	A	30 min.	79	172-74
			C	45 sec.	82	
			D	30 min.	80	
3g	CH(CH ₃) ₂	N-(CH ₂) ₃ -C	A	30 min.	79	168-69
			C	45 sec.	82	
			D	30 min.	80	
3h	CH(CH ₃)CH ₂ CH ₃	CH ₂ OCH ₂ C ₆ H ₅	A	40 min.	85	168-70
			C	45 sec.	91	
			D	25 min.	86	
3i	CH ₂ COOC(CH ₃) ₃	CH ₂ CH(CH ₃) ₂	A	35 min.	84	148-49
			C	35 sec.	88	

			D	30 min.	84
3j	$\text{CH}_2\text{CH}_2\text{CONH}(\text{Trt})$	$\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$	A	30 min.	72 164-65
			C	35 sec.	74
			D	30 min.	74
3k		$\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$	A	40 min.	71 167-69
			C	45 sec.	72
			D	25 min.	71
3l	$(\text{CH}_2)_3\text{NCN}(\text{NHPmc})$	$\text{CH}_2\text{CH}(\text{CH}_3)_2$	A	35 min.	82 182-83
			C	45 sec.	86
			D	30 min.	82
3m	$\text{CH}(\text{SCH}_2\text{C}_6\text{H}_5)\text{CH}_3$	$\text{CH}(\text{CH}_3)_2$	A	40 min.	79 154-55
			C	45 sec.	82
			D	25 min.	80
3n	$\text{CH}_2\text{OCH}_2\text{C}_6\text{H}_5$	$\text{CH}_2\text{C}_6\text{H}_5$	A	35 min.	82 188-89
			C	45 sec.	88
			D	30 min.	83
3o	$\text{CH}_2\text{OC}(\text{CH}_3)_3$	$\text{CH}_2\text{C}_6\text{H}_5$	A	40 min.	81 168-69
			C	30 sec.	86
			D	25 min.	80
3p	$\text{CH}(\text{CH}_3)\text{OC}(\text{CH}_3)_3$	$\text{CH}_2\text{CH}_2\text{COOCH}_2\text{C}_6\text{H}_5$	A	45 min.	76 139-40
			C	60 sec.	78
			D	20 min.	76
3q	$\text{CH}_2\text{C}_6\text{H}_4\text{OC}(\text{CH}_3)_3$	H	A	30 min.	88 123-25
			C	30 sec.	89
			D	20 min.	85

Method A : Reflux in toluene ; Method B : Microwave irradiation in toluene ; Method C : Microwave irradiation using solid powder ; Method D : Conversion by ultrasonication ; ^aIsolated yield after crystallization ; *Method B results are not shown in the table.

IR spectra of Fmoc-Ser(Bu^t)-Phe-CON₃ **2o**

IR spectra of Fmoc-Val-Pro-NCO **3g**

RP-HPLC of Fmoc-Val-Phe-CON₃ **2e** and RP-HPLC of Fmoc-Val-Phe-NCO **3e**

Waters C-18 bondapak (3.9 x 300 mm) column ; flow rate 1mL/min. ; eluent: MeCN: water (60 : 40 ; isocratic ; monitoring at 254 nm)

Analytical RP HPLC of Fmoc-Val-Phe-CON₃ 2e

Waters C-18 bondapak (3.9 x 300 mm) column ; flow rate 1mL/min. ; eluent: MeCN: water (60 : 40 ; isocratic ; monitoring at 254 nm)

Analytical RP HPLC of Fmoc-Val-Phe-NCO 3e

Monitoring of Curtius rearrangement of **2g** to **3g** using IR spectrum

Monitoring of Curtius rearrangement of **2g** to **3g** using IR spectra

¹H NMR spectra of Fmoc-Val-Ala-NCO **3b**

¹³C NMR spectra of Fmoc-Val-Ala-NCO **3b**

Data: <Untitled>.3 13 Aug 2004 17:40 Cal: angiotensin 23 Apr 2003 7:14
Kratos PCKompact SEQ V1.2.2: + Linear High, Power: 40, P.Ext. @ 490 (bin 67)

MALDI mass spectra of Fmoc-Val-Pro-Gly-NCO **4c**

%Int. 100% = 3.0 mV[sum= 118 mV] Profiles 1-40 Smooth Av 50

¹H NMR spectra of Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH 7a

¹³C NMR spectra of Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH **7a**

Current Data Parameters
 NAME 439836-1
 EXPNO 10
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040628
 Time 1.40
 INSTRUM AV400
 PROBHD 5 mm QNP 13C-1
 PULPROG zgpg30
 TD 32768
 SOLVENT DMSO
 NS 10000
 DS 0
 SWH 30120.482 Hz
 FIDRES 0.919204 Hz
 AQ 0.5439988 sec
 RG 512
 DE 16.600 usec
 TE 6.00 usec
 O1 0.0 K
 d1 3.00000000 sec
 d11 0.03000000 sec
 MCREST 0.00000000 sec
 MCWRR 0.01500000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 9.25 usec
 PL1 -2.00 dB
 SF01 100.6259487 MHz

===== CHANNEL f2 =====
 CDPORG2 waltz16
 NUC2 1H
 PCPD2 80.00 usec
 PL2 0.00 dB
 PL12 16.00 dB
 PL13 16.00 dB
 SF02 400.1320007 MHz

F2 - Processing parameters
 SI 131072
 SF 100.6127690 MHz
 MDW 0
 SSB 0
 LB 4.00 Hz
 GB 0
 PC 0.20

ID NMR plot parameters
 CX 20.00 cm
 CY 0.00 cm
 F1P 277.359 ppm
 F1 27905.81 Hz
 F2P -7.276 ppm
 F2 -732.06 Hz
 PPMCN 14.23173 ppm/cm
 HZCM 1431.89355 Hz/cm

ES MS of Fmoc-Val-Phe-ψ(NH-CO-NH)-Ala-OH 7a

page 1 of

Monday, January 19, 2004 12:15 PM

MultiView 1.5

Info for pane 2: 3012004

Period 1, Expt. 4; Mass range: 100.0 to 750.0 by 0.2 amu; Dwell: 0.8 ms; Pause: 2.0 ms

Acq. Time: Tue, Dec 30, 2003 at 8:01:37 AM; Samp. Name:

¹H NMR of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe 7e

Current Data Parameters
 NAME 439837
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040626
 Time 13:22
 INSTRUM AV400
 PROBHD 5 mm DUL 13C-1
 PULPROG zg30
 TD 32768
 SOLVENT DMSO
 NS 4
 DS 0
 SHW 8012.820 Hz
 FIDRES 0.244532 Hz
 AQ 2.0447731 sec
 RG 256
 DM 62.400 usec
 DE 6.00 usec
 TE 0.0 K
 D1 1.00000000 sec
 MCREST 0.00000000 sec
 MCHRG 0.01500000 sec

***** CHANNEL f1 *****
 NUC1 ¹H
 P1 12.00 usec
 PL1 0.00 dB
 SF01 400.1320007 MHz

F2 - Processing parameters
 S1 16384
 SF 400.1300000 MHz
 WDW EN
 SSB 0
 LB 0.50 Hz
 GB 0
 PC 0.50

1D NMR plot parameters
 CX 20.00 cm
 CY 0.00 cm
 FIP 12.156 ppm
 F1 4663.88 Hz
 F2P 0.040 ppm
 F2 15.97 Hz
 PPKCM 0.60579 ppm/cm
 HZCM 242.39577 Hz/cm

¹³C NMR spectra of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**

ES MS of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**

MultiView 1.5
3012006
Monday, January 19, 2004 12:15 PM
Period 1, Expt. 1; Mass range: 100.0 to 750.0 by 0.2 amu; Dwell: 0.8 ms; Pause: 2.0 ms
Acq. Time: Tue, Dec 30, 2003 at 8:07:26 AM; Samp.

¹³C NMR spectra of Tyr-Gly-Gly-ψ(NH-CO-NH)-Phe-Leu-OH 14

MALDI-mass of Tyr-Gly-ψ(NH-CO-NH)-Gly-Phe-Leu-OH 13

Multiple Spectra Report

Bruker Daltonics flexAnalysis

printed: 10/05/04 16:39:23

ES-MS of Tyr-Gly-ψ(NH-CO-NH)-Gly-ψ(NH-CO-NH)-Gly-ψ(NH-CO-NH)-Phe-ψ(NH-CO-NH)-Leu-OH **21**

Bruker Daltonics flexAnalysis

printed: 10/05/04 16:39:23

¹H NMR spectra of Val-Pro-Gly-ψ(NH-CO-NH)-Val-Gly-OH 22

MALDI-mass spectra of Pro-Gly-Val-Gly- ψ (NH-CO-NH)-Val-OH **23a**

Data: <Untitled>.7 23 Sep 2004 10:53 Cal: angior 23 Apr 2004 6:41
 Kratos PCKompact SEQ V1.2.2: + Linear High, Power: 40, P.Ext. @ 1312 (bin 111)

%Int. 100% = 48 mV[sum= 819 mV] Profiles 1-17 Smooth Av 100

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-R-(+)-1-phenylethylamine **25a**

Current Data Parameters
NAME 705-40963-1
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040220
Time 4.51
INSTRUM AV400
PROBHD 5 mm BBO BB-1H
PULPROG zgpg
TD 32768
SOLVENT DMSO
NS 4
DS 0
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 256
DM 62.400 usec
DE 6.00 usec
TE 0.0 K
U1 1.00000000 sec
MCREST 0.00000000 sec
MCWPK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 13.00 usec
PL1 0.00 dB
SF01 400.1332010 MHz

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.50 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 0.00 cm
FIP 12.034 ppm
F1 4815.25 Hz
F2 -0.240 ppm
F2 96.18 Hz
PPMCH 0.61373 ppm/cm
HZCM 245.57124 Hz/cm

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-R-(+)-1-phenylethylamine **25a**

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-S(-)-1-phenylethylamine **25b**

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-S-(-)-1-phenylethylamine **25b**

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-(R&S)-1-phenylethylamine **25c**

Current Data Parameters
 NAME 409563-III
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040220
 Time 4.54
 INSTRUM AV400
 PROBHD 5 mm BBO BB-1H
 PULPROG zg30
 TD 32768
 SOLVENT DMSO
 NS 4
 DS 0
 SFO1 8012.820 Hz
 FIDRES 0.244532 Hz
 AQ 2.0447731 sec
 RG 71.8
 DN 62.400 usec
 DE 6.00 usec
 TE 0.0 K
 O1 1.0000000 sec
 MCREST 0.0000000 sec
 MCWRK 0.0150000 sec
 ***** CHANNEL f1 *****
 NUC1 ¹H
 P1 13.00 usec
 PL1 0.00 dB
 SF01 400.1332010 MHz
 F2 - Processing parameters
 SI 16384
 SF 400.1300000 MHz
 WDW EM
 SSB 0
 LB 0.50 Hz
 GB 0
 PC 1.00
 1D NMR plot parameters
 CX 20.00 cm
 CY 0.00 cm
 FIP 12.034 ppm
 F1 4815.25 Hz
 F2P -0.240 ppm
 F2 -96.18 Hz
 PPMCH 0.61373 ppm/cm
 HZCM 245.57124 Hz/cm

¹H NMR spectra of Fmoc-Leu-Val-ψ(NH-CO-NH)-(R&S)-1-phenylethylamine **25c**

Current Data Parameters
NAME 409963-III
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040220
Time 4.54
INSTRUM AV400
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 32768
SOLVENT DMSO
NS 4
DS 0
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 71.8
DM 62.400 usec
DE 6.00 usec
TE 0.0 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCNRC 0.01500000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 13.00 usec
PL1 0.00 dB
SF01 400.1332010 MHz

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.50 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 0.00 cm
FIP 2.775 ppm
F1 1110.51 Hz
F2 433.07 Hz
PPMCH 0.08465 ppm/cm
HZCM 33.87189 Hz/cm

HR MS of Fmoc-Val-Ala-NCO **3b**

HR MS of Fmoc-Ala-Leu-NCO **3d**

HR MS of Fmoc-Ser(Bu^t)-Phe-ψ(NH-CO-NH)-Leu-OMe **7e**

HR MS of Fmoc-Leu-Val-ψ(NH-CO-NH)- S-(-)-1-phenylethylamine **25b**

HR MS of Fmoc-Val-Pro-Gly-ψ(NH-CO-NH)-Val-Gly-OH **22a**