Preparation and Synthetic Application of Novel Ketene Silyl Acetal of Methyl Trifluoropyruvate

*Go Takikawa, Toshimasa Katagiri and Kenji Uneyama**

Department of Applied Chemistry, Faculty of Engineering, Okayama University

3-1-1 Tsushima-naka, Okayama 700-8530, Japan

uneyamak@cc.okayama-u.ac.jp

General Procedures	S2
3,3,3-Trifluoro-2-[(4-methoxyphenyl)(trimethylsilyloxy)methyl]-2-trimethylsilyloxypropanoic acid methyl ester (10a)	S2
3,3,3-Trifluoro-2-[(3-chlorophenyl)(trimethylsilyloxy)methyl]-2-trimethylsilyloxypropanoic acid methyl ester (10b)	S2
3,3,3-Trifluoro-2-[(4-trifluoromethylphenyl)(trimethylsilyloxy)methyl]-2-trimethylsilyloxypropanoic acid methyl ester (10d)	S3
3,3,3-Trifluoro-2-[furyl(trimethylsilyloxy)methyl]-2-trimethylsilyloxypropanoic acid methyl ester (10e)	S3
3,3,3-Trifluoro-2-[(4-methylphenyl)(trimethylsilyloxy)methyl]-2-trimethylsilyloxypropanoic acid methyl ester (10f)	S4
3,3,3-Trifluoro-2-hydroxy-2-[(4-methoxyphenyl)hydroxymethyl]propanoic acid methyl ester (12a)	S4
2-[(3-Chlorophenyl)hydroxymethyl]-3,3,3-trifluoro-2-hydroxypropanoic acid methyl ester (12b)	S5
3,3,3-Trifluoro-2-hydroxy-2-[(4-trifluoromethylphenyl)hydroxymethyl]propanoic acid methyl ester (12d)	S5
3,3,3-Trifluoro-2-[(furyl)hydroxymethyl]-2-hydroxypropanoic acid methyl ester (12e)	S5
3,3,3-Trifluoro-2-hydroxy-2-[1'-(p-tolyl)hydroxymethyl]propanoic acid methyl ester (12f)	S6
2-Trifluoromethyl-2,3-dihydroxy-5-phenylpentanoic acid methyl ester (12g)	S6
2-Trifluoromethyl-2,3-dihydroxy-4-hexenoic acid methyl ester (12h)	S6
2-Hydroxy-3-oxo-2-trifluoromethyloctanoic acid methyl ester (15b)	S7

General Procedures

^1H NMR spectra were recorded at 300, 500 and 600 MHz, ^{19}F NMR was recorded at 282 and 470 MHz and ^{13}C NMR was recorded at 151 MHz. The chemical shifts are reported in δ (ppm) values relative to TMS (δ 0 ppm for ^1H NMR in CDCl_3) and C_6F_6 (δ 0 ppm for ^{19}F NMR in CDCl_3) as internal standards, respectively. Coupling constants are reported in hertz (Hz). All air and/or water-sensitive reactions were carried out under an argon atmosphere with dry, freshly distilled solvents using a standard syringe-cannula/septa technique. MTFP (**1**, 98% pure), was distilled from P_2O_5 . THF was distilled from sodium/benzophenone ketyl. *n*-Heptane and TMSCl were distilled from CaH_2 . CH_2Cl_2 was distilled from P_2O_5 . Aldehydes, acyl halides, ketones and benzaldehyde dimethyl acetal were purified by distillation. Other reagents were employed without further purification. Kanto Chemical Co., Inc. silica gel (Silica Gel 60N (spherical, neutral), 40-50 μm) was employed for the chromatography.

3,3,3-Trifluoro-2-[(4-methoxyphenyl)(trimethylsilanyloxy)methyl]-2-trimethylsilanyloxypropanoic acid methyl ester (10a).

97%; colorless oil; Diastereomeric mixture (63:37 by ^{19}F NMR); IR (neat) 2990, 2940, 2880, 1785, 1766, 1632, 1602, 1528, 1528, 1480, 1458 cm^{-1} ; ^1H NMR (300 MHz) δ -0.01 (s, 9H, major), 0.07 (s, 9H, minor), 0.06 (s, 9H, major), 0.21 (s, 9H, minor), 3.71 (s, 3H, major), 3.80 (s, 3H, major), 3.81 (s, 3H, minor), 3.86 (s, 3H, minor), 5.18 (s, 1H, minor), 5.20 (s, 1H, major), 6.80 (dd, $J = 1.8$ Hz, $J = 6.9$ Hz, 2H, major), 6.82 (d, $J = 8.7$ Hz, 2H, minor), 7.19 (dd, $J = 1.8$ Hz, $J = 6.9$ Hz, 2H, major), 7.27 (d, $J = 8.7$ Hz, 2H); ^{19}F NMR (282 MHz) δ 87.6 (s, 3F, minor), 89.9 (s, 3F, major); EI MS m/z (relative intensity) 423 ($\text{M}^+ - 15$, trace), 209 (100), 73 (44); Anal. Calcd for $\text{C}_{18}\text{H}_{29}\text{F}_3\text{O}_5\text{Si}_2$: C, 49.29; H, 6.66. Found; C, 49.46; H, 6.83.

3,3,3-Trifluoro-2-[(3-chlorophenyl)(trimethylsilanyloxy)methyl]-2-trimethylsilanyloxypropanoic acid methyl ester (10b).

86%; colorless oil; Diastereomeric mixture (61:39 by ^{19}F NMR); IR (neat) 2964, 2910, 1772, 1760, 1600, 1580, 1478, 1440 cm^{-1} ; ^1H NMR (300 MHz) δ -0.04 (s, 9H, minor), 0.02 (s, 9H, major), 0.06 (s, 9H, major), 0.20 (s, 9H, minor), 3.73 (s, 3H, major), 3.86 (s, 3H, minor), 5.18 (s, 1H, minor), 5.23 (s, 1H, major), 7.08-7.39 (m, 5H x 2, major and minor mixed); ^{19}F NMR (282 MHz) δ 87.7 (s, 3F, minor), 89.9 (s, 3F, major); EI MS m/z (relative intensity) 427 ($\text{M}^+ - 15$, trace), 302 (1), 213 (100), 73 (80); Anal. Calcd for $\text{C}_{17}\text{H}_{26}\text{ClF}_3\text{O}_4\text{Si}_2$: C, 46.09; H, 5.92. Found; C, 45.79; H, 6.16.

3,3,3-Trifluoro-2-[(4-trifluoromethylphenyl)(trimethylsilanyloxy)methyl]-2-trimethylsilanyloxypropanoic acid methyl ester (10d).

58%; colorless oil; Diastereomeric mixture (53:47 by ^{19}F NMR); IR (neat) 3000, 2948, 2897, 1958, 1783, 1635, 1458, 1437 cm^{-1} ; ^1H NMR (500 MHz) δ -0.06 (s, 9H, minor), 0.00 (s, 9H, major), 0.03 (s, 9H, major), 0.17 (s, 9H, minor), 3.74 (s, 3H, major), 3.87 (s, 3H, minor), 5.27 (s, 1H, minor), 5.30 (s, 1H, major), 7.41 (d, $J = 8.5$ Hz, 1H x 2, major and minor mixed), 7.47 (d, $J = 8.0$ Hz, 1H x 2, major and minor mixed), 7.55 (d, $J = 8.5$ Hz, 1H x 2, major and minor mixed), 7.56 (d, $J = 8.0$ Hz, 1H x 2, major and minor mixed); ^{19}F NMR (470 MHz) δ 88.4 (s, 3F, minor), 90.8 (s, 3F, major), 99.2 (s, 3F, major), 99.3 (s, 3F, minor); EI MS m/z (relative intensity) 461 ($\text{M}^+ - 15$, 2), 302 (5), 247 (100), 73 (55); Anal. Calcd for $\text{C}_{18}\text{H}_{26}\text{F}_6\text{O}_4\text{Si}_2$: C, 45.37; H, 5.50. Found; C, 45.25; H, 5.44.

3,3,3-Trifluoro-2-[furyl(trimethylsilanyloxy)methyl]-2-trimethylsilanyloxypropanoic acid methyl ester (10e).

50%; colorless oil; Diastereomeric mixture (64:36 by GC analysis); IR (neat) 2968, 2908, 1772, 1758, 1440 cm^{-1} ; ^1H NMR (600 MHz) δ -0.03 (s, 9H, major), 0.05 (s, 9H, minor), 0.11 (s, 9H, minor), 0.21 (s, 9H, major), 3.76 (s, 3H, minor), 3.84 (s, 3H, major), 5.28 (s, 1H, major), 5.33 (s, 1H, minor), 6.29-6.32 (m, 2H, minor), 6.35 (dd, $J = 1.8$ Hz, $J = 3.0$ Hz, 1H, major), 6.36 (d, $J = 3.0$ Hz, 1H, major), 7.31 (dd, $J = 0.6$ Hz, $J = 1.8$ Hz, 1H, minor), 7.37 (dd, $J = 0.6$ Hz, $J = 1.8$ Hz, 1H, major); ^{19}F NMR (282 MHz) δ

86.4 (s, 3F, major), 89.2 (s, 3F, minor); EI MS m/z (relative intensity) 383 (M^+ -15, 2), 302 (trace), 169 (100), 73 (59); Anal. Calcd for $C_{15}H_{25}F_3O_5Si_2$: C, 45.21; H, 6.32. Found; C, 44.91; H, 6.33.

3,3,3-Trifluoro-2-[(4-methylphenyl)(trimethylsilanyloxy)methyl]-2-trimethylsilanyloxypropanoic acid methyl ester (10f).

20%; colorless oil; Diastereomeric mixture (86:14 by ^{19}F NMR); IR (neat) 2980, 2950, 2928, 1780, 1638, 1526, 1450 cm^{-1} ; 1H NMR (300 MHz) δ -0.07 (s, 9H, major), 0.00 (s, 9H, minor), 0.05 (s, 9H, minor), 0.20 (s, 9H, major), 2.32 (s, 3H, minor), 2.35 (s, 3H, major), 3.72 (s, 3H, minor), 3.86 (s, 3H, major), 5.19 (s, 1H, major), 5.22 (s, 1H, minor), 7.05 (d, J = 8.1 Hz, 2H, minor), 7.09 (d, J = 8.1 Hz, 2H, major), 7.15 (d, J = 8.1 Hz, 2H, minor), 7.23 (d, J = 8.1 Hz, 2H, major); ^{19}F NMR (282 MHz) δ 87.6 (s, 3F, major), 89.8 (s, 3F, minor); EI MS m/z (relative intensity) 405 (M^+ -17, 2), 193 (100), 73 (32); Anal. Calcd for $C_{18}H_{29}F_3O_4Si_2$: C, 51.16; H, 6.92. Found; C, 50.95; H, 6.97.

3,3,3-Trifluoro-2-hydroxy-2-[(4-methoxyphenyl)hydroxymethyl]propanoic acid methyl ester (12a).

87%; oily solid; Diastereomeric mixture (52:48 by 1H NMR); mp 48-49 °C ; IR (neat) 3512, 3016, 2968, 2848, 1754, 1766, 1616, 1588, 1516, 1444, 1480, 1458 cm^{-1} ; 1H NMR (600 MHz) δ 2.63 (s, J = 7.8 Hz, 1H, minor), 2.91 (d, J = 7.2 Hz, 1H, major), 3.70 (s, 3H, major), 3.78 (s, 3H, major), 3.80 (s, 3H, minor), 3.97 (s, 3H, minor), 3.99 (s, 1H, major), 4.18 (s, 1H, minor), 5.15 (d, J = 7.2 Hz, 1H, minor), 5.20 (s, J = 7.8 Hz, 1H, major), 6.84 (dd, J = 1.8 Hz, J = 6.6 Hz, 2H, major), 6.88 (dd, J = 1.8 Hz, J = 6.6 Hz, 2H, minor), 7.24 (dd, J = 1.8 Hz, J = 6.6 Hz, 2H, major), 7.38 (d, J = 1.8 Hz, J = 6.6 Hz, 2H, minor); ^{19}F NMR (282 MHz) δ 87.81 (s, 3F, minor), 87.82 (s, 3F, major); EI MS m/z (relative intensity) 294 (M^+ , trace), 137 (100), 109 (33), 94 (23); 77 (22); Anal. Calcd for $C_{12}H_{13}F_3O_5$: C, 48.99; H, 4.45. Found; C, 48.97; H, 4.41.

2-[(3-Chlorophenyl)hydroxymethyl]-3,3,3-trifluoro-2-hydroxypropanoic acid methyl ester (12b).

79%; white powder; Diastereomeric mixture (64:36 by ^1H NMR); mp 106-108 °C; IR (neat) 3508, 2965, 1754, 1760, 1600, 15780, 1480, 1442 cm^{-1} ; ^1H NMR (300 MHz) δ 2.71 (d, J = 7.8 Hz, 1H, minor), 3.00 (d, J = 8.1 Hz, 1H, major), 3.75 (s, 3H, major), 3.98 (s, 1H, major), 3.99 (s, 3H, minor), 4.19 (s, 1H, minor), 5.17 (d, J = 7.2 Hz, 1H, minor), 5.23 (d, J = 7.8 Hz, 1H, major), 7.18-7.48 (m, 4H x 2, major and minor mixed); ^{19}F NMR (282 MHz) δ 87.88 (s, 3F, major), 87.90 (s, 3F, minor); EI MS m/z (relative intensity) 298 (M^+ , trace), 141 (100), 113 (36), 77 (69); Anal. Calcd for $\text{C}_{11}\text{H}_{10}\text{ClF}_3\text{O}_4$: C, 44.24; H, 3.38. Found; C, 44.04; H, 3.46.

3,3,3-Trifluoro-2-hydroxy-2-[(4-trifluoromethylphenyl)hydroxymethyl]propanoic acid methyl ester (12d).

68%; white powder; Diastereomeric mixture (67:33 by ^1H NMR); mp 108-110 °C; IR (neat) 3525, 3000, 1765, 1640, 1460 cm^{-1} ; ^1H NMR (500 MHz) δ 2.63 (d, J = 7.8 Hz, 1H, minor), 2.93 (d, J = 8.1 Hz, 1H, major), 3.75 (s, 3H, major), 3.96 (s, 1H, major), 4.01 (s, 3H, minor), 4.19 (s, 1H, minor), 5.27 (d, J = 7.8 Hz, 1H, minor), 5.33 (d, J = 8.1 Hz, 1H, major), 7.45-7.66 (m, 4H x 2, major and minor mixed); ^{19}F NMR (470 MHz) δ 87.88 (s, 3F, minor), 87.90 (s, 3F, major), 98.94 (s, 3F, major), 98.95 (s, 3F, minor); EI MS m/z (relative intensity) 313 (M^+ -19, 4), 175 (100), 158 (45), 127 (80); Anal. Calcd for $\text{C}_{12}\text{H}_{10}\text{F}_6\text{O}_4$: C, 43.39; H, 3.03. Found; C, 43.39; H, 3.25.

3,3,3-Trifluoro-2-[(furyl)hydroxymethyl]-2-hydroxypropanoic acid methyl ester (12e).

73%; brown oil; Diastereomeric mixture (63:37 by ^{19}F NMR); IR (neat) 3504, 2968, 1756, 1504, 1444 cm^{-1} ; ^1H NMR (300 MHz) δ 3.04 (br, 1H x 2, major and minor mixed), 3.81 (s, 3H, minor), 3.98 (s, 3H, major), 4.07 (s, 1H, minor), 4.27 (s, 1H, major), 5.23 (br, 1H, major), 5.30 (br, 1H, minor), 6.32-6.35 (m, 2H x 2, major and minor mixed), 6.37 (dd, J = 1.8 Hz, J = 3.3 Hz, 2H, major), 6.46 (dd, J = 0.3 Hz, J = 3.3 Hz, 1H, major), 7.35 (dd, J = 0.9 Hz, J = 1.8 Hz, 1H, minor), 7.31 (dd, J = 0.9 Hz, J = 1.8 Hz, 1H, minor); ^{19}F NMR (282 MHz) δ 86.9 (s, 3F, major), 87.8 (s, 3F, minor); EI MS m/z (relative

intensity) 254 (M^+ , trace), 177 (4), 97 (100), 69 (13); Anal. Calcd for $C_9H_9F_3O_5$: C, 42.53; H, 3.57. Found; C, 42.72; H, 3.68.

3,3,3-Trifluoro-2-hydroxy-2-[1'-(p-tolyl)hydroxymethyl]propanoic acid methyl ester (12f).

88%; white powder; Diastereomeric mixture (65:35 by 1H NMR); mp 80-82 °C; IR (neat) 3525, 3070, 2990, 1755, 1525, 1458 cm^{-1} ; 1H NMR (300 MHz) δ 2.33 (s, 3H, major), 2.35 (s, 3H, minor), 2.47 (d, J = 7.5 Hz, 1H, minor), 2.77 (d, J = 8.4 Hz, 1H, major), 3.72 (s, 3H, major), 3.90 (s, 1H, major), 4.00 (s, 3H, minor), 4.12 (s, 1H, minor), 5.17 (d, J = 7.2 Hz, 1H, minor), 5.23 (d, J = 8.1 Hz, 1H, major), 7.12-7.36 (m, 4H x 2, major and minor mixed); ^{19}F NMR (282 MHz) δ 87.80 (s, 3F, minor), 87.83 (s, 3F, major); EI MS m/z (relative intensity) 200 (M^+ -78, 11), 120 (100), 92 (32) 77 (14); Anal. Calcd for $C_{12}H_{13}F_3O_4$: C, 51.80; H, 4.71. Found; C, 51.70; H, 4.75.

2-Trifluoromethyl-2,3-dihydroxy-5-phenylpentanoic acid methyl ester (12g).

53%; colorless oil; Diastereomeric mixture (58:42 by ^{19}F NMR); mp 42-44 °C; IR (neat) 3524, 3032, 2968, 2872, 1744, 1606, 1500, 1444 cm^{-1} ; 1H NMR (500 MHz) δ 1.49-1.57 (m, 1H, major), 1.84 (d, J = 11.5 Hz, 1H, major), 1.85-1.93 (m, 1H x 2, major and minor mixed), 1.99-2.05 (m, 1H, minor), 2.06 (d, J = 9.0 Hz, 1H, minor), 2.70 (m, 1H x 2, major and minor mixed), 2.93 (m, 1H x 2, major and minor mixed), 3.86 (s, 3H, minor), 3.94 (s, 3H, major), 3.95 (s, 1H, minor), 4.03 (s, 1H, major), 4.12 (m, 1H x 2, major and minor mixed), 7.16-7.32 (m, 5H x 2, major and minor mixed); ^{19}F NMR (282 MHz) δ 87.9 (s, 3F, minor), 88.1 (s, 3F, major); EI MS m/z (relative intensity) 292 (M^+ , 4), 158 (29), 117 (55), 91 (100); Anal. Calcd for $C_{13}H_{15}F_3O_4$: C, 53.43; H, 5.17. Found; C, 53.28; H, 5.04.

2-Trifluoromethyl-2,3-dihydroxy-4-hexenoic acid methyl ester (12h).

64%; colorless oil; Diastereomeric mixture (58:42 by 1H NMR); IR (neat) 3516, 3000, 2955, 2898, 1765, 1690, 1460 cm^{-1} ; 1H NMR (300 MHz) δ 1.71 (ddd, J = 0.3 Hz, J = 1.8 Hz, J = 6.6 Hz, 2H, minor), 1.75 (dd, J = 1.8 Hz, J = 6.6 Hz, 2H, major), 2.04 (d, J = 8.4 Hz, 1H, major), 2.28 (d, J = 8.4 Hz, 1H,

minor), 3.87 (s, 3H, minor), 3.95 (s, 3H, major), 3.96 (s, 1H, minor), 4.03 (d, $J = 0.3$ Hz, 1H, major), 4.55 (d, $J = 8.4$ Hz, 1H, major), 4.58 (d, $J = 8.4$ Hz, 1H, minor), 5.42-5.50 (m, 1H, minor), 5.59-5.67 (m, 1H, major), 5.77-5.94 (m, 1H x 2, major and minor mixed); ^{19}F NMR (282 MHz) δ 87.6 (s, 3F, major), 88.8 (s, 3F, minor); EI MS m/z (relative intensity) 211 ($\text{M}^+ - 17$, trace), 138 (6), 103 (6), 71 (100), 69 (20); Anal. Calcd for $\text{C}_8\text{H}_{11}\text{F}_3\text{O}_4$: C, 42.11; H, 4.86. Found; C, 42.09; H, 4.61.

2-Hydroxy-3-oxo-2-trifluoromethyloctanoic acid methyl ester (15b).

72%; diastereomeric mixture (50:50 by ^{19}F NMR); colorless oil; IR (neat) 3005, 2968, 2880, 1795, 1785, 1480, 1460 cm^{-1} ; ^1H NMR (300 MHz) δ 0.90 (m, 3H), 1.34 (ddd, $J = 2.1$ Hz, $J = 7.2$ Hz, $J = 7.5$ Hz, 4H), 1.69 (m, 2H), 2.49 (td, $J = 1.5$ Hz, $J = 7.5$ Hz, 2H), 3.85 (s, 3H), 5.47 (q, $J = 7.2$ Hz, 1H); ^{19}F NMR (282 MHz) δ 88.1 (s, 3F), 88.2 (s, 3F); EI MS m/z (relative intensity) 237 ($\text{M}^+ - 19$, trace), 200 (20), 99 (100), 70 (63), 55 (41); Anal. Calcd for $\text{C}_{10}\text{H}_{15}\text{F}_3\text{O}_4$: C, 46.88; H, 5.90. Found; C, 46.61; H, 5.84.