Preparation of C-3,5-Acyl Furanoses via Highly Selective Intramolecular Acyl Migration

Feng Xu*, Bryon Simmons, Kimberly Savary, Chunhua Yang, and Robert A. Reamer,

Department of Process Research, Merck Research Laboratories,

Rahway, NJ 07065, USA

feng_xu@merck.com

Supporting Information

Table of Contents

1. Method for HRMS	Page S2
2. Through process procedures for preparation of 4a	Page S2
3. Analytical data for 4a-i, 10, 12	Page S2 –S7
4. General procedure for acyl migration	Page S7
5. Preparation of 6a	Page S7
6. Analytical data for 6a-i, 11, 13	Page S7-S12
7. Preparation of 8	Page S12

Method for HRMS: High resolution mass spectroscopy (HRMS) measurements were acquired on Micromass Ultima US Quadrupole Time of Flight (QTof) mass spectrometer. Electrospray ionization was used in the negative ion mode. Samples were dissolved in MeCN and injected into a carrier composed of 50:50 water/MeCN with 0.02% HOAc. Compounds were measured as deprotonated molecular ions, M-H, deprotonated water adducts, M-H+H₂O or acetate adducts, M+CH₃COO⁻. The internal lock mass was raffinose. Mass calibration was performed using NaOAc as the calibrant. The mass spectrometer was preformed at a resolution (M/ΔM) of 10,000.

Through process procedures for preparation of **4a**

To a yellow solution of **1** (100 g, 0.308 mol) and pyridine (74.4 mL, 0.924 mol) in 700 mL of MeCN at ambient temperature was added p-toluoyl chloride (92 mL, 0.6776 mol) dropwise over 30 min. During the addition, the reaction temperature was controlled below 45 °C by using external cooling. The reaction was aged at 50 °C for 16 h to achieve >99% conversion. Then, HClO₄ (60%, 60 mL, 0.924 mol) and H₂O (250 mL) were added. After 10 h age at 55 °C, the reaction solution was allowed to cool to 0-5 °C. A solution of periodic acid (86 g, 0.3696 mol) in 200 mL of H₂O was added dropwise over 1-2 h. The reaction was further aged for 30 minutes at 0 – 5 °C. iPrOAc (500 mL) and H₂O (200 mL) were added. The organic phase was separated and washed with 500 mL of H₂O, 400 mL of half saturated NaHCO₃, 300 mL of 5% sodium thiosulfate and 5% NaCl solution. The organic phase was concentrated in vacuo to give the crude product in about 150 mL in iPrOAc, while MeCN was also removed during the concentration. This crude product can be used directly for acyl migration (see procedure B for the preparation of **6a**). **4a** : **5** is typically about 9:1. Three-step assay yield for **4a** and **5** is 88%.

4a could also be purified through SiO₂ column chromatography eluted with EtOAc/hexane (25:75). ¹H NMR (CDCl₃, 400 MHz): δ 9.57 (s, 1 H), 8.23 (d, *J* = 0.6 Hz, 1 H), 7.89 (d, *J* = 8.2 Hz, 2 H), 7.84 (d, *J* = 8.27 Hz, 2 H), 7.33 (m, 4 H), 7.29 (m, 1 H), 7.24 (d, *J* = 8.2 Hz, 2 H), 7.23 (d, *J* = 8.1 Hz, 2H), 5.90 (dd, *J* = 0.6, 8.2 Hz, 1 H),

5.84 (ddd, $J = 2.4, 4.5, 8.2$ Hz, 1 H), 4.76 (dd, $J = 2.4, 12.6$ Hz, 1 H), 4.72 (d, $J = 11.4$ Hz, 1 H), 4.62 (d, $J = 11.4$ Hz, 1 H), 4.42 (dd, $J = 4.5, 12.6$ Hz, 1 H), 2.41 (s, 3 H), 2.40 (s, 3 H), 1.45 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 202.1, 166.3, 165.0, 159.3, 144.5, 144.1, 137.7, 130.2, 130.0, 129.5, 129.4, 128.7, 128.0, 127.5, 127.0, 126.6, 83.3, 73.8, 69.3, 67.0, 62.6, 21.9, 16.5.

HRMS: Calcd for $\text{C}_{30}\text{H}_{30}\text{O}_8\text{-H}+\text{H}_2\text{O}$ m/z 535.1968; Found 535.1953.

^1H NMR (CDCl_3 , 400 MHz): δ 9.58 (s, 1 H), 8.25 (s, 1 H), 8.01 (m, 2 H), 7.95 (m, 2 H), 7.56 (m, 2 H), 7.46 (m, 4 H), 7.2 (m, 5 H), 5.93 (d, $J = 8.9$ Hz, 1 H), 5.86 (ddd, $J = 2.4, 4.6, 8.9$ Hz, 1 H), 4.81 (dd, $J = 2.4, 12.7$ Hz, 1 H), 4.72 (d, $J = 11.3$ Hz, 1 H), 4.62 (d, $J = 11.3$ Hz, 1 H), 4.45 (dd, $J = 4.6, 12.7$ Hz, 1 H), 1.46 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 202.1, 166.3, 164.9, 159.3, 137.6, 133.7, 133.4, 130.1, 130.0, 129.7, 129.2, 128.8, 128.7, 128.1, 127.5, 83.3, 73.6, 69.5, 67.0, 62.7, 16.4.

4c plus about 10% deformylated product: 82% isolated yield for three-step through process.

^1H NMR (CDCl_3 , 400 MHz): δ 9.58 (s, 1 H), 8.25 (s, 1 H), 7.94 (d, $J = 8.5$ Hz, 2 H), 7.85 (d, $J = 8.5$ Hz, 2 H), 7.42 (d, $J = 8.5$ Hz, 2 H), 7.41 (d, $J = 8.5$ Hz, 2 H), 7.34 (m, 5 H), 5.90 (d, $J = 8.3$ Hz, 1 H), 5.82 (ddd, $J = 2.3, 4.7, 8.3$ Hz, 1 H), 4.82 (dd, $J = 2.3, 12.6$ Hz, 1 H), 4.72 (d, $J = 11.3$ Hz, 1 H), 4.62 (d, $J = 11.3$ Hz, 1 H), 4.42 (dd, $J = 4.7, 12.6$ Hz, 1 H), 1.46 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 201.9, 165.4, 164.1, 159.3, 140.4, 140.0, 137.5, 131.4, 131.3, 129.2, 129.1, 128.7, 128.1, 128.0, 127.6, 127.5, 83.2, 73.3, 69.7, 67.1, 62.8, 16.3.

HRMS: Calcd for $\text{C}_{28}\text{H}_{24}\text{Cl}_2\text{O}_8+\text{Ac}^-$ m/z 617.0981; Found 617.0961.

4f plus about 5% deformylated product: 85% isolated yield for three-step through process.

^1H NMR (CDCl_3 , 400 MHz): δ 9.69 (s, 1 H), 8.10 (s, 1 H), 7.99 (m, 2 H), 7.96 (m, 2 H), 7.56 (m, 2 H), 7.44 (m, 4 H), 7.28 (m, 2 H), 7.23 (m, 3 H), 5.86 (ddd, $J = 0.8, 2.8, 7.6$ Hz, 1 H), 5.79 (ddd, $J = 2.7, 4.7, 7.6$ Hz, 1 H), 4.88 (dd, $J = 2.7, 12.7$ Hz, 1 H), 4.76 (d, $J = 11.4$ Hz, 1 H), 4.57 (d, $J = 11.4$ Hz, 1 H), 4.52 (dd, $J = 4.7, 12.7$ Hz, 1 H), 4.15 (m, 1 H).
 ^{13}C NMR (CDCl_3 , 100 MHz): δ 199.6, 166.2, 165.3, 159.5, 136.0, 133.8, 133.5, 130.1, 129.9, 129.6, 129.2, 128.9, 128.84, 128.81, 128.73, 128.71, 81.1, 73.9, 69.7, 69.3, 62.3.

^1H NMR (CDCl_3 , 400 MHz): δ 9.48 (s, 1 H), 8.30 (s, 1 H), 7.95 (d, $J = 8.0$ Hz, 2 H), 7.88 (d, $J = 8.4$ Hz, 2 H), 7.42 (m, 4 H), 7.35 (m, 1 H), 7.25 (m, 4 H), 6.07 (d, $J = 9.6$ Hz, 1 H), 5.78 (ddd, $J = 2.0, 4.4, 9.6$ Hz, 1 H), 4.92 (d, $J = 11.2$ Hz, 1 H), 4.78 (dd, $J = 2.0, 12.5$ Hz, 1 H), 4.68 (d, $J = 11.2$ Hz, 1 H), 4.40 (dd, $J = 4.4, 12.5$ Hz, 1 H), 2.41 (s, 3 H), 2.40 (s, 3 H), 2.06 (m, 1 H), 1.79 (m, 1 H), 1.39 (m, 1 H), 1.29 (m, 2 H), 1.20 (m, 1 H), 0.88 (t, $J = 7.2$ Hz, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 203.7, 166.1, 164.5, 144.4, 143.9, 137.5, 129.9, 129.8, 129.3, 129.2, 128.5, 127.8, 127.6, 126.8, 126.3, 85.0, 74.5, 68.9, 66.2, 62.1, 30.5, 24.5, 23.1, 21.7, 13.8.

HRMS: Calcd for $\text{C}_{33}\text{H}_{36}\text{O}_8 + \text{Ac}^-$ m/z 619.2543; Found 619.2512.

^1H NMR (CDCl_3 , 400 MHz): δ 9.49 (s, 1 H), 7.90 (d, J = 8.1 Hz, 2 H), 7.87 (d, J = 8.2 Hz, 2 H), 7.37 (m, 4 H), 7.26 (m, 5 H), 5.50 (ddd, J = 2.8, 2.8, 9.4 Hz, 1 H), 4.94 (dd, J = 3.1, 12.6 Hz, 1 H), 4.87 (d, J = 11.4 Hz, 1 H), 4.69 (d, J = 11.4 Hz, 1 H), 4.61 (dd, J = 2.5, 12.6 Hz, 1 H), 4.19 (dd, J = 6.0, 9.4 Hz, 1 H), 3.41 (d, J = 6.0 Hz, 1 H), 2.42 (s, 3 H), 2.41 (s, 3 H), 2.32 (m, 1 H), 1.82 (m, 1 H), 1.49–1.20 (m, 4 H), 0.88 (t, J = 7.1 Hz, 3 H).
 ^{13}C NMR (CDCl_3 , 100 MHz): δ 205.5, 167.8, 165.0, 144.45, 144.37, 138.3, 130.14, 130.12, 129.5, 129.4, 128.6, 127.8, 127.7, 126.82, 126.79, 86.0, 74.3, 71.6, 66.6, 63.4, 30.9, 25.1, 23.5, 21.9, 14.0.

HRMS: Calcd for $\text{C}_{32}\text{H}_{36}\text{O}_7 + \text{Ac}^-$ m/z 591.2600; Found 591.2577.

^1H NMR (CDCl_3 , 400 MHz): δ 9.51 (s, 1H), 8.17 (d, J = 0.6 Hz, 1 H), 7.90 (d, J = 8.2 Hz, 2H), 7.85 (d, J = 8.2 Hz, 2 H), 7.33 (m, 4 H), 7.29 (m, 2 H), 7.23 (m, 4 H), 6.12 (dd, J = 0.6, 8.0 Hz, 1 H), 5.90 (dd, J = 11.1, 17.7 Hz, 1 H), 5.87 (m, 1 H), 5.59 (d, J = 11.1 Hz, 1 H), 5.55 (d, J = 17.7 Hz, 1 H), 4.76 (dd, J = 2.4, 12.5 Hz, 1 H), 4.75 (d, J = 11.6 Hz, 1 H), 4.73 (d, J = 11.6 Hz, 1 H), 4.43 (dd, J = 5.2, 12.5 Hz, 1 H), 2.4 (s, 6 H).
 ^{13}C NMR (CDCl_3 , 100 MHz): δ 198.2, 166.1, 164.5, 159.0, 144.2, 143.8, 137.4, 130.4, 129.8, 129.7, 129.2, 129.0, 128.4, 127.7, 127.2, 126.7, 126.4, 121.6, 86.6, 72.7, 69.0, 67.4, 62.3, 21.6.

HRMS: Calcd for $\text{C}_{31}\text{H}_{30}\text{O}_8 + \text{Ac}^-$ m/z 589.2074; Found 589.2079.

^1H NMR (CDCl_3 , 400 MHz): δ 9.68 (s, 1H), 7.99 (d, J = 0.7 Hz, 1 H), 7.90 (d, J = 8.2 Hz, 2 H), 7.85 (d, J = 8.2 Hz, 2 H), 7.49 (m, 2 H), 7.37 (m, 8 H), 7.24 (m, 4 H), 6.47 (d, J = 7.6 Hz, 1 H), 5.95 (ddd, J = 2.6, 5.3, 7.6 Hz, 1 H), 4.78 (d, J = 11.7 Hz, 1 H), 4.72 (d, J

= 11.7 Hz, 1 H), 4.63 (dd, J = 2.6, 12.5 Hz, 1 H), 4.36 (dd, J = 5.3, 12.5 Hz, 1 H), 2.40 (s, 6 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 197.9, 166.3, 165.0, 159.1, 144.4, 144.0, 137.6, 133.0, 130.2, 130.0, 129.5, 129.4, 129.27, 129.23, 128.7, 128.1, 127.8, 127.6, 127.1, 126.6, 87.4, 73.3, 69.5, 68.0, 62.8, 21.9.

10 plus about 15% deformedylated product: 94% isolated yield for three-step through process.

^1H NMR (CDCl_3 , 400 MHz): δ 9.81 (s, 1 H), 8.10 (s, 1 H), 7.86 (m, 4H), 7.40 (d, J = 8.7 Hz, 2 H), 7.26 (m, 7H), 5.83 (dd, J = 2.9, 7.2 Hz, 1 H), 5.74 (m, 1 H), 4.87 (dd, J = 2.6, 12.7 Hz, 1 H), 4.75 (d, J = 11.6 Hz, 1 H), 4.56 (d, J = 11.6 Hz, 1 H), 4.48 (dd, J = 4.8, 12.7 Hz, 1 H), 4.13 (d, J = 4.2 Hz, 1 H), 2.41 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 199.6, 166.2, 164.5, 159.4, 144.3, 140.4, 136.0, 131.4, 127.0, 129.4, 129.1, 128.9, 128.8, 128.7, 127.7, 126.8, 81.0, 73.9, 70.1, 69.3, 21.9.

12 plus about 15% deformedylated product was isolated.

^1H NMR (CDCl_3 , 400 MHz): δ 9.63 (s, 1 H), 8.09 (s, 1 H), 7.88 (m, 4 H), 7.60 – 7.20 (m, 9 H), 5.85 (dd, J = 5.2, 10.0 Hz, 1 H), 5.78 (m, 1 H), 4.82 (d, J = 11.7 Hz, 1 H), 4.63 (d, J = 11.7 Hz, 1 H), 4.53 (dd, J = 4.4, 12.0 Hz, 1 H), 4.39 (dd, J = 5.61, 12.0 Hz, 1 H), 4.18 (m, 1 H), 2.40 (s, 6 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 199.4, 165.9, 165.1, 159.5, 144.3, 143.9, 136.0, 129.9, 129.7, 129.2, 129.1, 128.7, 128.6, 128.4, 128.3, 126.2, 80.1, 73.5, 69.8, 69.2, 63.8, 22.6, 22.5.

General procedure for acyl migration

To a solution of **4a-h** (1 equiv) in an alcohol solvent (10 – 20 mL/g) under N₂ was added iPr₂NH (1.5 equiv.) at 0-5 °C. The reaction solution was aged at specified temperature till the reaction was completed. The residue upon concentration *in vacuo* was purified on column chromatography to afford the furanoses and pyranoses.

Preparation of 6a

Procedure A: A solution of **4a** (1.0 g, 1.98 mmol) and iPr₂NH (300 mg, 2.97 mmol) in MeOH (10 mL) was aged at 0 – 5 °C for 1 day. The crude was purified on SiO₂ column eluted with iPrOAc/hexane (3:7) to give **6a** (840 mg, 86% yield).

Procedure B: A solution of **4a** and **5** (**4a**:**5** = ca. 9:1) in iPrOAc, obtained from a through process (started with 0.154 mol of **1**), was concentrated *in vacuo* to about 150 mL. MeOH (400 mL) was added and the solution was cooled to 0 °C and diisopropylamine (7.8 g, 0.077 mol) was added over 10 min. The solution was aged at 0 °C for 2 days. Then, 2N HCl (160 mL) was added dropwise at 0 to 10 °C. iPrOAc (500 mL) and 250 mL of water were added and the organic phase was separated. The organic phase was washed with 150 mL of 10% NaHCO₃, followed by 150 mL of 10% NaCl. Then, the organic phase was concentrated *in vacuo* to about 300 mL. By HPLC: >98 A% conversion and 96% selectivity (**6a** vs. **7a**). 84% assay yield (63.4 g of **6a**, four steps starting from **1a**).

¹H NMR (CDCl₃, 400 MHz): δ The ratio of α:β isomers in CDCl₃ is about 5 to 2. For the α isomer: δ 7.92 (d, *J* = 8.1 Hz, 2 H), 7.88 (d, *J* = 8.8 Hz, 2 H), 7.39 – 7.18 (m, 9 H), 5.34 (d, *J* = 5.9 Hz, 1 H), 5.21 (s, 1 H), 4.73 (d, *J* = 11.0, 1 H), 4.70 – 4.62 (m, 2 H), 4.61 (d, *J* = 11.0 Hz, 1 H), 4.52 (dd, *J* = 5.1, 11.7 Hz, 1 H), 4.20 (s, br, 1 H), 2.42 (s, 3 H), 2.40 (s, 3 H), 1.63 (s, 3 H); for the β isomer: δ 7.95 – 7.87 (m, 4 H), 7.38 – 7.16 (m, 9 H), 5.66 (d, *J* = 7.3 Hz, 1 H), 5.36 (s, 1 H), 4.77 (d, *J* = 11.7 Hz, 1 H), 4.70 – 4.62 (m, 3 H), 4.51 (m, 1 H), 4.20 (s, br, 1 H), 2.42 (s, 3 H), 2.38 (s, 3 H), 1.55 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 166.7, 166.5, 166.3, 166.1, 144.6, 144.4, 144.0, 143.9, 137.8, 130.2, 130.1, 130.0, 129.4, 129.38, 129.3, 128.6, 128.5, 128.0, 127.5, 127.46, 127.3, 127.2, 127.1, 126.8, 126.5, 101.5, 101.3, 83.9, 80.6, 79.2, 78.6, 77.4, 76.9, 67.4, 66.2, 65.3, 64.2, 21.9, 21.78, 19.4, 15.8.

HRMS: Calcd for $\text{C}_{29}\text{H}_{30}\text{O}_7\text{-H}$ m/z 489.1913; Found 489.1916.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 10 to 6. For the α isomer: δ 8.05-7.98 (m, 4H), 7.60 – 7.29 (m, 11 H), 5.36 (d, J = 6.0 Hz, 1 H), 5.22 (s, 1 H), 4.72 (d, J = 11.2 Hz, 1 H), 4.73 – 4.64 (m, 2 H), 4.62 (d, J = 11.2 Hz, 1 H), 4.55 (dd, J = 4.8, 11.6 Hz, 1 H), 4.12 (br s, 1 H), 1.65 (s, 3 H); for the β isomer: δ 8.05-7.98 (m, 4H), 7.60 – 7.29 (m, 11 H), 5.68 (d, J = 7.6 Hz, 1 H), 5.38 (s, 1 H), 4.77 (d, J = 11.6 Hz, 1 H), 4.73 – 4.64 (m, 4 H), 3.05 (s, br, 1 H), 1.56 (s, 3 H).

Partial ^{13}C NMR (CDCl_3 , 100 MHz): δ 166.5, 166.2, 166.0, 137.7, 133.8, 133.7, 133.3, 133.2, 130.2, 130.1, 129.9, 129.3, 128.9, 128.7, 128.63, 128.57, 128.54, 128.0, 127.6, 127.5, 127.1, 101.5, 101.2, 83.9, 80.7, 79.1, 78.5, 77.7, 77.1, 67.5, 66.2, 65.5, 64.4, 19.5, 15.8.

HRMS: Calcd for $\text{C}_{27}\text{H}_{26}\text{O}_7\text{-H}$ m/z 461.1600; Found 461.1622.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 10 to 7. For the α isomer: δ 8.00-7.90 (m, 4H), 7.43 – 7.28 (m, 9 H), 5.32 (d, J = 5.6 Hz, 1 H), 5.25 (s, 1 H), 4.74 – 4.52 (m, 5 H), 4.11 (s, br, 1 H), 1.66 (s, 3 H); for the β isomer: δ 8.00-7.90 (m, 4H), 7.43 – 7.28 (m, 9 H), 5.66 (d, J = 7.6 Hz, 1 H), 5.42 (s, 1 H), 4.76 (d, J = 12.0 Hz, 1 H), 4.74 – 4.52 (m, 4 H), 4.30 (br s, 1 H), 1.58 (s, 3 H).

Partial ^{13}C NMR (CDCl_3 , 100 MHz): δ 165.7, 165.6, 165.4, 165.2, 140.4, 140.3, 139.9, 139.8, 137.6, 131.5, 131.4, 131.3, 129.10, 129.07, 129.0, 128.9, 128.7, 128.6, 128.3, 128.1, 128.0, 127.7, 127.6, 127.4, 127.0, 101.4, 101.0, 83.8, 80.7, 79.0, 78.3, 77.9, 77.3, 67.5, 66.2, 65.6, 64.6, 19.6, 15.7.

HRMS: Calcd for $\text{C}_{27}\text{H}_{24}\text{Cl}_2\text{O}_7\text{-H}$ m/z 529.0821; Found 529.0815.

^1H NMR (CDCl_3 , 600 MHz): The ratio of α : β isomers in CDCl_3 is about 4 to 1. For the α isomer: δ 7.37 - 7.31 (m, 5 H), 5.13 (d, J = 7.9 Hz, 1 H), 4.97 (d, J = 5.7 Hz, 1 H), 4.67 (d, J = 10.6 Hz, 1 H), 4.55 (d, J = 10.6 Hz, 1 H), 4.38 (m, 1 H), 4.27 (dd, J = 12.1, 3.8 Hz, 1 H), 4.19 (dd, J = 12.1, 4.9, 1 H), 3.95 (d, J = 8.3 Hz, 1 H), 1.55 (s, 3 H), 1.23 (s, 9 H), 1.13 (s, 9 H); for the β isomer (selected data) : δ 5.27 (br s, 1 H), 5.25 (d, J = 7.6 Hz, 1 H), 4.71 (d, J = 11.7 Hz, 1 H), 4.60 (d, J = 11.7 Hz, 1 H), 4.34 (m, 1 H), 2.78 (br s, 1 H), 1.45 (s, 3 H).

^{13}C NMR (CDCl_3 , 150 MHz): δ (α isomer) 178.15, 177.48, 137.49, 128.47, 127.93, 127.53, 100.86, 80.16, 78.96, 75.88, 67.06, 63.42, 38.84, 38.69, 27.19, 26.99, 19.20.

HRMS: Calcd for $\text{C}_{23}\text{H}_{34}\text{O}_7\text{-H}$ m/z 421.2226; Found 421.2206.

^1H NMR (CDCl_3 , 600 MHz): The initial ratio of α : β in CDCl_3 , was 2:3. Aging in CDCl_3 resulted in a α : β ratio of 5:3. For the α isomer: δ 8.07 (m, 2H), 7.97 (m, 2H), 7.61 (m, 1H), 7.56 (m, 1H), 7.47 (m, 2H), 7.40 (m, 2H), 7.29-7.22 (overlapping m, 5H), 5.50 (dd, J = 8.7, 4.2 Hz, 1H), 5.45 (dd, J = 6.0, 4.9 Hz, 1H), 4.70 (d, J = 11.7 Hz, 1H), 4.68 (m, 1H), 4.64 (d, J = 11.7 Hz, 1H), 4.59 (dd, J = 12.1, 3.8 Hz, 1H), 4.49 (dd, J = 12.1, 4.5 Hz, 1H), 4.27 (dd, J = 6.0, 4.5 Hz, 1H), 3.87 (d, J = 8.7 Hz, 1H); for the β

isomer: δ 8.08 (m, 2H), 8.04 (m, 2H), 7.60 (m, 1H), 7.54 (m, 1H), 7.45 (m, 2H), 7.39 (m, 2H), 7.29-7.22 (overlapping m, 5H), 5.59 (dd, J = 6.4, 4.9 Hz, 1H), 5.48 (dd, J = 3.4, 1.5 Hz, 1H), 4.69 (obscured m, 1H), 4.4 (s, 2H), 4.63, (dd, J = 12.1, 4.2 Hz, 1H), 4.58 (dd, J = 12.1, 5.3 Hz, 1H), 4.29 (dd, J = 4.9, 1.5 Hz, 1H), 3.01 (d, J = 3.4 Hz, 1H).

^{13}C NMR (CDCl_3 , 150 MHz): δ 166.4, 166.1, 165.9, 165.9, 137.4, 136.6, 133.6, 133.4, 133.2, 133.1, 129.9, 129.8, 129.7, 129.6, 129.6, 129.4, 129.2, 128.6, 128.5, 128.41, 128.39, 128.34, 128.29, 128.0, 127.8, 127.7, 100.9, 96.1, 81.4, 79.0, 78.5, 76.6, 73.8, 73.5, 73.1, 72.2, 65.0, 64.1.

HRMS: Calcd for $\text{C}_{26}\text{H}_{24}\text{O}_7\text{-H}$ m/z 447.1444; Found 447.1455.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 1 to 2.5. For the α isomer: δ 8.06 (m, 4H), 7.48 – 7.30 (m, 11 H), 5.61 (d, J = 2.4 Hz, 1 H), 5.48 (dd, J = 0.8, 3.6 Hz, 1 H), 4.79 (d, J = 12.0 Hz, 1 H), 4.74 – 4.66 (m, 3 H), 4.59 (dd, J = 5.6, 11.6 Hz, 1 H), 4.21 (s, 1 H), 2.90 (d, J = 3.6 Hz, 1 H); for the β isomer: δ 8.05 (m, 4H), 7.62 – 7.30 (m, 11 H), 5.53 (m, 2 H), 4.89 (d, J = 11.7 Hz, 1 H), 4.75 (d, J = 11.7 Hz, 1 H), 4.74 – 4.64 (m, 2 H), 4.38 (ddd, J = 2.9, 5.2, 8.1 Hz, 1 H), 4.13 (dd, J = 2.2, 3.7 Hz, 1 H), 3.97 (d, J = 9.5 Hz, 1 H).

Partial ^{13}C NMR (CDCl_3 , 100 MHz): δ 166.54, 166.08, 137.45, 133.84, 133.72, 133.28, 133.20, 130.04, 130.00, 128.88, 128.76, 128.73, 128.56, 128.51, 128.36, 128.15, 128.04, 101.93, 97.45, 87.49, 81.65, 81.41, 79.33, 78.36, 73.12, 72.32, 65.30, 64.63.

HRMS: Calcd for $\text{C}_{26}\text{H}_{24}\text{O}_7\text{-H}$ m/z 447.1444; Found 447.1455.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 1 to 1. δ 7.96 – 7.87 (m, 8 H), 7.42 – 7.14 (m, 18 H), 5.76 (d, $J = 7.3$ Hz, 1 H), 5.43 (d, $J = 5.9$ Hz, 1 H), 5.39 (s, 1 H), 5.35 (s, 1 H), 4.92 (d, $J = 11.0$ Hz, 1 H), 4.74 (d, $J = 11.0$ Hz, 1 H), 4.70 – 4.49 (m, 9 H), 4.26 (s, br, 1 H), 3.78 (s, br, 1 H), 2.42 (s, 6 H), 2.41 (s, 3 H), 2.37 (s, 3 H), 2.06 – 1.79 (m, 4 H), 1.58 – 1.27 (m, 8 H), 0.94 (t, $J = 7.3$ Hz, 3 H), 0.89 (t, $J = 7.3$ Hz, 3 H).

Partial ^{13}C NMR (CDCl_3 , 100 MHz): δ 166.8, 166.5, 166.0, 165.9, 144.5, 144.4, 143.9, 143.8, 137.7, 130.14, 130.08, 129.9, 129.4, 129.22, 129.18, 128.6, 128.53, 128.48, 127.9, 127.6, 127.5, 127.2, 127.0, 126.5, 101.4, 99.7, 85.8, 82.8, 79.1, 79.0, 77.7, 75.7, 74.3, 66.6, 66.0, 65.6, 64.2, 31.7, 29.6, 25.4, 25.3, 23.3, 23.1, 21.82, 21.79, 14.1.

HRMS: Calcd for $\text{C}_{32}\text{H}_{36}\text{O}_7\text{-H}$ m/z 531.2383; Found 531.2388.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 5 to 2. For the α isomer: δ 7.98 (d, $J = 8.5$ Hz, 2 H), 7.95 (d, $J = 8.1$ Hz, 2 H), 7.43 (d, $J = 8.5$ Hz, 2 H), 7.34 (m, 5 H), 7.11 (d, $J = 8.1$ Hz, 2 H), 5.61 (s, 1 H), 5.45 (d, $J = 3.3$ Hz, 1 H), 4.78 (d, $J = 11.6$ Hz, 1 H), 4.69 (m, 1 H), 4.68 (d, $J = 11.6$ Hz, 1 H), 4.65 (dd, $J = 4.0, 11.6$ Hz, 1 H), 4.56 (dd, $J = 5.6, 11.6$ Hz, 1 H), 4.18 (s, 1 H), 2.94 (s, br, 1 H), 2.38 (s, 3 H); for the β isomer: δ 7.94 (d, $J = 8.0$ Hz, 4 H), 7.45 (d, $J = 8.0$, 2 H), 7.34 (m, 5 H), 7.18 (d, $J = 8.0$ Hz, 1 H), 5.51 (m, 2 H), 4.86 (d, $J = 11.9$ Hz, 1 H), 4.75 (d, $J = 11.9$ Hz, 1 H), 4.69 (m, 1 H), 4.60 (m, 1 H), 4.35 (ddd, $J = 3.4, 3.4, 6.0$ Hz, 1 H), 4.11 (dd, $J = 3.3, 3.3$, 1 H), 3.98 (br d, $J = 8.7$ Hz, 1 H), 2.39 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 166.6, 165.2, 165.1, 144.0, 143.9, 140.4, 140.2, 137.4, 136.8, 131.42, 131.36, 130.1, 130.0, 129.3, 129.2, 129.12, 129.07, 128.9, 128.7, 128.6, 128.4, 128.1, 128.04, 127.79, 127.2, 101.9, 97.4, 87.4, 81.6, 81.4, 79.3, 78.6, 77.9, 73.1, 72.3, 65.1, 64.4, 21.8.

HRMS: Calcd for $\text{C}_{27}\text{H}_{25}\text{ClO}_7\text{-H}$ m/z 495.1211; Found 495.1229.

^1H NMR (CDCl_3 , 400 MHz): The ratio of α : β isomers in CDCl_3 is about 1 to 2. For the β isomer: δ 7.89 (m, 4 H), 7.40 – 7.30 (m, 5 H), 7.26 – 7.19 (m, 4 H), 5.64 (dd, J = 1.6, 4.0 Hz, 1 H), 5.60 (s, br, 1 H), 4.95 (d, J = 11.6 Hz, 1 H), 4.78 (m, 2 H), 4.55 (d, J = 5.8 Hz, 2 H), 4.12 (dd, J = 1.6, 4.0 Hz, 1 H), 4.54 (s, br, 1 H), 2.42 (s, 3 H), 2.41 (s, 3 H); for the α isomer: δ 7.90 (m, 4 H), 7.40 – 7.30 (m, 5 H), 7.26 – 7.19 (m, 4 H), 5.66 (dd, J = 1.3, 4.8 Hz, 1 H), 5.48 (s, 1 H), 4.86 – 4.69 (m, 4 H), 4.57 (dd, J = 5.2, 11.6 Hz, 1H), 4.16 (s, 1 H), 3.00 (s, br, 1 H), 2.41 (s, 3 H), 2.40 (s, 3 H).

^{13}C NMR (CDCl_3 , 100 MHz): δ 166.4, 165.9, 144.8, 144.0, 136.7, 130.00, 129.97, 129.5, 129.3, 128.9, 128.7, 128.6, 128.3, 128.2, 128.0, 127.2, 126.5, 102.2, 96.8, 87.1, 81.2, 78.9, 76.4, 75.8, 75.7, 73.5, 72.4, 63.8, 62.7, 21.91, 21.86.

HRMS: Calcd for $\text{C}_{28}\text{H}_{28}\text{O}_7\text{-H}$ m/z 475.1757; Found 475.1774.

Preparation of 8

To a *i*PrOAc solution of **6a** from above procedure was added Charcoal (Darco G60, 500g). The slurry was aged for 1 h at ambient temperature. Then, the slurry was filtered through Solka Floc (10 g) to remove charcoal. The organic phase was diluted with *i*PrOAc to 500 mL. Pd/C (1.5 kg, 10%) was charged. The reaction mixture was subjected to hydrogenolysis (45 psi H_2 , 50-60 °C) for 35 h. The reaction mixture was cooled to ambient temperature and was filtered through Solka Floc. The *i*PrOAc solution was concentrated to about 135 mL. The solution was heated to 50 °C and 90 mL of heptane was added in one portion. The solution was seeded with 2.5 g of **8** and allowed to age for 2 h at 50 °C. Heptane (270 mL) was added dropwise over 2.5 h. The slurry was aged at 50 °C for 1 h, then cooled to ambient temperature and aged for 2 h before filtration. The wet cake was displacement washed with 90 mL of 20% *i*PrOAc/heptane followed by 120

mL x 2 of 10% iPrOAc/heptane. The solids were suction dried under a N₂ atmosphere at ambient temperature to give 4.54 kg of **8**. 90 wt%. 66% isolated for overall five-step through process.

¹H NMR (CDCl₃, 400 MHz): The ratio of α:β isomers in CDCl₃ is about 5 to 1. For the α isomer: δ 7.95 – 7.90 (m, 4H), 7.26 (d, *J* = 8.0 Hz, 2 H), 7.17 (d, *J* = 8.0 Hz, 2 H), 5.53 (d, *J* = 7.2 Hz, 1 H), 5.22 (d, *J* = 2.8 Hz, 1 H), 4.65 – 4.49 (m, 3 H), 3.08 (d, *J* = 3.2 Hz, 1 H), 2.44 (s, 3 H), 2.38 (s, 3 H), 2.26 (s, 1 H), 1.44 (s, 3 H); for the β isomer: δ 7.95 – 7.90 (m, 4H), 7.27 (d, *J* = 8.0 Hz, 2 H), 7.22 (d, *J* = 8.0 Hz, 2 H), 5.16 (d, *J* = 5.6 Hz, 1 H), 5.12 (d, *J* = 5.6 Hz, 1 H), 4.66 – 4.49 (m, 3 H), 3.54 (d, *J* = 5.6 Hz, 1 H), 2.91 (s, 1 H), 2.43 (s, 3 H), 2.40 (s, 3 H), 1.44 (s, 3 H).

¹³C NMR (CDCl₃, 100 MHz): δ 166.6, 166.3, 165.9, 165.7, 144.6, 144.3, 143.8, 143.7, 129.9, 129.7, 129.3, 129.2, 129.1, 129.0, 127.0, 126.9, 126.4, 126.2, 102.9, 100.8, 79.8, 79.2, 78.7, 76.9, 76.5, 76.4, 65.5, 64.0, 23.5, 21.7, 21.6, 20.0.

HRMS: Calcd for C₂₂H₂₄O₇-H m/z 399.1444; Found 399.1453.