This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Prospective association of TV viewing with acute phase reactants and coagulation markers: English Longitudinal Study of Ageing

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1016/j.atherosclerosis.2015.02.009>

PUBLISHER

© Elsevier Ireland Ltd.

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Hamer, Mark, Lee Smith, and Emmanuel Stamatakis. 2019. "Prospective Association of TV Viewing with Acute Phase Reactants and Coagulation Markers: English Longitudinal Study of Ageing". figshare.
<https://hdl.handle.net/2134/19160>.

Prospective association of TV viewing with acute phase reactants and coagulation

markers: English Longitudinal Study of Ageing

Mark Hamer¹, PhD; Lee Smith^{1,2}, PhD; Emmanuel Stamatakis^{1,3,4,5} PhD

¹Physical Activity Research Group, Department of Epidemiology and Public Health, University College London, UK; ²Health Behaviour Research Centre, Department of Epidemiology and Public Health, University College London; ³Prevention Research Collaboration, Sydney School of Public Health, University of Sydney, Australia; ⁴Charles Perkins Centre, University of Sydney, Australia; ⁵Exercise and Sports Science, Faculty of Health Sciences, University of Sydney, Australia.

Correspondence to: Mark Hamer, PhD, Department of Epidemiology and Public Health, 1-19 Torrington Place, University College London, London, WC1E 6BT, UK.

E-mail: m.hamer@ucl.ac.uk; phone: +44 207 6795969; fax: +44 207 9168542

Word count = 2,697

Running head: Television viewing time and inflammation

Abstract

Objective: Inflammatory processes are putative mechanisms underlying the detrimental health effects of sedentary behaviour but no long-term prospective data are available. We examined the longitudinal association between TV viewing, physical activity and inflammatory markers over a 4-year follow-up period.

Methods: Participants were 3612 men and women (mean age 64.1 ± 8.2 years) from the English Longitudinal Study of Ageing. Self-reported daily TV viewing was measured at baseline and 2 years follow up. Inflammatory markers (serum high-sensitivity C-reactive protein [CRP], white blood cell count [WBC], and fibrinogen) were measured at baseline (2008/09) and 4 years follow-up (2012/13).

Results: On average, participants viewed TV for 5.1 ± 4.0 hrs/d, and there was an increase of 1.9 hr/wk TV viewing over 2 years. In linear models adjusted for covariates including physical activity, TV viewing was not associated with $\log_e$ CRP at follow-up ($B=0.004$, 95% CI, -0.001, 0.009, $p=0.09$) but was associated with WBC ($B=0.018$, 95% CI, 0.005, 0.031, $p=0.006$), and fibrinogen ($B=0.004$, 95% CI, 0.00, 0.008, $p=0.035$). In contrast, physical activity was inversely associated with CRP ($p=0.047$) and WBC ($p=0.026$), but not fibrinogen ($p=0.22$). An increase in TV viewing (of at least 1 hr/d) was associated with higher concentrations of CRP ($p=0.015$) and WBC ($p=0.05$) at follow up after adjustment for covariates and baseline TV viewing.

Conclusions: Physical activity and sedentary behaviour have contrasting associations with markers of low grade inflammation over 4 years of follow-up. These behaviors may be important in influencing the pro-inflammatory state seen with ageing.

Key words: Adiposity; Blood Coagulation; Exercise; Inflammation; Sedentary Lifestyle

Introduction

Pro-inflammatory status appears to increase with age, that may underlie biological mechanisms responsible for cardiovascular disease (CVD) and other age-related diseases.¹⁻⁴ Physical activity may have beneficial anti-inflammatory effects as evidence from epidemiological studies has consistently demonstrated an inverse association between physical activity and markers of low grade systemic inflammation.⁵⁻⁷ The anti-inflammatory effects of exercise may partly explain the well-documented cardio-protective effects of physical activity.⁸⁻¹⁰ In contrast, little is known about the association between sedentary behaviours and low grade inflammation.

Sedentary behaviour has been conceptualised as a distinct domain of behaviour, which may pose a risk to health independently of physical activity.¹¹ Sedentary behaviour has been consistently associated with risk of CVD in population cohort studies,¹²⁻¹⁵ although the mechanisms remain poorly understood. Television (TV) viewing is a major component of leisure sedentary time¹⁶, particularly in older adults¹⁷ and is distinct from other sedentary behaviours in that it is passive and encourages prolonged periods of sitting. Moreover, TV viewing is likely to influence other behaviour, for example, dietary habits by encouraging an increased consumption of energy-dense food.¹⁸ Thus, it remains unknown if excess TV viewing is simply a marker of energy imbalance and other confounding or if results are driven by actual postural effects (ie, prolonged sitting).

Several studies have demonstrated cross-sectional associations between TV viewing or objective sedentary time with markers of low grade inflammation.¹⁹⁻²² However, there are a paucity of longitudinal data. We recently demonstrated an association between TV viewing

(aged 23) and inflammatory markers at follow up (aged 44) in a birth cohort study but were unable to infer true changes over time since biomarkers were not collected at baseline.²³

The aim of this study was therefore to examine longitudinal associations between TV viewing, physical activity and inflammatory markers over a 4-year follow-up period in a cohort of older adults. Uniquely, we also explored associations between changes in TV viewing on subsequent inflammatory markers.

Materials and Methods

Study sample and procedures

The English Longitudinal Study of Ageing (ELSA) was initiated in 2002 containing a nationally representative sample of the English population living in households.²⁴ The ELSA cohort consists of men and women born on or before 29 February 1952. Multistage stratified probability sampling was used to select the sample with postcode sectors selected at the first stage and household addresses selected at the second stage. For the purposes of the present analyses, data collected at wave 4 (2008/9) were used as the baseline as this was the first occasion information on TV viewing was gathered. Clinical information was gathered by nurses in participants' homes at wave 4 and at follow-up (wave 6; 2012/13). In addition, data on TV viewing was also collected at wave 5 (2010/11). Participants gave full informed written consent to participate in the study. Ethical approval was obtained from the London Multi-center Research Ethics Committee, compliant with the Declaration of Helsinki.

Baseline TV viewing and physical activity

At waves 4 and 5 participants were asked to recall “How many hours of television do you watch on an ordinary day or evening, that is, Monday to Friday” and “How many hours of television do you normally watch in total over the weekend, that is, Saturday and Sunday.” Average daily time spent watching TV was calculated as $\{(weekday\ TV\ time \times 5) + (Weekend\ TV\ time)\}/7$. We calculated the 2 year change in TV viewing by subtracting total reported weekly TV viewing hours at wave 5 from wave 4.

Participants were asked three questions regarding how often they took part in vigorous, moderate- and low-intensity physical activity, using prompt cards with different activities to help them interpret physical activity intensities. Response options for each question were: more than once a week, once a week, one to three times a month and hardly ever/never. Physical activity was then categorised as: inactive (participants that reported no moderate or vigorous activity at least once a week); moderate activity at least once a week (but no vigorous); vigorous activity at least once a week. The physical activity and TV viewing measures used in ELSA have demonstrated excellent convergent validity in grading a plethora of psychosocial, physical and biochemical risk factors in previous work.²⁵⁻²⁸

Clinical assessments

Nurses collected anthropometric data (weight, height), and blood samples. Participants' body weight was measured using Tanita electronic scales without shoes and in light clothing, and height was measured using a Stadiometer with the Frankfort plane in the horizontal position. Body mass index (BMI) was calculated using the standard formulae [weight (kilograms)/height (meters) squared]. Blood samples were analyzed for several inflammatory marker categories, including acute phase reactants (high sensitivity C-reactive

protein [CRP], white blood cell count [WBC]) and coagulation products (fibrinogen). Blood analysis was carried out at the Royal Victoria Infirmary (Newcastle-upon-Tyne, UK). Detailed information on the technicalities of the blood analysis, the internal quality control, and the external quality assessment for the laboratory have been described elsewhere.²⁹

Covariates

Trained interviewers asked questions on cigarette smoking (current or non-smoker), alcohol intake (categorised as; at least five times a week, at least once a week, monthly, rarely/never), use of prescribed medication (including medication for diabetes, high blood pressure, cholesterol, blood thinning), chronic illness, and depressive symptoms (using the 8-item Centre of Epidemiological Studies Depression scale³⁰). Disability was assessed based on participants' responses to interviewer questions on perceived difficulties in six basic activities (e.g. difficulty dressing) and seven instrumental activities of daily living (e.g. difficulty preparing a hot meal).³¹ Participants reporting difficulties in one or more activities were considered to have some degree of disability.

Statistical analyses

Characteristics of the study population at baseline were described as categorical variables. Log transformations were used to normalise the distribution of CRP. We examined prospective associations between baseline TV and inflammatory markers (CRP, WBC, and fibrinogen) using linear regression models with TV time entered as a continuous independent variable. We tested for interactions with sex but none were noted. The models were adjusted for age, sex, baseline concentration of inflammatory marker, smoking, alcohol, medications, depressive symptoms, long standing illness, disability, body mass

index, and physical activity. These covariates were selected a priori based on previous literature.^{7,17,23,26} Similar models were run to examine the association between physical activity and inflammatory markers. In order to assess the association between changes in TV viewing on inflammatory markers we generated three categories (stable; reduction of at least 1 hr/d; increase of at least 1 hr/d). A change in 1hr/d TV was chosen as it roughly reflected the top and bottom quarter of the distribution. Using dummy variables (with the stable group designated as the referent category), TV change was regressed onto inflammatory markers at follow up, using the same covariates described above. All analyses were conducted using SPSS version 21.

Results

A sample size of 5463 provided complete baseline data although 1851 participants were lost to follow-up leaving a final analytic sample of 3612. Compared with the analytic sample, those lost to follow-up were slightly older (64.1 vs. 65.9 yrs, $p=0.001$), had higher baseline logCRP concentration (1.15 vs. 1.22, $p=0.001$), higher baseline TV viewing (5.1 vs. 5.5 hr/d, $p=0.001$), although there were no differences in sex distribution (% men; 45.0 vs 45.9, $p=0.53$) or body mass index (28.0 vs. 28.2 kg/m², $p=0.09$). A further 83 and 158 participants had missing data on WBC and fibrinogen, respectively, thus slightly different sample sizes were used in analysis of different inflammatory markers.

The baseline characteristics of the sample are shown in Table 1. High levels of TV viewing (>6 hr/d) were reported in 26.4% of participants and 15.0% of the sample reported no weekly physical activity of at least moderate intensity. Obesity was prevalent in 28.3% and clinically elevated CRP was observed in 31.2% of the sample.

Between baseline and follow-up, there was a small reduction in CRP (median [interquartile range]; 1.80 [2.88] vs. 1.50 [2.30] mg/L, $p=0.001$) and fibrinogen (mean $\pm$ SD; 3.33 ± 0.54 vs. 2.96 ± 0.52 g/L, $p=0.001$) but an increase in WBC (mean $\pm$ SD; 6.29 ± 2.0 vs. $6.42 \pm 1.94 \times 10^9$ cells/L, $p=0.001$). In models adjusted for age, sex, and baseline CRP, both TV viewing and physical activity were associated with logCRP at 4 years follow up (Table 2). In final adjusted models, only physical activity remained (inversely) associated with logCRP ($p=0.047$). Both TV viewing ($p=0.006$) and physical activity ($p=0.026$) remained associated with WBC in final adjusted models (Table 2). TV viewing remained associated with fibrinogen in final adjusted models ($p=0.035$) although physical activity did not ($p=0.22$) (Table 2). The results remained unchanged when we adjusted for waist circumference instead of BMI. For example, in final adjusted models TV viewing was not associated with log_eCRP ($p=0.08$) but was associated with WBC ($p=0.016$), and fibrinogen ($p=0.020$). In order to identify if one covariate in particular attenuated the effect estimates we added both BMI and physical activity separately into Model 1 before other covariates. However, this strategy caused minimal changes to the effect estimate leading us to conclude that possible confounding influences were explained evenly across all covariates, and not by one in particular.

We performed several sensitivity analyses. First, to explore possible reverse causation we removed participants ($n=802$) that reported any cardiovascular diseases (angina, myocardial infarction, congestive heart failure, heart murmur, arrhythmia) over the first four waves of ELSA. The results for TV viewing were unchanged in final adjusted models; CRP ($B=0.004$, 95% CI, -0.002, 0.009; $p=0.17$), WBC ($B=0.016$, 95% CI, 0.003, 0.030; $p=0.019$), fibrinogen ($B=0.004$, 95% CI, 0.000, 0.008; $p=0.06$). Second, we performed analyses to explore the

possible additive effects of physical activity and TV viewing. Participants were split into four groups based on combinations of high/low physical activity (vigorous at least once a week) and high/low TV (4hr/d cut off point). In high TV viewers, no differences were observed for any inflammatory markers between high or low physical activity groups. However, Active/high TV viewers demonstrated higher WBC compared to Active/ low TV viewers (adjusted B=0.23, 95% CI, 0.05, 0.40; $p=0.01$), presented in figure S1.

On average there was an increase of 1.9 hr/wk (SD, 31.5) TV viewing over 2 years. Approximately half of the sample (53.2%) remained relatively stable (less than 1 hr/d change), whilst 20.6% reported a substantial reduction of more than 1 hr/d and 26.2% an increase of more than 1 hr/d in TV viewing. An increase in TV viewing was associated with higher concentrations of CRP ($p=0.015$) and WBC ($p=0.05$) at follow up after adjustment for covariates and baseline TV viewing (Table 3).

Discussion

The aim of this study was to examine longitudinal associations between TV viewing (a major component of sedentary behaviour) and a range of inflammatory markers. Inflammatory processes are putative mechanisms underlying the detrimental health effects of sedentary behaviour but little long-term prospective data are available. We demonstrated an association between TV viewing and increased WBC (acute phase reactant) and fibrinogen (coagulation product) concentration at follow-up after adjustment for a range of clinical and behavioural covariates. In addition, an increase in TV viewing was associated with elevated CRP and WBC at follow up.

Several studies have previously demonstrated cross-sectional associations between TV viewing and markers of low grade inflammation.^{19,20} For example, in over 3,000 Australian adults TV time was associated with CRP and fibrinogen although after adjustment for waist circumference these relationships were attenuated to the null.¹⁹ In a small study of 285 adults, changes in sedentary time over 6 months were associated with CRP in women, although no associations were observed with other biomarkers, including adiponectin, soluble intracellular adhesion molecule-1, and interleukin-6.³² In a longitudinal cohort study baseline TV viewing time was not significantly associated with change in metabolic biomarkers over 5 years follow-up although inflammatory markers were not examined.³³ Nevertheless, increases in TV time over 5 years were associated with increases in waist circumference, diastolic blood pressure, and a clustered cardiometabolic risk score. In a British birth cohort study we demonstrated an association of TV viewing (aged 23) with CRP and fibrinogen at follow-up (aged 44),²³ although the effect estimates were attenuated after adjustments for BMI, especially so in participants with a BMI ≥ 25 kg/m². This is the first study to show associations between changes in TV viewing and inflammatory markers. Unlike previous work, our results were not largely attenuated by the inclusion of BMI or waist in the models. This is possibly explained by the older age of the present sample ; particularly in an elderly sample with onset of sarcopenia measures such as BMI are less reflective of adiposity.

Sedentary behaviours such as TV viewing are characterised by a lack of muscle contraction. Thus, the association between TV viewing and low grade inflammation is plausible since muscle contraction has been shown to stimulate the expression of various transcriptional co-activators such as PGC1 α , which lead to a transient release of myokines that may

promote systemic anti-inflammatory effects.³⁴ Other research has demonstrated that prolonged sitting is associated with the expression of various genes linked to inflammatory responses.³⁵ Adipose tissue inflammation might also explain links between sedentary behaviour and inflammation.³⁶ Indeed, reductions in central adiposity largely explain the changes in CRP observed after aerobic exercise training interventions.^{37,38} Thus, excess TV viewing may simply be a marker of chronic energy imbalance indicated by excess adiposity. We have recently demonstrated associations between objectively assessed sitting time and visceral adiposity, including liver fat,³⁹ which is believed to be a particularly important site for the production of pro-inflammatory markers.⁴⁰

The association between TV viewing and fibrinogen may also have relevance in terms of elevated risk of vascular conditions, particularly venous thrombosis. Our results support findings from a recent experimental trial that demonstrated increases in plasma fibrinogen with prolonged uninterrupted sitting that was attenuated with active breaks.⁴¹ Some epidemiological studies have demonstrated that associations between sedentary behaviour and CVD are attenuated in physically active participants.^{14,15} However, in our analysis to explore the possible additive effects of physical activity and TV viewing on inflammatory markers we did not observe any significant effect modification of physical activity (see figure S1). Physical activity was, however, rather crudely defined (vigorous activity at least once a week), thus future experimental studies are needed to define the dose of activity required to attenuate deleterious effects of prolonged sitting.

Our study has some notable strengths, including the longitudinal design to examine changes in TV time and biomarkers across 4 years; the use of a large national sample of community-dwelling men and women; and the inclusion of a wide range of potentially important

factors, including behavioral, social and clinical covariates. There are also some limitations. Our measure of sedentary behavior was limited to TV viewing thus our results cannot be generalized to total sedentary time or sitting. The questionnaires used to assess TV viewing in ELSA have not been validated and may have introduced an element of recall bias. Nevertheless, the TV measure has demonstrated excellent convergent validity in grading a plethora of psychosocial, physical and biochemical risk factors in previous work.^{26,27} Also, sedentary time questions that focus on TV viewing have demonstrated the strongest reliability and validity among non-occupational sedentary behaviour questions.⁴² It was not possible to adjust models for diet, as such data were not collected. However, recent data has suggested that snack food and TV viewing were independently associated with metabolic syndrome.⁴³ The analytic sample differed in some characteristics compared with those excluded that reduces the representativeness of our data. In particular, those excluded were slightly older and watched higher amounts of TV, thus the results may reflect a conservative estimate of the true associations. The stability of markers such as CRP over long term follow-up has been found to be similar to that of blood pressure and serum cholesterol,⁴⁴ which makes it suitable for use in risk prediction studies. Surprisingly, we observed slight reductions in two inflammatory markers over the 4 year follow-up period, although this might be accounted for by use of medication since 20.3% of the sample reported taking statins at baseline known to have anti-inflammatory properties.⁴⁵

In summary, TV viewing was associated with increased WBC and fibrinogen, but not CRP, over 4 years follow-up. For the first time, we show associations between changes in TV viewing and elevated inflammatory markers. Sedentary behaviours may be important in influencing the pro-inflammatory state seen with ageing.

Author contributions

M Hamer had full access to the data, and takes responsibility for the integrity and accuracy of the results. M Hamer drafted the paper, performed analyses and designed the study.

L Smith contributed to the concept and design of the study and critical revision of the manuscript.

E Stamatakis contributed to the concept and design of the study and critical revision of the manuscript.

Funding sources

The data were made available through the UK Data Archive. The English Longitudinal Study of Ageing (ELSA) was developed by a team of researchers based at University College London, the Institute of Fiscal Studies and the National Centre for Social Research. The funding is provided by the National Institute on Aging in the United States (grants 2RO1AG7644-01A1 and 2RO1AG017644) and a consortium of UK government departments coordinated by the Office for National Statistics. MH is supported by the British Heart Foundation (RE/10/005/28296). LS is supported by the National Institute for Health Research's School for Public Health Research. The funders had no role in the study design; in the collection, analysis and interpretation of data; in writing of the report; or in the decision to submit the paper for publication. The developers and funders of ELSA and the Archive do not bear any responsibility for the analyses or interpretations presented here.

Conflict of interest

None of the authors have any competing interests to declare.

Data sharing statement

Full ELSA data are available at the UK data archive <http://www.data-archive.ac.uk/>.

References

1. Vasto S, Carruba G, Lio D, Colonna-Romano G, Di Bona D, Candore G, Caruso C. Inflammation, ageing and cancer. *Mech Ageing Dev.* 2009;130:40-5.
2. Emerging Risk Factors Collaboration, Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, Collins R, Danesh J. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet.* 2010;375:132-40.
3. Interleukin-6 Receptor Mendelian Randomisation Analysis (IL6R MR) Consortium, Hingorani AD, Casas JP. The interleukin-6 receptor as a target for prevention of coronary heart disease: a Mendelian randomisation analysis. *Lancet.* 2012;379:1214-24.
4. Kaptoge S, Di Angelantonio E, Pennells L, Wood AM, White IR, Gao P, Walker M, Thompson A, Sarwar N, Caslake M, Butterworth AS, Amouyel P, Assmann G, Bakker SJ, Barr EL, Barrett-Connor E, Benjamin EJ, Björkelund C, Brenner H, Brunner E, et al. C-reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med.* 2012;367:1310-20.
5. Gleeson M, Bishop NC, Stensel DJ, Lindley MR, Mastana SS, Nimmo MA. The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease. *Nat Rev Immunol.* 2011;11:607-15.
6. Hamer M. The relative influence of fitness and fatness on inflammatory factors. *Prev Med.* 2007;44:3-11.
7. Hamer M, Sabia S, Batty GD, Shipley MJ, Tabák AG, Singh-Manoux A, Kivimaki M. Physical activity and inflammatory markers over 10 years: follow-up in men and women from the Whitehall II cohort study. *Circulation.* 2012;126:928-33.
8. Hamer M, Stamatakis E. Physical activity and risk of cardiovascular disease events events: inflammatory and metabolic mechanisms. *Med Sci Sports Exerc.* 2009;41:1206-11.
9. Mora S, Cook N, Buring JE, Ridker PM, Lee IM. Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms. *Circulation.* 2007;116:2110-2118.

10. Rana JS, Arsenault BJ, Després JP, Côté M, Talmud PJ, Ninio E, Wouter Jukema J, Wareham NJ, Kastelein JJ, Khaw KT, Matthijs Boekholdt S. Inflammatory biomarkers, physical activity, waist circumference, and risk of future coronary heart disease in healthy men and women. *Eur Heart J*. 2011;32:336-44.
11. Dempsey PC, Owen N, Biddle SJ, Dunstan DW. Managing sedentary behavior to reduce the risk of diabetes and cardiovascular disease. *Curr Diab Rep*. 2014 Sep;14(9):522.
12. Stamatakis E, Hamer M, Dunstan DW. Screen-based entertainment time, all-cause mortality, and cardiovascular events: population-based study with ongoing mortality and hospital events follow-up. *J Am Coll Cardiol*. 2011;57:292-9.
13. Dunstan DW, Barr EL, Healy GN, Salmon J, Shaw JE, Balkau B, Magliano DJ, Cameron AJ, Zimmet PZ, Owen N. Television viewing time and mortality: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation*. 2010;121:384-91.
14. Chomistek AK, Manson JE, Stefanick ML, Lu B, Sands-Lincoln M, Going SB, Garcia L, Allison MA, Sims ST, LaMonte MJ, Johnson KC, Eaton CB. Relationship of sedentary behavior and physical activity to incident cardiovascular disease: results from the Women's Health Initiative. *J Am Coll Cardiol*. 2013;61:2346-54.
15. Matthews CE, George SM, Moore SC, Bowles HR, Blair A, Park Y, Troiano RP, Hollenbeck A, Schatzkin A. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr*. 2012;95:437-45.
16. Scholes S, Coombs N, Pedisic Z, Mindell JS, Bauman A, Rowlands AV and Stamatakis E. Age- and sex-specific criterion validity of the health survey for England physical activity and sedentary behavior assessment questionnaire as compared with accelerometry. *Am J Epidemiol*. 2014;179:1493-502.
17. Stamatakis E, Davis M, Stathi A and Hamer M. Associations between multiple indicators of objectively-measured and self-reported sedentary behaviour and cardiometabolic risk in older adults. *Prev Med* 2012;54:82-87.

18. Harris JL, Bargh JA, Brownell KD. Priming effects of television food advertising on eating behavior. *Health Psychol.* 2009;28:404-13.
19. Howard BJ, Balkau B, Thorp AA, Magliano DJ, Shaw JE, Owen N, Dunstan DW. Associations of overall sitting time and TV viewing time with fibrinogen and C reactive protein: the AusDiab study. *Br J Sports Med.* 2014 Feb 18. doi: 10.1136/bjsports-2013-093014. [Epub ahead of print].
20. Pinto Pereira SM, Ki M, Power C. Sedentary behaviour and biomarkers for cardiovascular disease and diabetes in mid-life: the role of television-viewing and sitting at work. *PLoS One.* 2012;7:e31132.
21. Healy GN, Matthews CE, Dunstan DW, Winkler EA, Owen N. Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-06. *Eur Heart J* 2011;32:590-597.
22. Henson J, Yates T, Edwardson CL, Khunti K, Talbot D, Gray LJ, Leigh TM, Carter P, Davies MJ. Sedentary time and markers of chronic low-grade inflammation in a high risk population. *PLoS One.* 2013;8:e78350.
23. Stamatakis E, Hamer M, Mishra GD. Early adulthood television viewing and cardiometabolic risk profiles in early middle age: results from a population, prospective cohort study. *Diabetologia.* 2012;55:311-20.
24. Steptoe A, Breeze E, Banks J, Nazroo J. Chort Profile: The English Longitudinal Study of Ageing. *Int J Epidemiol* 2013; 42:1640-48.
25. Hamer M, Lavoie KL, Bacon SL. Taking up physical activity in later life and healthy ageing: the English longitudinal study of ageing. *Br J Sports Med* 2014; 48:239-243.
26. Hamer M, Stamatakis E. Screen-based sedentary behavior, physical activity, and muscle strength in the English longitudinal study of ageing. *PloS One* 2013; 8:e66222.
27. Hamer M, Stamatakis E. Prospective study of sedentary behavior, risk of depression, and cognitive impairment. *Med Sci Sports Exerc* 2014;46:718-23

28. Hamer M, de Oliveira C, Demakakos P. Non-Exercise Physical Activity and Survival: English Longitudinal Study of Ageing. *Am J Prev Med.* 2014;47:452-60.
29. Graig R, Deverill C, Pickering K. Quality control of blood, saliva and urine analytes. In: K. Spronston and J. Mindell, Editors, *Health survey for England 2004, Methodology and documentation Vol. 2*, The Information Centre, London, 2006, pp. 34–41.
30. Irwin M, Artin KH, Oxman MN. Screening for depression in the older adult: criterion validity of the 10-item Center for Epidemiological Studies Depression Scale (CES-D). *Arch Intern Med.* 1999;159:1701-4.
31. Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *The Gerontologist* 1970; 10:20-30.
32. Falconer CL, Cooper AR, Walhin JP, Thompson D, Page AS, Peters TJ, Montgomery AA, Sharp DJ, Dayan CM, Andrews RC. Sedentary time and markers of inflammation in people with newly diagnosed type 2 diabetes. *Nutr Metab Cardiovasc Dis.* 2014 Apr 26. pii: S0939-4753(14)00129-X. doi: 10.1016/j.numecd.2014.03.009. [Epub ahead of print]
33. Wijndaele K, Healy GN, Dunstan DW, Barnett AG, Salmon J, Shaw JE, Zimmet PZ, Owen N. Increased cardiometabolic risk is associated with increased TV viewing time. *Med Sci Sports Exerc.* 2010;42:1511-8.
34. Pedersen BK. Exercise-induced myokines and their role in chronic diseases. *Brain Behav Immun.* 2011;25: 811-16.
35. Latouche C, Jowett JB, Carey AL, Bertovic DA, Owen N, Dunstan DW, Kingwell BA. Effects of breaking up prolonged sitting on skeletal muscle gene expression. *J Appl Physiol.* 2013;114: 453-60.
36. Vieira VJ, Valentine RJ, Wilund KR, Antao N, Baynard T, Woods JA. Effects of exercise and low-fat diet on adipose tissue inflammation and metabolic complications in obese mice. *Am J Physiol Endocrinol Metab.* 2009;296: E1164-71.

37. Vieira VJ, Hu L, Valentine RJ, McAuley E, Evans EM, Baynard T, Woods JA. Reduction in trunk fat predicts cardiovascular exercise training-related reductions in C-reactive protein. *Brain Behav Immun* 2009;23:485-491.
38. Campbell PT, Campbell KL, Wener MH, Wood BL, Potter JD, McTiernan A, Ulrich CM. A yearlong exercise intervention decreases CRP among obese postmenopausal women. *Med Sci Sports Exerc.* 2009;41:1533-1539.
39. Smith L, Thomas EL, Bell JD, Hamer M. The association between objectively measured sitting and standing with body composition: a pilot study using MRI. *BMJ Open.* 2014;4:e005476.
40. Pou KM, Massaro JM, Hoffmann U, Vasan RS, Maurovich-Horvat P, Larson MG, Keaney JF Jr, Meigs JB, Lipinska I, Kathiresan S, Murabito JM, O'Donnell CJ, Benjamin EJ, Fox CS. Visceral and subcutaneous adipose tissue volumes are cross-sectionally related to markers of inflammation and oxidative stress: the Framingham Heart Study. *Circulation* 2007;116:1234-41.
41. Howard BJ, Fraser SF, Sethi P, Cerin E, Hamilton MT, Owen N, Dunstan DW, Kingwell BA. Impact on hemostatic parameters of interrupting sitting with intermittent activity. *Med Sci Sports Exerc.* 2013;45:1285-91.
42. Clark BK, Sugiyama T, Healy GN, Salmon J, Dunstan DW, Owen N. Validity and reliability of measures of television viewing time and other non-occupational sedentary behaviour of adults: a review. *Obes Rev* 2009; 10: 7-16.
43. Thorp AA, McNaughton SA, Owen N, Dunstan DW. Independent and joint associations of TV viewing time and snack food consumption with the metabolic syndrome and its components; a cross-sectional study in Australian adults. *Int J Behav Nutr Phys Act.* 2013 Aug 9;10:96.
44. Danesh J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, Rumley A, Lowe GD, Pepys MB, Gudnason V. C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med.* 2004;350: 1387-97.

45. Ridker PM, Lüscher TF. Anti-inflammatory therapies for cardiovascular disease. *Eur Heart J*. 2014;35:1782-1791.

Table 1. Descriptive characteristics of the sample at baseline (n=3612)

Variable	N (%)
<i>Age category</i>	
50 – 59	1170 (32.4)
60 – 69	1495 (41.4)
70 – 79	784 (21.7)
≥80	163 (4.5)
<i>Sex</i>	
Men	1626 (45.0)
Women	1986 (55.0)
<i>TV viewing</i>	
< 2hr/d	415 (11.5)
2 < 4 hr/d	1286 (35.6)
4 < 6 hr/d	957 (26.5)
≥ 6 hr/d	954 (26.4)
<i>Physical activity</i>	
Inactive	541 (15.0)
Moderate at least 1/wk	1809 (50.1)
Vigorous at least 1/wk	1262 (34.9)
<i>Body mass index category</i>	
15 < 25 kg/m ²	1006 (27.9)
25 < 30 kg/m ²	1584 (43.9)
≥ 30 kg/m ²	1022 (28.3)
<i>Smoking</i>	
None	3182 (88.1)
Current	430 (11.9)
<i>Alcohol intake</i>	
At least 5 times a week	857 (23.7)
at least once a week	1512 (41.9)
monthly	663 (18.4)
Rarely/never	580 (16.1)
<i>Depressive symptoms</i>	
CES-D score 0 – 3	3230 (89.4)
CES-D score ≥ 4	382 (10.6)
<i>Disabilities</i>	
None	2930 (81.1)
At least one	682 (18.9)
<i>Longstanding illness</i>	
None	1805 (50.0)
Any	1807 (50.0)
<i>Medications[†]</i>	
No	2233 (61.8)
Yes	1379 (38.2)
<i>C-reactive protein category</i>	
< 1 mg/L	1037 (28.7)
1 < 3 mg/L	1447 (40.1)
≥ 3 mg/L	1128 (31.2)

[†] including medication for diabetes, high blood pressure, cholesterol, blood thinning

Table 2. Linear regression analyses of TV viewing and physical activity on inflammatory markers at 4 years follow-up.

Baseline independent variable	Model 1 B (95% CIs)	Model 2 B (95% CIs)	Model 3 B (95% CIs)
<i>Log_e C-reactive protein (N=3612)</i>			
TV viewing	0.007 (0.002, 0.012)*	0.004 (0.00, 0.009)*	0.004 (-0.001, 0.009)
Physical activity	-0.058 (-0.087, -0.029)*	-0.032 (-0.062, -0.002)*	-0.031 (-0.061, 0.000)*
<i>White blood cell count (N=3529)</i>			
TV viewing	0.027 (0.014, 0.040)*	0.019 (0.006, 0.032)*	0.018 (0.005, 0.031)*
Physical activity	-0.17 (-0.25, -0.092)*	-0.099 (-0.18, -0.018)*	-0.092 (-0.17, -0.011)*
<i>Fibrinogen (N=3454)</i>			
TV viewing	0.006 (0.003, 0.010)*	0.004 (0.00, 0.008)*	0.004 (0.00, 0.008)*
Physical activity	-0.036 (-0.058, -0.014)*	-0.016 (-0.039, 0.007)	-0.015 (-0.038, 0.009)

B reflects unstandardised coefficient for inflammatory marker. *p<0.05

Model 1: adjusted for age, sex, inflammatory marker at baseline (CRP, or WBC, or fibrinogen).

Model 2: adjusted for age, sex, inflammatory marker at baseline, smoking, alcohol, depressive symptoms, long standing illness, disability (ADLs/IADLs), medications, body mass index.

Model 3: adjusted for all covariates in model 2 + mutually for TV time or physical activity.

Table 3. Regression of change in TV viewing on inflammatory markers at follow-up.

	Reduced TV viewing	Stable	Increased TV viewing
<i>Log_e C-reactive protein</i>			
Model 1	0.06 (-0.03, 0.13)	Reference	0.10 (0.05, 0.16)*
Model 2	0.07 (-0.01, 0.14)	Reference	0.07 (0.01, 0.12)*
<i>White blood cell count</i>			
Model 1	-0.013 (-0.22, 0.19)	Reference	0.24 (0.09, 0.39)*
Model 2	0.06 (-0.14, 0.25)	Reference	0.15 (0.00, 0.30)*
<i>Fibrinogen</i>			
Model 1	0.03 (-0.02, 0.09)	Reference	0.05 (0.01, 0.10)*
Model 2	0.05 (-0.01, 0.10)	Reference	0.03 (-0.01, 0.07)

Data reflect unstandardised coefficient for inflammatory marker. *p<0.05

Model 1: adjusted for age, sex, baseline TV time.

Model 2: adjusted for age, sex, baseline TV time, smoking, alcohol, depressive symptoms, long standing illness, disability (ADLs/IADLs), medications, body mass index, physical activity.