Supporting Information for:**Pt-Ag Catalyst System for Hydroarylations with Unactivated Arenes and Olefins**Dmitry Karshedt,^{†,‡} Alexis T. Bell,^{*,§,‡} and T. Don Tilley^{*,†,‡}

Department of Chemistry[†] and Department of Chemical Engineering,[§] University of California, Berkeley, Berkeley, CA 94720, and Chemical Sciences Division,[‡] Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720

Synthetic Methods. All experiments were conducted under a nitrogen atmosphere using standard Schlenk techniques or in a Vacuum Atmospheres drybox. Non-deuterated solvents were distilled under N₂ from appropriate drying agents and stored in PTFE-valved flasks. Deuterated solvents (Cambridge Isotopes) were dried with appropriate drying agents.

All olefins were purchased from Aldrich. Norbornene was purified by sublimation. Cyclopentene and cyclohexene were dried over sodium metal and vacuum-transferred onto activated molecular sieves before use. Ethylene, propylene, and 2-butene were used as received. AgOTf was purchased from Strem Chemicals and stored in the absence of light. The ligand precursor 2-(2-pyridyl)indole,¹ [(C₂H₄)PtCl(μ-Cl)]₂ (Zeise's dimer),² and (COD)PtCl₂³ were prepared as reported in the literature.

Analytical Methods. ¹H (500.1 MHz) and ¹³C{¹H} (124.7 MHz) NMR spectra were recorded using a Bruker DRX-500 spectrometer equipped with a 5 mm BBI probe. Spectra were recorded at room temperature and referenced to the residual protonated solvent for ¹H and to the solvent peak for ¹³C. Multiplets that appear as virtual triplets are denoted as "vt." Elemental analyses were performed by the College of Chemistry Microanalytical Laboratory at the University of California, Berkeley. Identities of organic products were confirmed by ¹H NMR (for

exo-phenylnorbornane),⁴ by ^1H and ^{13}C NMR (for tolylcyclohexanes),⁵ and by GCMS, using an Agilent Technologies 6890N GC system with an HP-5MS column.

2-(2-pyridyl)indolyl potassium. To a solution of 2-(2-pyridyl)indole (2.00 g, 10.3 mmol) in 15 mL of tetrahydrofuran, a suspension of KH (0.410 g, 10.2 mmol) in 5 mL tetrahydrofuran was transferred dropwise at room temperature. Stirring continued for 5 h, and the solvent was then removed. The resulting residue was washed with toluene and dried in vacuo at 70 °C for 16 h, providing a powdery light-green solid (1.87 g, 78% yield). ^1H NMR (CD_3CN): δ 6.63 (m, 1H), 6.74 (m, 1H), 6.86 (s, 1H), 6.98 (m, 1H), 7.39 (m, 2H), 7.60 (m, 1H), 7.91 (m, 1H), 8.42 (m, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_3CN): δ 97.4, 115.0, 117.0, 117.3, 118.8, 119.0, 120.2, 132.0, 135.8, 147.8, 148.8, 148.9, 158.6. Anal. Calcd. for $\text{C}_{13}\text{H}_9\text{N}_2\text{K}$: C, 67.21; H, 3.90; N, 12.06. Found: C, 67.02; H, 3.75; N, 11.80.

(NN)PtCl(C_2H_4) (1).⁶ To a suspension of Zeise's dimer (0.200 g, 0.340 mmol, 0.5 equiv) in 2 mL of benzene, 2-(2-pyridyl)indolyl potassium (0.158 g, 0.680 mmol) was added as a solid. The reaction mixture was stirred for 24 h, and the solvent was then removed. The resulting residue was extracted with dichloromethane (10 mL) and filtered. The solvent volume was then reduced to ca. 3 mL, and the solution was treated with 10 mL of pentane, leading to the precipitation of a red-orange solid. The solvent was decanted and the resulting solid was dried in vacuo (0.248 g, 81% yield). ^1H NMR (CD_2Cl_2): δ 4.20 (d with Pt satellites, $^2J = 15.0$ Hz, $J_{\text{Pt-H}} = 55.2$ Hz, 2H, vinylic), 4.51 (d with Pt satellites, $^2J = 15.0$ Hz, $J_{\text{Pt-H}} = 54.0$ Hz, 2H, vinylic), 6.90 (s, 1H), 7.00 (vt, $^2J = 6.0$ Hz, 1H), 7.11 (m, 2H), 7.39 (d, $^2J = 4.5$ Hz, 1H), 7.53 (d, $^2J = 6.0$ Hz, 1H), 7.73 (d, $^2J = 5.7$ Hz, 1H), 7.83 (vt, $^2J = 6.3$ Hz, 1H), 8.46 (d, $^2J = 6.6$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 104.0, 112.2, 116.2, 116.5, 119.7, 120.3, 120.4, 120.6, 120.7, 121.8, 124.0, 124.1, 139.0, 141.8, 142.7. Anal. Calcd. for $\text{C}_{15}\text{H}_{13}\text{ClN}_2\text{Pt}$: C, 39.87; H, 2.90; N, 6.20. Found: C, 39.61; H, 2.99; N, 5.86.

(COD)Pt(OTf)₂. To a solution of (COD)PtCl₂ (0.101 g, 0.270 mmol) in 2 mL of dichloromethane, AgOTf (0.139 g, 0.541 mmol, 2 equiv) was added as a solid. The reaction

mixture was stirred for 1 h in the absence of light and filtered, and the solvent was removed in vacuo to give a flaky colorless solid (0.153 g, 94% yield). ^1H NMR (CD_2Cl_2): δ 2.26 (d, $^2J = 9.0$ Hz, 4H, aliphatic), 2.85 (d, $^2J = 9.0$ Hz, 4H, aliphatic), 6.26 (s with Pt satellites, $J_{\text{Pt-H}} = 74.5$ Hz, 4H, vinylic). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 30.6 (s, aliphatic), 101.5 (s with Pt satellites, $J_{\text{Pt-C}} = 224.5$ Hz, vinylic), 119.2 (q, $J_{\text{F-C}} = 311.8$ Hz, triflate). Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{F}_6\text{O}_6\text{S}_2\text{Pt}$: C, 19.97; H, 2.01. Found: C, 20.26; H, 1.93.

General Procedure for Catalytic Runs. Reactions were conducted in 5 mm Wilmad NMR tubes equipped with J. Young teflon screw caps, which were heated in temperature-controlled oil baths. Samples were prepared by dissolving catalysts and solid or liquid olefins in the specified mixed solvent. Gases were condensed on a Schlenk line into J. Young NMR tubes containing catalysts dissolved in the specified mixed solvent. Reaction progress was monitored by ^1H NMR spectroscopy (a small amount of cyclohexane- d_{12} was added to all samples in order to obtain a lock signal). Product yields were quantified by GC, using a calculated response factor to account for the difference in ionization between the integration standard (naphthalene) and *exo*-phenylnorbornane.

References

- (1) Thummel, R. P.; Hedge, V. *J. Org. Chem.* **1989**, *54*, 1720.
- (2) Chatt, J.; Searle, M. L. *Inorg. Synth.* **1957**, *5*, 210.
- (3) Clark, H. C.; Manzer, L. E. *J. Organomet. Chem.* **1973**, *59*, 411.
- (4) Olah, G. A.; Lee, C. S.; Surya Prakash, G. K. *J. Org. Chem.* **1994**, *59*, 2590.
- (5) Mahindaratne, M. P. D.; Wimalasena, K. *J. Org. Chem.* **1998**, *63*, 2858.
- (6) Orientation of ligands in complex **1** was determined by a ^1H gradient NOESY experiment and by X-ray crystallography. These data will be published elsewhere.