Supplemental material for:

Pt-Catalyzed Pent-annulations from In Situ Generated Metallo-Carbenoids Utilizing Propargylic Esters

B. A. Bhanu Prasad, Francis K. Yoshimoto, and Richmond Sarpong*

Center for New Directions in Organic Synthesis, Department of Chemistry, University of California, Berkeley, California 94720

Email: rsarpong@berkeley.edu

Table of Contents:

Materials and Methods	S2
General Procedure for Synthesis of Propargylic Esters	S4
General Procedure for Pt-catalyzed Pent-annulations of Propargylic Esters	S10
Saponification Procedure	S17
Allylation Procedure	S17
³¹P NMR Experiments	S18
References	S19
Selected Spectral Data	S20-S38

Materials and Methods. All air or moisture sensitive reactions were conducted in flame-dried glassware under an atmosphere of nitrogen using dry, deoxygenated solvents. Toluene, methylene chloride, and triethylamine were distilled under nitrogen from calcium hydride immediately prior to use and tetrahydrofuran (THF) was distilled under nitrogen from sodium benzophenone ketyl. All reagents were purchased from Aldrich and Lancaster and used without further purification. Melting points were measured on a Büchi® melting point apparatus and are corrected using vanillin (mp 81-83 °C) as a standard. Reaction temperatures were controlled by an IKAmag temperature modulator. Thin-layer chromatography (TLC) was performed using E. Merck silica gel 60 F254 precoated plates (0.25 mm) and visualized by UV and anisaldehyde stain. Fisher silica gel 240-400 mesh (particle size 0.032-0.063) was used for flash chromatography. Pt-catalyzed reactions were performed in Schlenk flasks. ¹H and ¹³C NMR spectra were recorded on a Bruker AVB-400 (at 400 MHz and 100 MHz respectively) and on a Bruker AV-300 (at 300 MHz and 75 MHz respectively), in chloroform-*d* at 23 °C, unless otherwise stated. Chemical shifts were referenced to the residual chloroform-H peak, which was set at 7.26 ppm for ¹H and 77.20 ppm (center peak) for ¹³C spectra. ³¹P NMR spectra were recorded on a Bruker AVB-400 at 162 MHz in 1,2-dichloroethane-*d*₄ at 23 °C as well as at 100 °C by using phosphoric acid in chloroform-*d* as an external standard (0.00 ppm). Data for ¹H NMR are reported as follows: chemical shifts (δ ppm). Multiplicity, (s = singlet, d = doublet, t = triplet, q = quartet, dd = doublet of doublet, qd = quartet of doublet, m = multiplet, br = broad resonance), coupling constants (Hz) and integration. Data for ¹³C NMR are reported in terms of chemical shift. IR spectra were recorded on a Nicolet MAGNA-IR 850 spectrometer and are reported in frequency of absorption (cm⁻¹). High resolution mass spectral data were obtained from the University of California, Berkeley Mass Spectral Facility.

SM Table 1. Preparation of Propargylic Esters Starting from Ketones

$ \begin{array}{c} \text{R}^1-\text{C}(=\text{O})-\text{R}^2 + \text{H}-\text{C}\equiv\text{C}-\text{C}(=\text{O})-\text{OR}^3 \\ \xrightarrow[2. \text{Ac}_2\text{O or BzCl, Et}_3\text{N, DMAP, CH}_2\text{Cl}_2, 0^\circ\text{C}]{1. n\text{-BuLi, THF, } -78^\circ\text{C, 2 h}} \\ \text{R}^1-\text{C}(\text{OR}^2)(\text{R}^2)-\text{C}\equiv\text{C}-\text{CO}_2\text{R}^3 \end{array} $				
entry	ketone	product		yield (%) ^a
1			1a	71
2			1b	70
3			1c	68
4			1d	72
5			1e	70
6			1f	64
7			1g	68
8			1h	69
9			1i	62

^a Overall yield (over 2 steps)

SM Table 1 (continued). Preparation of Propargylic Esters Starting from Ketones

$$R^1-C(=O)-R^2 + H-C\equiv C-C(=O)OR^3 \xrightarrow[2. \text{Ac}_2\text{O, Et}_3\text{N, DMAP, CH}_2\text{Cl}_2, \text{rt}]{1. n\text{-BuLi, THF, -78 }^\circ\text{C, 2 h}} R^1-C(OAc)(R^2)-C\equiv C-OR^3$$

entry	ketone	product	yield (%) ^a
10			73
11			74
12			76

^a Overall yield (over 2 steps)**Representative Procedure for Synthesis of Propargylic Esters (SM Table 1, entry 2):**

The following procedure, which is a modification of a literature preparation, was used:¹

A flame-dried, round-bottom flask was charged with anhydrous THF (40 mL) and ethyl propiolate (1.22 g, 12.5 mmol). The solution was cooled to -78 °C and *n*-BuLi (4.8 mL, 12.0 mmol, 2.5 M in hexanes) was added slowly over 20 min. The solution was allowed to stir for 45 min at -78 °C, and then acetophenone (1.16 mL, 10.0 mmol) was added slowly over 15 min. The mixture was stirred for an additional 45 min at -78 °C, then the acetone bath was removed and the mixture was allowed to warm to 23 °C. After 30 min, sat. aqueous NH₄Cl solution (10 mL) was added slowly and stirring continued for 15 min. The mixture was diluted with EtOAc (40 mL) and washed with sat. aqueous NH₄Cl solution (30 mL). The aqueous layer was extracted with EtOAc (2 X 15 mL) and the combined organic layer was washed with brine and dried over MgSO₄. The crude propargyl alcohol was directly acetylated after evaporation of the solvent under reduced pressure.

To a solution of the crude propargyl alcohol in anhydrous methylene chloride (40 mL) was added triethylamine (4.17 mL, 30.0 mmol), acetic anhydride (1.18 mL, 12.5

mmol), and DMAP (0.061 g, 0.5 mmol) under a nitrogen atmosphere at 23 °C. The resultant mixture was stirred for 45 min then washed with aqueous 2 N HCl solution (2 X 15 mL), brine, dried over MgSO₄ and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (7:1 hexanes:EtOAc) to obtain the pure propargylic ester (1.84 g, 71% yield).

SM Table 1, 1a: Pale yellow oil (71% yield). *R*_F 0.53 (3:1 hexanes:EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 7.55-7.52 (m, 2H), 7.40-7.28 (m, 3H), 4.26 (q, *J* = 7.2 Hz, 2H), 2.09 (s, 3H), 1.92 (s, 3H), 1.33 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 168.4, 153.3, 141.0, 128.5, 128.3, 124.7, 85.5, 79.1, 74.7, 62.2, 31.1, 21.4, 14.0; IR (film) 2239, 1752, 1713, 1265, 1221 cm⁻¹; HRMS (EI⁺) calc'd for [C₁₅H₁₆O₄]⁺: *m/z*, 260.1048 found 260.1045.

SM Table 1, 1b: Faint yellow solid (70% yield). mp 61-63 °C; *R*_F 0.56 (3:1 hexanes:EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 7.54-7.52 (m, 2H), 7.39-7.29 (m, 3H), 3.80 (s, 3H), 2.09 (s, 3H), 1.92 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 168.5, 153.8, 141.0, 128.7, 128.4, 124.8, 86.2, 78.9, 74.8, 52.9, 31.3, 21.6; IR (film) 2239, 1751, 1717, 1266, 1222 cm⁻¹; HRMS (EI⁺) calc'd for [C₁₄H₁₄O₄]⁺: *m/z*, 246.0892 found 246.0895.

SM Table 1, 1c: Colorless dense oil (68% yield). *R*_F 0.63 (3:1 hexanes:EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 7.68-7.32 (m, 10H), 4.26 (q, *J* = 7.2 Hz, 2H), 2.08 (s, 3H), 1.32 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 164.2, 162.5, 153.5, 133.4, 130.7, 130.1, 130.0, 129.0, 128.8, 128.6, 128.5, 85.7, 79.5, 75.4, 62.4, 31.5, 14.3; IR (film) 2238, 1789, 1730, 1274, 1229 cm⁻¹; HRMS (EI⁺) calc'd for [C₂₀H₁₈O₄]⁺: *m/z*, 322.1205 found 322.1202.

SM Table 1, 1d: White solid (72% yield). mp 87-89 °C; R_F 0.50 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.50-7.47 (m, 4H), 7.37-7.30 (m, 6H), 4.24 (q, $J = 7.2$ Hz, 2H), 2.18 (s, 3H), 1.31 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.2, 153.4, 140.9, 128.6, 128.5, 126.4, 81.8, 81.6, 76.9, 62.3, 21.7, 14.2; IR (film) 2366, 1757, 1706, 1254, 1224 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{20}\text{H}_{18}\text{O}_4]^+$: m/z , 322.1205 found 322.1203.

SM Table 1, 1e: Colorless oil (70% yield). R_F 0.47 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.63-7.56 (m, 6H), 7.46-7.42 (m, 2H), 7.37-7.28 (m, 1H), 4.28 (q, $J = 7.2$ Hz, 2H), 2.12 (s, 3H), 1.97 (s, 3H), 1.34 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.7, 153.5, 141.4, 140.6, 128.9, 127.7, 127.5, 127.3, 125.4, 85.6, 79.3, 74.7, 62.4, 31.3, 21.7, 14.2; IR (film) 2232, 1752, 1712, 1260 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{21}\text{H}_{20}\text{O}_4]^+$: m/z , 336.1361 found 336.1357.

SM Table 1, 1f: Dense liquid (64% yield). R_F 0.46 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.46 (dd, $J = 6.8, 2.4$ Hz, 2H), 6.88 (dd, $J = 6.4, 2.0$ Hz, 2H), 4.24 (q, $J = 7.2$ Hz, 2H), 3.78 (s, 3H), 2.04 (s, 3H), 1.91 (s, 3H), 1.30 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.4, 153.3, 141.0, 128.5, 128.3, 124.7, 85.5, 79.1, 74.7, 62.2, 31.1, 21.4, 14.0; IR (film) 2230, 1750, 1713, 1252, 1220 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{16}\text{H}_{18}\text{O}_5]^+$: m/z , 290.1187 found 290.1189.

SM Table 1, 1g: Pale yellow oil (68% yield). R_F 0.55 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.45-7.27 (m, 6H), 7.15-7.13 (m, 2H), 6.92-6.90 (m, 1H), 5.07 (d, $J = 2$ Hz, 2H), 4.26 (q, $J = 7.2$ Hz, 2H), 2.07 (s, 3H), 1.90 (s, 3H), 1.32 (t, $J = 7.2$ Hz, 3H);

^{13}C NMR (100 MHz, CDCl_3) δ 168.6, 160.1, 153.5, 142.8, 136.9, 129.9, 128.8, 128.2, 127.8, 117.4, 115.2, 112.0, 85.6, 79.2, 74.7, 70.3, 62.3, 31.4, 21.6, 14.2; IR (film) 2235, 1748, 1706, 1258, 1220 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{22}\text{H}_{22}\text{O}_5]^+$: m/z , 366.1467 found 366.1466.

SM Table 1, 1h: Yellow dense oil (67% yield). mp 91-93 °C; R_F 0.52 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.93 (d, J = 8.4 Hz, 1H), 7.78 (d, J = 8.4 Hz, 2H), 7.74 (s, 1H), 7.69 (d, J = 7.6 Hz, 1H) 7.31 (m, 1H), 7.26-7.22 (m, 3H), 4.25 (q, J = 7.2 Hz, 2H), 2.33 (s, 3H), 2.09 (s, 3H), 2.04 (s, 3H), 1.32 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ ; 168.5, 153.3, 145.4, 135.7, 130.2, 129.2, 128.4, 127.1, 125.1, 124.2, 123.6, 122.2, 120.8, 114.0, 84.8, 78.1, 70.7, 62.5, 28.3, 21.7, 21.5, 14.2; IR (film) 2197, 1710, 1246 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{24}\text{H}_{23}\text{NO}_6\text{S}]^+$: m/z , 453.1246 found 453.1251.

SM Table 1, 1i: Yellow solid (62% yield). mp 118-120 °C; R_F 0.40 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 8.37-8.35 (m, 2H), 8.04-8.02 (m, 2H), 7.31 (t, J = 2.4 Hz, 1H), 7.10 (dd, J = 3.2, 2.4 Hz, 1H), 6.42 (dd, J = 3.2, 1.6 Hz, 1H), 4.24 (q, J = 7.2 Hz, 2H), 2.01 (s, 3H), 1.89 (s, 3H), 1.31 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.6, 153.3, 151.0, 144.2, 130.9, 128.4, 125.0, 121.7, 118.8, 112.9, 84.9, 77.8, 69.7, 62.5, 29.1, 21.7, 14.2; IR (film) 2196, 1708, 1533, 1350, 1183 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{19}\text{H}_{18}\text{N}_2\text{O}_8\text{S}]^+$: m/z , 434.0784 found 434.0785.

SM Table 1, 1j: Dense oil (69% yield). R_F 0.58 (hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.45-7.27 (m, 5H), 6.88 (d, J = 17.6 Hz, 1H), 6.30 (d, J = 16 Hz, 1H), 4.24 (q, J = 7.2 Hz, 2H), 2.07 (s, 3H), 1.85 (s, 3H), 1.32 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.9, 153.5, 135.7, 132.2, 128.8, 128.5, 128.3, 127.1, 85.2, 78.6, 73.3, 62.4,

28.0, 21.8, 14.2; IR (film) 2237, 1748, 1713, 1256, 1228 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{17}\text{H}_{18}\text{O}_4]^+$: m/z , 286.1205 found 286.1210.

SM Table 1, 1k: Colorless oil (68% yield). R_F 0.60 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 5.89 (m, 1H), 4.22 (q, $J = 7.2$ Hz, 2H), 2.56-2.36 (m, 4H), 2.04 (s, 3H), 1.96-1.88 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.7, 153.5, 142.5, 128.6, 85.9, 77.4, 72.6, 62.1, 32.5, 31.2, 26.7, 23.4, 21.5, 14.1; IR (film) 2235, 1750, 1714, 1248, 1225 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{14}\text{H}_{18}\text{O}_4]^+$: m/z , 250.1205 found 250.1209.

SM Table 1, 1l: Colorless oil (62% yield). R_F 0.67 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 6.05 (m, 1H), 4.23 (q, $J = 7.2$ Hz, 2H), 2.14-2.05 (m, 2H), 2.04 (s, 3H), 1.95-1.86 (m, 2H), 1.72 (s, 3H), 1.57-1.55 (m, 4H), 1.29 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.8, 153.6, 142.5, 128.7, 86.0, 72.7, 62.2, 34.1, 32.6, 31.2, 26.8, 23.4, 23.0, 21.6, 14.2; IR (film) 2243, 1748, 1708, 1258, 1221 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{20}\text{O}_4]^+$: m/z , 264.1361 found 264.1363.

SM Table 2: Pt-Catalyzed Pent-annulations of Propargylic Esters

entry	substrate	product ^a	time (h)	yield (%)
1	1a	6	8	62
2	1b	9	8	62
3	1c	10	6	72
4	1d	11	6	84
5	1e	12	8	71
6	1f	13	8	77
7	1g	14	6	61
8	1h	15	4	64
9	1i	16	3	77

^a Obtained as a mixture of regioisomers in some cases (see characterization data). Major shown here.

General Procedure for Pt-catalyzed Pent-annulations of Propargylic Esters:

A flame-dried 25 mL Schlenk flask equipped with a Teflon screw-top cap was charged with $\text{PtCl}_2(\text{PPh}_3)_2$ (0.1 mmol), propargylic ester (1.0 mmol), and PhIO (0.2 mmol) in anhydrous toluene (0.2 M). The tube was flushed well with nitrogen and capped tightly. The flask was placed in an oil bath which was heated to 100 °C (ca. 20 min) and held at this temperature for 3-8 h. At the completion of the reaction (determined by TLC analysis), the resultant mixture was filtered through a fritted funnel and washed thoroughly with acetone (10 mL). The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography to obtain the pure pent-annulated products.

SM Table 2, 6: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a colorless oil in 62% yield as a 17:3 inseparable mixture of olefin regioisomers. R_F 0.56 (3:1 hexanes:EtOAc); ^1H NMR (300 MHz, CDCl_3) δ 7.91 (m, 1H minor), 7.48 (d, J = 9.6 Hz, 1H), 7.35-7.29 (m, 2H, 1H from minor isomer), 7.226-7.15 (m, 4H, 2H from minor isomer), 4.84 (m, 1H), 4.34 (q, J = 7.2 Hz, 2H minor), 4.25-4.06 (m, 2H), 3.73-3.65 (q, J = 7.5 Hz, 1H minor), 2.34 (s, 3H minor), 2.27 (s, 3H), 2.02 (d, J = 2.1 Hz, 3H), 1.39 (t, J = 7.2 Hz, 3H minor), 1.36 (d, J = 7.5 Hz, 3H minor) 1.27 (t, J = 7.2 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 169.2, 168.5, 167.8, 167.0, 147.5, 143.1, 142.5, 136.3, 127.9, 127.3, 125.8, 125.6, 123.5, 122.7, 122.6, 119.3, 118.8, 61.5, 60.6, 53.3, 44.1, 21.1, 14.4, 14.3, 8.9, IR (film) 1763, 1745, 1210 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{16}\text{O}_4]^+$: m/z , 260.0747 found 260.0769.

SM Table 2, 9: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a dense oil in 62% yield as a 23:2 inseparable mixture of olefin regioisomers. R_F 0.50 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 7.91 (d, J = 7.2 Hz, 1H minor), 7.48-7.46 (m, 1H), 7.45-7.19 (m, 6H, 3H from minor isomer), 4.84 (m, 1H), 3.88 (s, 3H minor), 3.72 (s, 3H), 2.36 (s, 3H minor), 2.28 (s, 3H), 2.02 (d, J = 1.6 Hz, 3H), 1.36 (d, J = 7.6 Hz, 3H minor); ^{13}C NMR (100 MHz, $CDCl_3$) δ 169.8, 168.6, 147.4, 143.1, 136.2, 128.0, 125.8, 125.7, 123.6, 119.4, 53.1, 52.7, 21.0, 8.9; IR (film) 1761, 1740, 1206 cm^{-1} ; HRMS (EI^+) calc'd for $[C_{14}H_{14}O_4]^+$: m/z , 246.0892 found 246.0891.

SM Table 2, 10: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a clear oil in 72% yield as a 17:3 inseparable mixture of olefin regioisomers. R_F 0.66 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 8.24-8.19 (m, 2H, 1H from minor isomer), 7.99 (m, 1H minor), 7.66-7.62 (m, 2H, 1H from minor isomer), 7.55-7.49 (m, 4H, 2H from minor isomer), 7.44-7.34 (m, 4H, 2H from minor isomer), 7.30-7.26 (m, 4H, 2H from minor isomer), 5.03 (bm, 1H), 4.25-4.04 (m, 2H), 3.95 (q, J = 7.6 Hz, 1H minor), 2.12 (d, J = 2 Hz, 3H), 1.43 (d, J = 7.6 Hz, 3H minor), 1.18 (t, J = 7.2 Hz, 3H), 1.09 (t, J = 7.2 Hz, 3H minor); ^{13}C NMR (100 MHz, $CDCl_3$) δ 169.2, 164.1, 147.7, 143.1, 136.4, 133.8, 130.3, 129.3, 128.9, 128.7, 127.9, 125.9, 125.6, 123.6, 61.5, 53.4, 14.2, 9.0; IR (film) 1787, 1735, 1451, 1263 cm^{-1} ; HRMS (EI^+) calc'd for $[C_{20}H_{18}O_4]^+$: m/z , 322.1205 found 322.1199.

SM Table 2, 11: The reaction was heated at 100 °C for 6 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a clear oil in 84% yield as a 9:1 inseparable mixture of olefin regioisomers. R_F 0.46 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 7.99 (d, J = 6 Hz, 1H minor), 7.60 (d, J = 6.0 Hz, 1H), 7.56 (d, J = 6 Hz, 1H minor), 7.48 (t, J = 5.6 Hz, 2H), 7.41-7.26 (m, 8H, 4H from minor isomer), 7.21 (t, J = 6.0 Hz, 1H minor), 7.16 (m, 1H minor), 7.11 (m, 2H minor), 5.12 (s, 1H), 4.91 (s, 1H minor), 4.38 (q, J = 5.6 Hz, 2H minor), 4.31-4.18 (m, 2H minor), 2.22 (s, 3H), 2.18 (s, 3H minor), 1.42 (t, J = 6.0 Hz, 3H minor), 1.31 (t, J = 5.6 Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 168.8, 168.7, 153.4, 147.8, 136.4, 131.7, 128.9, 128.7, 128.2, 127.9, 126.5, 125.9, 124.1, 120.7, 61.6, 53.5, 21.1, 14.3; IR (film) 1763, 1736, 1367, 1206 cm^{-1} ; HRMS (EI^+) calc'd for $[C_{20}H_{18}O_4]^+$: m/z , 322.1205 found 322.1203.

SM Table 2, 12: The reaction was heated at 100 °C for 6 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a dense liquid in 77% yield as a 17:3 inseparable mixture of olefin regioisomers. R_F 0.54 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 8.23 (d, J = 1.6 Hz, 1H minor), 7.77 (s, 1H), 7.67-7.57 (m, 5H, 2H from minor isomer), 7.51-7.44 (m, 5H, 3H from minor isomer), 7.42-7.32 (m, 4H, 2H from minor isomer), 4.96 (br, 1H), 4.40 (q, J = 7.2 Hz, 2H minor), 4.32-4.16 (m, 2H), 3.78 (q, J = 7.6 Hz, 1H minor), 2.38 (s, 3H minor), 2.32 (s, 3H), 2.08 (d, J = 2.0 Hz, 3H), 1.44-1.41 (m, 6H minor), 1.31 (d, J

= 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.1, 168.4, 147.7, 142.3, 141.4, 138.8, 136.9, 128.9, 127.4, 127.2, 126.9, 125.3, 122.4, 119.6, 61.5, 53.3, 21.0, 14.3, 8.9; IR (film) 1763, 1735, 1202 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{21}\text{H}_{20}\text{O}_4]^+$: m/z , 336.1361 found 336.1361.

SM Table 2, 13: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a dense liquid in 71% yield as a 17:3 inseparable mixture of olefin regioisomers. R_F 0.45 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.53 (d, J = 2.4 Hz, 1H minor), 7.20 (d, J = 8.0 Hz, 1H minor), 7.12 (d, J = 8.4 Hz, 1H), 7.10 (m, 1H), 6.84 (dd, J = 8.0, 2.4 Hz, 1H), 6.79 (dd, J = 8.4, 2.4 Hz, 1H minor), 4.79 (br, 1H), 4.33 (q, J = 7.2 Hz, 2H minor), 4.27-4.11 (m, 2H), 3.85 (s, 3H minor), 3.80 (s, 3H minor), 3.63 (q, J = 7.6 Hz, 1H minor), 2.54 (s, 3H minor), 2.26 (s, 3H), 1.98 (d, J = 2.0 Hz, 3H), 1.38 (t, J = 7.2 Hz, 3H minor), 1.33 (d, J = 7.6 Hz, 3H minor) 1.27 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.1, 168.6, 159.2, 145.5, 137.8, 135.8, 119.7, 113.7, 112.8, 110.6, 61.4, 55.6, 53.1, 20.9, 14.2, 8.9; IR (film) 1761, 1735, 1239, 1206 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{16}\text{H}_{18}\text{O}_5]^+$: m/z , 290.1154 found 290.1156.

SM Table 2, 14: The reaction was heated at 100 °C for 6 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a clear oil in 61% yield as a 17:3 inseparable mixture of olefin regioisomers. R_F 0.51 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.82 (d, J =

7.2 Hz, 1H minor), 7.45-7.31 (m, 11H, 5H from minor isomer) 7.01-6.93 (m, 2H minor), 6.89 (d, $J = 2.0$ Hz, 1H), 6.81 (dd, $J = 8.4, 2.4$ Hz, 1H), 5.14 (s, 2H minor), 5.09 (s, 2H), 4.79 (bm, 1H), 4.33 (q, $J = 7.2$ Hz, 2H minor), 4.24-4.13 (m, 2H), 3.67 (q, $J = 6$ Hz, 1H minor), 2.34 (s, 3H minor), 2.28 (s, 3H minor), 1.99 (d, $J = 2.0$ Hz, 3H), 1.39 (t, $J = 7.2$ Hz, 3H minor), 1.35 (d, $J = 6$ Hz, 3H minor) 1.28 (t, $J = 5.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.3, 168.4, 159.2, 148.4, 144.5, 137.1, 128.7, 128.6, 128.1, 127.6, 125.4, 124.2, 111.4, 107.1, 70.4, 61.4, 52.6, 21.0, 14.3, 8.9; IR (film) 1763, 1734, 1197 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{22}\text{H}_{22}\text{O}_5]^+$: m/z , 366.1467 found 366.1457.

SM Table 2, 15: The reaction was heated at 100°C for 4 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (5:1 hexanes:EtOAc) to provide the desired indene as a dense oil in 64% yield as a 4:1 inseparable mixture of olefin regioisomers. R_F 0.41 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.95-7.93 (m, 2H, 1H from minor isomer), 7.87-7.85 (m, 4H, 2H from minor isomer), 7.62-7.60 (m, 2H, 1H from minor isomer), 7.34-7.23 (m, 8H, 4H from minor isomer), 4.95 (m, $J = 2$ Hz, 1H), 4.80 (d, $J = 8$ Hz, 1H minor), 4.36-4.23 (m, 2H), 4.17 (q, $J = 7.2$ Hz, 2H minor), 2.37 (s, 3H minor), 2.34 (s, 6H), 2.20 (d, $J = 2$ Hz, 3H), 2.09 (s, 3H minor), 1.33-1.28 (m, 9H, 6H from minor); ^{13}C NMR (100 MHz, CDCl_3) δ 169.5, 167.8, 145.3, 142.9, 138.4, 135.6, 135.3, 130.2, 130.0, 127.3, 127.3, 125.3, 124.3, 123.7, 119.4, 114.5, 62.0, 52.0, 21.7, 20.5, 14.4, 10.6; IR (film) 1746, 1652, 1370, 1225, 1174 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{24}\text{H}_{23}\text{NO}_6\text{S}]^+$: m/z , 453.1246 found 453.1248.

SM Table 2, 16: The reaction was heated at 100°C for 3 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (6:1 hexanes:EtOAc) to provide the

desired product as a dense liquid in 77% yield as a single olefin isomer. R_F 0.43 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 8.35-8.31 (m, 2H), 8.11-8.07 (m, 2H), 7.04 (d, J = 3.2 Hz, 1H), 6.32 (d, J = 3.2 Hz, 1H), 4.63 (m, 1H), 4.22-4.15 (m, 2H), 2.42 (s, 3H), 1.89 (d, J = 2.0 Hz, 3H), 1.26 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 169.4, 167.8, 150.8, 144.2, 143.3, 135.7, 130.0, 128.7, 124.7, 123.5, 123.4, 108.1, 62.0, 50.6, 20.5, 14.3, 10.1; IR (film) 1737, 1722, 1535, 1185 cm^{-1} ; HRMS (EI $^+$) calc'd for $[C_{19}H_{18}N_2O_8S]^+$: m/z , 434.0784 found 434.0786.

SM Table 3: Pt-Catalyzed Pent-annulations of Propargylic Esters

entry	substrate	product	time (h)	yield (%)
1 ^a	17	18	8	64
2 ^a	19	21b	8	73
3 ^b	20	22a 22b	8	78

^aProduct obtained as a single regioisomer. ^bProduct obtained as a 2:1 mixture of olefin regioisomers.

SM Table 3, 18: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 ml). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired cyclopentadiene as a pale yellow oil in 64% yield as a single isomer. R_F 0.5 (3:1 hexanes:EtOAc); 1H NMR (400 MHz, $CDCl_3$) δ 7.40-7.27 (m, 5H), 4.15 (q, J = 7.2 Hz,

2H), 3.34 (s, 2H), 2.27 (s, 3H), 1.92 (s, 3H), 1.14 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.4, 163.9, 149.8, 144.0, 135.9, 128.3, 128.1, 128.0, 125.9, 125.47, 60.1, 44.9, 20.8, 14.0, 11.52; IR (film) 1760, 1713, 1224 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{17}\text{H}_{18}\text{O}_4]^+$: m/z , 286.1205 found 286.1203.

SM Table 3, 21b: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired indene as a colorless oil in 73% yield as a single isomer. R_F 0.50 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 6.39 (m, 1H), 4.23-4.06 (m, 2H), 2.88-2.79 (m, 1H), 2.61-2.52 (m, 1H), 2.42-2.06 (m, 3H), 2.14 (s, 3H), 1.79 (m, 3H), 1.24 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 170.3, 169.6, 163.2, 147.8, 137.0, 122.6, 83.1, 59.6, 28.9, 28.1, 24.0, 21.1, 14.5, 12.3; IR (film) 1741, 1701, 1232 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{13}\text{H}_{18}\text{O}_4]^+$: m/z , 251.1238 found 251.1236.

SM Table 3, 22: The reaction was heated at 100 °C for 8 h, then filtered through a fritted funnel and washed with acetone (10 mL). The filtrate was concentrated under reduced pressure and purified by flash chromatography (7:1 hexanes:EtOAc) to provide the desired product as a dense liquid in 78% yield as a 2:1 mixture of olefin isomers. R_F 0.60 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 6.13 (m, 1H, minor), 4.23-4.03 (m, 4H, 2H from minor isomer), 2.96-2.92 (m, 2H), 2.81-2.62 (m, 4H), 2.30-2.24 (m, 2H), 2.28 (s, 3H), 2.11 (s, 3H minor), 2.08-1.94 (m, 2H), 1.79 (m, 2H), 1.74 (bs, 3H minor), 1.71 (bs, 3H), 1.61-1.59 (m, 3H), 1.51-1.39 (m, 1H), 1.25 (t, $J = 7.2$ Hz, 3H), 1.22 (t, $J = 7.2$ Hz, 3H minor), 1.19-1.09 (m, 1H), 0.86 (qd, $J = 12.8, 3.2$ Hz 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 170.5, 167.9, 163.4, 162.8, 161.1, 160.2, 152.9, 141.3, 139.9, 127.2, 125.1, 121.0, 77.9, 59.5, 49.4, 32.5, 28.4, 27.0, 26.8, 25.2, 23.5, 22.5, 22.4, 21.0, 20.8,

14.5, 14.4, 11.2, 8.7; IR (film) 1773, 1743, 1696, 1371, 1197, 1105 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{20}\text{O}_4]^+$: m/z , 264.1361 found 264.1366.

Procedure for Saponification of 6 (SM Table 2): To a solution of **6** (100 mg, 0.384 mmol) in anhydrous MeOH (3 mL) at 23 °C under N_2 atmosphere was added KOH (64.5 mg, 1.15 mmol). The resulting mixture was stirred at 23 °C for 2 h and then the MeOH was evaporated under reduced pressure. The concentrate was diluted with ether (10 mL) and poured into cold 1N HCl (20 mL). The aqueous layer was extracted with ether (2 X 10 mL) and the combined organic phase was washed with brine (15 mL) and dried over MgSO_4 . Concentration *in vacuo* afforded an oily product which was purified by flash column chromatography (7:1 hexanes:EtOAc) to give the β -hydroxyester **25** predominantly as its enol tautomer (79.7 mg, 95% yield) as a white solid. mp 53-55 °C; R_F 0.57 (3:1 hexanes:EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 11.07 (bs, 1H), 7.60 (d, J = 7.6 Hz, 1H), 7.28-7.24 (m, 2H), 7.14-7.10 (m, 1H), 4.42 (q, J = 7.2 Hz, 2H), 3.55 (q, J = 7.6 Hz, 1H), 1.45 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 184.6, 169.3, 139.7, 138.4, 127.3, 123.9, 122.7, 120.3, 103.6, 60.7, 43.1, 15.0, 14.5; IR (film) 1717, 1653, 1594, 1233, 1197 cm^{-1} ; HRMS (EI^+) calc'd for $[\text{C}_{13}\text{H}_{14}\text{O}_3]^+$: m/z , 218.0943 found 218.0940.

Procedure for Allylation of 6 (SM Table 2): To a cooled suspension (0 °C) of NaH (60% dispersion in oil, 32.1 mg, 0.790 mmol) in anhydrous THF (3 mL) under an atmosphere of N_2 was added a solution of **6** (158 mg, 0.607 mmol) in anhydrous THF (1 mL). The resulting gray suspension was stirred 20 min then allyl bromide (56.5 μL , 0.668 mmol) was added. Stirring was continued for 30 min at 0 °C then the ice-bath was removed and the mixture was stirred with warming to 23 °C for a further 3 h. The

mixture was cooled to 0 °C and quenched with water (15 mL). The solution was warmed to 23 °C and extracted with ether (3 X 10 mL). The combined ethereal phase was washed with brine (10 mL), dried over MgSO₄ and concentrated under reduced pressure. The crude oil was purified by flash column chromatography (7:1 hexanes:EtOAc) to yield **26** (150 mg, 82%) as a faint yellow oil and as a >95:5 inseparable mixture of regioisomers: R_F 0.62 (3:1 hexanes:EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 7.37 (d, *J* = 7.6 Hz, 1H), 7.31-7.26 (m, 1H), 7.22-7.17 (m, 2H), 5.46 (m, 1H), 5.01-4.95 (m, 1H), 4.91-4.88 (m, 1H), 4.17-4.05 (m, 2H), 2.90-2.80 (m, 2H), 2.62 (s, 3H), 1.94 (s, 3H), 1.17 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 171.3, 167.4, 149.7, 143.0, 140.3, 132.6, 127.9, 126.6, 125.6, 123.1, 119.3, 118.3, 61.5, 61.0, 38.5, 21.0, 14.1, 9.8; IR (film) 1772, 1728, 1241, 1196, 1763 cm⁻¹; HRMS (EI⁺) calc'd for [C₁₈H₂₀O₄]⁺: *m/z*, 300.1361 found 300.1357.

³¹P NMR Experiments:

In a preliminary study to identify the active catalyst for the pent-annulation of propargylic esters, we have undertaken a series of ³¹P NMR and preparative experiments at 23 °C as well as at 100 °C. Phosphoric acid was used as an external standard. Thus, a ³¹P NMR of 0.3 M solution of H₃PO₄ (85% solution in water) in CDCl₃ was recorded at 23 °C and the corresponding phosphorous peak of H₃PO₄ was calibrated to 0.00 ppm. Other ³¹P NMR experiments were carried out based on the calibrated value.

A 0.02 M solution of PtCl₂(PPh₃)₂ (9.48 mg,) in 1,2-dichloroethane-*d*₄ (600 μL) was placed in an oven-dried NMR tube, and the tube was sealed under vacuum using an NMR single port sealing manifold. A ³¹P NMR spectrum was recorded at 23 °C, then the NMR tube was heated for 8 h at 100 °C in an oil bath, at which time a ³¹P NMR spectrum was recorded at 23 °C as well as at 100 °C in the NMR instrument. In all the experiments, PtCl₂(PPh₃)₂ showed an apparent triplet centered at δ 15.7 ppm.

³¹P NMR for PtCl₂(PPh₃)₂: ³¹P NMR (162 MHz, CD₂Cl₂, 85% H₃PO₄ = 0.00 ppm, 23 °C) δ 15.70 (t, *J* = 1836 Hz) (see page 39).

A solution of PtCl₂(PPh₃)₂ (9.48 mg, 0.012 mmol), and PhIO (5.27 mg, 0.024 mmol) in 1,2-dichloroethane-*d*₄ (600 μL) was placed in an oven-dried NMR tube, and the

tube was sealed under vacuum using an NMR single port sealing manifold. A ^{31}P NMR spectrum was recorded at 23 °C, which showed a triplet centered at δ 15.7 ppm (corresponds to $\text{PtCl}_2(\text{PPh}_3)_2$). Then the sealed NMR tube was heated to 100 °C for 8 h in an oil bath, cooled to 23 °C and a ^{31}P NMR spectrum was recorded at 23 °C. The ^{31}P NMR of the $\text{PtCl}_2(\text{PPh}_3)_2$ and PhIO mixture showed five intense signals along with a less intense triplet centered at δ 15.7 ppm, which corresponds to the starting $\text{PtCl}_2(\text{PPh}_3)_2$. A similar ^{31}P NMR spectrum was observed at 100 °C in the NMR instrument.

^{31}P NMR for $[\text{PtCl}_2(\text{PPh}_3)_2$ and PhIO mixture]: ^{31}P NMR (162 MHz, CD_4Cl_2 , 85% H_3PO_4 = 0.00 ppm, 23 °C) δ 15.70 (t, J = 1836 Hz), 28.3, 37.9, 42.8, 43.9, 72.2 (see page 40).

References:

1. Midland, M. M.; Tramontano, A.; Cable, J. R. *J. Org. Chem.*, **1980**, *45*, 28-29.

**^{31}P NMR of
 $\text{PtCl}_2(\text{PPh}_3)_2$ in 1,2-DCE- d_4 at 23 °C**