Supporting Information

(for Marinescu, L.; Thinggaard, J.; Thomsen, I. B.; Bols; M. "Radical Azidonation of Aldehydes."

J. Org. Chem.)

Contents:

General procedures	S1
Physical (Chromatography, NMR, IR, MS) data of compounds:	S1-S5
¹³ C-NMR spectra for 4b , 4d , 4h and 4i	S6-S9
References	S10

General. Moisture sensitive reactions were carried out in flame-dried glassware under N₂ atmosphere in solvents dried according to standard procedures. Commercially available compounds were used without further purification. Evaporation was carried out on a rotatory evaporator with the temperature kept below 40°C. Columns were packed with silica gel 60 (230-400 mesh) as the stationary phase. TLC-plates (Merck, 60, F₂₅₄) were visualized by spraying with cerium sulfate (1%) and molybdic acid (1.5%) in 10% H₂SO₄ and heating until colored spots appeared. NMR-spectra were recorded on either a 200MHz or 400MHz (¹H) instrument. Mass spectra were run on a Electrospray LC-TOF instrument.

3-Phenylpropanoyl azide (**2a**)¹

¹H NMR (400 MHz, CDCl₃): δ 7.32–7.19 (m, 5H), 2.95 (t, 2H), 2.68 (t, 2H).

Decanoyl azide (**2b**)^{12c}

¹H NMR (400 MHz, CDCl₃): δ 1.62 (t, 2H), 1.24 (m, 14H), 0.88 (t, 3H).

IR (cm^{-1}): 2925 (strong), 2137 (medium N_3), 1714 (strong, C=O).

Octanoyl azide (2c)²

^1H NMR (400 MHz, CDCl_3): δ 1.58 (t, 2H), 1.22 (m, 10H), 0.79 (t, 3H).

IR (cm^{-1}): 2928 (strong), 2139 (medium N_3), 1710 (strong, C=O).

2-Phenylethyl isocyanate (3a)³

^1H NMR (400 MHz, CDCl_3): δ 7.2 – 7.04 (m, 5H), 3.22 (t, 2H), 2.62 (t, 2H).

Nonyl isocyanate (3b)⁴

^1H NMR (400 MHz, CDCl_3): δ 3.58 (t, 2H), 1.55 (m, 2H), 1.2 (m, 12H), 0.82 (t, 3H).

Heptyl isocyanate (3c)⁵

^1H NMR (400 MHz, CDCl_3): δ 3.17 (m, 2H), 1.56 (m, 2H), 1.21 (m, 12H), 0.81 (t, 3H).

Heptyl carbamoylazide (4c)^{12a}

Flash chromatography performed in pentane- CH_2Cl_2 1:1.

^1H NMR (400 MHz, CDCl_3): δ 5.28 (bs, 1H), 3.2 (q, 2H, $J = 8.0$ Hz), 1.52-1.44 (m, 2H), 1.29 -1.21 (m, 8H), 0.84 (t, 3H, $J = 7.2$ Hz).

MS (ES) m/z 207.1 [$\text{M} + \text{Na}^+$].

Nonyl carbamoylazide (4b)

Flash chromatography performed in pentane- CH_2Cl_2 1:1.

^1H NMR (400 MHz, CDCl_3): δ 5.3 (bs, 1H), 3.2 (q, 2H), 1.5 (m, 2H), 1.25 (m, 12H), 0.85 (t, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 156.617 (C=O), 41.344 (C_1), 32.015 (C_2), 29.702, 29.641, 29.404 (C_3 , C_4 , C_5), 26.885 (C_7), 22.824 (C_8), 14.228 (C_9)

IR (KBr, cm^{-1}): 3314 (NH), 2182, (medium, N_3), 1682 (strong, C=O), 1546 (N-C=O).

HRMS (ES): calculated for $\text{C}_{10}\text{H}_{20}\text{N}_4\text{ONa}$ [$\text{M} + \text{Na}^+$]: 235.1535, found: 235.1532.

2-phenylethyl carbamoylazide (4a)^{12b}

Flash chromatography performed in pentane-EtOAc 10:1.

^1H NMR (400 MHz, CDCl_3): δ 7.25-7.10 (m, 5H), 4.99 (bs, 1H), 3.42 (q, 2H, $J = 6.7$ Hz), 2.75 (t, 2H). ^{13}C NMR (100 MHz, CDCl_3): δ 154.960 (C=O), 51.825, 28.763 (CH_3).

MS (ES): m/z 213.1 [$\text{M} + \text{Na}^+$].

***t*-butyl carbamoylazide (4d)**

Flash chromatography performed in pentane-EtOAc 5:1.

^1H NMR (400 MHz, CDCl_3): δ 5.1 (bs, 1H), 1.3 (s, 9H).

^{13}C NMR (100 MHz, CDCl_3): δ 154.960 (C=O), 51.825, 28.763 (CH_3).

IR (KBr, cm^{-1}): 3320 (NH), 2133, (medium, N_3), 1702 (strong, C=O), 1529 (N-C=O).

HRMS (ES): calculated for $\text{C}_5\text{H}_{10}\text{N}_4\text{ONa}$ [$\text{M} + \text{Na}^+$]: 165.0752, found: 165.0759.

Cyclohexyl carbamoylazide (4e)^{12d}

Flash chromatography performed in pentane-EtOAc 10:1.

^1H NMR (400 MHz, CDCl_3): δ 5.03 (bs, 1H), 3.56 – 3.46 (m, 1H), 1.88 - 1.82 (m, 2H), 1.66 - 1.60 (m, 2H), 1.57 – 1.51 (m, 2H), 1.34 – 1.23 (m, 2H), 1.14 – 1.04 (m, 2H).

HRMS (ES): calculated for $\text{C}_7\text{H}_{12}\text{N}_4\text{ONa}$ [$\text{M} + \text{Na}^+$]: 191.0908, found: 191.0910.

Phenyl carbamoylazide (4f)^{12c}

Flash chromatography performed in pentane-diethylether 10:1.

¹H NMR (400 MHz, CDCl₃): δ 7.4 (m, 3H), 7.1 (m, 2H), 7 (bs, 1H)

HRMS (ES) C₇H₆N₄ONa [M + Na⁺]; calculated: 185.0439 found: 185.0443

4-Methoxyphenyl carbamoylazide (4g)

Flash chromatography performed in pentane-diethylether 10:1.

¹H NMR (400 MHz, CDCl₃): δ 7.3 (d, 2H, *J* = 8 Hz), 6.8 (d, 2H), 6.6 (s, 1H), 3.7 (s, 3H).

IR (cm⁻¹): 2144 (medium, N₃), 1675 (strong, C=O).

4-methylphenyl carbamoylazide (4h)

Flash chromatography performed in pentane-EtOAc 10:1.

¹H NMR (400 MHz, CDCl₃): δ 7.2 (d, 2H, *J* = 8.0 Hz), 7 (d, 2H, *J* = 8.0 Hz), 6.7 (bs, 1H), 2.2 (s, 3H).

¹³C NMR (100 MHz, CDCl₃): δ 154.235 (C=O), 134.6 (Ph), 129.883 (Ph), 119.653 (Ph), 21.035 (CH₃).

IR (KBr, cm⁻¹): 3276 (NH), 2143, (medium, N₃), 1682 (strong, C=O), 1538 (N-C=O).

HRMS (ES): calculated for C₈H₈N₄ONa [M + Na⁺]: 199.0596, found: 199.0594.

2-methylphenyl carbamoylazide (4i)

Flash chromatography performed in pentane-EtOAc 10:1.

¹H NMR (400 MHz, CDCl₃): δ 7.7 (d, 1H, *J* = 10.0 Hz), 7.18-7.02 (m, 3H), 6.5 (bs, 1H), 2.2 (s, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 154.487 (C=O), 135.028 (Ph), 130.822 (Ph), 127.211 (Ph), 122.165 (Ph), 17.785 (CH_3).

IR (KBr, cm^{-1}): 3321 (NH), 2147, (medium, N_3), 1679 (strong, C=O), 1550 (N-C=O).

HRMS (ES): calculated for $\text{C}_8\text{H}_8\text{N}_4\text{ONa}$ [$\text{M} + \text{Na}^+$]: 199.0596, found: 199.0596.

Hydrolysis of **4f**.

64 mg (0.4 mmole) phenyl carbamoylazide (**4f**) was stirred at room temperature with 1 mL NaOH (2M) and 1 mL dioxane for 30 min. Then 1 mL hydrochloric acid 6M was added and the mixture was stirred for another 30 min. Now 1 mL NaOH (2M) was added until basic pH and the mixture was extracted with diethylether. The organic layer was dried and evaporated under reduced pressure to give aniline **6**. Yield: 29 mg (78%).

N-butyl-*N*-nonyl urea (**7**).

To 63 mg (0.3 mmole) nonyl carbamoylazide **4b** was added slowly 5 mL (0.05 mole) butyl amine and the solution was allowed to reflux 1.5 h. Then 1.8 mL HCl (12 M) was added, and reaction mixture was extracted with diethylether. The organic extracts were dried over Na_2SO_4 and concentrated under reduced pressure. Yield: 73 mg (85%).

^1H NMR: (400 MHz, CDCl_3) δ 4.28 (bs, 1H), 3.11–3.05 (m, 4H), 1.42–1.38 (m, 4H), 1.27–1.16 (m, 14H), 0.87–0.79 (m, 6H).

HRMS (ES) calculated for $\text{C}_{14}\text{H}_{30}\text{N}_2\text{ONa}$ [$\text{M} + \text{Na}^+$]: 265.2255, found: 265.2249.

4b:

28. carbon2
exp1 s2pul1

SAMPLE		SPECIAL	
date	Sep 25 2003	temp	26.0
solvent	CDC13	gain	not used
file	exp	sp	0
sw	25188.9	pw90	10.800
at	1.199	alpha	9.550
np	60422	l1	11
fb	13800	n	n
ds	1.000	ds	y
nt	256	hs	nn
ct	256	hs	nn
tn	100	fn	1.00
tof	156106	sp	65536
tpwr	62	wp	-201.4
pw	5.400	rfl	21146.6
dn	H1	rfl	1517.7
dmf	yyv	lp	-73.1
dmf	yyv	lp	-206.2
dmf	yyv	wc	270
dmf	yyv	sc	36
dmf	yyv	vs	85
dmf	yyv	th	6
dmf	yyv	al	no
dmf	yyv	ph	6

28. carbon2

4d:

4h:

S8

4i:

References:

- ¹Stefanuti, I.; Smith, S. A.; Taylor, R. J. K. *Tetrahedron Lett.* **2000**, *41*, 3735-3738.
- ²Gumaste, V. K.; Bhawal, B. M.; Deshmukh, A. R. A. S. *Tetrahedron Lett.* **2002**, *43*, 1345-1346.
- ³Minisci, F.; Coppa, F.; Fontana, F.; *J.Chem.Soc.Chem.Comm.* **1994**, 679-680.
- ⁴Valenta, V.; Bartosova, M.; Protiva, M.; *Collect.Czech.Chem.Comm.* **1980**, *45*, 517-528.
- ⁵Plusquellec, D.; Roulleau, F.; Lefeuvre, M.; Brown, E.; *Tetrahedron*, **1990**, *46*, 465-474