Radicamines A and B: Synthesis and Revision of the Absolute Configuration

Chu-Yi Yu^{*a} and Mu-Hua Huang^{a,b}

*Contribution from ^aBeijing National Laboratory for Molecular Science (BNLMS), Laboratory for
Chemical Biology, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100080, China
and ^bGraduate University of the Chinese Academy of Sciences, Beijing 100049, China
yucy@iccas.ac.cn*

Supporting Information

Section A:	General Information	S-2
Section B:	Experimental Procedures	S-3
Section C:	¹H, ¹³C and 2D-NMR Spectrum.	S-13
Section D:	References	S-42

Section A: General Information

All reagents were used as received from commercial sources without further purification or prepared as described in the literature. Tetrahydrofuran was distilled from sodium and benzophenone immediately before use. Reactions were stirred using Teflon-coated magnetic stirring bars. Analytical TLC was performed with 0.20 mm silica gel 60F plates with 254 nm fluorescent indicator. TLC plates were visualized by ultraviolet light or by treatment with a spray of Pancaldi reagent $\{(\text{NH}_4)_6\text{MoO}_4, \text{Ce}(\text{SO}_4)_2, \text{H}_2\text{SO}_4, \text{H}_2\text{O}\}$ or a solution 0.5% ninhydrin in acetone. Chromatographic purification of products was carried out by flash column chromatography on silica gel (230-400 mesh). Acidic ion exchange chromatography was performed on Amberlite[®] IR-120 (H^+) or Dowex 50WX8-400, H^+ form. Melting points were determined using an electrothermal melting point apparatus. Both melting points and boiling points are uncorrected. Infrared spectra were recorded on a JASCO FT/IR-480 plus Fourier transform spectrometer. NMR spectra were measured in CDCl_3 (with TMS as internal standard) or D_2O on a Bruker AV300 (^1H at 300 MHz, ^{13}C at 75 MHz) or a Bruker AV600 (^1H at 600 MHz, ^{13}C at 150 MHz) magnetic resonance spectrometer. Chemical shifts (δ) are reported in ppm, and coupling constants (J) are in Hz. The following abbreviations were used to explain the multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet. High-resolution mass spectra (HRMS) were recorded on an APEXII FT-ICR (SIMS) mass spectrometer. Polarimetry was carried out using an Optical Activity AA-10R polarimeter and the measurements were made at the sodium D-line with a 0.5dm pathlength cell. Concentrations (c) are given in gram per 100mL.

Section B: Experimental Procedures

(3*S*,4*S*,5*R*)- 5-Hydroxy-3,4,6-tribenzyloxy-1-hexene 13¹

ⁿButyllithium (1.6 M in hexane, 34 ml, 54.5 mmol) was added dropwise to a stirred suspension of

methyltriphenylphosphonium bromide (19.6 g, 55 mmol) in dry THF (100 mL) at about 0°C (ice-bath) in the atmosphere of Argon. The ice-bath was then removed and the resulted yellowish mixture was stirred for another 4 hours at room temperature. The reaction mixture was cooled with ice-water bath again, to which a solution of 2,3,5-O-tribenzyl-D-xylofuranose **12** (7.70 g , 18.3 mmol) in THF (60 mL) was added dropwise. A cream-colored precipitate appeared, and the reaction mixture became difficult to stir. After stirring in room temperature for 12 hours, the mixture was heated for another 75 mins. The solvent was removed in vacuo, the residue was fully distributed between ethyl acetate and water. The combined ethyl acetate phase was dried and concentrated, and purified by flash column chromatography (silica gel, petroleum ether/AcOEt = 4:1) to give the title compound **13** as a light yellow oil (5.90 g , yield 77%).

$[\alpha]_D^{20} = -10.7$ (c= 0.56, CHCl₃) {for its enantiomer $[\alpha]_D^{25} = +5.8$ (c= 1.6, CHCl₃) in Pearson, W. H. ; Hines, J. V. *J. Org. Chem* **2000**, 65, 5785-5793, }. IR (cm⁻¹): 3447(w), 3064(w), 3030(w), 2864(w), 1507(m), 1496(s), 1091(vs), 1062(vs).ESI-MS: m/z 419.4 [M+H]⁺; 441.4 [M+Na]⁺. HRMS (SIMS): m/z 419.2218 [M+H]⁺, (C₂₇H₃₁O₄ requires 419.2217) .

**(3S,4R,5R)-5-Methanesulfonyloxy-3,4,6-tribenzyloxy-1-hexene 14 and
(2R,3R,4R)-1,3,4-tris(benzyloxy)-4-formylbutan-2-yl methanesulfonate 15**

Pyridine (10ml) was added to a solution of (3S,4S,5R)- 5-hydroxy-3,4,6-tribenzyloxy-1-hexene **13** (5.90 g, 14 mmol) in DCM (10 ml). The resulting solution was cooled with ice-bath and added dropwise with methanesulphonyl chloride (2.0 ml, 25.7 mmol). The reaction mixture was stirred at room temperature for about 10 hours. Then the solvent was evaporated in vacuo, the residue was purified by flash column chromatography (silica gel, petroleum ether/AcOEt = 4:1) to give the mesylate **14** as a light yellow oil (6.56 g, yield 94%).

$[\alpha]_D^{20} = +15.6$ (c= 0.64, CHCl₃). IR (cm⁻¹): 3063(w), 3031(w), 2979(w), 2936(w), 1496(w), 1454(m), 1358(vs), 1174(vs), 1115(s), 1088(s), 1066(s), 1028(s). ¹H-NMR (CDCl₃, 300 M) : δ (ppm): 7.34-7.24 (m, 15 H ,ArH) , 5.96-5.85 (m, 1 H, CH=) , 5.39 (dd, 2H, J = 9.3 and 9.6 Hz) ,

4.88 (dt, 1 H, $J = 2.9, 3.0$, and 8.9 Hz), 4.68 (ABQ, 2H, $J = 11.3$ Hz), 4.41 (ABQ, 2 H, $J = 11.7$ Hz), 4.38 (ABQ, 2 H, $J = 11.8$ Hz), 3.97 (dd, 1 H, $J = 4.3$ and 7.4 Hz), 3.67 (dd, 1 H, $J = 2.9$ and 11.4 Hz), 3.54 (dd, 1 H, $J = 5.8$ and 11.3 Hz), 2.90 (s, 3 H). ^{13}C -NMR (CDCl_3 , 75M): δ (ppm) 38.40 (CH_3), 69.19 (CH_2O), 70.53 (OCH_2), 73.29 (OCH_2), 75.35 (OCH_2), 79.37 (CHO), 80.03 (CHO), 81.86 (CHO), 119.98 ($=\text{CH}_2$), 127.82, 127.91, 127.97, 128.28, 128.38, 128.44, 128.50, 129.04, 129.79, 134.49 ($\text{CH}=\text{}$), 137.49 (C), 137.71 (C), 137.76 (C). ESI-MS: m/z 519.4 $[\text{M}+\text{Na}]^+$. HRMS (SIMS): m/z 497.1984 $[\text{M}+\text{H}]^+$ ($\text{C}_{28}\text{H}_{33}\text{O}_6\text{S}$, requires 497.1992); 519.1805 $[\text{M}+\text{Na}]^+$ ($\text{C}_{28}\text{H}_{33}\text{O}_6\text{SNa}$, requires 519.1811).

A solution of the alkene-mesylate **14** (550mg, 1.1mmol) in DCM (50mL) was cooled with cold acetone bath (-78°C) and bubbled with O_3 in O_2 atmosphere for 7 mins. TLC indicated the disappearance of the starting material. The unreacted O_3 was driven off by purging the system with N_2 . Dimethyl sulfide (0.24 ml) was then added to the reaction mixture. After stirring for 3 hours, the solvent was removed in vacuo. The light yellow residue (800mg) was purified by flash chromatography (silica gel, petroleum ether/AcOEt = 2:1) to give the title compound **15** as a colorless oil (415mg, yield 76%).

$[\alpha]_{\text{D}}^{20} = +30.3^0$ ($c = 1.1, \text{CHCl}_3$). IR(cm^{-1}): 3031(w), 2936(w), 2871(w), 1732(s), 1496(w), 1454(m), 1358(vs), 1175(vs), 1092(s). DEPT-135 (CDCl_3 , 75M): δ (ppm) positive: 201.82, 128.67, 128.58, 128.47, 128.42, 128.32, 128.07, 128.01, 81.49, 80.26, 77.34, 38.25. negative: 74.75, 73.45, 68.44. ^{13}C -NMR (CDCl_3 , 75M): δ (ppm) 201.82, 137.26, 136.80, 136.40, 128.67, 128.58, 128.47, 128.42, 128.32, 128.07, 128.01, 81.49, 80.26, 77.34, 74.75, 73.45, 68.44, 38.25. ESI-MS: m/z 521.4 $[\text{M}+\text{Na}]^+$. HRMS (SIMS): m/z 521.1609 $[\text{M}+\text{Na}]^+$ ($\text{C}_{27}\text{H}_{30}\text{O}_7\text{SNa}$, requires 521.1604).

(3*S*,4*S*,5*S*)-3,4-Dibenzyloxy-5-benzyloxymethyl-1-pyrroline-N-oxide **9**²

A solution of sodium carbonate (5.09 g, 48 mmol) in H_2O (7ml) was added to a suspension of

hydroxylamine hydrogen chloride (3.36 g, 48 mmol) in MeOH (20 ml). The resulting mixture (pH = 8.0) was added with a solution of the mesylate-aldehyde **15** (8.03 g, 16 mmol) in THF (6 ml) and MeOH (8 ml). The reaction mixture was stirred and heated (at 50-60 °C, oil bath) for 79 hours. Then water and AcOEt was added and the two phases were separated. The AcOEt phase was washed with water and the aqueous layer was extracted with AcOEt several times until TLC revealed no more products remaining in the aqueous solution. The combined organic phase was dried (Na₂SO₄) and concentrated. The residue was purified by flash column chromatography (silicon gel, petroleum ether/AcOEt: 2:1) to give the target nitron **9** as a white solid (4.03g, yield 60%).

m.p. 90-91 °C. $[\alpha]_D^{20} = +45^\circ$ (c = 0.4, CHCl₃) {Lit.^{2a} *ent*-**9**: m.p. 80-82 °C. $[\alpha]_D^{19} = -41.7^\circ$ (c = 1.00, CHCl₃); Lit.^{2b} *ent*-**9**: m.p. 91-93 °C. $[\alpha]_D^{19} = -42^\circ$ (c = 1.3, CHCl₃)}. IR (cm⁻¹): 3049(w), 2945, 2923, 2901, 2884, 2868, 2851(w), 1593 (s), 1551(s), 1496(s), 1452(s), 1361(s), 1247(vs), 1131(vs), 1247(vs), 1028 (vs). ¹H-NMR (CDCl₃): δ (ppm) 7.38– 7.26 (m, 15 H), 6.91 (d, *J*=1.9 Hz, 1 H), 4.69–4.67 (m, 1 H), 4.64–4.46 (m, 6 H), 4.39 (dd, *J*=3.2, 2.2 Hz, 1 H), 4.10–4.04 (m, 2 H), 3.78 (d, *J*=7.3 Hz, 1 H). ¹³C-NMR (CDCl₃, 75M): δ (ppm) 66.03 (CH₂O), 71.67, 71.91, 73.47, 80.30 (CHO), 82.74 (CHO), 127.70, 127.75, 127.92, 128.14, 128.17, 128.38, 128.55, 128.61, 133.02 (N=CH), 137.06 (C), 137.16 (C), 137.63 (C). [lit : ¹³C NMR: δ (ppm) 137.7 (s), 137.2 (s), 137.1 (s), 132.9 (d), 128.6, 128.5, 128.4, 128.1, 127.9 and 127.7 (d, 15 C), 82.7 (d), 80.3 (d), 77.4 (d), 73.9 (t), 71.9 (t), 71.6 (t), 66.1 (t).]. ESI-MS : m/z 440.3 [M+Na]⁺. HRMS (SIMS): m/z 418.2007 [M+H]⁺ (C₂₆H₂₈NO₄ requires 418.2013) .

(2*S*,3*S*,4*S*,5*S*)-2-(benzyloxymethyl)-3,4-dibenzyloxy-5-phenyl-N-Hydroxy-pyrrolidine **17**

To a solution of phenylmagnesium bromide in THF [prepared by heating Mg turnings (60 mg, 2.5 mmol) and bromobenzene (0.26 ml, 2.5 mmol) in hot dry THF] was added dropwise with a solution of cyclic nitron **9** (200 mg, 0.5 mmol) in THF (6 ml) at about 0 °C (ice-water bath). The reaction mixture was stirred overnight (about 12 h.), quenched with sat. aqueous NaHCO₃ solution and extracted with diethyl ether and AcOEt. The combined extracts were dried and solvents were

removed in vacuo. The residue was purified by flash column chromatography (silica gel, petroleum ether/AcOEt: 4/1) to give the title compound **17** as an colorless oil, which was dried under vacuum to give a white solid (216mg, yield 91%).

m.p. 81-82 °C. $[\alpha]_D^{25} = +16.8$ ($c=2.5$,CHCl₃). IR(cm⁻¹): 3343(w), 3107(w), 3087(w), 3063(w), 3031(w), 2971(w), 2870(w), 1604(w), 1595(w), 1496(w), 1472(w), 1454(m), 1365(w), 1098(s), 1076(vs). ¹H-NMR (CDCl₃): δ (ppm) : 3.69-3.75 (m, 1 H) , 3.78 (t , 1H, $J = 7.2$ Hz) , 3.89 (dd , 1 H , $J = 3.7$ and 8.7 Hz) , 4.06 (dd, 1H, $J = 3.4$ and 7.2 Hz) , 4.11 (dd , 1H, $J = 2.7$ and 3.2 Hz) , 4.23 (d, 1H, $J = 7.2$ Hz) , 4.33 (ABQ, 2 H , $J = 11.8$ Hz) , 4.55 (ABQ, 2H, $J = 12.0$ Hz) , 4.56 (ABQ, 2 H , $J = 12.1$ Hz) , 4.94 (s, 1H , OH) , 7.07-7.44 (m , 20 H) . ¹H-NMR (CDCl₃+D₂O): δ (ppm) : 3.73-3.77 (m, 1 H) , 3.80 (t , 1H, $J = 7.2$ Hz) , 3.93 (dd , 1 H , $J = 3.9$ and 8.9 Hz) , 4.09 (dd, 1H, $J = 3.4$ and 7.2 Hz) , 4.14 (dd , 1H, $J = 2.8$ and 3.2 Hz) , 4.26 (d, 1H, $J = 7.2$ Hz) , 4.37 (ABQ, 2 H , $J = 11.9$ Hz) , 4.60 (s, 2H) , 4.58 (ABQ, 2 H , $J = 12.0$ Hz) , 7.12-7.47 (m , 20 H) . DEPT-135 (CDCl₃, 75M) : δ (ppm) positive : 128.50, 128.38, 128.34, 128.23, 128.05, 127.78, 127.69, 127.67, 127.62, 127.56, 87.49, 83.68, 74.06, 68.93. negative : 73.44, 72.01, 71.66, 66.86. ¹³C-NMR (CDCl₃, 75M): δ (ppm) 139.36, 138.25, 138.11, 137.89, 128.49, 128.38, 128.34, 128.23, 128.05, 127.77, 127.69, 127.67, 127.61, 127.55, 87.50, 83.69, 74.07, 73.44, 72.01, 71.66, 68.93, 66.87. ESI-MS : m/z 496.5 [M+H]⁺ . HRMS (SIMS) : m/z 496.2485 [M+H]⁺ , (C₃₂H₃₃NO₄ requires 496.2482).

(2S,3S,4S,5S)-2-(hydroxymethyl)-5-phenylpyrrolidine-3,4-diol **18**

10 ml of MeOH and 1 ml of HCl (6N) was added to the solution of the hydroxylamine **17** (200 mg, 0.4 mmol) in DCM (10mL). To which Pd/C (10%, 60 mg) was added under stirring. After hydrogenation under atmospheric pressure for 20 hours, TLC (AcOEt/MeOH: 5/1) revealed the formation of a polar compound. The system was bubbled with N₂, and the Pd/C was filtered off.

After the solution was concentrated in vacuo, the residue was dissolved in MeOH and neutralized with con. aqueous ammonium solution. The resulting solution was concentrated in vacuo. The residue was dried under vacuum and purified by an acidic ion exchanger column (Dowex 5Wx8-400, H⁺ form, Aldrich, column size: 1.3x14cm), eluting with distilled water (50 ml) and then 1N NH₄OH, affording the title compound **18** as a white solid (72 mg, yield 85%).

$[\alpha]_D = -52^0$ (c=0.65, H₂O). ¹H-NMR (D₂O, 300M): δ (ppm) : 7.41 (m, 5H), 4.09 (dd, 1H, *J*=7.4 and 8.3 Hz), 3.93 (broad t, 2H), 3.72 (dd, 1H, *J*=4.7 and 11.6 Hz), 3.65 (dd, 1H, *J*=6.3 and 11.6 Hz), 3.22 (dd, 1H, *J*=6.3 and 11.5 Hz). ¹H-NMR (D₂O, 600M): δ (ppm) : 7.36 (broad d, 4H, H-2', H-3', H-5', H-6'), 7.32-7.27 (m, 1H, H-4'), 4.04 (dd, 1H, *J*=7.5 and 9.0 Hz, H-4), 3.88 (d, 1H, *J*=5.6 Hz, H-3 or H-5), 3.87 (d, 1H, *J*=7.4 Hz, H-5 or H-3), 3.67 (dd, 1H, *J*=4.5 and 11.6 Hz, H-1a), 3.61 (dd, 1H, *J*=6.5 and 11.6 Hz, H-1b), 3.17 (dd, 1H, *J*=2.1 and 6.8 Hz, H-2). DEPT-135 (D₂O, 75M) : δ (ppm) positive : 128.95, 128.23, 127.35, 82.26, 77.51, 64.00, 61.79. negative : 62.46. ¹³C-NMR (D₂O, 75M): δ (ppm) 139.73, 128.96, 128.23, 127.36, 82.27, 77.53, 64.01, 62.47, 61.81. ESI-MS : m/z 210.2 [M+H]⁺, 232.2 [M+Na]⁺. HRMS (SIMS) : m/z 210.1126 [M+H]⁺, (C₁₁H₁₆NO₃ requires 210.1125).

2-(benzyloxy)-4-bromo-1-methoxybenzene **23** ³

To a suspension of sodium hydride (0.95 g, 55% in mineral oil, 21.7 mmol) in THF (10 ml) was added a solution of 5-bromo-2-methoxyphenol **22** (4.0 g, 19.7 mmol) in THF (10 ml). TBAT (0.2 g, 0.5 mmol) was then added to the reaction mixture, followed by addition of a solution of benzyl bromide (2.6 ml, 21.7 mmol) in THF (5 ml). The reaction mixture was stirred at 50-60°C. When TLC indicated the disappearance of the starting material, the reaction mixture was cooled down to room temperature and was poured into ice-water. The white solids precipitating from the solution were collected, filtered and recrystallized from AcOEt/petroleum ether (1:1) to give **23** as colorless needles (3.40 g, yield 90%) .

m.p. 106-107 °C (in lit 70-71 °C). ¹H NMR (CDCl₃, 300 MHz): δ (ppm) 3.88 (3H, s), 5.14 (2H, s), 6.79 (1H, *J*=8.8 Hz), 7.06–7.09 (2H, m), 7.32–7.48 (5H, m). ¹³C NMR (CDCl₃, 75 MHz): δ (ppm) 56.2, 71.2, 112.6, 113.1, 117.3, 124.0, 127.4, 128.1, 128.6, 136.4, 149.0, 149.1;

(2*S*,3*S*,4*S*,5*S*)-2-(benzyloxymethyl)-3,4-dibenzyloxy-5-(3-benzyloxy-4-methoxyphenyl)-N-hydroxy-pyrrolidine **24**

A solution of 2-(benzyloxy)-4-bromo-1-methoxybenzene **23** (0.73 g, 2.5 mmol) in THF (4 ml) was added to a mixture of magnesium turnings (0.10 g, 4.2 mmol) in hot THF (4 ml). The mixture was heated to reflux in the atmosphere of N₂ until the magnesium disappeared and the grey solution formed. The Grignard reagent was cooled with an ice-water bath, and a solution of the cyclic nitrone **9** (300 mg, 0.72 mmol) in dry THF (5 ml) was added dropwise. The reaction mixture was stirred at room temperature overnight (about 12 h.), then quenched with sat. aqueous NaHCO₃ solution, extracted with diethyl ether and AcOEt. The combined organic phase was dried (anhydrous Na₂SO₄) and solvents were removed in vacuo. The residue was purified by flash column chromatography (silica gel, petroleum ether/ACOEt: 4/1) to give the title compound **24** as an colorless oil, which turned into a white solid (400mg, yield 89%) after being placed in a refrigerator.

[α]_D = -4.4 (c=0.46, CHCl₃). ¹H-NMR (CDCl₃, 300 M): δ (ppm): 3.70-3.75 (m, 1H), 3.79 (dd, 1H, *J*=7.2 and 9.1Hz), 3.91 (dd, 1H, *J*=4.2 and 8.9 Hz), 3.91 (s, 3H), 3.98 (dd, 1H, *J*=3.5 and 7.3 Hz), 4.12 (dd, 1H, *J*=3.2 and 3.0 Hz), 4.16 (d, 1H, *J*=7.3 Hz), 4.31 (ABQ, 2H, *J*=11.9 Hz), 4.58 (ABQ, 2H, *J*=11.9 Hz), 4.60 (s, 2H), 4.72 (s and b, 1H, OH), 5.11 (s, 3H), 6.88 (d, 1H, *J*=8.2 Hz), 6.99 (dd, 1H, *J*=1.8 and 8.2 Hz), 7.04 (d, 1H, *J*=1.8 Hz), 7.09-7.13 (m, 2H), 7.25-7.45 (m, 14H). ¹³C-NMR (CDCl₃, 75M): δ (ppm) 56.1 (OCH₃), 67.0, 68.7, 70.9, 71.7, 72.0, 73.5, 73.7, 83.7, 87.4, 111.6, 113.7, 121.5, 127.5, 127.6, 127.7, 127.8, 128.0, 128.2, 128.4, 128.4, 128.5, 131.8, 137.1, 137.9, 138.1, 138.2, 148.1, 148.4. Dept-135 (CDCl₃): δ (ppm) positive :56.1, 68.7, 73.6, 83.7, 87.4,

111.6, 113.7, 121.5, 127.5, 127.6, 127.7, 127.8, 128.0, 128.2, 128.4, 128.4, 128.5. negative: 67.0, 70.9, 71.7, 72.0, 73.5. HRMS (SIMS): m/z 632.3018 $[M+H]^+$, ($C_{40}H_{32}NO_6$ requires 632.3006).

(2*S*,3*S*,4*S*,5*S*)-2-(3-Hydroxy-4-methoxy-phenyl)-5-hydroxymethyl-pyrrolidine-3,4-diols **1**

MeOH (10 ml) and HCl (6 N, 1ml) were added to a solution of **24** (80 mg, 0.1 mmol) in DCM (10 m). To which Pd/C (10%, 60 mg) was added under stirring. After hydrogenation under atmospheric pressure for 20 hours, TLC (AcOEt/MeOH: 5/1) revealed the formation of a polar compound. The system was bubbled with N_2 , and the Pd/C was filtered off. After the solution was concentrated in vacuo, the residue (52 mg) was dissolved in MeOH and neutralized with con. aqueous ammonium solution. The resulting solution was concentrated in vacuo. The residue was dried under vacuum and purified by an acidic ion exchanger column (Dowex 5Wx8-400, H^+ form, Aldrich, column size: 1.3x14cm), eluting with distilled water (50 ml) and then 1N NH_4OH , affording the title compound **1** as a yellow syrup (28 mg, yield 87%).

$[\alpha]_D = -44.4$ ($c=0.45$, H_2O) {in the lit $[\alpha]_D = +43.7$ ($c=0.13$, H_2O)}. IR(cm^{-1}): 3209(vs), 1637(m), 1401(vs), 1276(w), 1252(w), 1132(w), 1037(w). 1H -NMR (D_2O , 300 M): δ (ppm): 6.99 (d, 1H, $J=8.1$ Hz), 6.91 (s, 1H), 6.88 (dd, 1H, $J=5.1$ Hz), 4.04 (dd, 1H, $J=7.5, 8.9$ Hz), 3.91 (t, 1H, $J=7.3$ Hz), 3.81 (s, 3H, OCH_3), 3.80 (d, 1H, $J=9.4$ Hz), 3.72 (dd, 1H, $J=4.7, 11.6$ Hz), 3.65 (dd, 1H, $J=6.4, 11.6$ Hz), 3.21 (dd, 1H, $J=6.4, 11.1$ Hz). 1H -NMR (D_2O , 600 M): δ (ppm): 6.91 (d, 1H, $J=8.2$ Hz, H-5'), 6.84 (s, 1H, H-2'), 6.83 (d, 1H, $J=8.5$ Hz), 3.97 (t, 1H, $J=7.8, 8.8$ Hz), 3.84 (t, 1H, $J=7.3$ Hz), 3.76-3.73 (d or t, 1H, H-5, overlap with OCH_3), 3.74 (s, 3H, OCH_3), 3.64 (dd, 1H, $J=4.4, 11.6$ Hz, H-1a), 3.58 (dd, 1H, $J=6.4, 11.5$ Hz, H-1b), 3.14 (dd, 1H, $J=6.5, 11.2$ Hz, H-2). DEPT 135 (D_2O , 75 MHz): δ (ppm): positive: 119.65, 114.44, 112.75, 81.92, 77.34, 63.51, 61.68, 56.08. negative: 62.38. ^{13}C -NMR (D_2O , 75 MHz): δ (ppm) 147.42, 145.43, 132.63, 119.66, 114.45, 112.77, 81.93,

77.36, 63.52, 62.39, 61.69, 56.10. HRMS (SIMS) : m/z 256.1185 [M+H]⁺, (C₁₂H₁₈NO₅ requires 256.1179).

The comparison of ¹³C-NMR of compound **1** (original lab note: **mhh-357-p**) with that of radicamine A isolated by Kusano (in *Chem. Pharm. Bull.* **2001**,49,1362)

Radicamine A (in D ₂ O) isolated by Kusano	1 (Mhh-357-p) (in D ₂ O)	Δ δ
64.69 (C-1)	62.39 (C-1)	2.30
64.15 (C-2)	61.69 (C-2)	2.46
79.62 (C-3)	77.36 (C-3)	2.26
84.16 (C-4)	81.93 (C-4)	2.23
65.97 (C-5)	63.52 (C-5)	2.45
134.90 (C-1')	132.63 (C-1')	2.27
116.95 (C-2')	114.45 (C-2')	2.50
147.81 (C-3')	145.43 (C-3')	2.38
149.97 (C-4')	147.42 (C-4')	2.55
115.23 (C-5')	112.77 (C-5')	2.46
122.35 (C-6')	119.66 (C-6')	2.69
58.55 (OCH ₃)	56.10 (OCH ₃)	2.45

(2S,3S,4S,5S)-2-(Benzyloxymethyl)-3,4-dibenzyloxy-5-(4-benzyloxyphenyl)-N-hydroxypyrrolidine 26

A mixture of 50 mg (2mmol) of Mg turnings and 500 mg (2mmol) of 4-benzyloxy-bromobenzene⁴ in THF (4 ml) was heated and refluxed for 30 mins. The solution changed from yellow to brown, and was then cooled with an ice-water bath. The cold Grignard reagent was added dropwise to a solution of the cyclic nitron **9** (200 mg, 0.5mmol) in THF (15 ml) at about 0°C (ice-water bath). The reaction mixture was stirred overnight (about 12h.), then quenched with sat. aqueous NaHCO₃ solution and extracted with diethyl ether. The combined extracts were dried (anhydrous Na₂SO₄) and solvents were removed in vacuo. The residue was purified by flash column chromatography (silica gel, petroleum ether/ACOEt: 4/1) to give the title compound **26** as a white solid (225 mg, yield 80%) .

m.p. : 129 °C. $[\alpha]_D^{19} = +18.0$ ($c=0.55$,CHCl₃). IR(cm⁻¹): 3136(s), 1613(w), 1518(w), 1454(w), 1401(vs), 1247(w), 1121(w), 1024(w). ¹H-NMR (CDCl₃, 300M): δ (ppm) : 3.66-3.71 (m, 1 H, H-2) , 3.75 (dd , 1H, $J = 7.0$ and 9.0 Hz , H-4) , 3.84 (dd , 1 H , $J = 3.9$, and 9.0 Hz, H-3) , 4.06-4.11 (m, 2 H, H-1) , 4.20 (d , 1H, $J = 7.0$ Hz, H-5) ,4.34 (ABQ, 2H, $J = 11.8$ and 19.0 Hz) , 4.56 (ABQ, 2 H , $J = 12.0$ Hz) , 4.55 (ABQ, 2 H , $J = 12.0$ Hz) , 5.05 (s, 2 H) , 6.94 (d, 2 H , $J = 8.7$ Hz) , 7.09 (dd, 2 H , $J = 3.6$, 2.3 and 7.3 Hz) , 7.21-7.45 (m , 20 H). ¹³C-NMR (CDCl₃, 75 MHz): δ (ppm) 66.8, 68.9, 70.0 , 71.7, 72.0 (CH₂O), 73.4 , 73.4 (NCH), 83.4 , 87.1(CHO), 114.7 , 127.5 , 127.6 , 127.7 , 128.0 , 128.0 , 128.2 , 128.3 , 128.4 , 128.6 , 129.8 (CH) , 137.0 , 137.8 , 138.0 , 138.1 , 158.5 . ESI-MS : m/z 624.3 [M+Na]⁺ , 602.3 [M+H]⁺. HRMS (SIMS): m/z 602.2899 [M+H]⁺ (C₃₉H₄₀NO₅ requires 602.2901).

(2S,3S,4S,5S)-2-(hydroxymethyl)-5-(4-hydroxyphenyl)pyrrolidine-3,4-diol 2

10 ml of acidic MeOH (200 mL/2 mL : MeOH/AcCl) was added to a solution of **26** (117 mg, 0.2 mmol) in DCM (5 ml). To which Pd/C (10%, 60 mg) was added under stirring. After hydrogenation under atmospheric pressure for 20 hours, TLC (AcOEt/MeOH: 5/1) revealed the formation of a polar compound. The system was bubbled with N₂, and the Pd/C was filtered off. After the solution was concentrated in vacuo, the residue (80 mg) was dissolved in MeOH and neutralized with con. aqueous ammonium solution. The resulting solution was concentrated in vacuo. The residue was dried under vacuum and purified by an acidic ion exchanger column (Dowex 5Wx8-400, H⁺ form, Aldrich, column size: 1.3x14cm), eluting with distilled water (50 ml) and then 1N NH₄OH, affording the title compound **2** as a yellow syrup (41 mg, yield 93%).

$[\alpha]_D^{30} = -72.7$ (c=0.17, H₂O) {in the lit $[\alpha]_D = +72.0$ (c=0.10, H₂O)}. IR(cm⁻¹): 3199(vs), 1617(s), 1512(m), 1400(vs), 1260(w), 1179(w), 1125(w), 1074(w). ¹H-NMR (D₂O, 300 M): δ (ppm) : 7.29 (d, 1H, *J*=8.6Hz), 6.87 (d, 1H, *J*=8.6 Hz), 4.10 (dd, 1H, *J*=7.5, 9.2 Hz, H-4), 3.93 (t, 1H, *J*=7.4 Hz, H-3), 3.90 (d, 1H, *J*=9.2 Hz, H-5), 3.74 (dd, 1H, *J*=4.6, 11.7 Hz, H-1a), 3.67 (dd, 1H, *J*=6.3, 11.7 Hz, H-1b), 3.28-3.22 (m, 1H, H-2). ¹H-NMR (D₂O, 600 M): δ (ppm) : 7.24 (d, 1H, *J*=8.4 Hz, H-2', H-6'), 6.87 (d, 1H, *J*=8.4 Hz, H-3', H-5'), 4.08 (dd, 1H, *J*=7.8 and 9.2 Hz, H-4), 3.91 (d, 1H, *J*=7.2 Hz, H-5), 3.90 (dd, 1H, *J*=3.7 and 7.4Hz, H-3), 3.70 (dd, 1H, *J*=4.3, 11.8 Hz, H-1a), 3.64 (dd, 1H, *J*=6.4, 11.8 Hz, H-1b), 3.24 (dd, 1H, *J*=6.8 and 11.2 Hz, H-2). DEPT135 (D₂O, 300 MHz): δ (ppm) : positive : 129.08, 115.79, 81.40, 76.98, 63.35, 61.59 . negative : 62.02. ¹³C-NMR (D₂O, 300 MHz): δ (ppm) 155.77, 130.48, 129.08, 115.79, 81.40, 76.98, 63.35, 62.02, 61.59. ESI-MS : m/z 226.2 [M+H]⁺. HRMS (SIMS): m/z 226.1074 [M+H]⁺ (C₁₁H₁₆NO₄ requires 226.1074).

The comparison of ¹³C-NMR of compound **2** (original lab note: **mhh-321-p**) with that of radicamine B isolated by Kusano⁵ (in *Chem. Pharm. Bull.* **2001**, 49, 1362)

Radicamine B isolated by Kusano (in D ₂ O)	2 (Mhh-321-p) (in D ₂ O)	$\Delta\delta$
64.38 (C-1)	62.05 (C-1)	2.33
64.12 (C-2)	61.64 (C-2)	2.48
79.35 (C-3)	77.03 (C-3)	2.32
83.72 (C-4)	81.45 (C-4)	2.27
65.86 (C-5)	63.39 (C-5)	2.47
131.78 (C-1')	130.26 (C-1')	1.52
131.68 (C-2')	129.10 (C-2')	2.58
118.34 (C-3')	115.82 (C-3')	2.52
158.30 (C-4')	155.79 (C-4')	2.51
118.34 (C-5')	115.82 (C-5')	2.52
131.68 (C-6')	129.10 (C-6')	2.58

Section C: ¹H, ¹³C and 2D-NMR Spectrum

nmh-276-p

¹H-NMR of 14

Current Data Parameters
NAME nmh-276-p
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20231119
Time 0.37
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.890 Hz
FIDRES 0.084180 Hz
AQ 5.3064680 sec
RG 101.6
DM 61.000 usec
DE 6.00 usec
TE 294.4 K
D1 2.00000000 sec

==== CHANNEL f1 =====
NUC1 1H
P1 8.30 usec
PL1 -1.00 dB
SFO1 300.1310034 MHz

F2 - Processing parameters
SI 32768
SF 300.1300143 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

ID NMR plot parameters
CX 20.00 cm
CY 6.00 cm
FLP 9.971 ppm
F1 2872.63 Hz
F2P -0.747 ppm
F2 -224.18 Hz
PRGM 0.51850 ppm/cm
HZCN 164.85555 Hz/cm

nmh-276-p

```

Current Data Parameters
NAME nmh-276-p 4
EXPNO 4
PROCNO 1
F2 - Acquisition Parameters
Date_ 20031119
Time 1.46
INSTRUM mv360
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 4
DS 4
SWH 17055.511 Hz
FIDRES 0.27463 Hz
AQ 1.8256000000
RG 4096
DN 27.800 UBB
DE 5.00 UBB
TE 295.3 K
D1 2.0000000000
d11 0.0000000000
d12 0.0000000000
===== CHANNEL f1 =====
NUC1 13C
P1 15.00
PL1 0.00 dB
SFO1 76.4705653 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2 10.00 UBB
PL2 -1.00 dB
PL3 18.00 dB
PL4 18.00 dB
SFO2 300.1312003 MHz
F2 - Processing parameters
SI 32768
SF 75.467450 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
1D NMR plot parameters
CX 20.00 cm
CY 6.68 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPM0 11.0000 ppm/cm
HZ0 14050 Hz/cm

```

nmh-392-p-13C

Current Data Parameters
NAME nmh-392-p-13C
EXPNO 16
PROCNO 1
F2 - Acquisition Parameters
Date_ 2004-05-23
Time 15:40
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 640
DS 4
SWH 17085.511 Hz
FIDRES 0.27463 Hz
AQ 1.024508 sec
RG 27.000 USBC
DE 5.00 USBC
TE 294.7 K
D1 2.00000000 sec
d11 0.00000000 sec
d12 0.00000000 sec
==== CHANNEL f1 =====
NUC1 13C
P1 19.40 USBC
PL1 -1.00 dB
SFO1 76.4705653 MHz
==== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 19.00 USBC
PL2 -1.00 dB
PL3 19.00 dB
PL4 19.00 dB
SFO2 300.1312603 MHz
F2 - Processing parameters
SI 32768
SF 75.4677450 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
F2 - NMR plot parameters
CX 20.00 cm
CY 3.68 cm
FIP 216.000 000
FL 18228.37 Hz
FEP -5.000 000
F2 -377.34 Hz
PPM0 11.00000 ppm/cm
HZ0 14080 Hz/cm

nmh-280-p-13C

0.000

¹³C-NMR of 9

Current Data Parameters
NAME nmh-280-p-13C
EXPNO 16
PROCNO 1
F2 - Acquisition Parameters
Date_ 2003130
Time 2.38
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 4
DS 4
FIDRES 17055.511 Hz
AQ 0.27463 Hz
RG 1.024500 960
FREQ 125.760 MHz
TE 300.2 K
D1 2.00000000 sec
d11 0.00000000 sec
d12 0.00000000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 16.40 uS
PL1 -1.00 dB
SFO1 76.4705653 MHz
===== CHANNEL f2 =====
COPROG2 wa1zg16
NUC2 1H
PCPR2 19.00 uS
PL2 -1.00 dB
PL3 19.00 dB
PL4 19.00 dB
SFO2 300.1312623 MHz
F2 - Processing parameters
SI 32768
SF 75.467513 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
F2 - NMR in 1D parameters
CX 20.00 cm
CY 10.00 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPH0 11.00000 ppm/cm
HZ0 140280 Hz/cm

nmh-292-p

Current Data Parameters
NAME nmh-292-p
EXPNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040126
Time 22.50
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.890 Hz
FIDRES 0.084180 Hz
AQ 5.3064660 sec
RG 408.4
DM 81.000 usec
DE 6.00 usec
TE 295.7 K
D1 2.00000000 sec

==== CHANNEL f1 =====
NUC1 1H
P1 8.30 usec
PL1 -1.00 dB
SFO1 300.1310054 MHz

F2 - Processing parameters
SI 32768
SF 300.1300063 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 6.00 cm
FLP 11.000 ppm
F1 3301.43 Hz
F2 -1.000 ppm
F2 -300.13 Hz
PRGM PRGM
ACQ 0.60000 ppm/cm
FID 180.07800 Hz/cm

0.107
0.000
0.109

7.4500
7.4298
7.4246
7.3672
7.3493
7.3381
7.3257
7.3124
7.2603
7.2522
7.2484
7.2376
7.2299
7.1013
7.0694
7.0769
4.5282
4.5683
4.5774
4.5656
4.5227
4.4626
4.3409
4.3321
4.2606
4.1305
4.1223
3.7691
3.7621

¹H-NMR of 17

nmh-282-p-13C

Current Data Parameters
NAME nmh-282-p-13C
EXPNO 28
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040228
Time 15:51
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 4
DS 4
SWH 17055.511 Hz
FIDRES 0.27463 Hz
AQ 1.0236500 sec
RG 27.000 USBC
DE 5.00 USBC
TE 291.2 K
D1 2.00000000 sec
d11 0.00000000 sec
d12 0.00000000 sec
==== CHANNEL f1 =====
NUC1 13C
P1 15.00 USBC
PL1 -1.00 dB
SFO1 76.4705653 MHz
==== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 19.00 USBC
PL2 -1.00 dB
PL3 19.00 dB
PL4 19.00 dB
SFO2 300.1312603 MHz
F2 - Processing parameters
SI 32768
SF 76.4676512 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
1D NMR plot parameters
CX 20.00 cm
CY 20.00 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPM0 11.00000 ppm/cm
HZ0 146280 Hz/cm

refined mnh-364-p D2O

Current Data Parameters

NAME	proton
EXPNO	32
PROCNO	1

F2 - Acquisition Parameters

Date_	20060620
Time	10.50
INSTRUM	cpd900
PROBHD	5 mm BBO 1H-8B
PULPROG	zgpg30
TD	65536
SOLVENT	D2O
NS	64
DS	4
SWH	6387.934 Hz
FIDRES	0.062213 Hz
AQ	0.0917909 sec
RG	71.8
DM	82.800 uMHD
DE	6.00 uMHD
TE	298.2 K
D1	2.00000000 sec
ACQRES	0.00000000 sec
MCPRG	0.01500000 sec

----- CHANNEL f1 -----

NUC1	1H
PL1	8.00 uMHD
PL2	-1.00 dB
RF1	600.1325000 MHz

F2 - Processing Parameters

SI	32768
SF	600.1325000 MHz
WDW	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

1D NMR plot parameters

CX	20.00 cm
CY	12.50 cm
FLP	7.814 ppm
F1	4888.12 Hz
F2P	2.910 ppm
F2	1746.43 Hz
NUC2	0.2617 ppm/ppm
NUC2N	147.13428 Hz/cm

nmh-384-p-13C

¹³C-NMR of 18

Current Data Parameters
NAME nmh-384-p
EXPNO 13
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040811
Time 5.48
INSTRUM mv360
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 268432
SOLVENT DMS
NS 500
DS 4
SWH 17055.511 Hz
FIDRES 0.27483 Hz
AQ 1.0255500 sec
RG 27.000 USBC
W 5.00 USBC
TE 300.2 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00000000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 15.00 USBC
PL1 -1.00 dB
SFO1 76.4705653 MHz
===== CHANNEL f2 =====
COPROG2 wa1zg16
NUC2 1H
PCPR2 19.00 USBC
PL2 -1.00 dB
PL3 19.00 dB
PL4 19.00 dB
SFO2 300.1312623 MHz
F2 - Processing parameters
SI 32768
SF 76.4677450 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
F2 NMR p101 Derivatives
GX 20.00 cm
EY 2.00 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPM0 11.00000 ppm/cm
HZ0 14050 Hz/cm

ppm

4.0492
4.0341
4.0367
4.0216
3.9855
3.9733
3.9610
3.9571
3.9697
3.9576
3.9503
3.9503
3.9213
3.9106
3.9020
3.5913
3.2376
3.1891
3.1780
3.1700
3.1665
3.1589

mth-364-p 020

4-555-141

¹H-NMR of 23

Current Data Parameters
NAME mhh-355-p
EXPNO 10
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040605
Time 3.07
INSTRUM ei300
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.890 Hz
FIDRES 0.084180 Hz
AQ 5.3064660 sec
RG 2031.2
DM 61.000 usec
DE 6.00 usec
TE 299.7 K
D1 2.00000000 sec
===== CHANNEL f1 =====
NUC1 1H
P1 8.30 usec
PL1 -1.00 dB
SFO1 300.1310034 MHz
F2 - Processing parameters
SI 32768
SF 300.1310032 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00
ID NMR plot parameters
CX 20.00 cm
CY 8.00 cm
FLP 7.863 ppm
F1 2369.86 Hz
F2 -0.227 ppm
F2 -68.18 Hz
PRGM 0.40950 ppm/cm
HZN 122.90493 Hz/cm

Integrat

92-S

nmh-395-p-13C

¹³C-NMR of 23

Current Data Parameters
NAME nmh-395-p
EXPNO 11
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040608
Time 5:30
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 4
DS 4
FIDRES 17855.511 Hz
AQ 0.27463 Hz
RG 1.024600000
IN 27.000 MHz
TE 300.2 K
F2 125.7600000 MHz
SFO1 76.4705653 MHz
SF2 0.0000000 MHz
===== CHANNEL f1 =====
NUC1 13C
P1 14.00 MHz
PL1 -1.00 dB
SFO1 76.4705653 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 00.00 MHz
PL2 -1.00 dB
PL3 18.00 dB
PL4 18.00 dB
SF02 300.1312000 MHz
F2 - Processing parameters
SI 32768
SF 75.467467 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
F2 - NMR unit parameters
CX 20.00 cm
CY 8.68 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPH0 11.00000 ppm/cm
HZ0 14000 Hz/cm

nmh-356-p4

¹H-NMR of 24

Current Data Parameters
NAME nmh-356-p4
EXNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040616
Time 4.10
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.850 Hz
FIDRES 0.084180 Hz
AQ 5.3064880 sec
RG 287.4
DM 81.000 usec
DE 6.00 usec
TE 299.8 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 8.30 usec
PL1 -1.00 dB
SFO1 300.1310534 MHz

F2 - Processing parameters
SI 32768
SF 300.1310534 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 6.00 cm
F1P 8.226 ppm
F1 2460.46 Hz
F2P -0.356 ppm
F2 -108.08 Hz
PRGM 1D
PCPN 0.43123 ppm/cm
PCN 108.10864 Hz/cm

mhh-3556-p4-13C

Current Data Parameters
NAME mhh-355-p4
EXPNO 12
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040616
Time 4.37
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 268360
SOLVENT CDCl3
NS 2048
DS 4
SWH 17085.611 Hz
FIDRES 0.27463 Hz
AQ 1.0245000 sec
RG 27.000 USBC
WV 5.00 USBC
TE 300.2 K
D1 2.00000000 sec
d11 0.00000000 sec
d12 0.00000000 sec
==== CHANNEL f1 =====
NUC1 13C
P1 15.00 USBC
PL1 -1.00 dB
SFO1 76.4705653 MHz
==== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2 10.00 USBC
PL2 -1.00 dB
PL3 18.00 dB
PL4 18.00 dB
SFO2 300.1312603 MHz
F2 - Processing parameters
SI 32768
SF 75.4677467 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
10 NMR plot parameters
CX 20.00 cm
CY 12.50 cm
FIP 216.000 000
FL 18228.37 Hz
FEP -5.000 000
F2 -377.34 Hz
PPM0 11.00000 ppm/cm
HZ0 14080.00 Hz/cm

nmh-471-p D2O

Current Data Parameters

NAME	proton
EXPNO	84
PROCNO	1

F2 - Acquisition Parameters

Date_	20060620
Time	14.45
INSTRUM	gvt000
PROBHD	5 mm BBO 1H-8B
PULPROG	zgpg30
TD	65536
SOLVENT	D2O
NS	16
DS	4
SWH	6387.934 Hz
FIDRES	0.062213 Hz
AQ	0.0917909 sec
RG	71.8
DM	82.800 uMHD
DE	6.00 uMHD
TE	298.0 K
D1	2.00000000 sec
ACQRES	0.00000000 sec
MCPRG	0.01500000 sec

CHANNEL f1

NUC1	1H
PL1	8.00 uMHD
PL2	-1.00 dB
RF01	600.1325000 MHz

F2 - Processing Parameters

SI	32768
SF	600.1325000 MHz
WDW	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

1D NMR plot parameters

EX	20.00 cm
CY	50.00 cm
FLP	7.264 ppm
F1	4383.80 Hz
F2	2.750 ppm
PC	1000.00 Hz
PC	0.22000 ppm/mm
PC	135.14776 Hz/cm

nmh-357-d-cosy

```

Current Data Parameters
NAME      MH357-d-13C
EXPNO     16
PROCNO    1
F2 - Acquisition Parameters
Date_     20040708
Time      3.18
INSTRUM   mv300
PROBHD    5 mm QNP 13C-1
PULPROG   zgpg30
TD        65536
DECR      0.000
SOLVENT   DMSO
NS         1280
DS         4
SWH        17055.511 Hz
FIDRES     0.27463 Hz
AQ         1.824600000
RG          27.800
AQ        5.0000000
TE         300.2 K
D1         2.000000000
d11        0.000000000
d12        0.000000000
===== CHANNEL f1 =====
NUC1       13C
P1         12.00
PL1        -1.00 dB
SFO1       75.476563 MHz
===== CHANNEL f2 =====
COPROG2    zgpg30
NUC2       1H
P2         12.00
PL2        -1.00 dB
PL3        18.00 dB
PL4        18.00 dB
SFO2       300.1312623 MHz
F2 - Processing parameters
SI          32768
SF          75.467450 MHz
WDW          EN
SSB          0
LB          1.00 Hz
GB          0
PC          1.40
=====
1D NMR plot parameters
CX          20.00 cm
CY          1.00 cm
F1          216.000 000
F2          18228.37 Hz
F3          -5.000 000
F4          -377.34 Hz
PPM01      11.00000 ppm/cm
HZ01       1830.14000 Hz/cm

```

NOESY of **1**

nmh-471-p D2O

NAME	Current Data Parameters
EXPNO	1
PROCNO	1
DATA_	EXPNO
TIME	15.42
PROBHD	5 mm BBI H-500
PULPROG	zgpg30
TD	131072
EXHAUST	1
NO	1
NUC1	13C
NUC2	1H
NUC3	1H
NUC4	1H
NUC5	1H
NUC6	1H
NUC7	1H
NUC8	1H
NUC9	1H
NUC10	1H
NUC11	1H
NUC12	1H
NUC13	1H
NUC14	1H
NUC15	1H
NUC16	1H
NUC17	1H
NUC18	1H
NUC19	1H
NUC20	1H
NUC21	1H
NUC22	1H
NUC23	1H
NUC24	1H
NUC25	1H
NUC26	1H
NUC27	1H
NUC28	1H
NUC29	1H
NUC30	1H
NUC31	1H
NUC32	1H
NUC33	1H
NUC34	1H
NUC35	1H
NUC36	1H
NUC37	1H
NUC38	1H
NUC39	1H
NUC40	1H
NUC41	1H
NUC42	1H
NUC43	1H
NUC44	1H
NUC45	1H
NUC46	1H
NUC47	1H
NUC48	1H
NUC49	1H
NUC50	1H
NUC51	1H
NUC52	1H
NUC53	1H
NUC54	1H
NUC55	1H
NUC56	1H
NUC57	1H
NUC58	1H
NUC59	1H
NUC60	1H
NUC61	1H
NUC62	1H
NUC63	1H
NUC64	1H
NUC65	1H
NUC66	1H
NUC67	1H
NUC68	1H
NUC69	1H
NUC70	1H
NUC71	1H
NUC72	1H
NUC73	1H
NUC74	1H
NUC75	1H
NUC76	1H
NUC77	1H
NUC78	1H
NUC79	1H
NUC80	1H
NUC81	1H
NUC82	1H
NUC83	1H
NUC84	1H
NUC85	1H
NUC86	1H
NUC87	1H
NUC88	1H
NUC89	1H
NUC90	1H
NUC91	1H
NUC92	1H
NUC93	1H
NUC94	1H
NUC95	1H
NUC96	1H
NUC97	1H
NUC98	1H
NUC99	1H
NUC100	1H

NOESY of 1

mhh-300-p2

0.0106
0.0000
0.0109

7.4233
7.4197
7.4173
7.4054
7.3932
7.3824
7.3687
7.3424
7.3340
7.3255
7.3134
7.3043
7.2547
7.2317
7.2236
6.9599
6.9309
5.0494
4.6154
4.6755
4.5538
4.5426
4.5133
4.4735
4.3486
4.3375
4.1906
4.1029
4.0940
3.7457
3.7156

ppm

¹H-NMR of 26

mhh-300-p

22.911
2.017
1.985
2.091
4.314
2.119
1.022
2.015
1.000
1.982

Integral

ppm

Current Data Parameters
NAME mhh-300-p2
EXNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040206
Time 1.37
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.890 Hz
FIDRES 0.084180 Hz
AQ 5.3064690 sec
RG 267.4
DM 81.000 usec
DE 6.00 usec
TE 293.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 8.30 usec
PL1 -1.00 dB
SFO1 300.1310034 MHz

F2 - Processing parameters
SI 32768
SF 300.1300081 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 7.00 cm
FLP 8.888 ppm
F1 2986.00 Hz
F2 -1.207 ppm
F2 -362.17 Hz
PRCM 0.55479 ppm/cm
HZCN 165.16873 Hz/cm

mhh-300-p2-13C

Current Data Parameters
 NAME mhh-300-p2
 EXPNO 20
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040206
 Time 1.58
 INSTRUM mv300
 PULPROG zgpg30
 PRCHRG 6 mm WIL 13C-1
 ZGPG30
 FIDRES 0.0003
 SOLVENT CDCl3
 NS 4
 DS 4
 SWH 17055.511 Hz
 FIDRES 0.27435 Hz
 AQ 1.023558 sec
 RG 327.60
 DN 27.800 uBBG
 DE 5.00 uBBG
 TE 293.0 K
 D1 2.0000000 sec
 d11 0.0000000 sec
 d12 0.0000000 sec
 ===== CHANNEL f1 =====
 NUC1 13C
 P1 15.00
 PL1 -1.00 dB
 SFO1 76.470563 MHz
 ===== CHANNEL f2 =====
 CDPRP2 na1216
 NUC2 1H
 PCPR2 50.00 uBBG
 PL2 -1.00 dB
 PL12 18.00 dB
 PL13 18.00 dB
 SFO2 300.1312623 MHz
 F2 - Processing parameters
 SI 32768
 SF 75.467511 MHz
 EN
 NDW 0
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40
 1D NMR plot parameters
 CX 20.00 cm
 CY 5.00 cm
 FIP 216.000 000
 F1 18228.37 Hz
 F2 -5.000 000
 F2 -377.34 Hz
 PPM0 11.0000 ppm/cm
 HZ0 14050 Hz/cm

nmh-321-p-13C

¹³C-NMR of 2

Current Data Parameters
NAME nmh-321-p-13C
EXPNO 16
PROCNO 1
F2 - Acquisition Parameters
Date_ 20040301
Time 13.38
INSTRUM mv300
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 65536
SOLVENT DMS
NS 128
DS 4
SWH 17085.511 Hz
FIDRES 0.27443 Hz
AQ 1.8236000 sec
RG 320
IN 27.000 MHz
TE 293.1 K
DE 5.00 MHz
JE 250.1 K
D1 2.0000000 sec
d11 0.0000000 sec
d12 0.0000000 sec
==== CHANNEL f1 =====
NUC1 13C
P1 15.00
PL1 0.00 dB
SFO1 76.4705653 MHz
==== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2 15.00 MHz
PL2 -1.00 dB
PL3 18.00 dB
PL4 18.00 dB
SFO2 300.1312603 MHz
F2 - Processing parameters
SI 32768
SF 75.467450 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
F2 - NMR p101 parameters
CX 20.00 cm
CY 2.00 cm
FIP 216.000 000
F1 18228.37 Hz
F2 -5.000 000
F2 -377.34 Hz
PPM0 11.00000 ppm/cm
HZ0 14080 Hz/cm

H-H COSY of 2

3.1383
3.1493
3.1576
3.1584

3.5746
3.5859
3.5936
3.6046
3.6343
3.6415
3.6534
3.6596
3.7896
3.8051
3.8376
3.8501
3.8525

3.9875
4.0010
4.0155

ppm

nmh-474-P-B D20 2005 June 1

3.1383
3.1493
3.1576
3.1584

3.5746
3.5936
3.5936
3.6046
3.6343
3.6415
3.6534
3.6596

3.7896
3.8051
3.8376
3.8501
3.8525

ppm

NOESY of 2

nmh-474-p-b D2O 2005 June 1

S-40

NOESY of 2

Section D: References

- (1) Calimente, D. ; Postema , M.H.D. *J.Org.Chem.* **1999**, *64*, 1770-1771.
- (2) (a) Cardona, F.; Faggi, E.; Liguori, F.; Cacciarini, M.; Goti, A. *Tetrahedron Lett.* **2003**, *44*, 2315-2318. (b) Desvergues, S. ; Py, S ; Vallée, Y. *J. Org. Chem.* **2005**, *70*, 1459-1462.
- (3) Yu Y.; Singh, S. K.; Liu, A.; Li,T.-K.; Liu L. F.; LaVoie E. J. *Bioorg. Med. Chem.* **2003**, *11*, 1475-1491.
- (4) Fuji , K.; Ichikawa , K.; Node , M.; Fujita , E. *J. Org. Chem.* **1979** , *44*, 1661-1664.
- (5) Shibano, M. ; Tsukamoto, D. ; Masuda, A. ; Tanaka, Y. ; Kusano, G. *Chem. Pharm. Bull.*, **2001**, *49*, 1362-1365.