

Rare Earth Starting Materials and Methodologies for Synthetic Chemistry

Fabrizio Ortu*



Cite This: *Chem. Rev.* 2022, 122, 6040–6116




Read Online

ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: The number of rare earth (RE) starting materials used in synthesis is staggering, ranging from simple binary metal-halide salts to borohydrides and “designer reagents” such as alkyl and organoaluminate complexes. This review collates the most important starting materials used in RE synthetic chemistry, including essential information on their preparations and uses in modern synthetic methodologies. The review is divided by starting material category and supporting ligands (*i.e.*, metals as synthetic precursors, halides, borohydrides, nitrogen donors, oxygen donors, triflates, and organometallic reagents), and in each section relevant synthetic methodologies and applications are discussed.



CONTENTS

1. Introduction	6040	9.3. Benzyls	6088
1.1. Motivation for the Review	6040	9.4. Organoaluminates	6090
1.2. Scope of the Review	6041	10. Conclusions and Future Perspectives	6095
1.3. Previous Reviews	6041	Author Information	6095
2. General Considerations on the Reactivity of REs	6042	Corresponding Author	6095
2.1. Salt Elimination/Metathesis	6042	Notes	6095
2.2. Metathesis	6043	Biography	6095
2.3. Protonolysis/Protolysis	6043	Acknowledgments	6096
2.4. Insertion/Oxidation	6043	Abbreviations	6096
3. Metals as Synthetic Precursors	6043	References	6096
3.1. Activation in Liquid Ammonia	6043	Note Added after ASAP Publication	6116
3.2. Insertion into C–X Bonds	6046		
3.3. Direct Metalation	6047		
3.4. Redox Transmetalation	6053		
3.5. Metal Vapor Synthesis	6059		
4. Halides	6061		
4.1. Divalent Ln Halides	6061		
4.2. Trivalent RE Halides	6067		
5. Borohydrides	6070		
5.1. Divalent Ln Borohydrides	6071		
5.2. Trivalent RE Borohydrides	6072		
6. Nitrogen Donors	6076		
6.1. Divalent Ln Silylamides	6076		
6.2. Trivalent RE Silylamides	6078		
6.3. Tetravalent Ce Amides	6080		
7. Oxygen Donors	6080		
7.1. Aryloxides	6080		
7.2. Ce(IV) Alkoxides and CAN	6081		
8. Triflates	6084		
9. Organometallic Reagents	6085		
9.1. Simple Hydrocarbyls (Me, ⁿ Bu, Ar)	6086		
9.2. α -Silyl-alkyls	6086		

1. INTRODUCTION

1.1. Motivation for the Review

The organometallic and coordination chemistry of the rare earth (RE) and lanthanoid (Ln) metals (Sc, Y, La, and Ce–Lu) was overshadowed for the best part of the 20th century by the popularity and wide applicability of transition-metal (TM) complexes. This historical disparity finds its roots in how research into the chemistry of REs and TMs developed through the centuries. While the first TM organometallic complexes were identified in the 19th century by pioneers of the ilk of Bunsen, Frankland, and Mond, attempts to stabilize RE–C bonds were largely unsuccessful until the 1950s.¹ Finally, in 1954 Wilkinson

Received: September 30, 2021

Published: January 31, 2022



and Birmingham succeeded in the stabilization of the first RE organometallic complexes, $\text{RE}(\text{Cp})_3$ ($\text{RE} = \text{Sc}, \text{Y}, \text{La}, \text{Ce}, \text{Pr}, \text{Nd}, \text{Sm}, \text{and Gd}$; $\text{Cp} = \{\text{C}_5\text{H}_5\}^-$),² over a century after Frankland's preparation of dialkylzinc species.³ Following Wilkinson and Birmingham's breakthrough, scientists started closing this historical 100 year gap with TM chemistry, and since then, RE organometallic and coordination chemistry has developed from mere curiosity to a burgeoning research field.^{4–9} A key component of these discoveries is the use of ligand architectures that can fulfill the electronic and steric demands of the large and electropositive RE ions. This step-change in RE chemistry was also propelled by the many advances in anaerobic manipulation techniques and ligand design, which in turn led to discoveries that challenged common assumptions and opened unexpected research avenues.^{7,10}

Crucially, these advances have largely been supported by the preparation of adequate starting materials for anaerobic synthesis and the development of innovative synthetic strategies. In particular, the number of RE starting materials is staggering, constituting an immense library that synthetic chemists can access to design their own unique methodologies. All this essential information is spread over 100 years' worth of literature and laboratory work, carried out by a huge number of researchers across the whole world. It is not uncommon for different research teams to devise their own unique methodologies and synthetic strategies, the nuances of which are often passed on as word of mouth—sometimes constituting a religious rite of passage for new starters. This review work provides a systematic account of the most important synthons used in RE synthetic chemistry, covering their preparations and applications as synthetic precursors. Despite the publication of several reviews and book chapters on many of the topics included in this work (*vide infra*, section 1.3), the last comprehensive account on the preparation of RE starting materials and their application in synthesis was provided by Edelman in 1997.⁹ Since then the field has seen an immense amount of progress and the number of new starting materials at the disposal of synthetic RE chemists has increased significantly. The aim is to compile an up-to-date account of the most important synthons and methodologies that constitute the synthetic toolbox for any researcher approaching RE chemistry, thus providing an essential point of reference especially for those at the beginning of their journey into the f-block world.

1.2. Scope of the Review

Throughout the review the terms “rare earth” and “lanthanoid” will be used. The term “rare earth” will be used to describe all group 3 and lanthanoid elements, with the exception of the highly radioactive and rare promethium. The abbreviation “RE” in chemical formulas is used to describe group 3 metals (Sc and Y) and lanthanoids (La–Lu) collectively, whereas the abbreviation Ln will be used whenever a specific reaction or methodology applies exclusively to La–Lu or divalent species.

The aim of this review is to illustrate essential aspects of synthetic methods applied to RE chemistry, focusing primarily on the preparation of starting materials and their most important applications as synthetic precursors. In order to make this work more focused, the starting materials and methods described in this review will primarily relate to anaerobic synthesis. The review is subdivided into starting material categories, and dedicated synthetic methods are discussed within each section: metals as synthetic precursors (section 3), halides (section 4), borohydrides (section 5), amides (section 6), oxygen donors

(section 7), triflates (section 8), and organometallics (section 9). With the exception of section 3, each section contains tables listing selected starting materials for each individual category of this review. In the case of organometallic reagents (section 9), Zimmermann and Anwander published a very comprehensive account in 2010,¹¹ so this review will provide a broad overview of the topic and illustrate the progress achieved over the past decade. This review work covers all relevant literature published up to December 2021.

1.3. Previous Reviews

Over the last 70 years a significant number of reviews have been written on the individual categories covered herein. Selected reviews which are relevant for this work are listed below:

- *Metals as synthetic precursors:*^{12,13}
 - Guo, Z.; Huo, R.; Tan, Y. Q.; Blair, V.; Deacon, G. B.; Junk, P. C. Synthesis of Reactive Rare Earth Complexes by Redox Transmetalation/Protolysis Reactions-A Simple and Convenient Method; 2020 (ref 12)
 - Cloke, F. G. N. Zero Oxidation State Compounds of Scandium, Yttrium, and the Lanthanides; 1993 (ref 13)
- *Halides:*^{14,15}
 - Taylor, M. D. Preparation of Anhydrous Lanthanoid Halides; 1962 (ref 14)
 - Meyer, G. The Rare Earth Elements – The Divalent State in Solid Rare Earth Metal Halides; 2012 (ref 15)
- *Borohydrides:*^{16–21}
 - Marks, T. J.; Kolb, J. R. Covalent Transition Metal, Lanthanide, and Actinide Tetrahydroborate Complexes; 1977 (ref 16)
 - Ephritikhine, M. Synthesis, Structure, and Reactions of Hydride, Borohydride, and Aluminohydride Compounds of the f-Elements; 1997 (ref 17)
 - Makhaev, V. D. Structural and Dynamic Properties of Tetrahydroborate Complexes; 2000 (ref 18)
 - Visseaux, M.; Bonnet, F. Borohydride Complexes of Rare Earths, and Their Applications in Various Organic Transformations; 2011 (ref 19)
 - Guillaume, S. M.; Maron, L.; Roesky, P. W. Catalytic Behavior of Rare-Earth Borohydride Complexes in Polymerization of Polar Monomers; 2014 (ref 20)
 - Paskevicius, M.; Jepsen, L. H.; Schouwink, P.; Černý, R.; Ravnsbæk, D. B.; Filinchuk, Y.; Dornheim, M.; Besenbacher, F.; Jensen, T. R. Metal Borohydrides and Derivatives—Synthesis, Structure and Properties; 2017 (ref 21)
- *Amides:*^{22–24}
 - Anwander, R. Lanthanide Amides; 2005 (ref 22)
 - Lappert, M. F.; Protchenko, A.; Power, P. P.; Seeber, A. Metal Amide Chemistry; 2009 (ref 23)
 - Goodwin, C. A. P.; Mills, D. P. Silylamides: Toward a Half-Century of Stabilizing Remarkable f-Element Chemistry; 2017 (ref 24)
- *Oxygen donors:*^{25–30}
 - Bradley, D. C.; Mehrotra R. C.; Gaurn D. P. Metal Akoxides; 1978 (ref 25)
 - Mehrotra, R. C.; Singh, A.; Tripathi, U. M. Recent Advances in Alkoxo and Aryloxo Chemistry of Scandium, Yttrium and Lanthanides; 1991 (ref 26)

- Bradley, D. C.; Mehrotra, R. C.; Rothwell, I. P.; Singh, A. Alkoxo and Aryloxo Derivatives of Metals; 2001 (ref 27)
- Anwander, R. Routes to Monomeric Lanthanide Alkoxides; 2005 (ref 28)
- Boyle, T. J.; Ottley, L. A. M. Advances in Structurally Characterized Lanthanide Alkoxide, Aryloxo and Silyloxo Compounds; 2008 (ref 29)
- Parmar, V.; Mills, D. P.; Winpenny, R. E. P. Mononuclear Dysprosium Alkoxide and Aryloxo Single-Molecule Magnets; 2021 (ref 30)
- *Triflates*.³¹
 - Lawrance, G. A. Coordinated Trifluoromethanesulfonate and Fluorosulfate; 1986 (ref 31)
- *Organometallics*.^{4,5,11,32–38}
 - Cotton, F. A. Alkyls and Aryls of Transition Metals; 1955 (ref 4)
 - Davidson, P. J.; Lappert, M. F.; Pearce, R. Metal σ -Hydrocarbyls, MR_n . Stoichiometry, Structures, Stabilities, and Thermal Decomposition Pathways; 1976 (ref 32)
 - Bochkarev, M. N.; Kalinina, G. S.; Bochkarev, L. N. Advances in the Chemistry of Organolanthanides; 1985 (ref 33)
 - Schumann, H.; Meese-Marktscheffel, J. A.; Esser, L. Synthesis, Structure, and Reactivity of Organometallic π -Complexes of the Rare Earths in the Oxidation State Ln^{3+} with Aromatic Ligands; 1995 (ref 34)
 - Cotton, S. A. Aspects of the Lanthanide-Carbon σ -Bond; 1997 (ref 5)
 - Deacon, G. B.; Forsyth, C. M.; Nickel, S. Bis-(pentafluorophenyl)mercury—a versatile synthon in organo-, organooxo-, and organoimido-lanthanoid chemistry; 2002 (ref 35)
 - Edelmann, F. T.; Freckmann, D. M. M.; Schumann, H. Synthesis and Structural Chemistry of Non-Cyclopentadienyl Organolanthanide Complexes; 2002 (ref 36)
 - Zimmermann, M.; Anwander, R. Homoleptic Rare-Earth Metal Complexes Containing $Ln-C$ σ -Bonds; 2010 (ref 11)
 - Lyubov, D.; Trifonov, A. A. $Ln(II)$ Alkyl Complexes: From Elusive Exotics to Catalytic Applications; 2021 (ref 37)
 - Izod, K. Alkyl, Carbonyl and Cyanide Complexes of the Group 3 Metals and Lanthanides; 2021 (ref 38)

A significant number of reviews have also been published on general synthetic methods in RE chemistry. Some of the most relevant for this work are listed below:^{6,8,40–43}

- Evans, W. J. The Organometallic Chemistry of the Lanthanide Elements in Low Oxidation States; 1987 (ref 39)
- Edelmann, F. T. Scandium, Yttrium, and the Lanthanide and Actinide Elements, Excluding their Zero Oxidation State Complexes; 1995 (ref 6)
- Edelmann, F. T. Lanthanides and Actinides; 1997 (ref 9)
- Anwander, R. Principles in Organolanthanide Chemistry; 1999 (ref 8)
- Anwander, R. Herrmann, W. A. Features of Organolanthanide Complexes; 2005 (ref 40)


- Liddle, S. T. Lanthanides: Organometallic Chemistry Fundamental Properties; 2012 (ref 41)
- Nicholas, H. M.; Mills, D. P. Lanthanides: Divalent Organometallic Chemistry; 2017 (ref 42)
- Ortu, F.; Mills, D. P. Low Coordinate Rare Earth and Actinide Complexes; 2019 (ref 7)
- Layfield, R. A. Lanthanides; 2021 (ref 43)

2. GENERAL CONSIDERATIONS ON THE REACTIVITY OF RES

Unlike TMs, RE metals are not involved in classic two-electron reactions like oxidative addition and reductive elimination. Therefore, synthetic strategies are mostly limited to a handful of key reactions: (1) *salt elimination*, (2) *metathesis*, (3) *protonolysis* (also referred to as *protolysis*), and (4) *insertion/oxidation*.^{9,44} This is not an exhaustive list, as other more specialized methodologies have been also implemented in RE chemistry and will be discussed in more detail in the various sections of this review.

2.1. Salt Elimination/Metathesis

In salt elimination/metathesis reactions, a RE salt precursor, LnX_2 or REX_3 ($X = \text{halide, borohydride, triflate, etc.}$), is reacted with a ligand transfer reagent $M(L)_n$ ($n = 1, 2$ depending on the oxidation state of the metal and formal charge of the ligand), producing a new RE complex and eliminating an inorganic salt (eq 1). Classic ligand transfer reagents are group 1 and group 2 salts of a variety of bases, such as hydrocarbyls, amides, alkoxides, and aryloxides. Additionally, the methodology entails the elimination of highly insoluble inorganic salts with very high lattice energies, thus providing a strong driving force for the reaction. For these reasons, this is one of the most popular methodologies used in RE synthetic chemistry and can be applied to the synthesis of both homoleptic and heteroleptic species, though the occurrence of salt occlusion can prevent the isolation of clean products.^{8,41}



The availability of different halide sources offers several possibilities in terms of lattice energy and solubility of the byproducts of salt elimination reactions. Estimates of lattice energies (eq 2) can be obtained from simple electrostatic considerations:

$$U \propto \frac{n \cdot |z^+| |z^-|}{r^+ + r^-} \quad (2)$$

where the lattice energy, U , is proportional to the number of ions (n) and their charges (z^+ and z^-) and inversely proportional to their ionic radii (r^+ and r^-), *i.e.*, the distance between the two charges.⁴⁵ Smaller ions will give the highest lattice energy values (LiF , 1034 kJ mol⁻¹), while larger ions will have the smallest lattice enthalpies (CsI , 613 kJ mol⁻¹) (Table 1).⁴⁶ Additionally, the relative strength of the RE–X bonds should also be taken into account: RE^{3+} ions are hard Lewis acids, and the bond strength with halides decreases descending the group. One additional consideration is the solubility of the alkali metal salts. Often salt elimination reactions have to be carried out in ethereal solvents because of the poor solubility of starting materials and reagents. However, alkali metal salt can have some solubility in certain organic solvents, which can have a detrimental effect on the outcome of salt elimination protocols and make purification procedures extremely challenging (Table 1).


Table 1. Lattice Energies (U)⁴⁶ and Solubilities of Selected Alkali Metal Halides in THF at 25 °C^{47–49}

	U (kJ mol ⁻¹)	solubility (mg/mL)
LiF	1034	–
LiCl	864	49.5
LiBr	820	388
LiI	764	552 ^a
NaCl	790	0.20
NaBr	754	0.15
NaI	705	29.98
KCl	720	0.30
KBr	691	0.06–0.1
KI	650	0.001

^aSolubility in diethyl ether.⁵⁰

2.2. Metathesis

Metathesis reactions produce a ligand exchange between two metals (eq 3), based on differences in their affinities for different donors. For example, this is a method that can be employed for the preparation of RE organometallics complexes (*i.e.*, transmetalation), such as the metathesis reaction between aryloxy complexes and Li{CH(SiMe₃)₂} to give [RE{CH(SiMe₃)₂}₃].⁵¹ This method is particularly useful for the preparation of homoleptic complexes, as the occurrence of ligand scrambling often prevents the isolation of heteroleptic species.



2.3. Protonolysis/Protolysis


RE starting materials can perform acid/base reactions with protic substrates, accompanied by the highly favorable elimination of volatile gases or liquids (eq 4). Such reactions are classically driven by differences in pK_a (Table 2), though

Table 2. pK_a Values of Selected Acids in H₂O^{7,11,52}

acid	pK_a
CH ₄	48
Ph ₃ CH	31.5
C ₅ H ₂ Me ₄	26.1 ^a
HN(SiMe ₃) ₂	25.8
HN(SiHMe ₂) ₂	22.6
C ₅ H ₆	15
HOC ₆ H ₃ Me _{3-2,4,6}	10.87
HOC ₆ H ₆	9.97
HN(Me)(C ₆ H ₅)	4.85
H ₂ NC ₆ H ₅	4.58

^aValue in DMSO.


other important factors can come into play (*e.g.*, saturation of metal coordination sphere or use of chelating ligands). However, simple acid/base considerations are often a good indication when planning a methodology that makes use of these reactions.



2.4. Insertion/Oxidation

RE metals can sometimes react directly with substrates, either by inserting into a polar bond (*e.g.*, a carbon–halogen bond, eq 5) or by reducing the target donor (eq 6). The former method has been used very rarely, and it mainly applies to divalent REs that are able to behave like *pseudo*-alkaline earth (AE) metals (section 3.2). Oxidation reactions instead are more versatile,

though their main applications are limited to Sm, Eu, and Yb; these are usually performed on protic substrates and accompanied by the elimination of hydrogen (sections 3.1 and 3.3). Both insertion and oxidation reactions can be sluggish with RE metals, and better results are often achieved when metals have been activated. Oxidation reactions have also been combined with transmetalation reactions by using redox transmetalating reagents, such as organomercurials (section 3.4).



3. METALS AS SYNTHETIC PRECURSORS

The use of REs in their metallic form as primary starting materials for synthesis has been investigated for over a century. For example, high-temperature reaction of RE powders with dry HCl or iodine is an established methodology for the preparation of solvent-free RECl₃, and most REs react readily with iodine or 1,2-diiodoethane in ethereal solvents to afford di- and trihalide species (*vide infra*, section 4.1). REs can also react directly with organic substrates (section 3.3), though often activation strategies have to be employed which circumvent the relative inertia of the metal. Examples of this approach are (1) activation in liquid ammonia (section 3.1), (2) activation with Hg or HgCl₂ (section 3.3), (3) redox transmetalation (section 3.4), and (4) metal vapor synthesis (section 3.5). Additionally, in some cases REs can also react directly with substrates in a fashion similar to the insertion of Mg into C–X bonds (section 3.2).

3.1. Activation in Liquid Ammonia

At the turn of the 20th century, the use of liquid ammonia as an alternative nonaqueous solvent sparked the interest of the scientific community,⁵³ leading to the development of new reaction apparatuses and experimental techniques which set criteria still used today for experimental work with ammoniacal solutions (Figure 1).^{54,55} Pioneering work by Joannis,⁵³ Mentrel,⁵⁶ Roederer,⁵⁷ and Moissan^{58–60} focused on the solubilization of alkali and AE metals and the properties of resulting solutions. Kraus further developed this new research field by studying various s-block metals in liquid ammonia and first proposed the presence of solvated electrons in such media.^{61–63} Around the same time, Cottrell focused his efforts on the dissolution of Mg and Ca and their reactivity with acetylene.⁶⁴ The first ammoniacal solutions of any f-element were first reported by Watt *et al.* in 1950, who described the use of liquid ammonia on various U and Th salts (*e.g.*, iodates, oxalates, peroxides).⁶⁵ Warf and Korst decided to extend this study to the classic divalent Lns, owing to their resemblance to the heavy AE elements.⁶⁶ The heavy AEs share many similarities with classic divalent Lns Eu, Sm, and Yb, such as the stability of the +2 oxidation state and ionic radii (Ca²⁺, 1.00 Å; Yb²⁺, 1.02 Å; Sm²⁺, 1.17 Å; Eu²⁺, 1.22 Å; Sr²⁺, 1.18 Å).⁶⁷ Because of this their coordination chemistry has often been developed in parallel.⁶⁸ By using freshly distilled ammonia, Warf and Korst were able to dissolve Eu and Yb at –78 °C, while no reaction was observed with Sm (Scheme 1 and Figure 1).

The nature of the cation–electron ion pair was further corroborated by EPR studies on Eu ammoniacal solutions performed by Catterall and Symons in 1964;^{69,70} additionally, the electronic spectra of Eu and Yb species were recorded a year





Figure 1. Illustration of basic setup for the preparation of ammoniacal solutions of divalent Lns. Alternatively, modification of Schlenk line manifolds can be implemented to optimize ammonia transfer and workup procedures.⁵⁵

Scheme 1. Synthesis of Ammonia Solution of Divalent Lns



later by Waugh and co-workers.⁷¹ Interestingly, Peer and Lagowski were able to dissolve various other Lns (Sm, Tb, Er, and Tm) in ammonia *via* co-condensation and recorded the electronic spectra of the resulting solutions; both Sm and Tb were identified to be in their divalent state, while definitive conclusions could not be drawn for Er and Tm.⁷²


Eu and Yb ammonia solutions are relatively stable when stored under anaerobic conditions. However, they decompose over time to generate the parent amide Ln(NH₂)₂, similar to what is observed with alkali and AE metals (Scheme 2).⁷³ In the case of Yb, depending on reaction conditions, degradation of ammoniacal solutions can also produce oxidized trivalent amide Yb(NH₂)₃.⁷³ Divalent Eu(NH₂)₂ is obtained also under ammonothermal reaction conditions (up to 5000 atm of NH₃), while for Yb the same reaction conditions generate pure trivalent Yb amide or the salt Na[Yb(NH₂)₄].⁷⁴ The use of ammonothermal conditions with Lns has limited synthetic utility and has attracted interest mostly for the preparation of new semiconductors;⁷⁵ also, supercritical ammonia (160 °C) has been used to prepare metal sulfide salts of Yb, [Yb(NH₃)₈]-[M(S₄)₂]-NH₃ (M = Cu, Ag), and La, [La(NH₃)₉][Cu(S₄)₂].⁷⁶ Additionally, Müller-Buschbaum and Quitmann reported the molecular structure of complex [Sm(NH₃)₉][Sm(Pyr)₆] (Pyr = pyrrolide, {C₄H₄N⁻}), obtained from the direct reaction of

pyrrole with Sm metal and pyrrole under solvothermal conditions, followed by treatment with liquid ammonia.⁷⁷

Recently, Kraus and co-workers investigated the preparation of Eu(II), Yb(II), and Ho(III) azides by reacting the pure metal with AgN₃ in liquid ammonia, leading to the structural identification of amino-adducts [Ho₂(μ-NH₂)₃(NH₃)₁₀]- (N₃)₃·(NH₃)_{1.25} and [Yb(NH₃)₈](N₃)₂.^{78,79}


The ready availability of soluble forms of divalent Lns gives access to the preparation of simple binary species. Such was the observation of Salot and Warf in 1968, who postulated the formation of YbI₂ from the reaction of Yb metal and NH₄I in liquid NH₃.⁸⁰ Shortly after, Howell and Pytlewski followed a similar synthetic protocol and reported the preparation of six unsolvated LnX₂ (Ln = Eu, Yb; X = Cl, Br, I) species (Scheme 3);⁸¹ this methodology has been used extensively by synthetic




Scheme 3. Synthesis of Eu and Yb Divalent Halides in Liquid NH₃⁸¹



chemists since then (*vide infra* for preparation and uses of LnX₂ precursors in synthesis, section 4.1).⁸² When Sm is employed, this methodology leads to the formation of trivalent species.⁸⁰

Scheme 2. Decomposition of Ammoniacal Solutions of Eu and Yb and Formation of Divalent and Trivalent Amides Eu(NH₂)₂, Yb(NH₂)₂, and Yb(NH₂)₃⁷³




Scheme 4. Synthesis of Divalent Ln Organometallic Complexes in Liquid NH₃^{83–87}Scheme 5. Reduction of 2,2'-Bipyridine and Phenantroline with Ln Ammoniacal Solutions^{88,89}Scheme 6. Synthesis of Ln(II) Aryloxy Complexes 9 and 10 from Ln Ammoniacal Solutions^{90,92}

Owing to the highly reducing nature of these ammoniacal Ln solutions, their synthetic utility was first explored in the 1960s. Fischer and Fischer reacted freshly distilled cyclopentadiene with an ammoniacal solution of europium or ytterbium, followed by sublimation to yield analytically pure metallocene Ln(Cp)₂ (**1-Ln**; Ln = Eu, Yb; Cp = {C₅H₅}⁻) (Scheme 4).⁸³ Similarly, Wayda *et al.* were able to synthesize their Cp* (Cp* = {C₅Me₅}⁻) analogues Ln(Cp*)₂(NH₃)_x (**2**), though only the ammoniate metallocene derivative [Yb(Cp*)₂(NH₃)(THF)] (**3**) was structurally authenticated (Scheme 4).⁸⁴ A similar approach has also been used for the preparation of other organometallic derivatives, such as Ln(COT) (COT = {C₈H₈}⁻)⁸⁵ and Eu(II) propynide Eu(C≡CCH₃)₂ (**4**).⁸⁶ Wayda *et al.* followed the original Ln(COT) synthesis developed by Thomas and Hayes⁸⁵ and were able to crystallize the Yb derivative by slow diffusion of pentane in a pyridine solution,

thus obtaining the piano-stool complex [Yb(COT)(py)₃] (5-3py, py = pyridine).⁸⁷ Around the same time of Fischer and Fischer's metallocene synthesis, Ln ammoniacal solutions were also used to for the reduction of 2,2'-bipyridine and phenantroline (Scheme 5), leading to the isolation of Ln(bipy)₄ (**6-Ln**; Ln = Eu, Yb; bipy = 2,2'-bipyridine) and Yb(phen)₄ (**7**, phen = phenantroline);⁸⁸ Pappalardo also reported the preparation of Yb(bipy)₃ (**8**) using a similar methodology.⁸⁹

Smith and co-workers also used Yb ammoniacal solutions to prepare the bis-aryloxy Yb complex [Yb(ODbmp)₂(THF)₃] (**9**, Scheme 6) (ODbmp = OC₆H₂tBu₂-2,6-Me-4),⁹⁰ which was originally prepared *via* protonolysis or redox transmetalation (*vide infra*, sections 3.4 and 7.1).⁹¹ Similarly, Evans *et al.* isolated the Eu²⁺ complex [Eu(ODb)₂(NCMe)₄] (**10**, ODb = OC₆H₃tBu₂-2,6) by dissolving a Eu ingot and 2,6-di-*tert*-butylphenol in liquid ammonia, followed by extractions with

Scheme 7. Reactivity of Yb and Eu with *N*-Heterocycles and Amines Using Ammoniacal Solutions or Ammonia-Activated Metals^{77,93–98,100}



acetonitrile (Scheme 6).⁹² However, when 2,6-diisopropylphenol (HODipp) is employed, the hydroxo-bridged Eu(II) complex $[\text{Eu}_4(\text{ODipp})_2(\mu\text{-ODipp})_4(\mu_3\text{-OH})_2(\text{NCMe})_6]$ is obtained; attempts by Evans *et al.* to use this methodology to obtain a hydroxo-free Eu(II) derivative were not successful.⁹²

This synthetic approach has been revisited more recently by Müller-Buschbaum and co-workers, who have used similar methodologies for the activation of a variety of nitrogen-containing heterocycles, *e.g.*, indole (11), carbazole (12), pyrazole (13), and pyrrole (14) (Scheme 7).^{77,93–98} These reactions in some cases are accompanied by concomitant formation of $\{\text{NH}_2\}^-$ from ammonolysis.⁹⁵ In their work, Müller-Buschbaum and co-workers carried out reactions in the absence of solvents and under solvothermal conditions, though in some cases low-temperature reactivity was also achieved, such as with the synthesis of $[\text{Yb}(\text{NH}_3)_8][\text{Yb}(\text{pyr})_6]$ (14c, pyr = pyrrolide, $\{\text{C}_4\text{H}_4\text{N}\}^-$).⁹⁷ This approach is particularly interesting for modern RE synthetic chemistry, as it can provide an access route to very challenging complexes avoiding the use of coordinating solvents.⁹⁸ In the case of the reaction between Yb and carbazole ($\text{C}_{12}\text{H}_8\text{NH}$, CarbH), *N*-phenylpiperazine (Phpip) was added to aid crystallization of the target complex $[\text{Yb}(\text{Carb})_2(\text{NH}_3)_4]$ (12, Carb = carbazolate, $\{\text{C}_{12}\text{H}_8\text{N}\}^-$), though this product cannot be separated from cocrystallized CarbH·Phpip.⁹⁴ It is also noteworthy that Müller-Buschbaum and co-workers extended this work to various divalent and trivalent REs by reacting metal powders with amines, either directly or in the presence of trace amounts of Hg under


solvothermal conditions (*vide infra*, sections 3.3 and 3.4).^{77,99} Remarkably, the use of ammoniacal Yb solutions in the presence of 2,2'-dipyridylamine afforded the isolation of the first RE molecular nitride $[\text{Yb}_3\text{N}(\text{dpa})_6][\text{Yb}(\text{dpa})_3]$ (15, dpa = 2,2'-dipyridylamide, $\{(\text{C}_6\text{H}_4\text{N})_2\text{N}\}^-$).¹⁰⁰

3.2. Insertion into C–X Bonds

Eu, Sm, and Yb resemble the chemistry of Sr and Ca respectively, owing in particular to the stability of the divalent state and ionic radii (see section 3.1). Organocalcium “heavy Grignard” reagents, $(\text{R})\text{CaX}$ (R = alkyl, aryl; X = halide), were originally investigated in the early 20th century as potential alternatives to classic organomagnesium analogues;^{101–103} additionally, several structurally authenticated examples have been reported by Westerhausen and co-workers over the last two decades,^{104,105} and recently this chemistry has also been extended to Sr and Ba.¹⁰⁶ Consequently, classic divalent Lns (Eu, Sm and Yb) have also been investigated for the preparation of *pseudo*-Grignard reagents, *i.e.* $(\text{R})\text{LnX}$ (R = alkyl, aryl; X = halide).^{107,108} Unlike for Mg, heavy group 2 Grignard reagents readily undergo Schlenk-type degradation affording dialkyl and dihalide species;^{104,105} therefore, owing to the similarities of classic divalent Lns with the heavy AE metals, this type of degradation is also a likely occurrence for a putative $(\text{R})\text{LnX}$ *pseudo*-Grignard derivative (Scheme 8).

D. F. Evans *et al.* first reported the reaction of Yb and Eu metal with alkyl and aryl iodides in THF at -20°C (Scheme 9).^{107,108} In the case of Yb, magnetic susceptibility measurements revealed

Scheme 8. Insertion Reaction of Ln Metal into C–X Bonds and Schlenk-Type Equilibrium



the presence of small quantities of Yb^{3+} ion (Yb^{3+} : $[\text{Xe}]4f^{13}$, $\mu_{\text{eff}} = 4.5 \mu_{\text{B}}$), with Yb^{2+} as the predominant ion (Yb^{2+} : $[\text{Xe}]4f^{14}$, $\mu_{\text{eff}} = 0 \mu_{\text{B}}$);¹⁰⁸ in the case of europium, the magnetic moment of (Ph)EuI was measured at $7.5 \mu_{\text{B}}$, close to the value for free Eu^{2+} (Eu^{2+} : $[\text{Xe}]4f^7$, $\mu_{\text{eff}} = 7.5 \mu_{\text{B}}$; Eu^{3+} : $[\text{Xe}]4f^6$, $\mu_{\text{eff}} = 1.5 \mu_{\text{B}}$).¹⁰⁸ In the case of Sm metal, reactivity with iodobenzene and iodoethane is sluggish, requiring higher temperatures compared to Eu and Yb and affording a much higher proportion of trivalent species.¹⁰⁸ Moreover, further reactivity tests with Gd and Er showed no reactivity, while La and Ce are converted into mixtures of trivalent species, *i.e.*, (R)REI₂, (R)₂REI, RE(R)₃, and REI₃.¹⁰⁸ Beletskaya and co-workers further investigated oxidative addition of Ce, Eu, Sm, and Yb with iodothiophenes and bromopentafluorobenzene in the presence of an initiator, *e.g.*, diiodoethane or dibromoethane (Scheme 9).¹⁰⁹ The authors reported the formation of organoderivatives (C₄H₃S)LnI (Ln = Ce, Eu, Sm and Yb) and (C₆F₅)LnBr (Ln = Eu, Sm and Yb), though their formation was confirmed only *via* reactivity with Ph₃SnCl and detection of corresponding Ph₃SnAr derivative (Ar = C₄H₃S, C₆F₅).¹⁰⁹ Ce has a very unstable divalent state ($\text{Ce}^{3+}/\text{Ce}^{2+}$: E = −3.2 V);^{110,111} therefore, it cannot be excluded that reactivity with iodothiophene produced trivalent (C₄H₃S)CeI₂ or (C₄H₃S)₂CeI rather than divalent (C₄H₃S)₂CeI, as both trivalent species can in turn generate Ph₃Sn(C₄H₃S) upon reaction with Ph₃SnCl.

(Ph)LnI (Ln = Eu, Yb) performs well as protonolysis reagent toward 3,5-diphenylpyrazole (HP₂pz) to give [Ln(Ph₂pz)(I)(THF)₄] (16-Ln, Ln = Eu, Yb; Ph₂pz = {3,5-Ph₂C₃HN₂}[−]; Scheme 10);¹¹² the same outcome was obtained also with freshly prepared (Me)LnI.¹¹³ Ali *et al.* further exploited the utility of (Ph)LnI by performing protonolysis reactions with a family of amidines, ArFormH (ArNCHNAr; Ar = C₆H₃Me₂-2,6 – Xyl, C₆H₂Me₃-2,4,6 – Mes, C₆H₃Pr₂-2,6 – Dipp), leading to the formation of divalent Ln complexes [Eu(DippForm)(I)(THF)₄] (17), [Eu(XylForm)(I)(DME)₂] (18), [Eu(XylForm)(I)(κ¹-DME)(μ₂:κ²O-DME)]_∞ (19), [Yb(DippForm)(I)(THF)₃] (20), [{Yb(MesForm)(μ-I)(THF)₂}₂] (21) and [{Yb(XylForm)(μ-I)(THF)₂}₂] (22) (Scheme 10); however, the authors also observed the formation of several trivalent derivatives during the isolation of these


species.¹¹³ To gain a better understating of the chemical behavior of these reagents, Junk and co-workers revisited the synthesis of putative (Ph)YbI first described by D. F. Evans and proposed the occurrence of Schlenk-type equilibria and disproportionation reactions (Scheme 10).^{112,113} The authors attempted to identify the components of these degradations and initially isolated [Yb(I)₂(THF)₄] and trivalent [Yb(Ph)₃(THF)₃]. Additionally, when phenyl iodide is reacted with Yb in DME, fractional crystallization affords [Yb(I)₂(DME)₃] and the mixed-valent separated ion pair complex [Yb(DME)₄][Yb(Ph)₄(DME)₂].¹¹² Subsequently, Junk and co-workers ascribed the formation of the trivalent Yb formamidinate species [Yb(DippForm)(I)₂(THF)₃] (23) and [Yb(XylForm)₂(I)(DME)] (24), together with the hydroxo-bridged Eu complex [{Eu(XylForm)(I)(μ-OH)(THF)₂}₂] (25), to the presence of other trivalent Yb derivatives, *e.g.*, (Ph)YbI₂ and (Ph)₂YbI (Scheme 11).¹¹³




The first structurally authenticated Ln *pseudo*-Grignard complexes [{Yb[C(SiMe₃)₃](μ-I)(OEt₂)₂] (26) and [{Yb[C(SiMe₂CH = CH₂)₃](μ-I)(OEt₂)₂] (27) were reported by Smith in 1994¹¹⁴ and 1996,¹¹⁵ obtained from the direct reaction of Yb powder with (SiMe₂R)₃Cl (R = Me, CH = CH₂) in diethyl ether (Scheme 12). Moreover, Niemeyer and Heckmann isolated aryl complex [Yb(C₆H₃Ph₂-2,6)(I)(THF)₃] (28) from the reaction between 2,6-Ph₂C₆H₃I and Yb powder, while use of Eu powder led to the isolation of bis-aryl complex [Eu(C₆H₃Ph₂-2,6)₂(THF)₂] (29) (Scheme 12).¹¹⁶ Such a difference in reactivity between Yb and Eu is an indication of the presence of a Schlenk-type equilibrium, which is more pronounced for the larger Eu²⁺ cation. Confirmation of this observation came also from the isolation of Eu₂(THF)₂ as a byproduct of the reaction. Conversely, treatment of Yb powder with neopentyl iodide, ^tBuCH₂I, leads to further oxidation to the trivalent Yb complex [Yb(CH₂^tBu)₃(THF)₂] (30) with concomitant formation of solvated YbI₂ (Scheme 12).¹¹⁷ Niemeyer proposed that the divalent *pseudo*-Grignard species (^tBuCH₂)YbI is initially formed, which then reacts with a further equivalent of ^tBuCH₂I to yield (^tBuCH₂)₂YbI₂; the latter species undergoes Schlenk-type equilibrium yielding 30 and YbI₃. Finally, YbI₃ reacts with excess Yb metal to form YbI₂.¹¹⁷

3.3. Direct Metalation


RE metals can react directly with protic substrates, thus providing a very efficient synthetic route which is accompanied by the evolution of hydrogen as a byproduct (Scheme 13).¹¹⁸ However, this is a relatively rare synthetic approach with limited scope, especially compared to other widely used synthetic

Scheme 9. Synthesis of Ln *pseudo*-Grignard Derivatives by D. F. Evans,^{107,108} Beletskaya,¹⁰⁹ and Co-workers




Scheme 10. Synthesis of Ln Formamidinate Complexes via Protonolysis Using (Ph)LnI^{112,113}Scheme 11. Schlenk-Type Equilibrium and Degradation of (Ph)₂YbI Proposed by Junk and Co-workers¹¹³Scheme 12. Synthesis of First Structurally Authenticated Ln *pseudo*-Grignard Species by Smith (26 and 27)^{114,115} and Niemeyer (28),¹¹⁶ Together with Synthesis of Diaryl Eu Complex 29¹¹⁶ and Trivalent Yb Derivative 30¹¹⁷

Scheme 13. Direct Reaction of RE Metals with Protic Substrates




methodologies such as salt metathesis, transamination, or transmetalation reactions; additionally, in some cases harsh reaction conditions and additives are required.

Evans and Pires de Matos reported independently the reactivity of Ln metals with alcohols.^{119–121} The reaction of Eu metal with 2-methoxyethanol proceeds smoothly at room temperature and affords polymeric [Eu(OCH₂CH₂OMe)₂]_n in good yields, which Evans and co-workers employed as protonolysis reagent toward substituted phenols HOC₆H₄Me_{2,6} (HOXyl) and HOC₆H₃ⁱPr_{2,6} (HODipp), affording the heteroleptic aryloxide complexes [Eu(μ₃:η²-

Scheme 14. Direct Reaction of Eu and Yb with Alcohols and Application in Protonolysis Reactions^{119–121}

$\text{OCH}_2\text{CH}_2\text{OMe}$) (η^2 - $\text{OCH}_2\text{CH}_2\text{OMe}$)($\text{OC}_6\text{H}_3\text{R}_2-2,6$)- $[\text{H}^+]_4$] (**31a**, $\text{R} = \text{Me}$; **31b**, $\text{R} = i\text{Pr}$) (Scheme 14).¹¹⁹ Pires de Matos and co-workers obtained $\text{Eu}(\text{OMe})_2$, $\text{Eu}(\text{OEt})_2$, $\text{Eu}(\text{O}^i\text{Pr})_2$, and $\text{Yb}(\text{OMe})_3$ by treating metal powders with methanol, ethanol, or isopropanol (Scheme 14);¹²¹ in the case of the reaction between Yb and isopropanol, activation with ammonia was required and the only isolated product was the cluster $[\text{Yb}_5\text{O}(\text{O}^i\text{Pr})_{13}]$ (**32**), previously prepared by Bradley and co-workers by using catalytic amounts of HgCl_2 in the reaction (*vide infra*, Scheme 16).¹²² In the same report, Carretas *et al.* illustrated the use of metal vapor synthesis (MVS) techniques for the preparation of the same alkoxides and reported an improvement of the overall yields in all cases.¹²¹ Moreover, Evans and Greci investigated the use of different solvents to facilitate the direct reaction of Eu metal with HOXyl and HODipp. When HOXyl was reacted with Eu ingots in *N*-methylimidazole ($\text{C}_3\text{H}_3\text{N}_2\text{-Me}$), the bridged aryloxide complex $[\text{Eu}_2(\text{OXyl})(\mu\text{-OXyl})_3(\text{C}_3\text{H}_3\text{N}_2\text{-Me})_5]$ (**33**) was obtained (Scheme 14); similarly, treatment of HODipp with Eu ingots in acetonitrile yielded the aryloxide complex $[\text{Eu}_2(\text{ODipp})_2(\mu\text{-ODipp})_2(\text{MeCN})_4(\mu\text{-NCMe})]$ (**34**).¹²⁰


Direct metalation reactions can also be facilitated by using small quantities of Hg or HgCl_2 . The proposed mechanism involves the amalgamation of metallic mercury with the RE metals, which are readily oxidized to their trivalent or divalent state upon reaction with protic substrates, accompanied by formation of hydrogen (Scheme 15).¹²³ When HgCl_2 is employed, metallic mercury is likely formed from the direct reaction with the substrate (usually amines or alcohols), leading to the formation of $\text{Hg}(\text{L})_2$ with the concomitant production of HCl. The transient Hg amide or alkoxide then reacts with the RE metal powder *via* redox transmetalation (*vide infra*), thus

Scheme 15. Direct Reaction between RE Metals and Protic Substrates with Catalytic Amounts of Hg or HgCl_2 


producing the desired RE derivative, $\text{RE}(\text{L})_3$ or $\text{Ln}(\text{L})_2$, and regenerating metallic Hg (Scheme 15).

Mazdiyasi and co-workers were the first to report the direct activation of isopropanol with RE metals in the presence of HgCl_2 (occasionally combined with $\text{Hg}(\text{OAc})_2$ or HgI_2) under reflux, and they discovered that the use of excess HgCl_2 led to the presence of RE chlorides in the final products.^{124,125} Deacon *et al.* subsequently demonstrated that solvated RECl_3 can indeed be prepared conveniently from metal powders and HgCl_2 (*vide infra*, section 4.2).¹²⁶ In Mazdiyasi's original reports, the authors identified the formation of the expected isopropoxide species $\text{RE}(\text{O}^i\text{Pr})_3$ for all metals but Eu, which formed the divalent $\text{Eu}(\text{O}^i\text{Pr})_2$ instead.^{124,125} These results were later revisited by Hubert-Pfalzgraf, Caulton, and Bradley, who studied the products of these direct reactions *via* single-crystal XRD studies and identified the formation of the oxo-bridged

Scheme 16. Examples of Activation with HgCl_2 and Hg: Formation of oxo-bridged clusters $[\text{RE}_5\text{O}(\text{O}^i\text{Pr})_{13}]$ (**32-RE**; RE = Y, Pr, Yb) via Direct Reaction between RE Metals and Isopropanol,^{122,124,125,127,128} and Synthesis of *bis*-Phospholyls **33-Ln**,¹²⁹ *bis*-Aryloxy **34-Ln**,¹³⁰ *ansa*-Metalloocene Complexes **35-Ln**,¹³¹ and *bis*-Gallyl Complex **36-Ln**.¹³²





Scheme 17. Synthesis of Aryloxides **37** and **38** via Direct Activation of HODpp Using RE(Hg) Amalgam^{134,135}



pentametallic clusters $[\text{RE}_5\text{O}(\text{O}^i\text{Pr})_{13}]$ (**32-RE**; RE = Y, Pr, Yb) as the main products, rather than the expected homoleptic isopropoxide species (Scheme 16).^{122,127,128} Nief and Mathey

used a similar method to synthesize $[\text{Ln}(\eta^5\text{-C}_4\text{H}_2\text{PPh}_2\text{-2,S})_2(\text{THF})_2]$ (**33-Ln**; Ln = Sm, Yb) and *bis*-aryloxy $[\text{Ln}(\eta^5\text{-C}_4\text{H}_2\text{AsPh}_2\text{-2,S})_2(\text{THF})_2]$ (**34-Ln**; Ln = Sm, Yb), obtained


Scheme 18. Synthesis of Heterobimetallic RE and AE Complexes Using Mercury Amalgam and Fluxes under Solvothermal Conditions¹⁴⁰Scheme 19. Examples of RE Pyrazolates Synthesized by Deacon and Co-workers Using Mercury Amalgam under Solvothermal Conditions^{141,143}

from the reaction of metal powders with *bis*-phospholyl or *bis*-arsolyl dimers in the presence of catalytic amounts of HgCl_2 (Scheme 16).^{129,130} Additionally, Edelman and Recknagel showed that this strategy can be used in the reaction of dimethylfulvene with Ln metal powders (Ln = Sm, Yb), generating *ansa*-metallocene complexes of formula $[\text{Ln}\{(\text{C}_5\text{H}_4)_2\text{C}_2\text{Me}_4\}(\text{THF})_2]$ (35-Ln) (Scheme 16).¹³¹ Finally, Roesky and co-workers synthesized the gallyl lanthanide complexes $[\text{Ln}\{\text{Ga}(\text{Dipp-Bian})\}_2(\text{THF})_4]$ (36-Ln; Ln = Sm, Eu, Yb; Dipp-Bian = 1,2-bis[(2,6-diisopropylphenyl)imino]-acenaphthene) from the reduction of dimeric precursor $[\{\text{Ga}(\text{Dipp-Bian})\}_2]$ with Ln powders activated with Hg.¹³²

This synthetic strategy was revisited by Deacon, Junk, and Müller-Buschbaum, who studied the activation of a variety of protic substrates such as substituted phenols and heterocyclic amines.^{98,133} In this new methodology, proligands are usually solids and the reactions are carried out without solvents and in the presence of Hg or HgCl_2 , operating under solvothermal conditions, *i.e.*, heating at temperatures higher than the melting point of the proligand. These types of reactions are usually carried out in partially evacuated sealed vessels (*e.g.*, Carius tubes or sealed glass ampules).^{98,133} Deacon and co-workers focused

their efforts on the use of sterically demanding aryloxides and substituted pyrazolates, comprising all of the RE metals in their methodologies. Because of the absence of coordinating solvents in these reactions, homoleptic species are usually obtained with various degrees of nuclearity depending on the ligands' steric features, ionic radii of the RE metals employed, reaction times, and crystal packing interactions in the solid state. When Eu or Yb are reacted with $\text{HOC}_6\text{H}_4\text{Ph}_2$ -2,6 (HODpp) directly, very little conversion is observed even after prolonged reaction time and under forceful conditions; when the same reactions are carried out in the presence of Hg (Scheme 17), the desired $\text{Ln}(\text{ODpp})_2$ species are generated, which can then be recrystallized from toluene to give dimeric $[\{\text{Eu}(\text{ODpp})\}(\mu\text{-ODpp})_3\text{Eu}]$ (37-Eu) and $[\{\text{Yb}(\text{ODpp})\}_2(\mu\text{-ODpp})_2]$ (37-Yb) (Scheme 18).¹³⁴ Similarly, when classic trivalent REs are employed, the expected tris-aryloxo complexes, $[\text{RE}(\text{ODpp})_3]$ (38-RE; RE = Y, La, Ce, Pr, Nd, Gd, Ho, Er, Lu), are obtained (Scheme 17); also in the case of RE^{3+} ions, direct reactivity in the absence of Hg leads to very little conversion into the desired products.¹²³ Deacon and Junk successfully extended this methodology to other REs (including Sc)¹³⁵ and aryloxides comprising different substitution patterns of the aryl groups,^{136–138} including buttressing

Scheme 20. Synthesis of RE Amides *via* Solvothermal Synthesis with Mercury Amalgam by Müller-Buschbaum and Co-workers^{146–148}



substituents in 3- and 5-positions.¹³⁵ Additionally, reactions can also be carried out in the presence of fluxes (e.g., 1,3,5-tri-*tert*-butylbenzene and 1,2,4,5-tetramethylbenzene) which increase the contact between metal and alcohol. This methodology has proven to be particularly useful for the preparation of heterobimetallic compounds and can be employed also for the synthesis of bimetallic systems incorporating alkali¹³⁹ and AE metals (39 and 40, Scheme 18).¹⁴⁰ One of the great advantages of these methodologies is that resulting complexes can spontaneously crystallize from the reaction mixtures in the Carius tubes upon cooling, together with avoiding the preparation of tailored starting materials and using coordinating solvents. However, in some cases recrystallizations have to be carried out, and owing to the insolubility of these species in nonpolar solvent media, these usually have to be performed at very high temperatures (e.g., toluene at 190 °C).¹⁴⁰


Reactivity of REs (Sc, La, Nd, Eu, Sm, Yb, and Lu) with bis-*tert*-butylpyrazole (^tBu₂pzH) in the presence of Hg and under solvothermal conditions delivers monomeric, [RE(^tBu₂pz)₃] (41-RE; RE = Sc, Sm), and dimeric species, [RE₂(^tBu₂pz)₆] (42-RE; RE = Nd, Sm, Lu).¹⁴¹ Similarly, 41-Sc can also be obtained *via* direct synthesis in the presence of mercury metal at temperatures between 270 and 300 °C.¹⁴² When divalent Eu and Yb are employed, the methodology produces slightly different outcomes, leading for example to the isolation of the mixed-valent Yb(II)/(III) complex [Yb₂(^tBu₂pz)₅] (43, Scheme 19).¹⁴³ Interestingly, these methodologies are affected by subtle variations in reaction conditions. For example, in the case of Eu heating of the reaction with ^tBu₂pzH at 220 °C for 24 h affords the monomeric complex [Eu(^tBu₂pz)₂] (though no structural validation has been provided for this conformation), while after heating for 15.5 h the tetrameric species [Eu₄(^tBu₂pz)₈] (44) is obtained instead (Scheme 19).¹⁴¹ Similar to the use of substituted phenols (*vide supra*, Scheme 17), this is a very effective methodology for the synthesis of homoleptic solvent-free RE complexes and has also been extended to other pyrazoles

featuring varying degrees of substitution.¹⁴⁴ Kempe and Deacon have also reported the direct reaction of Yb metal with substituted aminopyridines, leading to the formation of homoleptic divalent and trivalent Yb complexes depending on the ligand employed.¹⁴⁵

Müller-Buschbaum and co-workers extended this approach to other heterocyclic *N*-donors including 1*H*-1,2,3-benzotriazolo-[4,5-*b*]pyridine (45-RE, RE = La, Sm; 46-RE, RE = Y, Tb),¹⁴⁶ benzimidazole (47),¹⁴⁷ and unsubstituted pyrazole (48 and 49, Scheme 20).¹⁴⁸ All the reactions were performed in evacuated reaction vessels and under solvothermal conditions “in the melt”, with varying reaction temperatures depending on the substrates. The methods employed by Müller-Buschbaum are perfectly suited for the preparation of homoleptic derivatives with several REs, though in some cases proligands are also included in the resulting complexes, either trapped in the lattice through solid-state packing interactions or by coordinating directly to the metal centers (Scheme 20).

Another activation strategy consists of the use of iodine, which is a methodology commonly employed for the activation of Mg in the preparation of Grignard reagents;¹⁴⁹ REs can also react directly with I₂ to give REI₂ and REI₃ salts (*vide infra*, sections 4.1 and 4.2). Small amounts of iodine have been used by Junk, Deacon, and co-workers to activate a range of REs (Y, La, Nd, Sm, Eu, Dy, and Yb) and react them with variously substituted phenols and pyrazoles.^{150–153} The exact nature of the mechanism involved in this methodology has not been fully identified; however, Junk and Deacon invoked the formation of highly reactive divalent REI₂ salts, obtained from the comproportionation reaction between REI₃ and RE.¹⁵⁰ The transient divalent species is oxidized by protic proligand substrates, followed by Schlenk-type rearrangement to give homoleptic RE(L)₃ (L = alkoxide, pyrazolate) complexes together with the regeneration of REI₃ (Scheme 21).

Junk and Deacon have reported the reaction of Yb and Tb metal with variously substituted pyrazoles, C₃H₂N₂(CF₃)-1-

Scheme 21. Activation of REs with Iodine or REI₃ and Reaction with Protic Substrates


(C₄H₃S)-3 (HTfpz) and C₃H₂N₂-Ph-1-(C₄H₃S)-3 (HPhtpz) (Scheme 22).^{151,152} The former affords THF-solvated complex [Tb(Tfpz)₃(THF)₃] (**50**) when reacted with Tb filings and few crystals of iodine (*ca.* 8%), though the product is isolated in rather low yields. In the case of the reaction of HPhtpz with Yb, the intermediate [Yb(Phtpz)(I)(THF)₄] (**51**) is first isolated, which upon treatment with DME affords the *bis*-pyrazolate complex [Yb(Phtpz)₂(DME)₂] (**52**). Interestingly, this behavior is not observed in the reactions involving substituted phenols, where homoleptic species of the type [RE(OAr)₃(Slv)_{*n*}] (**53^{Ar}**-RE; OAr = ODipp, OMes, ODpp, ODbmp; Slv = THF, DME, diglyme, MeCN; *n* = 1–3)^{150,153} are usually obtained even with Yb.¹⁵⁰ Very recently, Junk, Deacon, and co-workers have also used a similar methodology for the preparation of homoleptic Ln(II) and heteroleptic RE(III) formamidinates.^{154,155} Bochkarev and co-workers have also shown that small amounts of preformed REI₃ (*ca.* 5 mol %) can be used in conjunction with metallic REs to give very similar results in the reaction with 1-phenyl-3-methyl-4-isobutyryl-5-pyrazolone (PMIP), to give dimeric complexes [{RE(PMIP)(μ-PMIP)₂}₂] (**54-RE**; RE = Y, Nd, Gd, Tb, Er, Tm, Lu) (Scheme 23).¹⁵⁶ Mashima *et al.* have also employed a similar methodology for the synthesis of RE(III) cyclooctatetraenyl complexes; in their synthetic strategy, RE metals are reacted with cyclooctatetraene in the presence of equimolar amounts of iodine, 1,2-dibromoethane, or Ph₃PCl₂ in hot THF, affording heteroleptic COT complexes with the formula [RE(COT)(I)(THF)_{*n*}] (RE = La, Ce, Pr, *n* = 3; RE = Nd, *n* = 2; Ln = Sm, *n* = 1), [{Sm(COT)(μ-X)(THF)₂}₂] (X = Cl, Br).¹⁵⁷

In some cases, Ln and RE metals can react directly without the aid of an activating agent, though this is a rare occurrence. For example, Müller-Buschbaum and Quitmann showed that carbazole reacts directly with Yb under solvothermal conditions (255 °C, carbazole mp 246 °C, Scheme 24) after a quick initiation at 280 °C, to form the homoleptic coordination polymer [Yb(Carb)₂]_∞ (**55**).¹⁵⁸ The outcome of this reaction is analogous to that obtained with Yb ammoniacal solutions of Yb, though requiring harsher conditions and long reaction times (*ca.*

48 days).⁹⁴ Anwander and co-workers have also shown that 3,4-dimethylpyrazole reacts with La powder at high temperatures (220 °C) under partial vacuum (Scheme 24), leading to the formation of coordination polymer [La(Me₂pz)₃]_∞ (**56**).¹⁵⁹ Similarly, some REs are also able to react directly with pyridylbenzimidazoles under solvothermal conditions and in the absence of Hg.¹⁴⁷ Additionally, in 2005 Junk and co-workers reported the direct reaction of La, Eu, and Yb with 2,6-dibenzylphenol (HODbp) carried out at 170 °C (Scheme 24).¹³⁶ The reactions with La and Eu proceed smoothly: in the case of La, the trivalent bimetallic complex [{La(ODbp)₂(μ-ODbp)₂}]₂ (**57**, ODbp = {O–C₆H₃(CH₂Ph)₂-2,6}⁻) is isolated, whereas the divalent species [{Eu(ODbp)(μ-ODbp)₂}]₂ (**58**) is obtained when Eu metal is employed.¹³⁶ Eu can also react directly with *N,N'*-bis(aryl)formamidines (ArFormH, Ar = Dipp, DF; DF = C₆H₃F₂-2,6) in acetonitrile to give complexes of formula [Eu(ArForm)₂(CH₃CN)₂] (**59**, Ar = Dipp; **60**, Ar = DF), while Yb requires the presence of catalytic amounts of Hg to obtain analogous complexes (Scheme 24).¹⁶⁰ Nief and Mathey also showed that Sm and Yb powders can react directly with biphospholyls and biarsolyls in THF at room temperature (Scheme 24), breaking the E–E (E = P, As) bond and forming *bis*-phospholyl and arsolyl complexes [Ln(η⁵-C₄Me₄P)₂(THF)₂] (**61-Ln**) and [Ln(η⁵-C₄Me₄As)₂(THF)₂] (**62-Ln**, Ln = Sm, Yb).^{129,130}

3.4. Redox Transmetalation

Direct reactivity of REs with proligands often requires prior activation of the metal or some other activation strategy (see sections 3.1, 3.2 and 3.3). However, there are some examples of direct reactivity in which the RE metal does not require activation and is involved in a redox exchange with another metal-containing species. Such species are typically transmetalating reagents, with organomercurials as the most popular choices (*e.g.*, HgPh₂, Hg(C₆F₅)₂, and Hg(CPh)₂);^{12,35,161–165} however, these methodologies have also been extended to include other redox active metals, such as Sn,^{90,166,167} Tl,^{91,168–174} Bi,^{175,176} and Ag.¹⁷⁷ The first reaction of an RE with an organomercurial reagent was reported in 1945 by Gilman and Jones, who reacted La metal with HgPh₂ though they were not able to identify the reactivity products.¹⁷⁸ Broadly speaking, these methodologies can be divided into two categories: (1) redox transmetalation (RT) reactions and (2) redox transmetalation combined with ligand exchange, also termed redox transmetalation protonolysis or protolysis (RTP) (Scheme 25).¹² Though initially developed for classic divalent Lns (Yb and Eu), RT and RTP reactions can also be used for the synthesis of trivalent RE derivatives.¹² RT/RTP reactions have been pioneered by Deacon and co-workers since the 1970s, who developed the first applications in RE coordination and

Scheme 22. Reactivity of Substituted Pyrazoles with Ln Metals (Ln = Yb, Tb) in the Presence of Iodine Reported by Deacon and Junk^{151,152}







Scheme 23. Reactivity of Iodine-Activated RE Metals with Alcohols by Junk, Deacon^{150,153} and Bochkarev¹⁵⁶Scheme 24. Direct Reactivity of RE Metals with Protic Substrates, *bis*-Phospholyls and *bis*-Aryloxy and in the Absence of Activators^{94,129,136,158-160}

Figure 2. Sketch of the apparatus originally developed by Deacon and co-workers for RT and RTP reactions with Hg and Tl reagents. *Note:* All joints are kept grease-free by using PTFE sleeves. Reproduced with permission from ref 179. Copyright 1990 Wiley.

organometallic chemistry and also engineered bespoke reaction apparatuses (Figure 2) for facile removal of hazardous byproducts (*i.e.*, Hg and Tl).¹⁷⁹ Their implementation in synthetic protocols has given access to complexes supported by a wide array of ligands such as cyclopentadienyl,^{168,169,172,173,180} carboranes,¹⁸¹ monodentate amides,^{90,182-185} bidentate amides,^{160,186-191} pyrazolates,^{166,174,175} aryloxides,^{91,170,171,183,192} thiolates,^{193,194} and *N*-heterocyclic carbenes (NHCs).^{195,196} These methodologies have been covered extensively in other review articles;^{12,35} therefore, this section will focus on key examples that best demonstrate their applications in RE chemistry.

Scheme 25. RT and RTP Reactions of RE Metals with Hg, Tl, Sn, Bi, and Ag Reagents





Scheme 26. RT Reactions of Organomercurial Reagents with Divalent and Trivalent Lns^{161–165,193,194}Scheme 27. RTP Reactions of RE Metals and Organomercurials with Substituted Phenols^{91,112,123,135,153,183,198,199}

addition of small quantities of Hg, I₂, and REI₃). Deacon pioneered the use of Hg reagents in RT and RTP reactions and reported the preparation of several organoytterbium and organoeuropium complexes, Ln(R)₂ (Ln = Eu, Yb; **64-Ln**, R = CCPh; **65-Ln**, CC^tBu; **66-Ln**, Ph; **67-Ln** C₆F₅; **68-Ln** o-HC₆F₄; **68-Ln** p-HC₆F₄), by reacting metal powder with the corresponding organomercurial reagent in ethereal solvents (Scheme 26).^{161–165} While Yb reacts smoothly with organomercurials, in the case of Eu, addition of Hg metal is necessary for the reaction to take place. Interestingly, reactivity of Sm powder with Hg(R)₂ reagents is less straightforward; when Hg(C₆F₅)₂ is employed, Sm(C₆F₅)₂ or Sm(C₆F₅)₃ cannot be isolated and various decomposition products are obtained instead, e.g., SmF₂/SmF₃, Sm(C₆F₅)(F)₂, and fluorohydrocarbons,^{164,197} and similar issues are encountered when using other trivalent Lns. Bochkarev and co-workers were the first to report the synthesis of [Er(Ph)₃(THF)₃] (**69-Er**) and [Tm(Ph)₃(THF)₃] (**69-Tm**) from the reaction of metal powders and HgPh₂ in the presence of small quantities corresponding triiodides (ErI₃ and TmI₃), though the isolation of such species is extremely challenging.¹⁷⁶ Despite the difficulties in isolating

putative RE(R)₃ (R = CCPh, Ph, C₆F₅) via RT, reactivity *in situ* of trivalent REs with organomercurials is used extensively as a first step in RTP reactions with various protic substrates (*vide infra*). Nonetheless, RT reactions with fluorinated thiols proceed smoothly to give fluorinated thiolate complexes [Ln(S-C₆F₅)₃(Slv)_x] (**70a-Ln**; Ln = Ho, Er, Yb; Slv = THF, DME, C₅H₅N; x = 2–3) and [{Ln(S-C₆F₅)₂(μ-S-C₆F₅)-(THF)_x}₂] (**70b-Ln**; Ln = Ce, Sm; x = 1, 2), which can also be obtained using RTP methodologies involving Hg(C₆F₅)₂ (Scheme 26).^{193,194}

Deacon and co-workers have shown that Hg-mediated RTP reactions are extremely effective for the preparation of RE(II) and RE(III) complexes with various aryloxy (OAr) ligands (Scheme 27).^{91,112,123,135,153,183,198,199} These reactions are typically carried out at room temperature with relative short reaction times, and the resulting complexes are usually crystallized as THF adducts, [RE(OAr)_n(THF)_m] (**71^{Ar}-RE**, n = 2, RE = Sm, Eu, Yb; **72^{Ar}-RE**, n = 3, RE = Y, La, Pr, Nd, Sm, Eu, Gd, Dy, Er, Yb, Lu; m = 0–3), though in some cases solvent-free monomeric complexes can also be isolated, e.g. [Nd(ODpp)₃] (**72^{Dpp}-Nd**, Scheme 27).²⁰⁰ The outcome of these RTP



Scheme 28. Examples of RTP Methodologies Involving Organomercurials and Cyclopentadienyl Ligands^{180,201–203}Scheme 29. Synthesis of Pyrazolate^{174,206–208} and Amidinate^{160,186,188–191,211} RE Complexes via RTP Reactions with Organomercurials

reactions can be affected by the nature of the RE involved, the choice of organomercurial reagents, and aryloxy ligand. In the case of Yb and Eu, the corresponding divalent complexes **71^{Ar}-RE** are usually obtained, which can be crystallized as THF adducts.²⁰⁰ Interestingly, the use of different $\text{Hg}(\text{R})_2$ reagents can selectively switch oxidation of Yb metal between Yb(II) and Yb(III); this was demonstrated by Deacon *et al.* in the reaction with 3,5-substituted 2,6-diphenylphenols: reactivity between Yb powder, 2,6-diphenyl-3,5-dimethylphenol and HgPh_2 leads to the Yb(II) complex **71^{Ar}-Yb**, whereas when $\text{Hg}(\text{C}_6\text{F}_5)_2$ is used the trivalent complex **72^{Ar}-RE** is isolated instead.¹³⁵ In some cases isolating pure divalent or trivalent complexes can be problematic, like in the case of RTP reactions with $\text{Hg}(\text{C}_6\text{F}_5)_2$ and the phenol $\text{HOC}_6\text{H}_2-2,6\text{-}^i\text{Bu}-4\text{-OMe}$ (HOAr^{OMe}). When Yb is employed, the divalent complex $[\text{Yb}(\text{OAr}^{\text{OMe}})_2(\text{THF})_3]$ (**71^{OMe}-RE**) is obtained in very good yields, while in the case of Sm the corresponding divalent and trivalent complexes $[\text{Sm}(\text{OAr}^{\text{OMe}})_2(\text{THF})_3]$ (**71^{OMe}-RE**) and $[\text{Sm}(\text{OAr}^{\text{OMe}})_3(\text{THF})_3]$ (**72^{OMe}-RE**) are obtained, with the latter as the major product.¹⁹⁹ RE-fluoride species can also be obtained because of the decomposition of RE-(C_6F_5) intermediates with certain supporting aryloxides.¹⁹⁹ To solve this problem, Deacon and co-workers replaced $\text{Hg}(\text{C}_6\text{F}_5)_2$ with $\text{Hg}(\text{CCPh})_2$ in their methodology and obtained the target homoleptic species **71^{Ar}-RE** or **72^{Ar}-RE**.¹⁹⁹

Some of the earliest applications of RTP methodologies were the preparation of Cp derivatives, comprising both divalent $\text{Ln}(\text{Cp})_2$ and trivalent $\text{Ln}(\text{Cp})_3$ complexes (Scheme 28).¹⁹⁷ These methodologies have also been extended to substituted Cp ligands, such as $\{\text{C}_5\text{H}_4\text{Me}\}^-$ (Cp^{Me}),¹⁹⁷ Cp^* ,¹⁸⁰ and $\{\text{C}_5\text{H}_4\text{PPh}_2\}^-$ (Cp^{PPh_2}).¹⁷⁰ Deacon and co-workers found that when $\text{Yb}(\text{C}_6\text{F}_5)_2$ was used as protonolysis reagent with HCp^{Me} the reaction afforded an explosive solid, likely because of the formation of unstable fluorinated species.¹⁹⁷ Therefore, the use of fluorine-free organomercurials is often preferable for these methodologies. RTP reactions are also capable of producing homoleptic $\text{Ln}(\text{Cp})_2$ complexes with extremely bulky ligands,



such as $\{\text{C}_5\text{HPh}_4\}^-$ (Cp^{Ph_4}) and $\{\text{C}_5\text{Ph}_5\}^-$ (Cp^{Ph_5}), e.g. $[\text{Ln}(\text{Cp}^{\text{Ph}_5})_2]$ (**73-Ln**; RE = Sm, Eu, Yb),^{201,202} $[\text{Yb}(\text{Cp}^{\text{Ph}_4})_2(\text{THF})_2]$.^{202,203} This is a particularly remarkable application as classic metathetical reactivity between alkali metal Cp salts and RE halides is often not suitable to obtain Cp derivatives with high steric congestions. Interestingly, Jaroschik, Deacon, and co-workers occasionally used also heteroleptic organomercurials, $\text{Hg}(\text{Ph})(\text{C}_6\text{F}_5)$ and $\text{Hg}(\text{Ph})(\text{CCPh})$, for some of these reactions.^{201,204,205}

Another successful application of RTP reactions with REs and organomercurials involves the use of mono- and multidentate N-donors. A large number of protic substrates have been employed, with silylamines, arylamines, formamidines, and pyrroles among the most popular.¹² This is a particularly important use of RTP methodologies owing to the prominent role played by N-donors in RE coordination chemistry.^{7,24} Also in this case, the main products of these reactions are homoleptic bis-amide complexes, both with mono- and multidentate donors. Deacon and co-workers have applied this methodology to several substituted pyrazolates and showed that this approach is applicable to both divalent Lns (Eu and Yb) and trivalent REs (Scheme 29).^{174,206–208} The choice of mercurial reagent can be crucial for the outcome of these reactions. When HgPh_2 is reacted with Yb and proligand in a 1:2 stoichiometry, the target bis-pyrazolates $\text{Yb}(\text{R}_2\text{pz})_2$ (**74^R-Yb**) are usually obtained.¹⁷⁴ However, when $\text{Hg}(\text{C}_6\text{F}_5)_2$ is employed, the reaction leads to oxidation to Yb(III) and formation of $\text{Yb}(\text{R}_2\text{pz})_3$ (**75^R-Yb**);^{174,209,210} it is noteworthy that short reaction times tend to favor the formation of **74^R-Yb** over **75^R-Yb**.³⁵ Interestingly, direct protonolysis between $\text{Yb}(\text{C}_6\text{F}_5)_2$ and Ph_2pzH proceeds smoothly to give $\text{Yb}(\text{Ph}_2\text{pz})_2$ (**75^{Ph}_2}-Yb**).¹⁷⁴ The use of RTP reactions is exemplified by the extensive work done by Junk, Deacon, and co-workers in the synthesis of RE formamidinate (ArForm ; Ar = Dipp, *p*-Tol, *o*-(CF_3) C_6H_4 , DF, 2,3,4,5- $\text{F}_4\text{C}_6\text{H}$ -TF, *o*- FC_6H_4 -F) complexes (Scheme 29).^{160,186,188–191,211} For these ligand systems, reactivity conditions and stoichiometries can be tailored to obtain either divalent or trivalent

Scheme 30. RTP and Direct RT Reactions with Sterically Demanding Amides^{182,212,213}Scheme 31. Synthesis of Cp,^{168,169,172,173,215} Pyrazolate,^{174,206} and Aryloxo,^{91,170,198} Ln Complexes via RT Reactions with Tl(I) Reagents

formamidate complexes, “RE(ArForm)₂” (76^{Ar}-RE) and “RE(ArForm)₃” (77^{Ar}-RE), irrespective of the choice of organomercurial agent, although $\text{Hg}(\text{C}_6\text{F}_5)_2$ is usually the oxidant of choice. It should be noted that a large excess of metal filings is used for Yb to afford clean conversion to divalent 76^{Ar}-Yb . Reactions are usually carried out in THF, and the resulting complexes are normally isolated as ethereal adducts (e.g., THF, DME, diglyme) upon recrystallization, with varying coordinated solvents depending on steric congestion around the metal center and differences in ionic radii. As previously discussed, fluorinated mercurial reagents can lead to the formation of undesired byproducts through fluoride abstraction and oxidation of the RE metal. Junk and co-workers have been able to take advantage of this type of reactivity to selectively synthesize heteroleptic complexes of general formula RE(ArForm)₂(X) (X = F, Br), such as $[\text{La}(\text{DippForm})_2(\text{F})(\text{THF})]^{211}$ and $[\text{RE}(\text{DippForm})_2(\text{Br})(\text{THF})]$ (RE = La, Nd, Yb),¹⁸⁸ the latter obtained by employing $\text{Hg}(2\text{-BrC}_5\text{F}_4)_2$.

Deacon and co-workers demonstrated that RTP methodologies can be of great utility for the stabilization of complexes with sterically demanding amides. Amines, $\text{HN}(\text{SiMe}_3)_2$, and $\text{HN}(\text{SiMe}_3)(\text{Dipp})$ react smoothly with Sm, Eu, and Yb in THF in the presence of HgPh_2 , to give Ln(II) bis-amide complexes $[\text{Ln}\{\text{N}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$ ($78\text{-Ln}\cdot 2\text{THF}$) and $[\text{Ln}\{\text{N}(\text{SiMe}_3)(\text{Dipp})\}_2(\text{THF})_2]$ (79-Ln).²¹² Deacon, Jones, and co-workers have also shown that RTP methodologies can afford complexes with very low coordination numbers, reporting the synthesis of 3-coordinate complex $[\text{Yb}\{\text{N}(\text{Dipp})(\text{Mes})\}_2(\text{THF})]$ (80) from Yb metal, HgPh_2 , and $\text{HN}(\text{Dipp})(\text{Mes})$ (Scheme 30).¹⁸² Niemeyer and Hauber reacted Eu and Yb powders with triazene $\text{HN}_3(\text{Dmp})(\text{Tph})$ ($\text{Dmp} = \text{C}_6\text{H}_3\text{Me}_2-2,6$; $\text{Tph} = \text{C}_6\text{H}_4\text{Tripp}-2$; $\text{Tripp} = \text{C}_6\text{H}_2\text{Pr}_3-2,4,6$) and $\text{Hg}(\text{C}_6\text{F}_5)_2$, obtaining the heteroleptic 3-coordinate complexes $[\text{Ln}\{\text{N}_3(\text{Dmp})(\text{Tph})\}(\text{C}_6\text{F}_5)]$ (81-Ln ; Ln = Eu, Yb).²¹³ Additionally, Jones and co-workers showed that heteroleptic organomercurials of the type “Hg(L)(X)” (L = ligand, X = halide) can be employed as direct

Scheme 32. RT Reactions with Ag(I)-NHC Reagents by Roesky and Co-workers^{195,196}Scheme 33. Use of Sn(II) and Sn(IV) Reagents in RT Reactions^{90,166,167}

RT reagents (Scheme 30).¹⁸² Reaction of $\text{Hg}(\text{L}^\dagger)(\text{I})$ ($\text{L}^\dagger = \text{N}(\text{Ar})(\text{SiMe}_3)$; $\text{Ar} = \text{C}_6\text{H}_2\text{Pr}\{\text{C}(\text{H})\text{Ph}_2\}_2\text{-4,2,6}$) with metal powders yields two different results: with Yb, the homoleptic 2-coordinate complex $[\text{Yb}(\text{L}^\dagger)_2]$ (**82**) is obtained, with concomitant formation of $[\text{Yb}(\text{I})_2(\text{THF})_2]$; in the case of Eu, the halide-bridged dimer $[\{\text{Eu}(\text{L}^\dagger)(\mu\text{-I})(\text{THF})_2\}_2]$ (**83**) is obtained instead. Analogous reactions with Sm and Tm were also attempted, but without success.

The use of Tl reagent in RT methodologies is also very well-established. Tl^+ is not as oxidizing as Hg^{2+} ($E^{\text{Tl}^+/\text{Tl}} = -0.34$ V; $E^{\text{Hg}^{2+}/\text{Hg}} = 0.85$ V),²¹⁴ thus allowing for more control of the reactivity particularly in the case of divalent Lns. Additionally, suitable transmetalation reagents “Tl(L)” are easily accessible by reacting $\text{Tl}(\text{OEt})$ with protic substrates such as cyclopentadienes,^{168,169,172,173,215} pyrazoles,^{174,206} and phenols.^{91,170,198} Methodologies are similar to those used for organomercurial RT and RTP reactions, requiring usually ethereal solvents (THF, DME), though pyridine can also be used in some cases.¹⁷⁹ Additionally, small aliquots of mercury can also be added to these reactions to aid reactivity.¹⁷⁹ The first compounds to be synthesized with these Tl(I) transmetalation reagents were Cp complexes, $\text{Ln}(\text{Cp}^{\text{R}})_2$ (**1-Ln**, $\text{Cp}^{\text{R}} = \text{Cp}$; **84-Ln**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{Me}}$; **85-Ln**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{PPh}_2}$; RE = Eu, Yb) and $\text{Ln}(\text{Cp}^{\text{R}})_3$ (**86-Ln**, $\text{Cp}^{\text{R}} = \text{Cp}$; **87-Ln**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{Me}}$; RE = Ce, Nd, Sm, Gd, Er, Yb) (Scheme 31).^{168,169,172,173,215} Despite the toxicity of Tl, Tl(I) reagents offer some advantages with respect

to organomercurials, owing to their higher chemical and thermal stability and tolerance toward a wider variety of solvents (e.g., pyridine and MeCN). It is noteworthy that the synthesis of divalent derivatives with Yb, **1-Yb** and **84-Yb**, often requires large excess of metal powder to avoid formation of trivalent **86-Yb** or **87-Yb**; however, in the case of Sm, only trivalent complexes **86-Sm** and **87-Sm** are obtained even in the presence of excess metal.^{169,215} Deacon and co-workers monitored the reaction between Yb powder and $\text{Tl}(\text{Cp})$ by IR spectroscopy, which revealed that the reaction proceeds first with the formation of **84-Yb** followed by reduction with the excess Yb metal to yield divalent **1-Yb**.¹⁷⁹ Deacon^{170,198} and Lappert⁹¹ reported also the application of Tl(I) RT reactions to the synthesis of sterically encumbered aryloxides, $[\text{Yb}(\text{OAr})_2(\text{THF})_3]$ (**53^{Ar}-Yb**) and $\text{Yb}(\text{OAr})_3$ (**88-Yb**) ($\text{Ar} = \text{C}_6\text{H}_3\text{tBu}_2\text{-2,6}$, $\text{C}_6\text{H}_2\text{tBu}_2\text{-2,6-Me-4}$, $\text{C}_6\text{H}_2\text{tBu}_3\text{-2,4,6}$) (Scheme 31).⁹¹ Similar to what was reported for the synthesis of Cp complexes, a large excess of Yb is required to access divalent derivatives. In addition, pyrazolate complexes, $\text{RE}(\text{R}_2\text{pz})_2$ (**74^R-Ln**, R = Me, Ph), are readily accessible for Eu and Yb, while reaction of Sm powder with $\text{Tl}(\text{Ph}_2\text{pz})$ affords trivalent $\text{Sm}(\text{Ph}_2\text{pz})_3$ (**75^{Ph2}-Sm**) even under strict stoichiometric control (Scheme 31).^{174,206}

Bi and Ag reagents, BiPh_3 , $[\text{Bi}(\text{C}_6\text{F}_5)_3]$, and $[\text{Ag}(\text{C}_6\text{F}_5)(\text{py})]$, have also been used as alternatives to Hg and Tl in transmetalation reactions, though their application so far has been more limited compared to Hg and Tl reagents.^{175–177}




Figure 3. Examples of MVS reactors employed by Cloke (left) and DeKock (right). Parts in the reactor used by Cloke are A, coolant level; B, glass reaction vessel; C, gasket; D, insulated container for coolant; E, ground flange seating for Viton O-ring; F, coolant drain; G, ligand vapor inlet, “gassing”; H, electron-beam furnace; I, outlet to trap and diffusion pump; J, metal sample; K, gutter for collection of products; L, vessel for product receipt. Panel on the left reproduced with permission from ref 217. Copyright 1981 Royal Society of Chemistry Publishing. Panel on the right reproduced from ref 222. Copyright 1978 American Chemical Society.


BiPh₃ was used by Bochkarev *et al.* to prepare [Er(Ph)₃(THF)₃] in good yields from Er powder and with a small quantity of ErI₃ (4%).¹⁷⁶ More recently, Deacon, Junk, and co-workers have demonstrated that [Bi(C₆F₅)₃]¹⁷⁵ and [Ag(C₆F₅)(py)]¹⁷⁷ can be used in RTP reactions for the synthesis of RE pyrazolate and formamidinate complexes,¹⁵⁴ thus proving that these species could provide an attractive alternative to the more toxic Hg and Tl reagents. Roesky and co-workers have also shown that RT reactions with Ag(I)-NHC reagents can be employed for the synthesis of heteroleptic Ln-NHC complexes (Scheme 32).^{195,196} In these methodologies, Ag salts “(NHC)Ag(I)” [NHC = 1,3-bis(R)imidazolin-2-ylidene; R = Me (IMe₂), Mes (IMes), and Dipp (IDipp)] are reacted with Sm, Eu or Yb powders in THF at room temperature, and the outcome of these reactions is dictated by the steric demands of the NHC ligand employed and the ionic radii of the metal centers. With the most sterically demanding IDipp, the products of this reaction are free carbene and [Ln(I)₂(THF)_n] (RE = Eu, Yb).¹⁹⁵ On the other hand, when the smaller IMes ligand is employed, the heteroleptic NHC complexes [Ln(IMes)(I)₂(THF)₃] (89-Ln; Ln = Eu, Yb) are obtained.¹⁹⁵ Finally, the reaction between (IMe₂)Ag(I) and Yb affords the bis-NHC adduct [Yb(IMe₂)₂(I)₂(THF)] (90), while the tetra-NHC adducts [Ln(IMe₂)₄(I)₂] (91-Ln; Ln = Sm, Eu) are obtained with the larger divalent lanthanoids.¹⁹⁶

Sn reagents have also been employed in RT reactions. This methodology was first introduced into RE chemistry by Lappert and co-workers who reacted Yb powder with [Sn{N(SiMe₃)₂}₂] in THF under reflux and obtained the bis-silylamide complex

[Yb{N(SiMe₃)₂}₂(DME)] (78-Yb-DME) upon crystallization from DME (Scheme 33).⁹⁰ Lappert’s methodology involved a Sn(II)-to-Sn(0) reduction, whereas Deacon and co-workers later employed Sn(IV) reagents—SnMe₃(Ph₂pz), SnMe₃(OAr) (Ar = C₆H₂⁴Bu₂-2,6-Me-4), and SnMe₂(Ph₂pz)₂—for the synthesis of RE pyrazolate and aryloxides, *i.e.*, RE(Ph₂pz)₂ (74^{Ph₂}-RE), RE(Ph₂pz)₃ (75^{Ph₂}-RE), RE(OAr)₂ (53^{tBuMe}-Ln), and RE(OAr₃) (88^{tBuMe}-Ln, Scheme 33).^{166,167} While the first two reagents afford a one-electron reduction, Sn(IV)/(III),¹⁶⁷ forming Sn₂Me₆ as byproduct, the latter behaves as a two-electron oxidizing agent generating putative “SnMe₂” (Scheme 33).¹⁶⁶

3.5. Metal Vapor Synthesis

The use of MVS techniques is not a very common methodology in RE coordination and organometallic chemistry and has been employed by very few research teams across the world.¹³ However, its application has led to some remarkable results such as the first synthesis of [Sm(Cp*)₂(THF)₂] (probably one of the most iconic and well-studied f-element complexes ever reported),²¹⁶ the isolation of the first zerovalent RE molecular species [RE(C₆H₃⁴Bu₃-1,3,5)₂] (RE = Y, La, Pr, Nd, Sm, Gd, Tb, Dy, Ho, Er, Lu),^{217–219} and the first examples of Sc(II) and Sc(I) complexes, [RE(C₆H₃⁴Bu₃-1,3,5){1-CH₂C(Me)₂-3,5-^tBu₂C₆H₃}(H)]²²⁰ and [Sc(η⁵-^tBu₂C₂P₃)₂](μ-η⁶:η⁶-^tBu₃C₃P₃)].²²¹ With this technique, metal vapor is generated at low temperatures (−196 °C) under high vacuum and condensed with a substrate (Figure 3), and the resulting

Scheme 34. Examples of RE Complexes Obtained *via* Metal Vapor Co-condensation Methods^{216–219,221,222,226,229}Table 3. Selected Examples of Solvent-free and Solvated LnI₂ Salts Used in Synthesis^a

	Cl	Br	I
Nd			NdI ₂ ²⁴⁴
Sm	SmCl ₂ ²⁵⁹ <i>SmCl₂(THF)_x</i> ²⁵⁰	SmBr ₂ ²⁶⁰ <i>SmBr₂(THF)_x</i> ^{250,261}	[Nd(I) ₂ (THF) ₅] ²⁵⁴ SmI ₂ ²⁶² [Sm(I) ₂ (THF) ₂] ²⁴⁵ [Sm(I) ₂ (THF) ₃] ²⁴⁹
Eu	EuCl ₂ ²⁶³	EuBr ₂ ^{241,263,264} [Eu(Br) ₂ (THF) ₂] ²⁴⁷	EuI ₂ ²⁴¹ [Eu(I) ₂ (THF) ₂] ²⁴⁵
Dy			DyI ₂ ²⁴⁴ [Dy(I) ₂ (THF) ₅] ²⁵⁴ [Dy(I) ₂ (DME) ₃] ²⁵⁴
Tm			TmI ₂ ²⁴⁴ [Tm(I) ₂ (DME) ₃] ^{252,253} [Tm(I) ₂ (THF) ₅] ^{252,253}
Yb	YbCl ₂ ^{259,265,266}	YbBr ₂ ^{266,267} [Yb(Br) ₂ (THF) ₂] ²⁴⁷	YbI ₂ ^{268,269} [Yb(I) ₂ (THF) ₂] ²⁴⁵ [Yb(I) ₂ (THF) ₄] ²⁷⁰

^aCompounds in italics have not been isolated.

reaction mixture is then extracted in an organic solvent for recrystallization.^{13,39,222}

The first reports on the reaction between RE metal vapors and organic substrates (*e.g.*, 1-hexyne, 3-hexyne, 1,3-butadiene, 2,4-pentanedione, and cyclooctatetraene) were presented by Blackborow,²²³ Evans,^{224,225} and DeKock.^{222,226} By reacting RE metal atoms (RE = La, Ce, Nd, Er) with C₈H₈ (Scheme 34), DeKock and co-workers isolated a family of asymmetric complexes of formula [RE(COT)(THF)₂][RE(COT)₂] (92-RE; RE = La, Ce, Nd, Er), while reactivity with Yb afforded “Yb(COT)” (5), previously obtained from reaction in liquid

NH₃ (*vide supra*, Scheme 4).^{222,226} Evans and co-workers successfully applied this technique to the preparation of [Sm(Cp^{*})₂(THF)₂] (93·2THF) and [Sm(Cp^{Me⁴Et})₂(THF)₂] (94·2THF; Cp^{Me⁴Et} = {C₅EtMe₄}⁻) by reacting Sm atoms with functionalized cyclopentadienes (Scheme 34).^{216,227} In general, products obtained from reactivity of RE atoms with unsaturated substrates show magnetic properties that are in agreement with the presence of oxidized RE ions (either +2 or +3) despite the formal 0 oxidation state,¹³ such as in the diazabutadiene complexes RE[{N(^tBu)CH₂]₂]₃ (95-RE; RE = Y, Nd, Sm, Yb) reported by Cloke and co-workers (Scheme 34).²²⁸ On the

other hand, reaction of RE atoms with 1,3,5-*tri*-tertbutylbenzene, 1,3,5-^tBu₃C₆H₃, generates sandwich complexes of general formula [RE(C₆H₃^tBu₃-1,3,5)₂] (**96-RE**; RE = Sc, Y, La, Pr, Nd, Sm, Gd, Tb, Dy, Ho, Er, Lu) where magnetic characterization reveals that the RE metals are unequivocally in a zerovalent state (Scheme 34).^{217–219} When the same reaction is performed with Sc atoms, together with the Sc(0) complex [Sc(C₆H₃^tBu₃-1,3,5)₂] (**96-Sc**) a C–H activation product is also isolated, [Sc(C₆H₃^tBu₃-1,3,5){1-CH₂C(Me)2-3,5-^tBu₂C₆H₃}(H)] (**97**), in which the metal center is formally in the +2 oxidation state (Scheme 34).²²⁰ Cloke and co-workers extended this synthetic methodology to cyclization reactions involving phosphalkyne ^tBuCP. Trimerization of ^tBuCP promoted by Sc atoms generated the triphosphabenzene ring C₃P₃^tBu₃, together with formation of phospholyl ligand {C₂P₃^tBu₂}[−] and assembling of the double-decker Sc(I) complex [{Sc(η⁵-C₂P₃^tBu₂)₂(μ-η⁶:η⁶-C₃P₃^tBu₃)] (**98**, Scheme 34).²²¹ A by-product of the same reaction is also the scandocene complex [Sc(η⁵-C₃P₂^tBu₃)₂] (**99**), which was at the time only the second Sc(II) complex ever reported.²²⁹

4. HALIDES


RE halides are very common starting materials in RE coordination and organometallic chemistry. Divalent salts used in synthetic chemistry are usually diiodides of classic divalent Lns (Sm, Eu, and Yb), with the addition of Nd, Dy, and Tm. Both divalent and trivalent salts, LnI₂ and REX₃ are classically employed in salt metathesis reactions and with a variety of ligand sets. In this section some examples of their reactivity will be covered.

4.1. Divalent Ln Halides

Divalent halides of the Lns, LnX₂ (Table 3), have been known for over a century, obtained primarily as diiodides.²³⁰ However, fluorides, chlorides, and bromides are also reported for classic divalent Lns (Sm, Eu, and Yb) with the addition in some cases of Nd, Dy, and Tm.^{6,9} Noticeably, ScI₂ has also been reported, though it contains impurities of ScI₃ and is best described as Sc_{0.9}I₂.^{15,231,232} Out of all these halides, only a handful of them are of some synthetic utility in coordination and organometallic chemistry. This is due to the different physicochemical behavior of binary halides: for dihalides of Nd, Sm, Eu, Dy, Tm, and Yb the metal is in a true divalent state similar to group 2 metals (RE²⁺), while in the case of the other REs these could be best described as trivalent metals with an electron residing in the 5d band, *i.e.*, RE³⁺(e[−]) (3d band in the case of Sc and 4d band in the case of Y).¹⁵ Nonetheless, NdI₂, DyI₂, and TmI₂ are very reducing and difficult to handle, which poses challenges for their application in synthetic chemistry. In this section we will cover dihalides employed in synthetic chemistry, focusing primarily on the iodides as they are the ones that have found most widespread applications in synthetic laboratories.⁴²



Solvent-free dihalides can be obtained using a variety of methodologies which usually entail solid-state high-temperature methods.^{15,233–236} These are convenient starting materials whenever ethereal solvents are avoided, either for stabilization of complexes with low coordination numbers or because of potential side reactivity.^{7,42} Solid-state methodologies to obtain solvent-free REX₂ salts comprise (1) metallothermic reduction (Wohler method, Scheme 35, A);^{15,237} (2) comproportionation of trivalent halides with metallic RE (Scheme 35, B);^{238,239} (3) reduction of trivalent halides with hydrogen (Scheme 35, C);^{230,240} (4) reaction of RE metal with HgX₂, where X is usually

Scheme 35. Preparation of Solvent-free RE Dihalides



an iodide (Scheme 35, D);²³⁹ (5) thermal decomposition (Scheme 35, E); and (6) direct reaction between elemental halogen and RE metal (Scheme 35, F).^{15,236,241} Additionally, solvent-free EuI₂ and YbI₂ can be obtained from the reaction of metal powders with NH₄I in liquid ammonia (Scheme 35, G).^{82,241}

With the exception of liquid ammonia synthesis (method G, Scheme 35), these methods usually require specialized equipment classically used for solid-state synthesis and are unsuitable for standard synthetic laboratories.¹⁵ Bochkarev and Evans were able to adapt solid-state methods originally developed by Corbett, Kruse, and others and devised an approach which can be used in normal laboratory conditions (method F, Scheme 35).^{242–244} Bochkarev and Fagin initially reported the synthesis of NdI₂, DyI₂, and TmI₂ using a simple apparatus consisting of two thick-walled glass ampoules connected to a vacuum manifold (Figure 4).^{242,243} In this methodology, after charging ampule A with metal shavings and iodine, the system is thoroughly



1. Vacuum (D, 10^{−2} mbar), system sealed
2. Preheating (B), 200–300 °C
3. A transferred into B through C
4. 400–500 °C (B), 5 minutes

Figure 4. Schematic representation of the apparatus for the synthesis of solvent-free LnI₂ salts used by Bochkarev and co-workers. A and B, thick-walled ampoules; C, rubber tubing; D, rubber tubing to vacuum line.^{242,243} Adapted with permission from ref 243. Copyright 2003 Elsevier.

evacuated and then sealed off. Following this, the starting materials are gradually dispensed from ampule A to ampule B (while A is kept at room temperature throughout the operation, B has been preheated to 200–300 °C with a gas burner), followed by heating at 400–500 °C for a maximum of 5 min, generating the desired LnI₂ with no further purifications. The authors also noted that after initiation molten metal was formed, thus indicating that the reaction core reached temperatures around 1500 °C.^{242,243} In Bochkarev's original method the ampules are reported to be made of thick-walled borosilicate glass; however, owing to the high temperatures necessary for the initiation of the reaction and the heat subsequently generated, it is recommended that quartz reactors be used for this setup instead.

Evans and co-workers developed a more sophisticated apparatus which can produce LnI₂ salts in large quantities (up to 50 g) and with more controlled conditions (Figure 5).²⁴⁴ In





Figure 5. Apparatus for the synthesis of solvent-free LnI₂ salts used by Evans and co-workers. A, quartz tube; B, O-ring joint; C, connection to vacuum line; D, quartz addition tubes; E, quartz crucible; F, addition funnel; G, furnace. Reproduced from ref 244. Copyright 2003 American Chemical Society.

Evans' method, Ln metal powders (40 mesh) and iodine are placed in separate addition funnels (F) and the whole system is kept under static vacuum. The reactor is then heated to 450 °C inside a furnace, and the starting materials are added gradually into a quartz crucible (E) placed at the bottom of the reactor. After an initiation period, the temperature is raised above the melting point of the relative LnI₂ (Ln = Nd 562 °C, Sm 520 °C, Eu 510 °C, Dy 659 °C, Tm 756 °C, Yb 772 °C),⁹ and the reagents are added portion-wise over a period of 2 h, giving the desired LnI₂ material.

Solvated iodide salts, [Ln(I)₂(Slv)_x] (**100-Ln**; Slv = THF, DME; *x* = 2–5) of divalent Lns can also be obtained *via* standard solution methods (Scheme 36). For the most stable divalent Lns (Sm, Eu, and Yb), the main methodology was developed by Kagan and co-workers for the synthesis of [Sm(I)₂(THF)₂] (**100-Sm·2THF**),²⁴⁵ which consists of direct reaction of Ln metal powder or chips with 1,2-diiodoethane in THF.^{246–248} Evans and co-workers crystallographically characterized the 7-coordinate THF adduct [Sm(I)₂(THF)₅],²⁴⁹ but the bis-THF adduct **100-Sm·2THF** is the predominant species after standard workups following Kagan's methodology. In a similar fashion, Watson *et al.* reported the synthesis of [Eu(Br)₂(THF)₂] (**101-Eu·2THF**) and [Yb(Br)₂(THF)₂] (**101-Yb·2THF**), obtained from the reaction between Ln metal powders and 1,2-dibromoethane;²⁴⁷ this methodology cannot be applied to the

Scheme 36. Synthesis of Solvated LnX₂ Salts 100-Ln·2THF (Ln = Sm, Eu, Yb),^{246–248} 101-Ln·2THF (Ln = Eu, Yb),²⁴⁷ 100-Tm·5THF and 100-Tm·3DME^{252,253}



synthesis of solvated SmBr₂, which is obtained from the reduction of SmBr₃ or substitution of SmI₂.²⁵⁰ Solvated SmI₂ can also be obtained from the comproportionation reaction of SmI₃ with Sm metal;²⁵¹ this method has also been successfully applied to the preparation of solvated TmI₂, which can be then isolated as either THF or DME adduct, [Tm(I)₂(THF)₅] (**100-Tm·5THF**) and [Tm(I)₂(DME)₃] (**100-Tm·3DME**).^{252,253} Solvent-free NdI₂ and DyI₂ can also be converted into THF or DME adducts, [Ln(I)₂(THF)₅] (**100-Nd·5THF**) and [Dy(I)₂(DME)₃] (**100-Dy·3DME**), though manipulations have to be carried out at low temperature owing to the propensity of these species to react with ethereal solvents.²⁵⁴ Adducts of SmI₂ and YbI₂ with *N*-heterocycles (pyridine, lutidine, and 4-*tert*-butylpyridine) have also been reported by Sella and Maunder,²⁵⁵ while Wakatsuki and Hou isolated also the hexamethylphosphoramide (HMPA) adducts [Sm(I)₂(HMPA)₄] and [I]₂[Yb(HMPA)₄(THF)₂].²⁵⁶ Finally, crown ether adducts of various LnI₂ salts have also been reported: Xémard *et al.* reacted **100-Tm·3DME** with 18-crown-6 to obtain the adduct [Tm(I)₂(18-crown-6)],²⁵⁷ while Huh *et al.* successfully attempted the encapsulation of LnI₂ salts with 2.2.2-cryptand in DMF to give cationic adducts [I]₂[Ln(crypt)(DMF)_n] (Ln = Sm, Eu, *n* = 2; Ln = Yb, *n* = 1; crypt = 2.2.2-cryptand).²⁵⁸

LnI₂ salts and their adducts are commonly used in salt metathesis reactions and are particularly effective for the synthesis of divalent Sm, Eu, and Yb complexes with a variety of supporting ligands (*vide infra*, Figure 6). Most of the complexes obtained with these starting materials are homoleptic derivatives, with varying coordination numbers depending on the presence of coordinated solvents, ligand denticity, and steric properties. Salt elimination reactions have been employed for several decades in the preparation of cyclopentadienyl derivatives, such as the archetypal [Ln(Cp*)₂(THF)₂] (**93-Ln**; Ln = Sm, Yb; Slv = Et₂O, THF).^{247,271} Metallocene-type Ln(Cp)₂ complexes can be obtained by reacting group 1 cyclopentadienyl salts with SmI₂, EuI₂, and YbI₂ (**93-Yb** can also be obtained from YbCl₂)²⁷² and are isolated either as solvent-free or solvated adducts of formula [Ln(Cp^R)₂(Slv)_x] (*x* = 0–2; **1**, Cp^R = Cp; **73-Ln**, Cp^R = Cp^{iPr5}; **93-Ln** Cp^R = Cp*; **102^R-Ln**, Cp', Cp'', Cp''', Cp^{Naph}, Cp^{SiPh3}, Cp^{SiPh2Me}, Cp^{tt}, Cp^{ttt}, Cp^{iPr4}, Cp^{Bn5})^{273–279} (Cp' = {C₅H₃(SiMe₃)₂-1,3}⁻, Cp'' = {C₅H₃(SiMe₃)₂-1,3}⁻, Cp''' = {C₅H₃(SiMe₃)₃-1,2,4}⁻, Cp^{tt} = {C₅H₃^{tt}Bu₂-1,3}⁻, Cp^{ttt} = {C₅H₃^{ttt}Bu₃-1,3}⁻, Cp^{Bn5} = {C₅(CH₂Ph)₅}⁻) (Scheme 37). These methodologies are more problematic with the highly reducing salts NdI₂, DyI₂, and TmI₂. When TmI₂ or DyI₂ are reacted with KCp* or KCp'' under nitrogen, N₂-activation products [{Tm(Cp*)₂}(μ-N₂)] (**103**)²⁸⁰ and [{Dy-





Figure 6. Selected examples of ligands used in salt elimination reactions with Ln diiodides.



Scheme 37. Reactivity of LnI_2 ($\text{Ln} = \text{Nd, Sm, Eu, Dy, Tm, Yb}$) with Cyclopentadienes and Cyclopentadienyl Metal Salts^{247,271,273–279,282,283}



$(\text{Cp}^*)_2\text{Ln}(\mu\text{-N}_2)]$ (**104**)²⁷⁷ are obtained, highlighting the reducing nature of these starting materials. The direct reaction of NdI_2 , DyI_2 , and TmI_2 with cyclopentadiene (Scheme 37) was also reported by Bochkarev and co-workers, who observed oxidation of the metal to Ln^{3+} and formation of monoring complexes $[\text{Ln}(\text{Cp})(\text{I})_2(\text{THF})_3]$ (**105-Ln**) with Nd and Dy, whereas the metallocene-type complex $[\text{Tm}(\text{Cp})_2(\text{I})(\text{THF})_2]$ (**106**) was obtained from the reaction with TmI_3 , with concomitant formation of LnI_3 .²⁸¹ When HCp^* was reacted with NdI_2 and DyI_2 , products of the reaction were metallocenes $[\text{Ln}(\text{Cp}^*)_2(\text{I})(\text{THF})]$ (**107-Ln**, $\text{Ln} = \text{Nd, Dy}$) (Scheme 37).²⁸¹ Nief and Evans were able to obtain Tm metallocenes $[\text{Tm}(\text{Cp}^R)(\text{THF})_x]$ (**108^R**, $\text{Cp}^R = \text{Cp}^{\prime\prime}, \text{Cp}^{\prime\prime\prime}, \text{Cp}^{\text{tt}}, \text{Cp}^{\text{ttt}}$; $x = 0, 1$) via salt elimination reaction between TmI_2 and group 1 salts of sterically demanding Cp ligands.^{277,282,283}


The same methodologies have also been extended to the synthesis of phospholyl and arsolyl derivatives. Nief and co-

workers were able to obtain several Tm(II) metallocene-type complexes, with varying degrees of substitution on the phosphorus or arsenic heterocycles (Scheme 38), i.e., $[\text{Tm}\{\eta^5\text{-C}_4\text{H}_2\text{P}^t\text{Bu}_2\text{-2,5}\}_2(\text{THF})]$ (**109**),²⁸⁴ $[\text{Tm}\{\eta^5\text{-C}_4\text{H}_2\text{P}(\text{SiMe}_3)_2\text{-2,5}\}_2(\text{THF})]$ (**110**),²⁸⁴ $[\text{Tm}\{\eta^5\text{-C}_4\text{P}^t\text{Bu}_2\text{-2,5-Me}_2\text{-3,4}\}_2(\text{THF})_x]$ (**111**, $x = 0, 1$),^{285,286} $[\text{Tm}\{\eta^5\text{-C}_4\text{E}(\text{SiMe}_3)_2\text{-2,5-Me}_2\text{-3,4}\}_2]$ (**112^E**, $\text{E} = \text{P, As}$),²⁸⁵ and $[\text{Tm}\{\eta^5\text{-C}_4\text{PMe}_4\}_2]$ (**113**). SmI_2 and YbI_2 have also been used for the preparation of bis-phospholyl (**114^P**–**116^P**) and arsolyl complexes (**114^{As}**–**116^{As}**);^{283,286} interestingly, in some cases with Sm and Tm the resulting phospholyl complexes are obtained as dimeric structures in which the lone pair of the phosphorus atom within the phospholyl ring bridges between two metallocene moieties, i.e., $[\text{Ln}\{\eta^5\text{-C}_4\text{P}(\text{R})\}\{\mu\text{-}\eta^5\text{-C}_4\text{H}_2\text{P}(\text{R})\}]_2$ (**117**, $\text{Ln} = \text{Sm, R} = {}^t\text{Bu}_2\text{Me}_2$; **118**, $\text{Ln} = \text{Sm, R} = (\text{SiMe}_3)_2\text{Me}_2$; **119**, $\text{Ln} = \text{Tm, R} = {}^t\text{Bu}_2$).^{283,286}

Scheme 38. Synthesis of Sm, Tm, and Yb Bis-phosphohyl Complexes^{284–286}Scheme 39. Reactivity of LnI₂ Salts with Monodentate Silylamides and Aryloxides, Leading to the Isolation of Bis-amides, Tris-amides, and Dinitrogen Activation Products

The discussion on cyclopentadienyl, phosphohyl, and arsoly derivatives of Ln(II) metals provides a good snapshot of the possibilities offered by the species; classic divalent Lns (with the addition of Tm) are able to preserve the divalent state in the final products, while NdI₂ and DyI₂ are too reducing. As mentioned

above, LnI₂ salts can also be used in salt elimination reactions with a variety of ligand transfer reagents. Nitrogen donors are among the most popular ligand choices in RE chemistry, particularly monodentate silylamide {N(SiMe₃)₂}⁻ (Scheme 39). The homoleptic complexes [Yb{N(SiMe₃)₂}₂(Et₂O)₂]

Scheme 40. Selected Examples of Heteroleptic Ln(II) Complexes Obtained *via* Salt Elimination Reactions from LnI₂ Salts

(78-Yb·2OEt₂) and [Eu{N(SiMe₃)₂}₂(DME)₂] (78-Eu·2DME) were first obtained by Andersen and co-workers from the reaction between solvent-free LnI₂ and two equivalents of Na[N(SiMe₃)₂] in ethereal solvents (Et₂O, THF or DME).^{272,287} Similarly, Evans *et al.* reported the synthesis of [Sm{N(SiMe₃)₂}₂(THF)₂] (78-Sm·2THF) from [Sm(I)₂(THF)₂] and Na[N(SiMe₃)₂] in THF.²⁸⁸ These salt elimination protocols cannot be applied to Tm(II), Dy(II), and Nd(II) because of their high reducing nature. In the case of Dy and Tm, dinitrogen activation products [Ln{N(SiMe₃)₂}₂(THF)₂(μ-N₂)] (120-Ln; Ln = Dy, Tm) are obtained;²⁸⁹ when TmI₂ is employed, Evans and co-workers isolated a purple solid which they assigned as putative Tm{N(SiMe₃)₂}₂(THF)_x (78-Tm); however, no structural information has yet been reported.²⁸⁹ DyI₂ is much more reducing than TmI₂, and as a result, the solutions obtained from the reaction with Na[N(SiMe₃)₂] are highly temperature-unstable, affording dinitrogen activation product 120-Dy even at -78 °C. It is noteworthy that the analogous Nd complex [Nd{N(SiMe₃)₂}₂(THF)₂(μ-N₂)] (120-Nd) has been obtained only from the reduction of Ln/group 1 “ate” complexes, rather than through a transient Nd(II) complex, “Nd{N(SiMe₃)₂}₂”.²⁹⁰ Additionally, Evans and co-workers applied a similar synthetic strategy to the synthesis of aryloxide complexes [Ln(OC₆H₂^tBu₂-2,6)₂(THF)₂(μ-N₂)] (121-Ln; Ln = Nd, Dy), which were isolated from the reaction between LnI₂ and K(OC₆H₂^tBu₂-2,6).^{289,291} Mills and co-workers were able to isolate the first Tm(II) bis-amide complex, [Tm{N(SiⁱPr)₂}₂]

(122-Tm), which was obtained from the reaction between TmI₂ and K[N(SiⁱPr)₂] in benzene;²⁹² similar protocols were also used by the same authors for the synthesis of the analogous [Sm{N(SiⁱPr)₂}₂] (122-Sm),²⁹³ [Eu{N(SiⁱPr)₂}₂] (122-Eu), and [Yb{N(SiⁱPr)₂}₂] (122-Yb).²⁹² When the less sterically demanding silylamide ligand {N(Si^tBuMe₂)}⁻ is employed, the solvated species [Ln{N(Si^tBuMe₂)₂}₂(THF)₂] (123-Ln; Ln = Sm, Yb) are obtained instead (Scheme 39). LnI₂ salts can also be used for the preparation of heterobimetallic “ate” complexes of general formula LnM{N(SiMe₃)₂}₃ (124^M-Ln; M = alkali metal; Ln = Eu, Sm, Yb); the methodology normally consists of the reaction of three equivalents of ligand transfer reagent with LnI₂ (Scheme 39) and has also been used to synthesize separated ion pair complexes [K(L)_n][Ln{N(Si^tBuMe₂)₂}₃] (125-Ln, Ln = Sm, Eu, Tm, Yb, L = 2.2.2-cryptand, n = 1; 126-Ln, Ln = Sm, Eu, L = C₇H₈, n = 2; 127-Ln = Sm, L = DME, n = 3).²⁹⁴

A multitude of other ligands have also been employed in salt elimination reactions with LnI₂ salts, which include aromatic ligands (Figure 6, A, B)^{295,296} mono- and multidentate dentate alkyls (C-H),^{114,191,297–304} multidentate N-donors (I-M),^{299,305–310} phosphides (N),³¹¹ silanides (O),^{312,313} and gallyls.^{132,314} A major challenge in Ln(II) chemistry is the stabilization of heteroleptic complexes of the type “Ln(L)(I)” directly from salt elimination reactions, owing to the large steric demands of divalent Lns and the tendency in some cases to rearrange to homoleptic Ln(L)₂ and LnI₂. This interest originates from the possibility of further functionalizing the complexes by substituting the iodide ligand *via* metathetical

Table 4. Anhydrous and THF Adducts of RE Trihalides^a

	Cl	Br	I
Sc	ScCl ₃ ^{339,355} [Sc(Cl) ₃ (THF) ₃] ³⁵⁶ <i>ScCl₃(THF)_{1.5}</i> ³³⁹	ScBr ₃ ^{339,355} [Sc(Br) ₃ (THF) ₃] ³⁵⁷ <i>ScBr₂(THF)_{2.5}</i> ³³⁹	ScI ₃ ^{231,232} <i>ScI₃(THF)₃</i> ³⁵⁸
Y	YCl ₃ ³⁴⁰ [Y(Cl)(μ-Cl) ₂ (THF) ₂] _n ³⁵⁹ [Y(Cl) ₃ (THF) _{3.5}] ³⁵⁹ <i>YCl₃(THF)₃</i> ^{339,360}	YBr ₃ ³⁴⁹ <i>YBr₃(THF)₂</i> ³⁶¹ <i>YBr₃(THF)₃</i> ³³⁴	YI ₃ ^{233,362,363} [Y(I) ₃ (THF) _{3.5}] ³³⁵
La	LaCl ₃ ^{14,348} [La(Cl)(μ-Cl) ₂ (THF) ₂] _n ³⁶⁴ <i>LaCl₃(THF)₃</i> ³⁶⁰ <i>LaCl₃(THF)₄</i> ^{364,365}	LaBr ₃ ^{14,348} [La(Br) ₃ (THF) ₄] ³⁴³ <i>LaBr₃(THF)₂</i> ³⁶⁶ <i>LaBr₃(THF)₃</i> ³⁶⁷	LaI ₃ ^{14,233,348} [La(I) ₃ (THF) _{3.5}] ³⁴² [La(I) ₃ (THF) ₄] ^{335,368}
Ce	CeCl ₃ ^{14,348} [Ce(Cl)(μ-Cl) ₂ (THF) ₂] _n ³⁶⁹ <i>CeCl₃(THF)₃</i> ^{370,371}	CeBr ₃ ³⁵¹ [Ce(Br) ₃ (THF) ₄] ³⁷²	CeI ₃ ^{14,233,348} [Ce(I) ₃ (THF) ₄] ^{341,373}
Pr	PrCl ₃ ^{14,348} [Pr(Cl)(μ-Cl) ₂ (THF) ₂] _n ^{364,374} <i>PrCl₃(THF)₃</i> ³⁷⁵	PrBr ₃ ³⁵¹ [Pr(Br) ₃ (THF) ₄] ^{344,351}	PrI ₃ ^{14,233,348} [Pr(I) ₃ (THF) ₄] ^{335,341}
Nd	NdCl ₃ ^{14,339,348} [Nd(Cl)(μ-Cl) ₂ (THF) ₂] _n ³⁶⁹ [Nd(Cl) ₃ (THF) ₄] ³⁷⁶ <i>NdCl₃(THF)₃</i> ³⁷⁵ <i>NdCl₃(THF)_{2.5}</i> ³³⁹	NdBr ₃ ³³⁹ [Nd(Br) ₃ (THF) ₄] ³⁵⁷	NdI ₃ ^{14,233,348} [Nd(I) ₃ (THF) _{3.5}] ^{335,341} [Nd(I) ₃ (THF) ₄] ³⁷⁷ <i>NdI₃(THF)₃</i> ³⁷⁸
Sm	SmCl ₃ ^{14,348} [Sm(Cl) ₃ (THF) ₄] ³⁷⁹ <i>SmCl₃(THF)₂</i> ³⁸⁰ <i>SmCl₃(THF)₃</i> ³⁷⁵	SmBr ₃ ^{14,348} [Sm(Br) ₃ (THF) ₄] ^{344,351}	SmI ₃ ^{14,233,348} [Sm(I) ₃ (THF) ₃] ^{381,382} [Sm(I) ₃ (THF) _{3.5}] ^{341,383} <i>SmI₂(THF)₂</i> ³⁸⁴
Eu	EuCl ₃ ^{14,348} [Eu(Cl) ₃ (THF) ₄] ³⁸⁵	EuBr ₃ ^{14,348} [Eu(Br) ₃ (THF) _{3.5}] ³⁴⁴	EuI ₃ ³⁸⁶ <i>EuI₃(THF)_{3.5}</i> ³⁴¹
Gd	GdCl ₃ ³³⁹ [Gd(Cl) ₃ (THF) _{3.5}] ³⁶⁴ [Gd(Cl) ₃ (THF) ₄] ³⁷⁴ <i>GdCl₃(THF)₂</i> ³⁸⁷ <i>GdCl₃(THF)₃</i> ¹⁷¹	GdBr ₃ ³³⁹ <i>GdBr₃(THF)_{3.5}</i> ³³⁹	GdI ₃ ^{14,233,348} [Gd(I) ₃ (THF) _{3.5}] ³³⁵
Tb	TbCl ₃ ³⁴⁰ [Tb(Cl) ₃ (THF) _{3.5}] ³⁶⁹ <i>TbCl₃(THF)₃</i> ^{375,388}	TbBr ₃ ³⁵¹	TbI ₃ ^{14,233,348} [Tb(I) ₃ (THF) _{3.5}] ³⁴¹
Dy	DyCl ₃ ³⁴⁰ [Dy(Cl) ₃ (THF) _{3.5}] ³⁸⁹ <i>DyCl₃(THF)₃</i> ^{390,391} [Dy(Cl) ₂ (THF) ₅][BPh ₄] ³⁹²	DyBr ₃ ³⁴⁹	DyI ₃ ^{14,233,348} <i>DyI₃(THF)_{3.5}</i> ^{335,341}
Ho	HoCl ₃ ³³⁹ [Ho(Cl) ₃ (THF) _{3.5}] ³⁹³ <i>HoCl₃(THF)₃</i> ³⁹¹ <i>HoCl₃(THF)_{2.5}</i> ³⁶⁴	HoBr ₃ ³⁴⁹	HoI ₃ ^{14,233,348} <i>HoI₃(THF)_{3.5}</i> ³⁴¹
Er	ErCl ₃ ³⁴⁰ [Er(Cl) ₃ (THF) _{3.5}] ³⁶⁴ <i>ErCl₃(THF)₂</i> ³⁹⁴ <i>ErCl₃(THF)₃</i> ³⁷⁵	ErBr ₃ ³⁴⁹	ErI ₃ ^{14,233,348} [Er(I) ₃ (THF) ₃] ³⁹⁵ <i>ErI₃(THF)_{3.5}</i> ^{335,341}
Tm	TmCl ₃ ³⁵¹ <i>TmCl₃(THF)₃</i> ³⁹⁰ <i>TmCl₃(THF)_{2.7}</i> ³⁶⁴	TmBr ₃ ³⁴⁹	TmI ₃ ^{14,233,348} <i>TmI₃(THF)_{3.5}</i> ^{335,341}
Yb	YbCl ₃ ^{14,348} [Yb(Cl) ₃ (THF) ₃] ¹⁷¹ [Yb(Cl) ₃ (THF) _{3.5}] ³⁹⁶ <i>YbCl₃(THF)₂</i> ³⁹⁴	YbBr ₃ ^{14,348}	YbI ₃ ^{14,348} [Yb(I) ₃ (THF) ₃] ³⁹⁷ [Yb(I) ₃ (THF) _{3.5}] ³⁹⁸
Lu	LuCl ₃ ³³⁹ [Lu(Cl) ₃ (THF) ₃] ^{339,364,399} <i>LuCl₃(THF)₂</i> ³⁶⁶	LuBr ₃ ³⁴⁹	LuI ₃ ^{14,233,348} <i>LuI₃(THF)₄</i> ⁴⁰⁰

^aSolvated salts in italics have not been structurally authenticated.

reactivity. Multidentate donors such as amidinates (I), guanidinates (J), β -diketimines (BDI, K, M), tris-pyrazolyloborates (Tp, L), and bis-iminophosphorano-methanide (E) have been successfully employed to deliver such species.


Lappert and co-workers reported the preparation of the heteroleptic complexes $[\{Yb(R^vBDI^R)(\eta-I)(THF)\}_2]$ (**128**, BDI = β -diketiminate; R = Dipp, R' = Me; **129**, R = SiMe₃, R' = Ph) from $K(R^vBDI^R)$ and YbI_2 in THF (Scheme 40);³¹⁵ Jones *et al.* also reported the analogous complex $[\{Yb(t^BuBDI^{Dipp})(\eta-I)(THF)\}_2]$ (**130**) obtained with similar methodologies.³¹⁶ Chen and co-workers reacted potassium salts of tethered BDI ligands, $\{(R)C(NDipp)CHC(R)NCH_2CH_2N(Me)CH_2CH_2NMe_3\}^-$ (R^vBDI^{N2} , R = Me, t^Bu), with $[Yb(I)_2(THF)_2]$ and $[Sm(I)_2(THF)_2]$ to give dimeric heteroleptic complexes $[\{Ln(R^vBDI^{N2})(\eta-I)(THF)\}_2]$.^{317–320} Amidinate and guanidinate potassium salts afford similar results; interestingly, the flanking aryl groups can also interact with the metal center, thus supporting a switch in the coordination mode from bidentate to monodentate, such as in the dimeric complexes $[\{Yb(Giso)(\eta-I)\}_2]$ (**131**, Giso = $\{(DippN)_2CN(C_6H_{11})_2\}^-$),³²¹ $[\{Sm(Piso)(\eta-I)\}_2]$ (**132**, Piso = $\{(DippN)_2CN^tBu\}^-$),³²² $[\{Yb(Priso)(\eta-I)(THF)\}_2]$ (**133**, Priso = $\{(DippN)_2CN^iPr_2\}^-$),³²¹ and monomeric Sm complex $[Sm(Piso)(I)(THF)_2]$ (**134**, Scheme 40).³²³ Furthermore, Roesky and co-workers reacted the potassium salt $K(H-BIPM)$ ($H-BIPM = \{CH(PPh_2NSiMe_3)_2\}^-$) with $[Ln(I)_2(THF)_2]$ (Ln = Sm, Eu) in THF (Scheme 40) to give dimeric $[\{Ln(H-BIPM)(\mu-I)(THF)\}_2]$ (**135-Ln**, Ln = Sm, Eu) and monomeric $[Yb(H-BIPM)(I)(THF)_2]$ (**136**).^{324,325} Finally, Takats and co-workers were able to isolate the heteroleptic complexes $[Ln(Tp^{tBu,Me})(I)]$ (**137-Ln**, Ln = Sm, Yb) from facile reaction between $K(Tp^{tBu,Me})$ and LnI_2 in THF at room temperature;³⁰⁹ remarkably, the authors were also able to extend this chemistry to Tm(II) and isolated complex $[Tm(Tp^{tBu,Me})(I)]$ (**137-Tm**), which is a very rare example of a heteroleptic Tm(II) species (Scheme 40).³²⁶

4.2. Trivalent RE Halides

Trivalent halides (Table 4) are the most common starting materials used in RE synthetic chemistry, particularly for salt elimination reactions; the only exceptions are trifluorides, which are very rarely employed in organometallic and coordination chemistry methodologies.^{327,328} These materials are usually commercially available, but the cost of anhydrous salts can be significant; additionally, a high degree of purity is required for these starting materials, which cannot always be guaranteed from commercial sources. There are also some special circumstances in which a knowledge of these basic methodologies can be extremely useful, as shown by Chilton and co-workers in the preparation of single molecule magnets with isotopically pure ^{164}Dy ;^{329,330} the only commercially available starting material containing pure ^{164}Dy is $^{164}Dy_2O_3$, which has to be converted into $^{164}DyCl_3$ for the preparation of organometallic complexes.^{329,330} Additionally, owing to the similarities between early Lns and trivalent actinides (Ans), knowledge of these preparative methods can be used to develop synthetic protocols with RE surrogates that can be transferred to the An family, which is particularly important when working with scarce and highly hazardous transuranic elements.^{331–335}

Trivalent salts can be obtained from the direct reaction of the metal with either halide or hydrogen halides (A and B, Scheme 41), though this type of approach can be challenging for at least one of the following factors: (1) reactions need to be carried out

Scheme 41. Main Synthetic Strategies for the Preparation of Trivalent RE Halides



at high temperatures above the melting point of the REX_3 salt (usually above 700 °C); (2) the halide source needs to be of very high purity (particularly if using HCl); (3) anhydrous halides can react with the reaction vessels at high temperatures.²³⁵ This method can be problematic particularly for $RECl_3$, while the preparation of tribromides³³⁴ and triiodides³³⁵ via direct reaction of the metal with bromine or iodine under inert atmosphere is more straightforward, as both materials can be obtained with very high purity.^{9,233} A more convenient method consists of the reaction of the RE oxides, RE_2O_3 , with NH_4X (X = Cl, Br, I) at high temperatures. Reed *et al.* originally developed this methodology as a solid-state synthesis (“dry method”) for $RECl_3$ and $REBr_3$,^{336,337} which was later applied for the preparation of REI_3 by Young and Hastings (B, Scheme 41).³³⁸ However, this approach can also be implemented in synthetic laboratories as a “wet method” by dissolving the RE oxides in acid (HCl and HBr) and in the presence of NH_4X (X = Cl, Br). This methodology has been refined over the years and has now become a routine approach for the preparation of anhydrous RE chlorides and bromides in many synthetic laboratories.^{339,340} The synthesis of RE tribromides³³⁴ and triiodides³³⁵ is also easily achieved via direct reaction of the metal with bromine or iodine under inert atmosphere in ethereal solvents to yield solvated adducts.^{334,335} This is a particularly efficient method for the synthesis of triiodides, which can be prepared in very large scales as ethereal adducts $[RE(I)_3(Slv)_x]$ (Slv = Et₂O, THF; x = 3, 3.5, 4);^{335,341} Evans and co-workers have also reported the preparation of pyridine adducts,³⁴² while Deacon and Petricěk have obtained adducts of various halides with DME and glyme.^{343,344} An alternative approach is based on the use of HgX_2 salts with metallic REs (D, Scheme 40), analogous to the method described previously for divalent Lns (*vide supra*, section 4.1).^{233,345,346}

In Reed's original methodology, RE_2O_3 and NH_4Cl are mixed by heating with a Bunsen burner, forming RE chlorides as ammonium salts adducts, $(NH_4)_2RECl_5$ or $(NH_4)_3RECl_6$,³⁴⁰ dispersed in an NH_4Cl matrix;³³⁶ the quantity of ammonium salt is crucial for the success of this methodology, as incorrect stoichiometries could lead to the formation of decomposition products such as oxychlorides “REOCl”. The resulting mixture is then transferred into the bulb (A) of the apparatus shown in Figure 7.³⁴⁷ A is connected to a bent delivery tube (Figure 7, C) equipped with a collection bulb at the end (Figure 7, E); the apparatus is connected to a high-vacuum line via a trap (Figure 7, F) positioned at the end of the apparatus. Once the reactor has been charged with the $RECl_3/NH_4Cl$ mixture, the furnace is heated to 300 °C for up to 30 h, affording anhydrous $RECl_3$ upon sublimation of NH_4Cl . It is important to avoid higher temperatures at least at the initial stages of the final purification




Figure 7. Apparatus for the “dry method” synthesis of anhydrous trivalent RE chlorides.³⁴⁷ A, pyrex flask; B, furnace; C, delivery tube (internal diameter 28 mm); D, ground glass connection; E, ammonium chloride receiving bulb (500 mL); F, trap (connected *via* rubber stopper); G–G', galvanized-iron cans; H, asbestos; I, asbestos boards; J, asbestos inner cover; K, heating unit (six Westinghouse No. 299-425 space heaters, 110 V and 220 W capacity); L, thermometer; M, pyrex thermometer cover; N, high-vacuum pump connection. Reproduced with permission from ref 347. Copyright 1939 Wiley.

step, as unreacted RE_2O_3 can react with $(\text{NH}_4)_3\text{RECl}_6$ at 320°C to form YOCl , NH_3 , and H_2O .³³⁷ A similar apparatus was developed by Taylor and Carter, and later refined by Kutscher and Schneider, for drying hydrated salts $\text{REX}_3(\text{H}_2\text{O})_x$ ($X = \text{Cl}, \text{Br}, \text{I}$) with ammonium halides NH_4X ($X = \text{Cl}, \text{Br}, \text{I}$).^{14,348} Other iterations have also been reported by Corbett, Meyer, and Edelmann, and the method has also been applied to the synthesis of REBr_3 .^{240,349}


A more convenient approach toward the preparation of REX_3 salts ($X = \text{Cl}, \text{Br}$) is the so-called “wet method”. Different from the previously described “dry method”, RE_2O_3 (oxides containing tetravalent metals can also be used, *e.g.*, CeO_2 , Pr_2O_7 , and Tb_4O_7) is first dissolved in a concentrated HCl or HBr solution containing excess NH_4X (Scheme 42),^{339,340,350,351} The solution is boiled to dryness (125°C) under a stream of dry air or nitrogen, and the fumes are passed through a trap containing a 10% NaOH solution. The resulting white material, $(\text{NH}_4)_2\text{REX}_5(\text{H}_2\text{O})_x$ is finely ground and transferred into a sublimation apparatus analogous to those illustrated by Meyer and Corbett.^{240,349} The final stage of this protocol is analogous to the one previously described in the “dry method”, which entails drying the crude material under reduced pressure. Different temperature programs are reported in the

literature for this procedure; according to Marks and Diaconescu, the raw material should be heated at 200°C maximum to remove excess water first, followed by removal of excess NH_4X at higher temperatures ($T > 300^\circ\text{C}$).^{339,351}


Anhydrous RECl_3 can also be obtained by dehydration of $\text{RECl}_3(\text{H}_2\text{O})_x$. One method is briefly described above, which uses NH_4X ($X = \text{Cl}, \text{Br}, \text{I}$) as a drying agent and can be applied to other halides.³⁵² Another typical methodology employed for drying $\text{RECl}_3(\text{H}_2\text{O})_x$ involves the use of SOCl_2 and was originally reported by Freeman and Smith.³⁵² In their methodology, $\text{RECl}_3(\text{H}_2\text{O})_x$ is converted into a fine powder, transferred into a flask with excess SOCl_2 , and then refluxed gently until dehydration is complete; particular care should be taken to avoid overheating, as decomposition of SOCl_2 could lead to the formation of undesired oxychlorides. The time required to achieve complete dehydration increases going across the Ln family, with 1 h required for LaCl_3 and 110 h required for ErCl_3 ; this is likely due to the increased Lewis acidity of the metals as a consequence of the higher charge density of the smaller Lns.^{9,352} It should be noted that anhydrous RE halides are not particularly soluble, and in many instances these have to be converted into solvated forms. Such procedures can be very time-consuming; therefore, simple and quick methodologies that deliver these species as ethereal adducts can be particularly desirable (*vide infra*).

High-temperature methods are used for the preparation of unsolvated REX_3 with $\text{Hg}(\text{II})$ halides, but these methodologies have several drawbacks (*e.g.*, bespoke solid-state reactors, removal of Hg by distillation, and purification of REX_3 by sublimation) which make it impractical for most synthetic laboratories. However, Deacon and co-workers showed that standard solution synthesis can also be used for laboratory-scale preparation of solvated RECl_3 ; this methodology can also be adapted to other halides and is carried out using the same apparatus employed for other RT and RTP reactions (*vide supra*, Figure 2).^{126,345} In Deacon's method, a solution of REX_3 in THF is produced which allows removal of Hg residues *via* filtration (Scheme 43). The compounds obtained through this method are solvated species, which were originally reported by the authors as $\text{YbCl}_3(\text{THF})_3$, $\text{YbBr}_3(\text{THF})_3$, $\text{YbI}_3(\text{THF})_3$, $\text{SmI}_3(\text{THF})_3$ and $\text{ErCl}_3(\text{THF})_{3.5}$.³⁴⁵ An alternative method was also reported by Deacon *et al.* in which RE powders are treated with hexachloroethane in THF, to give THF adducts $[\text{RE}(\text{Cl})_3(\text{THF})_n]$ (137-RE; RE = La, Nd, Sm, $n = 2$; RE = Gd,

Scheme 42. Synthesis of Anhydrous RECl_3 and Apparatus for the “Wet Method” Synthesis^{339,351}



Scheme 43. Synthesis of Solvated Trivalent RE Halides via RT Reactions with HgX₂ or Direct Reaction with Hexachloroethane or Trimethylsilylchloride




Yb, $n = 3$; RE = Er, $n = 3.5$) (Scheme 43);¹⁷¹ additionally, the authors reported the structure of dimeric $[\{\text{Yb}(\text{Cl})_2(\mu\text{-Cl})(\text{THF})_2\}_2]$ (138), which was obtained upon treatment of $[\text{Yb}(\text{Cl})_3(\text{THF})_3]$ (137-Yb·3THF) with pentane over several months. It is noteworthy that there are several different reports in the literature regarding the number of THF molecules present in solvated RECl₃ salts. This is due to the variety of methods and conditions employed for their preparation, which lead to the isolation of these adducts as either neutral molecular complexes—monomeric $[\text{RE}(\text{Cl})_3(\text{THF})_3]$ (137-RE·3THF) or polymeric $[\text{RE}(\text{Cl})_3(\text{THF})_2]_n$ (137-RE·2THF)—or separated ion pair species $[\text{RE}(\text{Cl})_2(\text{THF})_4][\text{RE}(\text{Cl})_4(\text{THF})_2]$, which are usually simplified as $[\text{RE}(\text{Cl})_3(\text{THF})_{3.5}]$ (137-RE·3.5THF). The adducts reported by Deacon and co-workers were obtained by Soxhlet extraction,¹⁷¹ while other THF adducts are obtained by heating RECl₃ in THF followed by simple removal of solvent, and their formula is usually reported as RECl₃(THF)₂.³³⁹ Finally, Wu *et al.* reported the direct reaction of RE powders with trimethylsilyl chloride and methanol at room temperature (Scheme 43).³⁵³ These reactions produce the desired RE chlorides in quantitative yields for all the metals tried in this protocol (Y, La, Ce, Pr, Sm, Dy, and Yb); additionally, different silanes (dimethylchlorosilane and silicon tetrachloride) and alcohols can be employed (ethanol, propanol, and 1-pentanol).

Solvated RE triiodides can be obtained *via* various solution methods, including (1) RT reaction of RE metal with HgI₂ (*vide supra*, Scheme 43);³⁴⁵ (2) reaction of RE metal with iodoethane or diiodoethane in THF;^{343,354} (3) reaction of RE metal with iodine in isopropanol;³³⁵ and (4) reaction of RE metal with iodine in THF.³³⁵ The last method was developed by Izod *et al.* and provides a very clean and consistent route toward obtaining

triiodides of all REs (Scheme 44).³³⁵ In this method, iodine is added slowly to RE metal chips in THF, and the crude material is then baked to remove excess iodine, followed by Soxhlet extraction to afford solvated $[\text{RE}(\text{I})_3(\text{THF})_x]$ (138-RE; $x = 3.5$ or 4 for La–Nd; Table 1).³³⁵ La Pierre and co-workers reported an analogous methodology in which the initial step is performed in diethyl ether; resulting diethyl ether adducts $[\text{RE}(\text{I})_3(\text{Et}_2\text{O})_3]$ (139-RE; RE = La, Ce, Nd, Pr, Sm, Gd, Tb, Dy, Ho, Er, Tm) were also isolated, and several of them were structurally authenticated by the authors (Scheme 44).³⁴¹ It is noteworthy that these etherates easily lose solvent under reduced pressure, so the final solvent content and composition can vary.³⁴¹

In principle, salt elimination reactions with REX₃ salts can afford both homoleptic, “RE(L)₃”, and heteroleptic complexes, “RE(L)₂X” and “RE(L)X₂” (Scheme 45). The outcome of these


Scheme 45. Schematic Representation of Salt Elimination Reactions with REX₃ Salts




reactions is largely dictated by the electronic and steric features of the ligands employed, and the alkali metal and halide source chosen for the reaction (*vide supra*, section 2.1). The ligands used in salt elimination reactions with LnX₃ (*vide supra*, Figure 6) have also been used in analogous chemistry with trivalent REs, for which the ligand scope is far greater owing to the greater stability of most RE³⁺ ions.

Because of all the possible combinations of RE trihalides, ligand transfer reagents, stoichiometric ratios and reaction conditions, the number of salt elimination applications of REX₃ salts is enormous, and a full account is beyond the scope of this work. Herein, some key examples are illustrated of their use with cyclopentadienyl salts to form homoleptic, RE(Cp^R)₃, and heteroleptic complexes, RE(Cp^R)₂X and RE(Cp^R)X₂ (Scheme 46). In 1954 Wilkinson and Birmingham used anhydrous RECl₃ (RE = Sc, T, La, Ce, Pr, Nd, Sm, and Gd) to synthesize RE(Cp)₃ (140-RE) complexes *via* salt elimination with Na(Cp), thus pioneering a synthetic route which quickly became a gold-standard for RE synthetic chemists.^{2,401} Group 1 salts of a variety of substituted Cp ligands have been used in salt metathesis reactions with RE halides; in some cases, group 2 (Be, Mg) and Tl(I) reagents have also been employed.³²⁷ Salt elimination reactions between RECl₃ and Na(Cp) or K(Cp) salts are usually carried out in THF or Et₂O and under reflux conditions.^{2,401} The same strategy is applicable to substituted Cps, though it becomes increasingly difficult to displace all three halides as the steric demands of the ligands increase; RECl₃ are the salts that

Scheme 44. Synthesis of THF (138-RE) and Et₂O Adducts (139-RE) or REI₃



Scheme 46. Selected Examples of Salt Elimination Reactions of REX_3 with Cp Salts

usually give the best results when attempting to isolate homoleptic $\text{RE}(\text{Cp}^R)_3$ compounds with the smaller Cp ligands, e.g., Cp^R (**141-RE**),^{402–407} Cp^{Me} (**142-RE**),⁴⁰³ Cp^{Me} (**143-RE**),^{403,408–410} Cp^{t} (**144-RE**),^{411,412} and Cp^{t} (**145-RE**).⁴¹³ Heteroleptic species can be obtained by altering the stoichiometric ratio of these reactions, thereby using a 2:1 ratio for metallocene-type complexes, $\text{RE}(\text{Cp}^R)_2\text{X}$,^{282,283,414–418} and 1:1 ratio for monoring complexes, $\text{RE}(\text{Cp}^R)\text{X}_2$ (Scheme 46).^{354,419,420} Metallocene-type complexes are particularly desirable as they provide a fine control of the coordination sphere of the metal center, with the additional possibility of functionalizing the complexes by displacing the halide ligand.^{282,283,414–418} However, often alkali metal salts are occluded in the resulting complexes; this occurrence can sometimes be avoided by using different combinations of alkali metal salts. Lappert *et al.* reported the reactivity of RECl_3 ($\text{RE} = \text{Sc}, \text{Y}, \text{La}, \text{Ce}, \text{Pr}, \text{Nd}, \text{Yb}$) with two equivalents of LiCp^R , in the attempt to obtain metallocene type complexes $\text{RE}(\text{Cp}^R)_2\text{Cl}$.⁴²¹ However, the reactivity resulted in the formation of LiCl adducts of formula $[\text{RE}(\text{Cp}^R)_2(\mu\text{-Cl})_2\text{Li}(\text{THF})_2]$, which can be converted into the desired species, in this case dimers $[\{\text{RE}(\text{Cp}^R)_2(\mu\text{-Cl})\}_2]$, upon recrystallization or sublimation.⁴²¹ Xie and co-workers showed that by using NaCp^R the alkali metal-free compounds $[\{\text{RE}(\text{Cp}^R)_2(\mu\text{-Cl})\}_2]$ are obtained instead.⁴²² Nonetheless, it is not always possible to avoid alkali metal occlusion in salt elimination reaction, as shown with the synthesis of heteroleptic $\text{RE}(\text{III})$ metallocenes with Cp^* as supporting ligand. With the exception of $[\text{Sc}(\text{Cp}^*)_2(\text{Cl})]$ (**146-Sc**)⁴²³ and $[\text{Lu}(\text{Cp}^*)_2(\text{Cl})(\text{THF})]$ (**146-Lu**),⁴²⁴ all the complexes obtained from salt elimination reactions between RECl_3 and $\text{M}(\text{Cp}^*)$ afford MCl adducts $[\text{RE}(\text{Cp}^*)_2(\mu\text{-Cl})_2\text{M}(\text{THF})_2]$ (**147^M-RE**; $\text{Slv} = \text{Et}_2\text{O}, \text{THF}, \text{DME}$; $\text{M} = \text{Li}, \text{Na}, \text{K}$; $\text{RE} = \text{Y},$ ⁴²⁵ $\text{Ce},$ ^{426,427} $\text{La},$ ⁴²⁸ $\text{Pr},$ ⁴²⁸ $\text{Nd},$ ⁴²⁹ $\text{Sm},$ ⁴³⁰ $\text{Gd},$ ⁴²⁸ $\text{Tb},$ ⁴²⁸ $\text{Dy},$ ^{428,431} $\text{Ho},$ ⁴²⁸ $\text{Er},$ ⁴²⁸ $\text{Tm},$ ^{428,432} $\text{Yb},$ ^{428,433} Lu ⁴³⁴) (Scheme

46). Despite the drawbacks of these salt elimination reactions, “ate” complexes are still excellent starting materials which can be further functionalized and converted into very useful reagents. Nonetheless, Meng *et al.* demonstrated that with the use of different RE starting materials it is possible to avoid alkali metal occlusion and were able to obtain $[\text{Dy}(\text{Cp}^*)_2(\text{X})(\text{THF})]$ (**148^X**; $\text{X} = \text{Br}, \text{I}$) from the reaction between KCp^* and DyBr_3 or DyI_3 in THF.⁴³¹ It is also possible to laboriously convert **147^M-RE** into monomeric complexes $[\text{RE}(\text{Cp}^*)_2(\text{Cl})(\text{THF})]$ or solvent-free $[\text{RE}(\text{Cp}^*)_2(\text{Cl})]_n$, usually *via* sublimation under high vacuum and subsequent recrystallization.^{427,432} Monomeric solvent-free $\text{RE}(\text{III})$ metallocenes $[\text{RE}(\text{Cp}^R)\text{X}]$ can be obtained *via* salt metathesis by employing bulkier Cp ligands, such as Cp^{tBu} (**148^X-RE**; $\text{X} = \text{Cl}, \text{Br}, \text{I}$; $\text{RE} = \text{Y}, \text{La-Lu}$ (except Eu and Tb)) and $\text{Cp}^{i\text{Pr}4\text{R}}$ (**149^R-RE**; $\text{R} = \text{H}, \text{Me}, \text{Et}, i\text{Pr}$), with anhydrous REX_3 ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) (Scheme 46).^{282,283,414–418}

5. BOROHYDRIDES

RE borohydrides (Table 5) have gained increasing popularity over the last three decades.^{17–19} These synthons have found numerous applications in salt metathesis reactions, especially for the stabilization of heteroleptic compounds of formulas $\text{Ln}(\text{L})(\text{BH}_4)$, $\text{RE}(\text{L})(\text{BH}_4)_2$, and $\text{RE}(\text{L})_2(\text{BH}_4)$. Part of this interest is due to their accessibility and ease of preparation, combined with their versatile coordination chemistry. The borohydride ligand displays various binding modes, mainly acting as mono-, bi-, or tridentate donor (Figure 8), and provides a certain degree of flexibility in salt elimination reactions compared to RE halides.^{16,17} Though the BH_4^- ligand can act as a *pseudo*-halide, it can also occupy multiple coordination sites when binding in a k^2 - or k^3 -fashion.¹⁹ Furthermore, BH_4^- is effectively a masked hydride which can readily decompose into H^- and “ BH_3 ”,¹⁹ or can be removed with hydride-abstracting agents.^{435–438} It is noteworthy that

Table 5. Anhydrous and THF Adducts of RE Borohydrides^a

	Ln(BH ₄) ₂	RE(BH ₄) ₃
Sc		Sc(BH ₄) ₃ ⁴⁷⁸ [Sc(BH ₄) ₃ (THF) ₂] ^{479,480} <i>Sc(BH₄)₃(THF)</i> ⁴⁵⁷ <i>Sc(BH₄)₃(THF)_{1.5}</i> ⁴⁶⁵ <i>Sc(BH₄)₃(THF)₂</i> ⁴⁷⁹
Y		Y(BH ₄) ₃ ^{460,481,482} [Y(BH ₄) ₃ (THF) ₃] ^{436,483} [Y(BH ₄) ₂ (THF) ₄][Y(BH ₄) ₄] ⁴⁸⁴ <i>Y(BH₄)₃(THF)</i> ⁴⁴⁷ [Y(BH ₄) ₂ (THF) ₅][BPh ₄] ⁴⁷²
La		La(BH ₄) ₃ ⁴⁸⁵ [La(BH ₄) ₃ (THF) _{3.5}] ^b [La(BH ₄) ₃ (THF) ₄] ⁴³⁵ <i>La(BH₄)₃(THF)₃</i> ^{448,454,486} <i>La(BH₄)₃(THF)₂</i> ⁴⁸⁷
Ce		Ce(BH ₄) ₃ ^{488,489} [Ce(BH ₄) ₃ (THF) _{3.5}] ^{435,453} <i>Ce(BH₄)₃(THF)₃</i> ⁴⁴⁸ [Ce(BH ₄) ₂ (THF) ₅][BPh ₄] ⁴⁷³
Pr		Pr(BH ₄) ₃ ^{489,490} [Pr(BH ₄) ₃ (THF) _{3.5}] ⁴⁹¹ <i>Pr(BH₄)₃(THF)₃</i> ⁴⁴⁸
Nd		Nd(BH ₄) ₃ ³³⁹ [Nd(BH ₄) ₃ (THF) _{3.5}] ^{437,491} [Nd(BH ₄) ₃ (THF) ₃] ^{448,452} [Nd(BH ₄) ₂ (THF) ₅][BPh ₄] ⁴⁷² [Nd(BH ₄) ₂ (THF) ₅] [B(C ₆ F ₅) ₄] ⁴⁵⁶
Sm	Sm(BH ₄) ₂ ^{461,462,492,493} [Sm(BH ₄) ₂ (THF) ₂] _∞ ⁴⁴⁶	Sm(BH ₄) ₃ ⁴⁸⁷ [Sm(BH ₄) ₃ (THF) ₃] ^{448,486,491} [Sm(BH ₄) ₂ (THF) ₅][BPh ₄] ⁴⁷²
Eu	Eu(BH ₄) ₂ ^{461,462,493} [Eu(BH ₄) ₂ (THF) ₂] _∞ ⁴⁴¹ [Eu(BH ₄)(THF) ₅][BPh ₄] _∞ ⁴⁴¹	[Eu(BH ₄) ₃ (THF) ₃] ⁴⁹¹
Gd		Gd(BH ₄) ₃ ^{460–462} [Gd(BH ₄) ₃ (THF) ₃] ^{448,487} <i>Gd(BH₄)₃(THF)₂</i> ⁴⁸⁷ <i>Gd(BH₄)₃(THF)</i> ⁴⁴⁷
Tb		Tb(BH ₄) ₃ ^{459,461} [Tb(BH ₄) ₃ (THF) ₃] ^{436,448} <i>Tb(BH₄)₃(THF)</i> ⁴⁴⁷
Dy		Dy(BH ₄) ₃ ^{460,462} [Dy(BH ₄) ₃ (THF) ₃] ^{438,448,486} [Dy(BH ₄) ₂ (THF) ₅][BPh ₄] ^{474,475}
Ho		Ho(BH ₄) ₃ ^{461,462} <i>Ho(BH₄)₃(THF)₃</i> ⁴⁴⁸ <i>Ho(BH₄)₃(THF)</i> ⁴⁴⁷
Er		Er(BH ₄) ₃ ^{462,494} [Er(BH ₄) ₃ (THF) ₃] ^{448,487,491}
Tm	[Tm(BH ₄) ₂ (DME) ₂] ⁴⁴²	Tm(BH ₄) ₃ ^{461,462} <i>Tm(BH₄)₃(THF)₃</i> ^{442,448} <i>Tm(BH₄)₃(THF)</i> ⁴⁴⁷
Yb	Yb(BH ₄) ₂ ⁴⁶² [Yb(BH ₄) ₂ (THF) ₂] ⁴⁴¹ [Yb(BH ₄)(THF) ₅][BPh ₄] ⁴⁴¹	Yb(BH ₄) ₃ ^{462,487} [Yb(BH ₄) ₃ (THF) ₃] ^{448,491}
Lu		Lu(BH ₄) ₃ ^{462,487} [Lu(BH ₄) ₃ (THF) ₃] ^{448,486,487} <i>Lu(BH₄)₃(THF)</i> ⁴⁴⁷


^aCompounds in italics have not been structurally authenticated.^bCSD entry 1198893.



Figure 8. Typical binding modes of the borohydride ligand in metal complexes.

borohydride compounds have more covalent character compared to halides; for this reason, Ln(BH₄)₂ and RE(BH₄)₃ have a relatively higher solubility than their halide counterparts.¹⁹ Crucially, the formation of higher aggregates and “ate” complexes are less likely when these materials are employed in salt elimination reactions instead of REX₃, thus making them excellent candidates for obtaining discrete molecular entities.^{19,20}

5.1. Divalent Ln Borohydrides

The syntheses of divalent Ln borohydrides [Ln(BH₄)₂(MeCN)_n] (Ln = Eu, n = 2; Ln = Yb, n = 4) and [Ln(BH₄)₂(py)_n] (Ln = Yb, n = 4; Ln = Eu, n = 1.8) were first reported by Shore and co-workers in 1991.⁴³⁹ To obtain these materials, the authors first prepared ammoniacal solutions of LnCl₂ by reacting metal powders with NH₄Cl in liquid ammonia; removal of ammonia afforded solid ammoniacal salts LnCl₂(NH₃)_x which were converted into MeCN or pyridine solvates, LnCl₂(Slv)_x (Slv = MeCN, pyridine), and then reacted with two equivalents of NaBH₄ (Scheme 47, A).⁴³⁹ Makhaev and Borisov were able to obtain Sm(BH₄)₂, Eu(BH₄)₂, and Yb(BH₄)₂ from the thermal decomposition of the corresponding NaLn(BH₄)₄ precursor (Scheme 47, B);⁴⁴⁰ interestingly, in the case of Yb it was also possible to obtain the divalent species from thermal decomposition of Yb(BH₄)₃(THF)₂.⁴⁴⁰ A more convenient approach was devised by Visseaux and co-workers, who showed that treatment of [Sm(BH₄)₃(THF)₃] with Sm metal in THF at room temperature can afford [Sm(BH₄)₂(THF)₂] (150-Sm·2THF) in very good yields (Scheme 47, C).¹⁹ The same method was applied by Roesky and co-workers to synthesize the Yb analogue [Yb(BH₄)₂(THF)₂] (150-Yb). [Eu(BH₄)₂(THF)₂]_∞ (150-Eu·2THF) was also prepared by Roesky and co-workers either *via* salt metathesis reaction between [Eu(I)₂(THF)₂] and NaBH₄ or from the reaction between EuCl₃ and NaBH₄ with the concomitant formation of H₂ and “BH₃” (Scheme 47, D and E).⁴⁴¹ Finally, Visseaux, Nief, and co-workers obtained [Tm(BH₄)₂(DME)₂] (150-Yb·2THF) from direct salt metathesis between [Tm(I)₂(DME)₃] and KBH₄ (Scheme 47, F) or by reduction of Tm(III) precursor [Tm(BH₄)₃(THF)₃] with KC₈ (Scheme 47, G).⁴⁴² Interestingly, the authors noted that comproportionation reaction between [Tm(BH₄)₃(THF)₃] and Tm did not yield a reduced Tm(II) species.



Divalent borohydrides have found application as salt metathesis precursors with various ligands, including BDIs (151-Ln; Ln = Sm, Eu, Yb),⁴⁴³ bis-iminophosphorano-methanides,^{441,444} tris-pyrazolylborate,⁴⁴² substituted pyrrol-yls,⁴⁴⁵ and cyclopentadienyls.⁴⁴⁶ Roesky and co-workers reacted K(H-BIPM) with [Ln(BH₄)₂(THF)₂] (Ln = Eu, Yb) to give heteroleptic borohydride complexes of formula [Ln(H-BIPM)(BH₄)(THF)₂] (152-Ln; Ln = Eu, Yb) (Scheme 48).⁴⁴¹ The dual behavior of these reagents (*pseudo*-halide and masked hydride) was highlighted by the same authors who synthesized cationic [Eu(BH₄)(THF)₅][BPh₄]_∞ (153) and [Yb(BH₄)(THF)₅][BPh₄] (154) by reacting [Ln(BH₄)₂(THF)₂] with

Scheme 47. Synthesis of Divalent Ln(BH₄)₂ Complexes^{19,439–442}Scheme 48. Selected Applications of Ln(BH₄)₂ Starting Materials^{441–443,446}

$[\text{Me}_3\text{NH}][\text{BPh}_4]$ (Scheme 48);⁴⁴¹ in this reaction the hydride is abstracted to form H₂ with concomitant formation of “BH₃” and NEt₃. Finally, Momin *et al.* used salt metathesis to synthesize the dimeric Sm(II) monoring complex $[\{\text{Sm}(\text{Cp}^*)_2(\mu\text{-BH}_4)(\text{THF})_2\}_2]$ (155) and $[\text{Tm}(\text{Tp}^{\text{tBu,Me}})(\text{BH}_4)]$ (156) (Scheme 48), which have both been used as initiators for the polymerization of ϵ -caprolactone.^{442,446}


5.2. Trivalent RE Borohydrides

Trivalent RE borohydrides, RE(BH₄)₃, are known for all the RE metals (Table 5 and Scheme 49). With the exception of Sc, Ce, and Eu, these species were first obtained as THF solvated salts RE(BH₄)₃(THF)₃ from the reaction of RE(OMe)₃ with B₂H₆ in THF at room temperature (Scheme 49, A).^{447,448} A more convenient approach is based on simple metathetical reactivity between RECl₃ and MBH₄ (M = Li, Na, K) in THF. This method was first attempted by Rossmannith and Muckenhuber in

Scheme 49. Synthesis of $\text{RE}(\text{BH}_4)_3(\text{THF})_n$ Complexes *via* Solution Methods and Synthesis of Solvent-free $\text{Ln}(\text{BH}_4)_2$ and $\text{RE}(\text{BH}_4)_3$ *via* Ball-Milling

Scheme 50. Selected Examples of Salt Elimination Reactions between $\text{RE}(\text{BH}_4)_3$ Reagents and Cyclopentadienyl^{435,438,452,465–468} and Phospholy⁴³⁷ Ligand Transfer Reagents


1959, but in their methodology they could only obtain partial conversion to mixed borohydride-chloride salts $\text{RE}(\text{BH}_4)_2\text{Cl}$.^{449,450} Moreover, Andersen reported that reactivity of LiBH_4 with NdCl_3 was sluggish and did not afford the desired $\text{Nd}(\text{BH}_4)_3$ product.⁴⁵¹ However, when these reactions are carried out under reflux for at least 48 h a good conversion to solvated $\text{RE}(\text{BH}_4)_3$ complexes is obtained, as demonstrated by Ephritikhine and co-workers with the preparation of $[\text{Nd}(\text{BH}_4)_3(\text{THF})_3]$ (**157-Nd·3THF**, Scheme 49, B),⁴⁵² and this method has been successfully applied to the synthesis of trisborohydrides with all REs.^{443,452–457} Traces of halides are often present in the final products despite very long reaction times and use of large excess of MBH_4 .⁴⁵⁵ Mills and co-workers developed this methodology further by using triiodide RE salts with a large excess of KBH_4 (Scheme 49, C) and were able to identify halide impurities *via* X-ray studies;^{435–437} the percentage of iodide

($\text{BH}_4)_3(\text{THF})_3$) (**157-Nd·3THF**, Scheme 49, B),⁴⁵² and this method has been successfully applied to the synthesis of trisborohydrides with all REs.^{443,452–457} Traces of halides are often present in the final products despite very long reaction times and use of large excess of MBH_4 .⁴⁵⁵ Mills and co-workers developed this methodology further by using triiodide RE salts with a large excess of KBH_4 (Scheme 49, C) and were able to identify halide impurities *via* X-ray studies;^{435–437} the percentage of iodide


Scheme 51. Synthesis of Mixed Allyl-borohydride Complexes⁴⁶⁹ and Protonolysis Reactions with Ammonium Salts^{456,472–475}

present in $[\text{La}(\text{BH}_4)_3(\text{THF})_4]^{435}$ (**157-La-4THF**) was measured at 3%, whereas the halide content in $[\text{Y}(\text{BH}_4)_3(\text{THF})_3]$ (**157-Y-3THF**) and $[\text{Tb}(\text{BH}_4)_3(\text{THF})_3]$ (**157-Yb-3THF**) was significantly higher (ca. 15%).⁴³⁶ Adducts with other Lewis bases are also reported, including pyridine and NH_3 . Finally, solvent-free $\text{RE}(\text{BH}_4)_3$ of several REs (Sm, Gd, Tb, Dy, Er, Tm, and Yb) have been obtained by Hauback, Jensen, and co-workers *via* ball-milling of RECl_3 with LiBH_4 ,^{458–461} including divalent Sm- $(\text{BH}_4)_2$ and Yb- $(\text{BH}_4)_2$ (**Scheme 49, D**). Solvent-free $\text{RE}(\text{BH}_4)_3$ can also be obtained from the reaction of REH_3 with $\text{H}_3\text{B}\cdot\text{SMe}_2$, both with dry (ball-milling) and solution methods.⁴⁶²

Much like REX_3 salts, trivalent RE borohydrides are very efficient reagents for salt metathesis reactions, and they have been used extensively in cyclopentadienyl chemistry for the synthesis of monoring and metallocene-type complexes (**Scheme 50**). It is noteworthy that RE(III) metallocenes have also been synthesized by reacting Cp salts with $\text{RE}(\text{BH}_4)_3$ generated *in situ* from RECl_3 and NBH_4 .⁴⁶³ Monoring complexes, $\text{RE}(\text{Cp})\text{X}_2$, can be particularly challenging to isolate when using halide starting materials, while they are usually readily accessible with borohydride reagents owing to the stabilizing properties of the borohydride ligand and its flexible coordination modes.⁴⁶⁴ This is exemplified by the variety of coordination motifs exhibited by this family of complexes: (1) monoring complexes, e.g. $[\text{RE}(\text{Cp}^{\text{R}})(\text{BH}_4)_2(\text{THF})]$ (**158**, $\text{Cp}^{\text{R}} = \text{Cp}^*$, RE = Sc;⁴⁶⁵ **159**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{iPr}^5}$; RE = Dy⁴³⁸) and $[\text{RE}(\text{Cp}^{\text{R}})(\text{BH}_4)_2(\text{THF})_2]$ (**160-RE**, $\text{Cp}^{\text{R}} = \text{Cp}^*$, RE = La,⁴⁶⁶ Nd;^{452,467} **161**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{ttt}}$, RE = La⁴³⁵); (2) borohydride-bridged dimers, e.g. $[\{\text{Ce}(\text{Cp}^{\text{ttt}})(\mu\text{-BH}_4)_2(\text{THF})\}_2]$ (**162**);⁴³⁵ (3) clusters, e.g. $[\{\text{La}(\text{Cp}^{\text{ttt}})(\text{BH}_4)_2\}_6]$ (**163**)⁴³⁵ (**Scheme 50**). Metallocene-type complexes, $[\text{RE}(\text{Cp}^{\text{R}})_2(\text{BH}_4)]$, can be ob-


tained with several supporting Cp ligands (**164-RE**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{iPr}^4}$, RE = Nd, Sm,⁴⁶⁸ **165**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{m}}$, RE = La;⁴³⁵ **166-RE**, $\text{Cp}^{\text{R}} = \text{Cp}^{\text{ttt}}$, RE = La, Dy, Tm);⁴³⁵ in a similar vein to heteroleptic halide-Cp RE complexes, the formation of dimeric species is observed when steric hindrance of the ligands is reduced, e.g., $[\{\text{RE}(\text{Cp}^{\text{tt}})(\mu\text{-BH}_4)\}_2]$ (**167-RE**; RE = La, Ce).⁴³⁵ Unlike with halide-based starting materials, “ate” complex formation is usually not an issue in these methodologies. However, when similar reactions were carried out by Liu *et al.* between phospholyl $\text{K}(\text{Htp})$ ($\text{Htp} = \{\text{C}_4\text{H}_2\text{P}^t\text{Bu}_2\text{-2,5}\}^-$) and various RE borohydrides (RE = La, Ce, Nd, Sm), the leading to the isolation of “ate” complexes $[\{\text{RE}(\text{Htp})_2(\mu\text{-BH}_4)_2\text{K}(\mu\text{-DME})_2\}_2]$ (**168-RE**; RE = La, Ce) and $[\text{La}(\text{Htp})(\mu\text{-Htp})(\mu\text{-BH}_4)_2\text{K}(\text{OEt}_2)(\text{THF})]_{\infty}$ (**169**) (**Scheme 50**) were isolated.⁴³⁷ Nonetheless, the authors were able to avoid salt occlusion by performing the reactions in $^n\text{Bu}_2\text{O}$ under reflux, which gave borohydride-bridged dimers $[\{\text{RE}(\text{Htp})_2(\mu\text{-BH}_4)\}_2]$ (**170-RE**; RE = La, Ce, Nd, Sm).⁴³⁷

While RE halides can be converted into alkyls and allyls from the reaction with organolithium and Grignard reagents, analogous reactivity of RE borohydrides is not as straightforward. Visseaux and co-workers reacted $\text{Nd}(\text{BH}_4)_3$ and $\text{Sm}(\text{BH}_4)_3$ with Grignard reagents and could not obtain pure products.⁴⁶⁹ Nonetheless, reactions with half an equivalent of $\text{Mg}(\text{C}_3\text{H}_5)_2$ generated the heteroleptic allyl derivatives $[\text{RE}(\text{BH}_4)_2(\text{C}_3\text{H}_5)(\text{THF})_3]$ (**171-RE**; RE = Nd, Sm) in crystalline form and with excellent yields (**Scheme 51**).⁴⁶⁹ In addition to this, RE borohydrides can be used in protonolysis reactions, similar to their divalent analogues (*vide supra*, **Scheme 50**). This particular application was pioneered by Ephritikhine and co-workers, inspired by their previous work with An borohydrides,

Scheme 52. Synthesis of Ln(II) Silylamides 78-Ln and 176-Ln^{90,162,499–501,503,505,506}Table 6. Divalent and Trivalent RE Silylamides and Tetravalent Ce Amides Used as Synthetic Precursors^a

	Ln{N(SiR ₃) ₂ } ₂	RE{N(SiR ₃) ₂ } ₃	Ce(NR ₂) ₄
Sc		[Sc{N(SiMe ₃) ₂ } ₃] ^{539,540} [Sc{N(SiHMe ₂) ₂ } ₃ (THF)] ⁴⁹⁷	
Y		[Y{N(SiMe ₃) ₂ } ₃] ⁵⁴¹ [Y{N(SiMe ₃) ₂ } ₃ (THF)] ^b [Y{N(SiHMe ₂) ₂ } ₂ {μ-N(SiHMe ₂) ₂ } ₂] ⁵⁴² Y{N(SiMe ₃) ₂ } ₃ (THF) ₂ ⁵⁴³	
La		La{N(SiMe ₃) ₂ } ₃ ^{342,495,496,544} [La{N(SiHMe ₂) ₂ } ₂ {μ-N(SiHMe ₂) ₂ } ₂] ^{497,517} [La{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁴⁹⁷	
Ce		[Ce{N(SiMe ₃) ₂ } ₃] ^{495,496,514,545} [Ce{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁴⁶	[Ce{N(SiHMe ₂) ₂ } ₄] ⁵³⁵ [Ce(N ⁱ Pr ₂) ₄] ⁵³⁶
Pr		[Pr{N(SiMe ₃) ₂ } ₃] ^{495,496} [Pr{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁴⁶	
Nd		[Nd{N(SiMe ₃) ₂ } ₃] ^{495,496,547} [Nd{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁴³	
Sm	[Sm{N(SiMe ₃) ₂ } ₂ (THF) ₂] ²⁸⁸ [Sm ₃ {N(SiHMe ₂) ₂ } ₆ (THF) ₂] ⁵⁰⁵	[Sm{N(SiMe ₃) ₂ } ₃] ^{495,496,548} [Sm{N(SiMe ₃) ₂ } ₃ (THF)] ⁵⁴⁹ [Sm{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁵⁰	
Eu	Eu{N(SiMe ₃) ₂ } ₂ (Et ₂ O) ₂ ^{502,551} [Eu{N(SiMe ₃) ₂ } ₂ (THF) ₂] ^{502,551} [Eu{N(SiMe ₃) ₂ } ₂ (DME) ₂] ^{500,502} [Eu ₃ {N(SiHMe ₂) ₂ } ₆ (THF) ₂] ⁵⁰¹	[Eu{N(SiMe ₃) ₂ } ₃] ^{495,496,540} [Eu{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁰¹	
Gd		Gd{N(SiMe ₃) ₂ } ₃ ^{495,496} [Gd{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵¹⁹	
Tb		[Tb{N(SiMe ₃) ₂ } ₃] ^{495,496,514}	
Dy		[Dy{N(SiMe ₃) ₂ } ₃] ^{543,552}	
Ho		Ho{N(SiMe ₃) ₂ } ₃ ^{495,496} [Ho{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵⁵³	
Er		[Er{N(SiMe ₃) ₂ } ₃] ⁵⁴³ Er{N(SiHMe ₂) ₂ } ₃ ⁴⁹⁷	
Tm	Tm{N(SiMe ₃) ₂ } ₂ (THF) _x ²⁹⁰	[Tm{N(SiMe ₃) ₂ } ₃] ⁵⁴³	
Yb	Yb{N(SiMe ₃) ₂ } ₂ (Et ₂ O) ₂ ⁵⁰⁰ [Yb{N(SiMe ₃) ₂ } ₂ (THF) ₂] ^{212,554–556} Yb{N(SiMe ₃) ₂ } ₂ (DME) ₂ ⁵⁰⁰ [Yb{N(SiMe ₃) ₂ } ₂] ₂ ⁵⁰³ [Yb{N(SiMe ₃) ₂ } ₂ (BPh ₄)] ⁵¹¹ [Yb ₃ {N(SiHMe ₂) ₂ } ₆ (THF) ₂] ⁵⁵⁰	[Yb{N(SiMe ₃) ₂ } ₃] ^{495,496,557} [Yb{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁵¹⁸	
Lu		[Lu{N(SiMe ₃) ₂ } ₃] ^{495,496,558} [Lu{N(SiHMe ₂) ₂ } ₃ (THF) ₂] ⁴⁹⁷	

^aCompounds in italics have not been structurally authenticated. ^bCSD entry 2056064.

Scheme 53. Selected Examples of Protonolysis Reactivity of 78-Ln with Amides, Alcohols, and Ammonium Salts ^{191,507508–510}

which can form cationic species upon treatment with ammonium salts.⁴⁷⁰ Accordingly, Guillaume *et al.* synthesized the dimeric COT complex $[\{\text{Nd}(\text{COT})(\mu\text{-BH}_4)(\text{THF})\}_2]$ (172) *via* salt elimination reaction between $\text{K}_2(\text{COT})$ and $\text{Nd}(\text{BH}_4)_3(\text{THF})_3$ and then reacted it with $[\text{Et}_3\text{NH}][\text{BPh}_4]$ to give the cationic complex $[\text{Nd}(\text{COT})(\text{THF})_4][\text{BPh}_4]$ (173) (Scheme 51).⁴⁷¹ Furthermore, $\text{RE}(\text{BH}_4)_3$ starting materials can also be converted cleanly into cationic separated ion pair complexes $[\text{RE}(\text{BH}_4)_2(\text{THF})_5][\text{BPh}_4]$ (174-RE; RE = Y,⁴⁷² La,⁴⁷² Ce,⁴⁷³ Nd,^{472,473} Sm,⁴⁷² Dy)^{474,475} upon reaction with $[\text{Et}_3\text{NH}][\text{BPh}_4]$ (Scheme 51). Visseaux and co-workers obtained also the analogous complex $[\text{Nd}(\text{BH}_4)_2(\text{THF})_5][\text{B}(\text{C}_6\text{F}_5)_4]$ (175) from the reaction between $[\text{Me}_2\text{PhNH}][\text{B}(\text{C}_6\text{F}_5)_4]$ and $[\text{Nd}(\text{BH}_4)_3(\text{THF})_3]$.⁴⁵⁶ Other reagents that have been used with borohydride complexes for the formation of cations are highly electrophilic hydride abstracting reagents $[\text{Ph}_3\text{C}][\text{B}(\text{C}_6\text{F}_5)_4]$ ⁴³⁶ and $[(\text{Me}_3\text{Si})_2\text{H}][\text{B}(\text{C}_6\text{F}_5)_4]$.^{438,476,477}



6. NITROGEN DONORS

Amide ligands are ubiquitous in RE chemistry and have found a multitude of applications.^{22–24} Among these, silylamides are probably the most popular ligand class; Bradley *et al.* pioneered the use of the bis(trimethylsilyl)amide ligand $\{\text{N}(\text{SiMe}_3)_2\}^-$ in RE coordination chemistry in the early 1970s,^{495,496} and their work became a centerpiece of modern RE and f-element synthetic chemistry.²⁴ More recently, the conjugated base of tetramethylsilazane, $\{\text{N}(\text{SiHMe}_2)_2\}^-$, has also become a very important ligand in RE chemistry.⁴⁹⁷ Together with possessing a rich coordination chemistry, silylamides are also excellent Brønsted bases and are therefore extremely useful for protonolysis reactivity.^{22–24} Because of this, RE silylamides have now become essential starting materials for synthetic chemistry. Other RE amides have also been used in protonolysis

reactions (e.g., $\text{Ln}(\text{NH}_2)_2$ and $\text{RE}(\text{NH}_2)_3$,⁴⁹⁸ $\text{RE}(\text{NMe}_2)_3$, $\text{RE}(\text{N}^{\text{iPr}2})_3$, and $\text{RE}(\text{NCy}_2)_3$),^{22–24} but these will not be covered in this review as their uses in RE chemistry are limited compared to the vast applications of silylamides.

6.1. Divalent Ln Silylamides

The first Ln(II) silylamides, $\text{Ln}\{\text{N}(\text{SiMe}_3)_2\}_2$ (Ln = Yb, Eu), were originally synthesized by Tilley *et al.* *via* salt elimination reactions between $\text{Na}[\text{N}(\text{SiMe}_3)_2]$ and EuI_2 or YbI_2 (divalent salts were both been obtained from reactions in liquid ammonia) and were isolated as Et_2O or DME adducts, $[\text{Ln}\{\text{N}(\text{SiMe}_3)_2\}_2(\text{Slv})_2]$ (78-Ln; Ln = Yb, Eu; Slv = Et_2O , DME) (Scheme 52 and Table 6).^{499–501} It is noteworthy that 78-Eu·2THF was originally obtained from the reduction of $\text{Eu}\{\text{N}(\text{SiMe}_3)_2\}_2\text{Cl}$ with sodium naphthalene; however, current salt elimination methodologies are far more convenient.⁵⁰² The Sm analogue $[\text{Sm}\{\text{N}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$ (78-Sm·2THF) was obtained by Evans and co-workers with similar methodologies using $[\text{Sm}(\text{I})_2(\text{THF})_2]$ and $\text{Na}[\text{N}(\text{SiMe}_3)_2]$.²⁸⁸ 78-Yb can also be synthesized *via* (1) RT reaction of Yb metal with $[\text{Sn}\{\text{N}(\text{SiMe}_3)_2\}_2]$,⁹⁰ (2) RTP reaction of Yb metal with HgPh_2 and $\text{HN}(\text{SiMe}_3)_2$,¹⁶² and (3) protonolysis between $\text{Yb}(\text{Bn})_2$ (Bn = CH_2Ph) and $\text{HN}(\text{SiMe}_3)_2$ ⁵⁰³ (Scheme 52; see section 3.4 for details on RT and RTP methods). All these alternative methods are highly desirable to avoid the presence of alkali metal or halide impurities. Additionally, the protonolysis route affords 78-Yb as a solvent-free species, $[\text{Yb}\{\text{N}(\text{SiMe}_3)_2\}_2]$, without the need to use labor-intensive desolvation protocols.⁵⁰⁴ The Tm analogue $\text{Tm}\{\text{N}(\text{SiMe}_3)_2\}_2$ has been obtained as a transient species by Evans and co-workers, but its isolation and full characterization have not been achieved to date.²⁹⁰


Scheme 54. Redox Reactivity of 78-Ln and 176-Ln with Pb(Cp*)₂⁵¹²Scheme 55. Synthesis of Trivalent Silylamides 185-RE and 186-RE^{495–497,513–517,520}

Anwander and co-workers were also able to introduce the smaller silylamide $\{\text{N}(\text{SiHMe}_2)_2\}^-$ to divalent Ln chemistry.⁵⁰⁵ In their work, they observed that salt metathesis reaction between $[\text{Sm}(\text{I})_2(\text{THF})_2]$ and $\text{Li}[\text{N}(\text{SiHMe}_2)_2]$ could not deliver clean products, so they decided to explore a protonolysis route.⁵⁰⁵ $78\text{-Sm} \cdot 2\text{THF}$ was an ideal candidate as precursor for this strategy, owing to the favorable difference in $\text{p}K_a$ between $\text{HN}(\text{SiHMe}_2)_2$ and $\text{HN}(\text{SiMe}_3)_2$ (22.6 and 25.8 respectively).⁵⁰⁵ The reaction of $78\text{-Sm} \cdot 2\text{THF}$ with $\text{HN}(\text{SiHMe}_2)_2$ afforded oligomeric silylamide $[\text{Sm}_3\{\text{N}(\text{SiHMe}_2)_2\}_6(\text{THF})_2]$ (176-Sm),⁵⁰⁵ and the same method has been used for the synthesis of $[\text{Yb}_3\{\text{N}(\text{SiHMe}_2)_2\}_6(\text{THF})_2]$ (176-Yb) and $[\text{Eu}_3\{\text{N}(\text{SiHMe}_2)_2\}_6(\text{THF})_2]$ (176-Eu) (Scheme S2 and Table 6).⁵⁰¹

Amides 78-Ln have been employed extensively as protonolysis reagents with various ligand systems, particularly multidentate amines and alcohols. 78-Yb reacts with $^{\text{Me}}\text{B-DI}^{\text{Dipp}}\text{-H}$ in toluene under reflux (Scheme S3) to give the heteroleptic tetrameric complex $[\{\text{Yb}(\text{Me}^e\text{BDI}^{\text{Dipp}})\{\text{N}(\text{SiMe}_3)_2\}_2\}_4]$ (177), which was further converted into the parent dimeric hydride $[\{\text{Yb}(\text{Me}^e\text{BDI}^{\text{Dipp}})(\mu\text{-H})\}_2]$ upon treatment with PhSiH_3 .⁵⁰⁷ Junk and Cole followed a similar approach with the synthesis of the sterically hindered bis-formamidate complex $[\text{Sm}(\text{DippForm})_2(\text{THF})_2]$ ($76^{\text{Dipp}}\text{-Sm}$), which in addition was obtained *via* salt elimination and RTP methodologies (*vide supra*, Scheme 29).¹⁹¹ Moreover, Shi *et al.* treated the terphenyl-aniline $\text{H}_2\text{N}(\text{C}_6\text{H}_3\text{Xyl-2,6})$ with $78\text{-Yb} \cdot 2\text{THF}$ (Scheme S3), yielding the dimeric heteroleptic amide complex

$[\{\text{Yb}\{\text{NH}(\text{C}_6\text{H}_3\text{Xyl-2,6})\}\{\mu\text{-N}(\text{SiMe}_3)_2\}\}_2]$ (178).⁵⁰⁸ Another very effective use of these silylamides is the synthesis of aryloxy complexes. This methodology can provide an excellent route toward low-coordinate complexes, especially because of the possibility of operating in the absence of ethereal solvents. Lappert and co-workers isolated the dimeric complex $[\{\text{Yb}(\text{OAr})(\mu\text{-OAr})\}_2]$ (179 ; $\text{Ar} = \text{C}_6\text{H}_2(\text{tBu})_2\text{-2,6-Me-4}$) from the reaction between 78-Yb and two equivalents of $\text{HOC}_6\text{H}_2(\text{tBu})_2\text{-2,6-Me-4}$ in hexane (Scheme S3).⁵⁰⁹ By using the highly sterically demanding ligand $\{\text{OC}_6\text{H}_3\text{Tripp}_2\text{-2,6}\}^-$, Zhao *et al.* stabilized the bis-aryloxy complex $[\text{Sm}(\text{OC}_6\text{H}_3\text{Tripp}_2\text{-2,6})_2]$ (180) from the reaction between the parent phenol, $\text{HOC}_6\text{H}_3\text{Tripp}_2\text{-2,6}$, and $78\text{-Sm} \cdot 2\text{THF}$ (Scheme S3).⁵¹⁰ The ability of silylamides to act as Brønsted bases has also been exploited by Deacon and co-workers in the reaction of $78\text{-Yb} \cdot 2\text{THF}$ with the ammonium salt $[\text{Me}_3\text{NH}][\text{BPh}_4]$ (Scheme S3), which generates the *pseudo*-metallocene cationic complex $[\text{Yb}\{\text{N}(\text{SiMe}_3)_2\}\{\eta^6\text{-Ph}\}_2\text{BPh}_3](\text{THF})_n$ (181 ; $n = 0, 1$).⁵¹¹ 181 is itself an excellent synthetic precursor for protonolysis reactivity, as demonstrated by the same authors in the reaction with $^t\text{Bu}_2\text{pzH}$ and subsequent formation of the pyrazolate complex $[\text{Yb}(\text{tBu}_2\text{pz})\{\eta^6\text{-Ph}\}_2\text{BPh}_2](\text{THF})$.⁵¹¹

Another interesting application of 78-Ln was demonstrated by Anwander and co-workers with the synthesis of heteroleptic mono-Cp complexes using the lead reagent $\text{Pb}(\text{Cp}^*)_2$ (Scheme S4).⁵¹² In this methodology, $78\text{-Ln} \cdot 2\text{THF}$ and 176-Ln are reacted with 0.5 equiv of $\text{Pb}(\text{Cp}^*)_2$, causing oxidation of the Ln(II) center to Ln(III) and formation of piano-stool complexes

Scheme 56. Selected Examples of Reactivity of 185-RE and 186-RE^{521,525–529}

$[\text{Ln}(\text{Cp}^*)\{\text{N}(\text{SiMe}_3)_2\}_2]$ (**182-Ln**; Ln = Sm, Yb) and $[\text{Ln}(\text{Cp}^*)\{\text{N}(\text{SiHMe}_2)_2\}_2(\text{THF})]$ (**183-Ln**; Ln = Sm, Yb), with concomitant formation of metallic Pb.⁵¹² Conversely, when 78-Eu·2THF is employed, the metallocene complex $[\text{Eu}(\text{Cp}^*)_2\{\text{N}(\text{SiMe}_3)_2\}(\text{THF})]$ (**184**) is isolated.


6.2. Trivalent RE Silylamides

RE(III) silylamides $[\text{RE}\{\text{N}(\text{SiMe}_3)_2\}_3]$ (**185-RE**) were originally synthesized by Bradley *et al.* in 1972 *via* salt elimination reactions between RE trichlorides and $\text{Li}\{\text{N}(\text{SiMe}_3)_2\}$ in THF (Scheme 55 and Table 6).^{495,496} The methodology developed by Bradley is still widely used, but it can suffer from salt occlusion or “ate” complex formation. However, the use of heavier alkali metal salts and RE triiodides can suppress salt occlusion, and recrystallization or sublimation can be used for further purifications.^{24,342} Alternative procedures have also been used, such as (1) salt elimination reaction between $\text{RE}(\text{OTf})_3$ and $\text{Na}\{\text{N}(\text{SiMe}_3)_2\}$ ^{513–515} and (2) salt elimination reaction between benzyl-potassium reagents and RECl_3 followed by protonolysis with $\text{HN}(\text{SiMe}_3)_2$ (Scheme 55).^{513,516} The dimethylsilyl analogues $[\text{RE}\{\text{N}(\text{SiHMe}_2)_2\}_3(\text{THF})_n]$ (**186-RE**· n THF; RE = Sc, n = 1; RE = Y, La–Lu, n = 2) can be prepared following similar salt elimination methods using RE chlorides and group 1 transfer reagents $\text{M}\{\text{N}(\text{SiHMe}_2)_2\}$ (Scheme 55), though reactions involving chloride salts and $\text{K}\{\text{N}(\text{SiHMe}_2)_2\}$ lead to the isolation of products containing halide impurities.⁴⁹⁷ **185-La** can also be used as a starting material for the preparation of solvent-free **186-La** *via* protonolysis, similar to the strategy used for obtaining their divalent congeners **176-Ln** (Scheme 55).^{501,517–519} Moreover, Dietrich *et al.* obtained **186-Y** from the protonolysis reaction between $[\text{Y}(\text{Me})_3]_n$ and $\text{HN}(\text{SiHMe}_2)_2$ (Scheme 55; *vide infra*, section 9.1).⁵²⁰ Finally, Anwander and co-workers attempted to


prepare **186-RE** using $\text{RE}(\text{OTf})_3$ and $\text{M}\{\text{N}(\text{SiHMe}_2)_2\}$ (M = Na, K); however, they could not obtain the desired products in pure form because of the contamination of $\text{M}(\text{OTf})$.⁴⁹⁷

The applications of RE silylamides in protonolysis reactivity are numerous, and a full account is beyond the scope of this review.^{22–24} Some representative examples will be listed here to outline their main applications. Lappert and co-workers used **185-RE** (RE = Sc, Y, La Pr, Nd, Dy, Ho, Er, Yb) in protonolysis reactions with HODbmp, obtaining homoleptic aryloxo complexes $[\text{RE}(\text{ODbmp})_3]$ (**187-RE**; RE = Sc, Y, La Pr, Nd, Dy, Ho, Er, Yb) (Scheme 56).⁵²¹ This method has found many applications, particularly for obtaining RE aryloxides (especially whenever targeting solvent-free derivatives) and to exclude the possibility of halide and alkali metal contamination.^{522–524} Additionally, these protonolysis methodologies can be employed toward the synthesis of complexes supported by multidentate oxygen donors. Examples of this synthetic approach have been shown by Schelter and co-workers with the preparation of $[\text{RE}(\text{TriNOx}^{\text{OMe}})(\text{THF})]$ (**188-RE**; RE = Nd, Dy; $\text{TriNOx}^{\text{OMe}} = \{[(2\text{-}^t\text{BuNO})(5\text{-OMe})\text{-C}_6\text{H}_3\text{CH}_2]_3\text{N}\}^{3-}$) from the deprotonation of parent hydroxylamine $\text{H}_3\text{TriNOx}^{\text{OMe}}$ with **185-Nd** and **185-Dy** (Scheme 56),⁵²⁵ or by Dong and Robinson with the use of **186-RE** (RE = Y, La) for the synthesis of the heteroleptic complexes $[\text{La}\{\kappa^3\text{N}, \text{O}, \text{O}'\text{-N}(\text{C}_6\text{H}_2^t\text{Bu}_2\text{-3,5-O-2})_2(\text{CH}_2\text{Ph})\}\{\text{N}(\text{SiMe}_3)_2\}(\text{THF})_2]$ (**189**) and $[\text{Y}\{\mu\text{-}\kappa^3\text{N}, \text{O}, \text{O}'\text{-N}(\text{C}_6\text{H}_2^t\text{Bu}_2\text{-3,5-O-2})_2(\text{CH}_2\text{Ph})\}\{\text{N}(\text{SiMe}_3)_2\}_2]$ (**190**) (Scheme 56).⁵²⁶ Similarly, **185-RE** can also be used for the deprotonation of amines to target the isolation of solvent-free amides. Evans and co-workers reacted H_2NDipp with **185-RE** (RE = Y, Yb), which led to the deprotonation of the aniline and formation of the target anilido complexes $\text{RE}(\text{HNDipp})_3$ (**191-RE**; RE = Y, Yb), though only

Scheme 57. Deprotonation of Multidentate Amines with 185-RE and 186-RE^{531–533} and Insertion Reactivity of 185-Ce with Carbodiimides^{524,534}



Scheme 58. Synthesis of Tetravalent Ce(IV) Amides 201 and 202 and Protonolysis Reactivity^{535–537}




191-Y was structurally authenticated, revealing a dimeric arrangement in the solid state (Scheme 56).⁵²⁷ Anwander *et al.* have also shown that **186-Y** and **185-La** are very effective deprotonating agents toward cyclopentadienes, obtaining a range of heteroleptic metallocene-silylamide complexes, e.g. [RE(Cp*)₂{N(SiHMe₂)₂}₂] (**183-RE**; RE = Y, Lu), [Y(Cp*)₂{N(SiHMe₂)₂}₂] (**192**), [Y(Cp^{tet})₂{N(SiHMe₂)₂}₂] (**193**; Cp^{tet} = {C₅HMe₄}⁻), and [La(Cp^{Ph4})₂{N(SiHMe₂)₂}₂] (**194**) (Scheme 56).^{528,529} Interestingly, Teuben and co-workers could not obtain clean products when reacting **185-RE** (RE = Y, La, Ce) with HCp*, thus highlighting an advantage in using {N(HSiMe₂)₂}⁻ over {N(SiMe₃)₂}⁻ as a base for this type of reaction.⁵³⁰

This strategy is also very effective for obtaining heteroleptic complexes supported by multidentate amides. Vitanova *et al.* reacted Me^cBDI^{Mes}-H with **185-La** and **186-RE** (RE = Y, La) and

obtained the heteroleptic complexes [La(Me^cBDI^{Mes}){N(SiMe₃)₂}₂] (**195**) and [RE(Me^cBDI^{Mes}){N(SiHMe₂)₂}₂] (**196-RE**; RE = Y, La) (Scheme 57).⁵³¹ A similar strategy was adopted by Luo and co-workers to synthesize the heteroleptic species [Y{(NDipp)₂CPh}{N(SiHMe₂)₂(THF)}] (**197**) and [RE{(NXyl)₂CPh}{N(SiHMe₂)₂(THF)_n}] (**198-RE**; RE = Sc, Y; n = 0, 1) (Scheme 57).⁵³² Moreover, Roesky and co-workers deprotonated the chiral amidine (*S,S*)-*N,N*-bis(1-phenylethyl)-benzamidine, (*S*)-HPEBA-H, with **185-Lu**, affording the bisamidinate complex [Lu{(S)-PEBA}₂{N(SiMe₃)₂}] (**199**, Scheme 57).⁵³³ Another approach toward the stabilization of heteroleptic silylamide complexes was demonstrated by Schelter and co-workers with the preparation of heteroleptic guanidinate complexes [Ce{(NR)₂CN(SiMe₃)₂}{N(SiMe₃)₂}] (**200^R**; R = ⁱPr, Cy); in their study, they reacted **185-RE** with carbodiimides

Scheme 59. Synthetic Strategies for the Synthesis of Divalent and Trivalent RE Aryloxides



R–N=C=N–R (R = ⁱPr, Cy) leading to insertion into one of the C–N bonds (Scheme 57).^{524,534}

6.3. Tetravalent Ce Amides

Amide ligands can function as supporting ligands for Ce(IV) complexes, and some of these species are now routinely used as Ce(IV) starting materials in protonolysis reactions. Two of the most common Ce(IV) amide starting materials are complexes [Ce{N(SiHMe₂)₂}]₄ (**201**)⁵³⁵ and [Ce(NⁱPr₄)₄] (**202**).⁵³⁶ Both of these species are obtained from oxidation of Ce(III) precursors, **186**-Ce and [Ce(NⁱPr₄)₄Li], respectively, using PhICl₂, Ph₃CCl, or C₂Cl₆ (Scheme 58). Both **201** and **202** can react with protic substrates to generate new Ce(IV) complexes (Scheme 58), as shown by Kim *et al.* with the synthesis of the 8-coordinate complexes [Ce{Ph[C₆H₄N(^tBu)(O)-2]₂}]₂ (**203**)⁵³⁷ and [Ce{O[CH₂-C₆H₄N(^tBu)(O)-2]₂}]₂ (**204**)⁵³⁸ and by Schneider *et al.* with the stabilization of the tetramethylguanidinate (TMG) complex [Ce(TM₃G)₃(μ-TMG)]₂ (**205**) (Scheme 58).⁵³⁶

7. OXYGEN DONORS

The RE metals are very electropositive and highly oxophilic and therefore form very stable complexes with O-based donors.^{26,28,44,559} The coordination chemistry of aliphatic alkoxides and aryloxides with the REs is extremely well-developed, and throughout this review various methodologies which are employed for the synthesis of Ln(OR)₂ and RE(OR)₃ derivatives are discussed (see section 3). Alkoxides have attracted a lot of interest because of their high volatility, which makes them very desirable precursors for the fabrication of new materials.²⁹ In principle, these derivatives can be used as starting materials for protonolysis reactivity by matching them with substrates with higher pK_a (*vide supra*, Table 2); however, their applications are limited compared to other protonolysis reagents illustrated in this review (*e.g.*, amides and alkyls). Nonetheless, aryloxide RE complexes have found applications as protonolysis or metathesis reagents for the synthesis of challenging coordination complexes and organometallic derivatives, and Ce(IV) alkoxides have also been used in protonolysis and metathesis reactions.^{560–562} Additionally, RE inorganic derivatives containing oxygen donors have also found several applications in synthesis, particularly triflates and nitrates. For

the purpose of this review we will focus on the following reagents: (1) aryloxides (section 7.1) and (2) Ce(IV) alkoxides and CAN (section 7.2). Triflates are discussed in a separate section (section 8).

7.1. Aryloxides

The synthesis of RE aryloxides [RE(ODbmp)₃] (**72^{Dbmp}-RE**) was first reported by Lappert and co-workers in 1983, based on protonolysis of substituted phenol HODbmp with trisilylamide precursors **185-RE** (A, Scheme 59).⁵²¹ This method is still widely applied to RE aryloxide synthesis and has been used also for the synthesis of very sterically congested systems which cannot be obtained using other synthetic strategies, *e.g.* [Y(OC₆H₂Ad₂-2,6-^tBu-4)₃].⁵⁶³ Alcoholysis with Ln(II) silylamides can also be used to obtain divalent Ln alkoxides, and by using solvent-free **78-Yb** it was possible to obtain the homoleptic aryloxide **179** (A, Scheme 59; *vide supra*, Scheme 53).^{504,509} Other approaches, some of which are illustrated in detail in other sections of this review, include (1) salt elimination between REX₃ and MOAr (B, Scheme 59);^{270,522,564} (2) ammoniacal synthesis (C, Scheme 59; *vide supra*, Scheme 9);^{90,92,565} (3) direct reaction with metals (D, Scheme 59; *vide supra*, Scheme 14);¹²⁰ (4) direct reaction with metals in the presence of activators, *e.g.* Hg, HgCl₂, I₂ (E, Scheme 59; *vide supra*, Schemes 17 and 23);^{134,135,150,153,566} (5) RTP with organomercurials and HOAr (F, Scheme 59; *vide supra*, Scheme 27);^{112,123,135,153,183,198,199} and (6) RT with Tl(OAr) reagents (G, Scheme 59; *vide supra*, Scheme 31).^{91,170,198}

RE aryloxides (Table 7) have found extensive usage in protonolysis reactions with various substrates, particularly with multidentate donors containing aryloxide functionalities.^{567–570} An example of this is the alcoholysis reaction of β-ketoimines (Scheme 60) with **72^{Dbmp}-RE** (RE = Y, La, Nd, Sm, Yb).^{567–569} β-Ketoimine 1-phenyl-3-N-(*p*-methoxyphenylimino)-1-butanone reacts smoothly in a 2:1 fashion with **72^{Dbmp}-RE** (RE = Y, Nd) in THF at room temperature to give heteroleptic complexes of formula [RE{N(C₆H₄OMe-3)C(Me)CHC(Ph)-O}₂(ODbmp)] (**206-RE**; RE = Y, Nd).⁵⁶⁷ In a similar fashion, Shen and co-workers were able to obtain a series of bimetallic RE complexes supported by a phenyl-bridged BDI ligand (**207-RE**; RE = Y, La, Nd, Sm, Yb)⁵⁶⁸ and a tridentate β-ketoiminate

Table 7. Selected Divalent and Trivalent RE Aryloxides and Ce(IV) Alkoxides Used as Synthetic Precursors^a

	Ln(OAr) ₂	RE(OAr) ₃	Ce(OR) ₄
Sc		[Sc(ODbmp) ₃] ⁵²¹	
Y		[Y(ODbmp) ₃] ^{521,585}	
La		[La(ODbmp) ₃] ^{521,586}	
		[La(ODb) ₃ (MeCN)] ¹⁵³	
Ce		[Ce(ODbmp) ₃] ^{521,585}	Ce(OMe) ₄ ^{576,577,579}
		[Ce(ODb) ₃] ^{587,588}	Ce(OEt) ₄ ^{576,577}
		[Ce(ODb) ₃ (MeCN)] ⁵⁸⁷	Ce(O ⁱ Pr) ₄ ^{576,577,579,580}
			[{Ce(O ^t Bu) ₃ (μ-O ^t Bu)} ₂] ⁵⁸⁹
			[Ce(O ^t Bu) ₄ (py) ₂] ^{583,590}
			Ce(O ^t Bu) ₄ ⁵⁸¹
			Ce(O ^t Bu) ₃ (NO ₃) ⁵⁸¹
			Ce(O ^t Bu) ₂ (NO ₃) ₂ ⁵⁸¹
			Ce(O ^t Bu)(NO ₃) ₃ ⁵⁸¹
			[{Ce(OCH ₂ ^t Bu) ₄ } ₃] ⁵⁸²
Pr		<i>Pr(ODbmp)</i> ₃ ⁵²¹	
		[Pr(ODb) ₃] ⁵⁹¹	
Nd		<i>Nd(ODbmp)</i> ₃ ⁵²¹	
		[Nd(ODb) ₃] ⁵⁹¹	
Sm	[Sm(ODbmp) ₂ (THF) ₃] ^{592–595}	[Sm(ODbmp) ₃ (THF)] ⁵⁹⁶	
		[Sm(ODbmp) ₃ (MeCN) ₂] ¹⁵³	
		[Sm(ODb) ₃] ⁵⁸⁸	
Eu	[Eu(ODbmp) ₂ (THF) ₃] ^{153,270}		
	[Eu(ODb) ₂ (THF) ₂] ⁵⁹⁷		
Gd		[La(ODbmp) ₃] ^{521,586}	
Tb		[Tb(ODipp) ₃ (THF) ₂] ⁵⁶⁶	
Dy		Dy(ODbmp) ₃ ⁵²¹	
		[Dy(ODb) ₃] ⁴⁹⁸	
		[Dy(ODb) ₃ (THF)] ⁴⁹⁸	
		[Dy(ODb) ₃ (py)] ⁴⁹⁸	
		[Dy(ODb) ₃ (NH ₃)] ⁴⁹⁸	
Ho		Ho(ODbmp) ₃ ⁵²¹	
Er		[Er(ODbmp) ₃] ⁵⁹⁸	
		[Er(Dbmp) ₃ (THF)] ⁵⁶⁴	
		[Er(ODb) ₃] ⁵⁸⁸	
		[Er(ODb) ₃ (py) ₂]	
Tm		[Tb(OC ₆ H ₃ ^t Bu ₃ -2,4,6)] ⁵⁹⁹	
Yb	[{Yb(ODbmp)(μ-ODbmp)} ₂] ^{504,509}	[Yb(ODbmp) ₃ (THF)] ¹⁷⁰	
	[Yb(ODbmp) ₂ (THF) ₂] ⁹¹	[Yb(ODbmp) ₃ (MeCN)] ⁶⁰²	
	[Yb(ODbmp) ₂ (THF) ₃] ⁶⁰⁰	[Yb(ODb) ₃] ⁵⁸⁸	
	[Yb(ODbmp) ₂ (OEt ₂) ₂] ⁹¹		
	<i>Yb(ODb)₂(THF)₂</i> ⁶⁰¹		
Lu		Lu(ODbmp) ₃ ⁵²¹	
		[Lu(ODb) ₃] ⁵⁸⁸	

^aCompounds in italics have not been structurally authenticated.




containing an additional aryloxo functionality (**208-RE**; RE = Y, Nd, Sm, Yb).⁵⁶⁹

A particularly important application of **72^{Dbmp}-RE** is their use as synthetic precursors for the preparation of new RE organometallic complexes. In this type of methodology, the aryloxo donor is reacted with an organolithium reagent, resulting in transmetalation and formation of insoluble Li(OAr) as a byproduct. This strategy was applied successfully to the synthesis of heteroleptic aryloxo-Cp RE complexes by Teuben and Watkin, who reacted **72^{Db}-RE** (RE = Ce, Sm) with one equivalent of Li(Cp*) to give the monoring complex [RE-(Cp*)(ODb)₂(THF)_n] (**209-RE**; RE = Ce, n = 0; RE = Sm, n = 1) (Scheme 61).^{571,572} Additionally, several σ-bonded alkyl complexes have been obtained with this methodology, using alkyl ligands stabilized by silyl substituents on the β-position.

Lappert originally reported the reaction of **72^{Db}-La** and **72^{Db}-Sm** with Li{CH(SiMe₃)₂}, which gave homoleptic tris-alkyls [RE{CH(SiMe₃)₂}₃] (**210-RE**; RE = La, Sm).⁵¹ This method was then extended to the synthesis of mid- and late-Lns and is now the main methodology used for the preparation of this class of compounds (Scheme 61).^{573,574}


7.2. Ce(IV) Alkoxides and CAN

The coordination chemistry of tetravalent cerium has attracted a lot of interest, especially because of the use of CAN as a selective oxidizing agent in organic transformations.⁵⁷⁵ However, advances in the coordination chemistry of Ce(IV) have been hampered by the lack of well-defined and hydrocarbon-soluble starting materials, especially compared to the ample variety of compounds available for trivalent REs. A strong electron-donating environment is usually required to stabilize Ce⁴⁺, and

Scheme 60. Application of $72^{\text{Dbmp}}\text{-RE}$ as Precursor in Protonolysis Reactions with Various β -Ketoimines^{567–569}Scheme 61. Use of 72^{Db}-RE in Metathesis Reactions with Organolithium Reagents^{571–574}Scheme 62. Preparation of Ce(IV) Alkoxides *via* the “Ammonia Method”^{576,577} and “CAN Method”^{579–581} and Synthesis of 211 and 212 *via* Alcoholysis with Silylamide Complex 201⁵⁸³

for this reason there has been a lot of progress in developing Ce synthons containing oxygen donors.⁵⁵⁹ Bradley and co-workers reported the preparation of $\text{Ce}(\text{OR})_4$ (Et, ⁱPr, ⁿPr, ⁿBu) compounds by passing ammonia in an alcoholic solution of dipyridinium Ce hexachloride, $[\text{C}_5\text{H}_5\text{N-H}]_2[\text{CeCl}_6]$ (“ammonia method”, Scheme 62, A).^{576–578} Gardeff and co-workers

showed that CAN can also be used as a starting material in a very similar methodology for the preparation of $\text{Ce}(\text{OMe})_4$ (“modified ammonia method”, Scheme 62, B).⁵⁷⁹ Alternatively, CAN is dissolved in an alcohol of choice and then reacted with NaOMe, thus allowing for an easier control of reaction conditions and stoichiometry (“CAN method”, Scheme 62,

Scheme 63. Reactivity of Ce(IV) Alkoxides with Protic Substrates and Cp Reagents^{560–562}Table 8. RE Triflate Salts Used in Anaerobic Synthesis^a

	Ln(OTf)_2	RE(OTf)_3	RE(OTf)_4
Sc		Sc(OTf)_3 ⁶⁰³	
Y		Y(OTf)_3 ^{603,605,616}	
La		La(OTf)_3 ^{605,616,627}	
Ce		Ce(OTf)_3 ⁴¹³	Ce(OTf)_4 ^{614,615}
Pr		Pr(OTf)_3 ⁶⁰⁴	
Nd		Nd(OTf)_3 ^{605,616}	
Sm	Sm(OTf)_2 ⁶¹⁰	Sm(OTf)_3 ^{605,628}	
	$\text{Sm(OTf)}_2(\text{DME})_2$ ⁶⁰⁹		
Eu	$\text{Eu(OTf)}_2(\text{H}_2\text{O})$ ⁶⁰⁶	Eu(OTf)_3 ^{605,621}	
	$\text{Eu(OTf)}_2(\text{MeCN})_2$ ⁶¹²		
Gd		Gd(OTf)_3 ⁶¹⁶	
Tb		Tb(OTf)_3 ⁶²⁹	
Dy		Dy(OTf)_3 ⁶³⁰	
Ho		Ho(OTf)_3 ⁶³⁰	
Er		Er(OTf)_3 ^{605,616}	
Tm	$[\text{Tm(OTf)}_2(\text{DME})_2]_\infty$ ⁶¹³	Tm(OTf)_3 ⁶²⁹	
Yb	Yb(OTf)_2 ^{610,611}	Yb(OTf)_3 ⁶⁰⁵	
	$\text{Yb(OTf)}_2(\text{THF})_2$ ⁶¹¹		
Lu		Lu(OTf)_3 ⁶⁰⁴	


^aCompounds in italics have not been isolated.

C); the authors also noted that $\text{Ce(O}^i\text{Pr)}_4$ obtained with the original “ammonia method” is more difficult to purify, in comparison with the “CAN method” which allows for quick reaction times and easier purification procedures. Additionally, the reactions can be carried out in ethereal solvents, such as THF or DME.^{579,580} Evans *et al.* used Gardeff’s method to obtain $\text{Ce(O}^i\text{Bu)}_4$, together with other mixed nitrate-alkoxide Ce(IV) salts, e.g. $\text{Ce(O}^i\text{Bu)}_n(\text{NO}_3)_{4-n}$ ($n = 1-4$).⁵⁸¹ More recently Anwander and co-workers have used the CAN method to obtain

$\text{Ce(OCH}_2\text{Bu)}_4$, which can also be synthesized from alcoholysis with silylamide **201** and crystallizes as the trimer $[\{\text{Ce(OCH}_2\text{Bu)}_4\}_3]$ (**211**).⁵⁸² Finally, Schelter and Anwander converted Ce(IV) silylamide **201** into $[\text{Ce(O}^i\text{Bu)}_4(\text{py})_2]$ (**212**) via fast alcoholysis with ${}^t\text{BuOH}$ (Scheme 62, D).⁵⁸³

Ce(IV) alkoxides (Table 7) have been applied as protonolysis reagents for the synthesis of Ce(IV) complexes, as demonstrated by Kim *et al.* with the synthesis of 8-coordinate complex $[\text{Ce(Harene-TriNOx)}_2]$ (**213**, HareneTriNOx = $\{[(2-{}^t\text{BuNO})-$

Scheme 64. Preparation of Divalent and Trivalent RE Triflate Salts




$\text{C}_6\text{H}_4)_2[(2\text{-}^t\text{BuNOH})\text{C}_6\text{H}_4]\text{C}_6\text{H}_3\}^{2-}$ (Scheme 63).⁵⁶⁰ Gulino *et al.* attempted to use $\text{Ce}(\text{O}^i\text{Pr})_4$ in salt elimination reactions with $\text{Tl}(\text{Cp})$ or $\text{Mg}(\text{Cp})_2$ without success, observing reduction to Ce(III) instead.⁵⁶¹ Nonetheless, when the tin reagent R_3SnCp ($\text{R} = \text{Me}, ^n\text{Bu}$) was employed, they were able to isolate $\text{Ce}(\text{Cp})_3(\text{O}^i\text{Pr})$ (**214**), though structural authentication was not obtained (Scheme 63).⁵⁶¹ A different approach was followed by Evans *et al.*, who reacted mixed Ce(IV) nitrate-alkoxides $\text{Ce}(\text{O}^t\text{Bu})_2(\text{NO}_3)_2$ and $\text{Ce}(\text{O}^t\text{Bu})(\text{NO}_3)_3$ with $\text{Na}(\text{Cp})$. In the case of the reaction between $\text{Ce}(\text{O}^t\text{Bu})_2(\text{NO}_3)_2$ and two equivalents of $\text{Na}(\text{Cp})$, the metallocene Ce(IV) complex $\text{Cp}(\text{Cp})_2(\text{O}^t\text{Bu})_2$ (**215**) was obtained in excellent yields but could not be structurally authenticated (Scheme 63).⁵⁶² However, when $\text{Ce}(\text{O}^t\text{Bu})(\text{NO}_3)_3$ was reacted with three equivalents of $\text{Na}(\text{Cp})$, a mixture containing **215** and $[\text{Cp}(\text{Cp})_3(\text{O}^t\text{Bu})]$ (**216**) was obtained, and the latter was structurally characterized (Scheme 63).⁵⁶² Finally, Gordon and co-workers used the same methodology to isolate metallocene derivatives $[\text{Ce}(\text{Cp}')_2(\text{O}^t\text{Bu})_2]$ (**217**) and $[\text{Ce}(\text{Cp}'')_2(\text{O}^t\text{Bu})_2]$ (**218**) (Scheme 63).⁵⁸⁴

8. TRIFLATES

Triflate (OTf , $\{\text{CF}_3\text{SO}_3\}^-$) salts are a very popular choice of starting materials in RE synthetic chemistry (Table 8). These salts are used as more soluble replacements of halides in salt elimination reactions.^{9,31} Additionally, triflates are better leaving groups and can be easily displaced from the metal coordination sphere.^{9,31} Similar to RECl_3 , $\text{RE}(\text{OTf})_3$ salts are usually commercially available both in anhydrous and hydrate form. However, they can also be conveniently made from the direct reaction of the oxide (RE_2O_3) with triflic acid (HOTf), followed by thorough desiccation under reduced pressure; reports vary from a minimum of 4 h to several days (Scheme 64, A).^{9,603–606} This method was originally reported by Massaux and is the most common strategy for obtaining anhydrous $\text{RE}(\text{OTf})_3$ salts.⁶⁰⁵

Triflate salts of divalent Sm, Eu, and Yb are obtained *via* various methodologies. $\text{Sm}(\text{OTf})_2$ can be prepared from direct salt elimination reaction of $\text{K}(\text{OTf})$ with SmI_2 (Scheme 64, B)⁶⁰⁷ or by reducing $\text{Sm}(\text{OTf})_3$ with Sm metal in the presence of iodine⁶⁰⁸ or mercury⁶⁰⁹ (Scheme 64, C). Nief and co-workers obtained poor results when performing this reaction in THF, whereas they achieved excellent conversions when using DME, ultimately leading to the isolation of the DME adduct $\text{Sm}(\text{OTf})_2(\text{DME})_2$.⁶⁰⁹ Another method for the preparation of $\text{Ln}(\text{OTf})_2$ is the reduction of $\text{Ln}(\text{OTf})_3$ with Grignard reagents, which has also been successfully employed for the preparation of $\text{Yb}(\text{OTf})_2(\text{THF})_3$ (Scheme 64, D).⁶¹⁰ The DME-solvate analogue, $\text{Yb}(\text{OTf})_2(\text{DME})$, has also been obtained from the reaction of HOTf with $[\text{Yb}\{\text{N}(\text{SiMe}_3)_2\}_2(\text{OEt}_2)_2]$.⁶¹¹

$\text{Eu}(\text{OTf})_2$ can be obtained by treating EuCO_3 with excess triflic acid, affording monohydrate $\text{Eu}(\text{OTf})_2(\text{H}_2\text{O})$ upon drying *in vacuo* at 90 °C overnight (Scheme 64, E).⁶⁰⁶ Alternatively, water-free $\text{Eu}(\text{OTf})_2$ can be obtained with the method described by Krossing and co-workers, which starts from hydrated $\text{Eu}(\text{OTf})_3(\text{H}_2\text{O})_x$.⁶¹² Eu(III) triflate is dissolved in triethylorthoformate and heated at 50 °C for 1 h, followed by treatment with MeCN and subsequent crystallization of $\text{Eu}(\text{OTf})_3(\text{MeCN})_2$ at room temperature. The MeCN solution of $\text{Eu}(\text{OTf})_3(\text{MeCN})_2$ is then passed through a column filled with Zn/Hg amalgam to give the final product $\text{Eu}(\text{OTf})_2(\text{MeCN})_2$ (Scheme 64, F).⁶¹² Outside the triad of classic Ln^{2+} ions, only Tm(II) forms a stable triflate salt, $[\text{Tm}(\text{OTf})_2(\text{DME})_2]_\infty$.⁶¹³ This was reported by Nocton and co-workers in 2017 and was obtained *via* reduction of $\text{Tm}(\text{OTf})_3$ with KC_8 in DME at $-35 \text{ }^\circ\text{C}$ (Scheme 64, G).⁶¹³ Finally, tetravalent $\text{Ce}(\text{OTf})_4$ can be obtained from the reaction between CAN, K_2CO_3 , and HOTf.⁶¹⁴ However, Berthet and co-workers reported that anhydrous material cannot be obtained following this methodology because drying above 100 °C results in reduction to Ce(III).⁶¹⁵ The authors were able to dry commercial $\text{Ce}(\text{OTf})_4(\text{H}_2\text{O})_x$ by treating it with trifluorome-

Scheme 65. Selected Applications of Triflate Salts for the Synthesis of Cyclopentadienyl, COT, and Allyl RE Complexes^{622–626}

thanesulfonic anhydride, TfOTf, followed by drying at 100 °C for 15 h.⁶¹⁵

RE(III) triflates have numerous applications as salt metathesis reagents, with a scope very similar to RE halides including numerous ligands, such as cyclopentadienyls, allyls, and a variety of mono- and multidentate donors.^{616–621} Therefore, only some selected applications are listed in this section to give an overview of the type of coordination chemistry accessible with these synthons. Because triflates are very good leaving groups, they have been used for the synthesis of sterically congested RE(Cp^R)₃ systems, such as [La(Cpⁿ)₃] (**219**) and [RE(Cp^{tt})₃] (**220-RE**; RE = La, Ce, Pr),⁶²² whose synthesis is typically more challenging when using halide precursors.⁴¹³ For example, **219** can be obtained from the reaction between LaCl₃ and K(Cpⁿ) requiring first treatment in THF at room temperature for 24 h (to form transient La(Cpⁿ)₂Cl), followed by reflux in toluene for an additional 24 h; whereas with La(OTf)₃ and Mg(Cpⁿ)₂ the reaction only requires stirring at room temperature in THF overnight.⁶²² Triflate salts have also been used for the synthesis of heteroleptic open metallocene complexes (e.g., [Ce(Cp^{tt})₂(OTf)], **221**), though in some cases alkali metal salts are occluded in the products resulting in “ate” complex formation, as observed by Lappert and co-workers with the isolation of [{Nd(Cpⁿ)₂(μ-OTf)₂Li]₂] (**222**).⁶²³ From the reaction of K₂COT with Y(OTf)₃, Edelman and Kilmann obtained dimeric [{Y(COT)(μ-OTf)(THF)}₂] (**223**) (Scheme 65),⁶²⁴ which can then be functionalized further by substituting the triflate anion with various ancillary ligands *via* salt elimination. Moreover, John and co-workers have shown that triflate salts of various REs (Ce, Nd, Eu, Tb, and Yb) can react with the potassium salts of substituted allyls (Scheme 65) to form homoleptic allyl derivatives [RE{C₃H₃(SiMe₃)₂-1,3}₃(THF)] (**224-RE**; RE = Ce, Nd, Tb).⁶²⁵ The same metathetical protocol is also effective for Yb(OTf)₂ (Scheme 65) and the authors were able to obtain the bis-allyl complexes [Yb{C₃H₃(SiMe₃)-1-R-3}₂(THF)_n] (**225^R**; R = H, SiMe₃,

SiPh₃; n = 1, 2).⁶²⁶ This is noteworthy, as there are very few applications in the literature of Ln(II) triflates as synthetic precursors in anaerobic synthesis.

9. ORGANOMETALLIC REAGENTS

Organometallic compounds of the REs are generally very reactive because of the high polarization of metal–carbon interactions. This is a problem typical of σ-bonded organometallic compounds but is also true for π-bonded organometallics, such as Cp, allyl, and COT derivatives. Additionally, RE organometallics can easily undergo β-hydride elimination and intramolecular C–H activation reactions. All these issues, together with the high electropositive character and large ionic radii of the REs, make the stabilization of organometallic species extremely challenging. However, over the years there has been a lot of progress in the stabilization of relatively simple RE organometallics which can function as synthons for the preparation of new compounds, particularly as reagents in protonolysis reactions. Simple σ-bonded RE alkyls (Me, ⁿBu) and aryls (Ph, C₆F₅) have been well-investigated (section 9.1 and Table 9), though such compounds tend to be extremely reactive and often thermally sensitive. In this section Ln and RE aryls will not be discussed in detail because an extensive account of their preparation and application is provided in section 3 (*vide supra*). Alkyl complexes can be further stabilized by substituting the central carbon donor with silyl substituents, and this strategy has been used very effectively for the preparation of RE complexes α-silyl ligands {CH₂SiMe₂R}[−] (R = Me, Ph) and {CH(SiMe₃)₂}[−]; such complexes are now common synthetic precursors in RE organometallic chemistry (section 9.2 and Table 10). Moreover, RE benzyl (Bn, {CH₂Ph}[−]) complexes constitute another important class of organometallic starting materials (section 9.3 and Table 11); benzyl ligands are nominally alkyl-type donors, though they ligate the metal center in a multihapto fashion. Finally, organoaluminate, Ln(AlR₄)₂ and RE(AlR₄)₃ (R = Me, Et), are special types of organometallic

Table 9. Selected Simple RE Hydrocarbyls Used as Starting Materials^a

RE	"RE(R) ₃ "
Sc	<i>[Sc(Me)₃]_n</i> ⁶³³ <i>Sc(μ-Me)₆Li₃(THF)_{1,2}</i> ⁶³⁹ <i>[Sc₂(μ-Me)₆Li₃(Et₂O)₃(THF)₂]⁶³⁹</i> <i>Sc(μ-ⁿBu)₆Li₃(THF)_x</i> ⁶⁴⁰
Y	<i>[Y(Me)₃]_n</i> ^{631,632} <i>[Y₂(μ-Me)₆Li₃(Et₂O)₂(THF)₃]⁶³⁹</i> <i>Y(μ-Me)₆Li₃(THF)_{1,3}</i> ⁶³⁹ <i>Y(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸ <i>"LaMe₃"</i> ⁶³²
La	<i>La(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸ <i>La(μ-ⁿBu)₆Li₃(THF)_x</i> ⁶⁴⁰
Ce	<i>[Ce(μ-Me)₆Li₃(TMEDA)₃]⁶⁴⁰</i> <i>[Ce(μ-ⁿBu)₄Li₂(TMEDA)₂]⁶⁴⁰</i> <i>Ce(μ-ⁿBu)₆Li₃(THF)_x</i> ⁶⁴⁰
Pr	<i>Pr(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸
Nd	<i>Nd(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸
Sm	<i>Sm(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸
Eu	
Gd	<i>Gd(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸ <i>Gd(μ-Me)₆Li₃(THF)_{2,4}</i> ⁶³⁹
Tb	<i>[Tb₂(μ-Me)₆Li₃(Et₂O)₂(THF)₃]⁶³⁹</i> <i>Tb(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸ <i>Tb(μ-Me)₆Li₃(THF)_{1,5}</i> ⁶³⁹
Dy	<i>Dy(μ-Me)₆Li₃(TMEDA)₃</i> ⁶³⁸ <i>Dy(μ-Me)₆Li₃(THF)_{1,65}</i> ⁶³⁹
Ho	<i>HoMe₃·[Ho(Me)₃]_n</i> ⁶³⁴ <i>[Ho(μ-Me)₆Li₃(TMEDA)₃]⁶³⁸</i> <i>Ho(μ-Me)₆Li₃(THF)_{1,2}</i> ⁶³⁹
Er	<i>[Er(μ-Me)₆Li₃(TMEDA)₃]^{635,636}</i> <i>Er(μ-Me)₆Li₃(THF)₂</i> ⁶³⁹
Tm	<i>Tm(μ-Me)₆Li₃(TMEDA)₂</i> ⁶³⁸ <i>Tm(μ-Me)₆Li₃(THF)_{1,7}</i> ⁶³⁹
Yb	<i>[Yb(Me)₂]_n</i> ⁵⁰³ <i>[Yb₂(μ-Me)₆Li₃(Et₂O)₃(THF)₂]⁶³⁹</i> <i>Yb(μ-Me)₆Li₃(THF)⁶³⁹</i>
Lu	<i>[Lu(Me)₃]_n</i> ^{631,632} <i>[Lu(μ-Me)₆Li₃(DME)₃]^{635,637,638}</i> <i>Lu(μ-Me)₆Li₃(THF)⁶³⁹</i> <i>[Lu(μ-ⁿBu)₄Li₂(TMEDA)₂]⁶⁴⁰</i> <i>[Lu(μ-ⁿBu)₆Li₃(THF)₄]⁶⁴⁰</i>

^aCompounds in italics have not been structurally authenticated.

reagents which can be effectively considered as masked alkyls stabilized by the formation of an adduct with AlR₃ (section 9.4 and Table 12). Most of these synthetic precursors have been the subject of a recent comprehensive review by Zimmermann and Anwander;¹¹ therefore, this section will aim to provide a broad overview of these reagents, with some highlights on the emergence of new reagents and new applications.

9.1. Simple Hydrocarbyls (Me, ⁿBu, Ar)

Salt metathesis reactivity is the most common methodology employed for the synthesis of binary alkyl and aryl complexes (Scheme 66, A), though there are some notable exceptions such as (1) RT reactions with organomercurials and organobismuth reagents for the synthesis of aryl complexes (Scheme 66, B; *vide supra*, section 3.4),¹² (2) donor cleavage reactions from organoaluminate precursors (Scheme 66, C),⁶³¹ and (3) transmetalation reactions between amide precursors and MeLi (Scheme 66, D).⁵⁰³ Saran and co-workers reported the reaction

of MeLi and PhLi with RECl₃ (Sc, Y, La, and Pr) in 1970.⁶³² In the case of reactivity with PhLi, the authors identified Sc(Ph)₃ and Y(Ph)₃ *via* IR spectroscopy and elemental analysis, though with La and Pr they identified the formation of "ate" complex LiLa(Ph)₄ from analogous reactivity. However, the formulas of the methyl derivatives "RE(Me)₃" (226-RE; RE = Sc, Y, La) obtained from the reactions between RECl₃ and MeLi could not be unequivocally identified.⁶³² Neutral methyl complexes have been obtained with Ho, Lu, and Y by Anwander and co-workers, isolated as polymeric [RE(Me)₃]_n (226-RE; RE = Sc,⁶³³ Y,⁶³¹ Ho,⁶³⁴ Lu⁶³¹) from the donor-cleavage reaction of aluminate precursors (Scheme 66; *vide infra*, section 9.4). Schumann reacted RECl₃ with six equivalents of MeLi in the presence of TMEDA and obtained a series of hexamethyl "ate" complexes of formula [RE(μ-Me)₆Li₃(TMEDA)₃] (227-RE; RE = Y, La, Pr, Nd, Sm, Gd, Tb, Dy, Ho, Er, Tm, and Lu).^{635–638} Furthermore, Okuda and co-workers carried out an in-depth study on the stability and reactivity of 227-RE and were also able to obtain the pentametallic complexes [RE₂(μ-Me)₆Li₃(Et₂O)_n(THF)_m] (RE = Sc, n = 3, m = 2; RE = Y, Tb, n = 2, m = 3).⁶³⁹ Anwander and co-workers recently developed analogous chemistry with Ce, and in addition they extend this approach to the preparation of butyl derivatives [RE(μ-ⁿBu)₄Li₂(THF)₄] (228-RE; RE = Sc, Y, La, Ce, Lu) and [RE(μ-ⁿBu)₄Li₂(TMEDA)₂] (229-RE; RE = Ce, Lu).⁶⁴⁰ Finally, Anwander and co-workers reported the synthesis and reactivity of the neutral methyl complex [Yb(Me)₂]_n (230), which is the only example of methyl derivative with a divalent Ln; 230 was obtained *via* transmetalation reaction between bis-silylamide precursors and MeLi.⁵⁰³

RE alkyl complexes (Table 9) are excellent protonolysis reagents owing to the very favorable pK_a (*vide supra*, Table 2) and react promptly with Brønsted acids. Anwander and co-workers treated 226-Y with HN(SiMe₃)₂, HN(SiHMe₂)₂, and HOCH^tBu₂ (Scheme 67), obtaining the *tris*-amide complexes 185-Y and 186-Y and alkoxide derivative Y(OCH^tBu₂)₃ (231).^{542,631} Interestingly, reactivity of 226-RE with HN-(SiMe₃)(Dipp) leads to the deprotonation of one methyl ligand and formation of methylidene bridged cluster [RE₃{N(SiMe₃)-(Dipp)}₃(μ₂-Me)₃(μ₃-Me)(THF)₃] (232-RE; RE = Y, Ho, Lu) (Scheme 67).⁶³⁴ In a similar fashion, Berger *et al.* investigated the reactivity of 229-Ce with HOCH^tBu₂, which led to the isolation of the cluster [Ce₂Li₃(OCH₂^tBu)₉(HOCH₂^tBu)₂(THF)].⁶⁴⁰ Kramer *et al.* reacted the THF adducts of hexamethyl complexes 227-RE with five equivalents [Et₃NH]-[BPh₄] (Scheme 68), affording cationic complexes [RE(Me)-(THF)_n][BPh₄]₂ (233-RE; RE = Sc, Tm, n = 5; RE = Y, Gd–Er, Yb, Lu, n = 6).⁶³⁹ As would be expected, 230 is also an excellent protonolysis reagent and reacts smoothly with HTP^{tBu,Me} (Scheme 68) to give the terminal methyl complex [Yb-(TP^{tBu,Me})(Me)(THF)] (234).⁵⁰³

9.2. α-Silyl-alkyls

RE(III) complexes with α-silyl-alkyl ligands (Table 10) were first reported by Lappert and Pearce; in their seminal report, the authors obtained [RE(CH₂SiMe₃)₃(THF)₂] (235-RE; RE = Sc, Y) from the reaction of RECl₃(THF)_x with Li{CH₂SiMe₃} in a mixture of hexane and ether (Scheme 69, A).⁶⁴¹ This salt elimination method has since been applied to obtain analogous complexes with most of the trivalent REs,^{642–645} with the exception of the larger Lns La–Nd. The larger {CH(SiMe₃)₂}[−] ligand was originally introduced in RE chemistry by Lappert and Barker in 1974, who obtained Sc and Y complexes [RE{CH-

Table 10. Selected α -Silyl-alkyl RE Complexes Used as Starting Materials^a

RE	Ln(CR ₃) ₂	RE(CR ₃) ₃
Sc		[Sc(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{641,656,657} [Sc(CH ₂ SiMe ₂ Ph) ₃ (THF) ₂] ⁶⁵⁸ <i>Sc{CH(SiMe₃)₂}</i> ⁶⁴⁶
Y		[Y(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{641–643,659,660} [Y(CH ₂ SiMe ₃) ₃ (THF) ₃] ⁶⁴⁵ <i>[Y{CH(SiMe₃)₂}]⁵¹</i>
La		<i>La(CH₂SiMe₃)₃(THF)₃</i> ⁶⁶¹ <i>[La{CH(SiMe₃)₂}]⁵¹</i>
Ce		<i>[Ce{CH(SiMe₃)₂}]⁶⁶²</i>
Pr		
Nd		
Sm	[Sm{C(SiMe ₃) ₃ }] ⁶⁶³	[Sm(CH ₂ SiMe ₃) ₃ (THF) ₃] ⁶⁴³ <i>[Sm{CH(SiMe₃)₂}]⁵¹</i>
Eu	[Eu{C(SiMe ₃) ₃ }] ¹¹⁵	
Gd		[Gd(CH ₂ SiMe ₃) ₃ (THF) ₂] ⁶⁶⁴
Tb		[Tb(CH ₂ SiMe ₃) ₃ (THF) ₂] ⁶⁶⁰
Dy		[Dy(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{660,665}
Ho		[Ho(CH ₂ SiMe ₃) ₃ (THF) ₂] ⁶⁶⁰
Er		[Er(CH ₂ SiMe ₃) ₃ (THF) ₂] ⁶⁴³ <i>[Er{CH(SiMe₃)₂}]⁵⁹⁸</i>
Tm		[Tm(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{660,666}
Yb	<i>Yb{CH(SiMe₃)₂}(Et₂O)₂</i> ²⁹⁹ <i>[Yb{C(SiMe₃)₃}]¹¹⁴</i>	[Yb(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{664,667}
Lu		[Lu(CH ₂ SiMe ₃) ₃ (THF) ₂] ^{642,643,656} [Lu(CH ₂ SiMe ₃) ₃ (py) ₂] ⁶⁶⁸ [Lu(CH ₂ SiMe ₃) ₃ (THF)(DME)] ⁶⁶⁹ <i>[Lu{CH(SiMe₃)₂}]⁵⁷³</i>

^aCompounds in italics have not been structurally authenticated.

(SiMe₃)₂}₃] (**210-RE**) *via* salt elimination reactions, though no structural authentication was provided (Scheme 69, A).⁶⁴⁶ However, it has been amply demonstrated that salt elimination reactions can often lead to the formation of “ate” complexes with these systems.⁶⁴⁷ Nonetheless, salt occlusion can be avoided by following the transmetalation route devised by Lappert and co-workers, in which aryloxide complexes [RE(ODb)₃] (**72^{Db}-RE**) are treated with Li{CH(SiMe₃)₂} to give neutral *tris*-alkyl complexes **210-RE** (Scheme 69, B).⁵¹ It is important to note that these ligands are not ideal for stabilizing neutral *bis*-alkyl Ln(II) complexes; Lappert and co-workers obtained [Yb{CH(SiMe₃)₂}₂(OEt₂)₂] (**236**) *via* both analogous salt metathesis and transmetalation methodologies, but no structural information has been reported to validate these results (Scheme 69, C and D).²⁹⁹ It is also noteworthy that other ligand variations have been introduced, particularly the very sterically demanding {C(SiMe₃)₃}[−], which has enabled the stabilization and structural authentication of neutral homoleptic *bis*-alkyl complexes [Ln{C(SiMe₃)₃}₃] (Ln = Sm, Eu, Yb), which were obtained *via* salt elimination reactions.¹¹⁴

The applications of both **235-RE** and **210-RE** as synthetic precursors in protonolysis reactions are numerous and well-established.¹¹ **235-RE** is a particularly useful reagent owing to the production of volatile SiMe₄ as byproduct upon treatment with Brønsted acids. This strategy has been applied to produce complexes with a plethora of supporting ligands, and their utility as synthetic precursors has been extensively reviewed before.¹¹ Therefore, only some representative examples are discussed herein. Okuda and co-workers showed that by treating **235-RE** (RE = Sc, Y, Lu) with [Me₂PhNH][B(C₆F₅)₄] or [Et₃NH][BPh₄] the cationic complexes [RE(CH₂SiMe₃)₂(THF)_n][BR₄]

(**237^{BPh₄}-RE**: R = BPh₄, RE = Sc, Y, Lu; **237^{BARF}-RE**: R = C₆F₅, RE = Y, Lu; n = 3, 4) are produced (Scheme 70).^{648,649} When at least two equivalents of Brønsted acid are employed, the dicationic complexes [RE(CH₂SiMe₃)(THF)_n][BPh₄]₂ (**238-RE**: RE = Y, n = 5; RE = Lu, n = 4) are obtained (Scheme 70).^{648,649} Interestingly, **235-RE** react also with Lewis acids (*e.g.*, BPh₃, B(C₆F₅)₃, AlR₃) to give analogous cationic complexes upon abstraction of one or more alkyl ligands.^{648,649} The efficacy of **235-RE** as a protonolysis reagent has also been demonstrated by Li *et al.* with the reaction of **235-Sc** with a series of substituted cyclopentadienes (HCp, HCp^{Me}, HCp^{tet}, HCp^{*}, and HCp^{*'}) to produce half-sandwich dialkyl complexes [Sc(Cp^R)(CH₂SiMe₃)₂(THF)] [Cp^R = Cp (**239**), Cp^{Me} (**240**), Cp^{tet} (**241**), Cp^{*} (**242**), Cp^{*'} (**243**) - Cp^{tet} = {C₅HMe₄}[−], Cp^{*'} = {C₅Me₄SiMe₃}[−]] (Scheme 70).⁶⁵⁰ This approach can be extended to different REs, as demonstrated by Hou and co-workers with the preparation of **244-RE** (RE = Sc, Y, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu).⁶⁵¹ The use of **210-RE** in similar alkane elimination reactions has also been explored, but these applications are more limited compared to **235-RE** owing to the kinetic inertia provided by the more sterically demanding {CH(SiMe₃)₂}[−] ligand.¹¹ Nonetheless, an important application of these reagents has been demonstrated by Kempe with the preparation of heterobimetallic RE-TM species *via* alkane elimination reactions between RE alkyls and TM hydride precursors.^{652–654} **210-RE** (RE = La, Sm, and Lu) reacts with [RE(Cp)₂(H)] in benzene at room temperature to give the tetrametallic complexes [RE{Re(Cp)₂}₃] (**244-RE**; RE = La, Sm, Lu);⁶⁵⁵ these species are extremely sensitive, which according to the authors is the reason behind the low yields of these reactions. The choice of reagent and solvent conditions in

Table 11. Selected Benzyl Complexes of Divalent Lns and Trivalent REs^a

RE	Ln(Bn) ₂	RE(Bn) ₃
Sc		[Sc(Bn) ₃ (THF) ₂] ⁶⁷⁷ [Sc(Bn) ₃ (THF) ₃] ⁶⁷⁷ <i>Sc(Bn^{NMe2})₃</i> ⁶⁷⁰
Y		[Y(Bn) ₃ (THF) ₃] ^{673,685} [Y(Bn ^{NMe2}) ₃] ⁶⁷¹ [Y(Bn) ₂ (1)(THF) ₃] ⁶⁷¹
La		[La(Bn) ₃ (THF) ₃] ⁶⁷⁵ [La(Bn ^{tBu}) ₃ (THF) ₃] ⁶⁷⁵ <i>La(Bn^{Me})₃(THF)₃</i> ⁶⁷³ [La(Bn ^{NMe2}) ₃] ^{671,690}
Ce		[Ce(Bn) ₃ (THF) ₃] ⁶⁷⁸
Pr		[Pr(Bn) ₃ (THF) ₃] ⁶⁷⁸ <i>Pr(Bn^{NMe2})₃</i> ⁶⁷²
Nd		[Nd(Bn) ₃ (THF) ₃] ⁶⁷⁸ [Nd(Bn ^{NMe2}) ₃] ⁶⁷⁶
Sm	<i>[Sm(Bn)₂]_n</i> ⁶⁸¹ [Sm ₂ (Bn) ₄ (THF)(THP) ₂] _∞ ⁶⁸¹ [Sm{CH(C ₆ H ₄ NMe ₂)(SiMe ₃) ₂ (THF) ₂ }] ⁶⁸⁰	[Sm(Bn) ₃ (THF) ₃] ⁶⁷⁸ [Sm(Bn ^{NMe2}) ₃] ^{672,689}
Eu	<i>[Eu(Bn)₂]_n</i> ⁶⁸¹ [Eu ₄ (Bn) ₈ (THF) ₂] ⁶⁸¹ <i>Eu{CH(C₆H₄NMe₂)(SiMe₃)₂(THF)₂}</i> ⁶⁷³	
Gd		[Gd(Bn) ₃ (THF) ₃] ^{339,678} <i>Gd(Bn^{NMe2})₃</i> ⁶⁷²
Tb		[Dy(Bn) ₃ (THF) ₃] ⁶⁷⁸
Dy		[Dy(Bn ^{NMe2}) ₃] ⁶⁷³
Ho		[Ho(Bn) ₃ (THF) ₃] ³³⁹ [Ho(Bn ^{NMe2}) ₃] ⁶⁷³
Er		[Er(Bn) ₃ (THF) ₂] ³³⁹ [Er(Bn) ₂ (THF) ₃] ^{339,678} [Er(Bn) ₂ (1)(THF) ₃] ⁶⁷⁹
Tm		
Yb	<i>[Yb(Bn)₂]_n</i> ⁶⁸¹ [Yb(Bn) ₂ (DME) ₂] ⁶⁸¹ [Yb(Bn) ₂ (THP) ₄] ⁶⁸¹ [Yb{CH(C ₆ H ₄ NMe ₂)(SiMe ₃) ₂ (THF) ₂ }] ⁶⁸⁰	[Yb(Bn ^{NMe2}) ₃] ⁶⁷²
Lu		[Lu(Bn) ₃ (THF) ₂] ⁶⁷⁷ [Lu(Bn) ₃ (THF) ₃] ⁶⁷⁷ <i>Lu(Bn^{NMe2})₃</i> ⁶⁷²

^aCompounds in italics have not been structurally authenticated.

Kempe's methodology is crucial, as the presence of THF leads to undesired side-reactivity. Because of this, only unsolvated **210-RE** can be employed.

9.3. Benzyls

A range of benzyl (Bn, {CH₂Ar}⁻) complexes have been reported with trivalent REs, which comprise (1) unsubstituted benzyls, (2) substituted benzyls, and (3) bidentate benzyls (Table 11). Compared to other hydrocarbyls, benzyls offer the advantage of acting as multihapto ligands and can delocalize the negative charge from the benzylic position into the phenyl ring; because of this they impart greater stability to resulting RE complexes while still preserving the basicity (or nucleophilicity) of alkyl fragments and are therefore very useful synthetic precursors for protonolysis reactivity. Ln(Bn)₂ and RE(Bn)₃ are obtained *via* salt elimination reactions between LnI₂ or REX₃ and benzylpotassium salts, though occasionally Li salts are also employed. That was indeed the case when Mazer first reported in 1978 the reaction between ScCl₃ and three equivalents of Li(Bn^{NMe2}) (Bn^{NMe2} = {CH₂[C₆H₄(NMe₂)₂]}⁻), leading to

the formation of *tris*-benzyl complex Sc(Bn^{NMe2})₃ (**245-Sc**) (Scheme 71, A).⁶⁷⁰ Workup of the reaction following Mazer's method can be frustrated by the solubility of the byproduct (LiCl), so more convenient routes entail the use of benzylpotassium salts. Harder^{671–673} and then Hou⁶⁷⁴ obtained benzyl complexes **245-RE** (RE = Y, La, Pr, Nd, Sm, Gd, Dy, Ho, Lu) by employing K(Bn^{NMe2}) with either RECl₃ (RE = Y, Nd, Gd, Sm, Dy, Ho, Yb, Lu—both anhydrous and solvated)^{671–673} or [RE(Br)₃(THF)₄] (RE = La, Pr, Nd) (Scheme 71, A).⁶⁷⁴ RE complexes with monodentate benzyl ligands were first reported by Hessen and co-workers, who prepared [La(Bn)₃(THF)₃] (**246-La**) and [La(Bn^{Me})₃(THF)₃] (**247**; Bn^{Me} = {CH₂(C₆H₄Me-4)}⁻) from [La(Br)₃(THF)₄] and KBn^R in THF at 0 °C (Scheme 71, B).⁶⁷⁵ In a similar vein, Harder reacted [La(Br)₃(THF)₄] with K(Bn^{tBu}) (Bn^{tBu} = {CH₂(C₆H₄^tBu-4)}⁻) in THF at -50 °C to give [La(Bn^{tBu})₃(THF)₃] (**248**).⁶⁷⁶ RECl₃ precursors are also a good match with benzylpotassium for the preparation of **246-RE** *via* salt metathesis, as demonstrated by Harder,⁶⁷⁶ Roesky,⁶⁷⁷ and

Table 12. Selected Alkylaluminum Complexes of Divalent Lns and Trivalent REs^a

RE	Ln(AlR ₄) ₂	RE(AlR ₄) ₃
Sc		[Sc(AlMe ₄) ₃]·Al ₂ Me ₆ ⁶³³
Y		Y(AlMe ₄) ₃ ^{631,691}
		[Y(AlMe ₄) ₃]·Al ₂ Me ₆ ⁶⁹¹
La		[La(AlMe ₄) ₃] ⁶⁹²
		[La(AlEt ₄) ₃] ⁷⁰⁴
Ce		[Ce(AlMe ₄) ₃] ^{518,693}
Pr		[Pr(AlMe ₄) ₃] ⁶⁹²
Nd		[Nd(AlMe ₄) ₃] ⁶⁹¹
		[Nd(AlMe ₄) ₃]·Al ₂ Me ₆ ⁶⁹⁴
Sm	<i>[Sm(AlMe₄)₂]_n</i> ^{556,696}	[Sm(AlMe ₄) ₃] ⁶⁹²
	<i>[Sm(AlEt₄)₂]_∞</i> ⁵⁵⁶	
	<i>[Sm(AlEt₄)₂(THF)₂]⁶⁹⁶</i>	
Eu	<i>[Eu(AlMe₄)₂]_n</i> ⁵¹⁸	
	<i>[Eu(AlEt₄)₂]_∞</i> ^{518,705}	
Gd		[Gd(AlMe ₄) ₃] ⁷⁰⁶
Tb		[Tb(AlMe ₄) ₃] ⁷⁰⁶
Dy		[Dy(AlMe ₄) ₃] ⁷⁰⁷
Ho		[Ho(AlMe ₄) ₃] ⁶⁹³
Er		[Er(AlMe ₄) ₃] ⁶⁹³
Tm		[Tm(AlMe ₄) ₃] ⁵¹⁸
Yb	<i>[Yb(AlMe₄)₂]_n</i> ^{556,705}	[Yb(AlMe ₄) ₃] ⁵¹⁸
	<i>[Yb(AlEt₄)₂]_∞</i> ⁷⁰⁵	[Yb(AlMe ₄) ₃]·Al ₂ Me ₆ ⁵¹⁸
	<i>[Yb(AlEt₄)₂(THF)₂]⁶⁹⁶</i>	
Lu		[Lu(AlMe ₄) ₃] ⁶⁹²


^aCompounds in italics have not been structurally authenticated.




Diaconescu.³³⁹ Interestingly, Diaconescu and co-workers observed that **246-Ho** and **246-Er** could be obtained in good yields starting from corresponding RECl₃; however, bromide salts were required to obtain benzyl complexes of larger metals, *i.e.*, Nd and Gd.³³⁹ Liddle and co-workers reported the convenient synthesis of **246-RE** (RE = Y, La–Sm, Gd, Dy, Er) from [RE(I)₃(THF)_n] (**138-RE**) and K(Bn) in THF at 0 °C

(Scheme 71, C).⁶⁷⁸ It is noteworthy that attempts to obtain a Yb(III) analogue led to the isolation of the mixed-valent Yb(II)/Yb(III) complex [Yb(Bn)(THF)₅][Yb(Bn)₄(THF)₂].⁶⁷⁸ Additionally, Liddle and co-workers reported the heteroleptic benzyl complexes [RE(Bn)₂(I)(THF)₃] (**249-RE**; RE = Y, Er) from the reaction of **138-Y** and **138-Er** with two equivalents of K(Bn) (Scheme 71, C).⁶⁷⁹ Finally, another class of benzyl complexes has been reported by Schmidt and Behrle using *N,N*-dimethylbenzylamine (DBA) metalated in the α -position.⁵¹⁶ Analogously to the other methodologies employed for the synthesis of **245-RE** and **246-RE**, Schmidt and Behrle reacted three equivalents of K(DBA) with RECl₃ (RE = Y, La, Ce, Nd, Sm, Gd) in THF at –50 °C (Scheme 71, D), obtaining [RE(DBA)₃] (**250-RE**; RE = Y, La, Ce, Nd, Sm, Gd).⁵¹⁶ X-ray studies of **250-RE** reveal that DBA ligands coordinate the metal centers in an η^4 -fashion.⁵¹⁶

Benzyl derivatives of divalent Lns have been reported by Harder and Anwander. Harder and co-workers obtained Sm, Eu, and Yb benzyls [Ln{CH(C₆H₄NMe₂)(SiMe₃)₂(THF)₂}] (**251-Ln**; Ln = Sm, Eu, Yb) by reacting LnI₂ with K{CH(C₆H₄NMe₂)(SiMe₃)₂} (Scheme 72).⁶⁸⁰ This bidentate benzyl ligand imparts greater stability to the complexes owing to the stabilization of the carbanion by the α -silyl substituent. Recently, Anwander and co-workers have reported the synthesis of [Ln(Bn)₂]_n (**252-Ln**; Ln = Sm, Eu, Yb) from [Ln(I)₂(THF)₂] and two equivalents of K(Bn) (Scheme 72).⁶⁸¹ The amorphous **252-Ln** can be recrystallized in the presence of donors to give discrete molecular species, *i.e.*, [Eu₄(Bn)₈(THF)₂] (**253**), [Sm₂(Bn)₄(THF)(THP)₂]_∞ (**254**), [Yb(Bn)₂(THP)₄] (**252-Yb·4THP**), and [Yb(Bn)₂(DME)₂] (**252-Yb·2DME**).⁶⁸¹

RE benzyl complexes have been used as synthetic precursors in protonolysis reactions with a range of substrates. Hessen and co-workers treated **246** and **247-La** with either one or two equivalents of [MePh₂NH][BPh₄] and obtained the corresponding monocationic or dicationic complex, *i.e.*, [La(Bn)₂(THF)₄][BPh₄]₂ (**255**), [La(Bn^{Me})₂(THF)₄][BPh₄], [La(Bn)(THF)₆][BPh₄]₂ (**256**), and [La(Bn^{Me})(THF)₆][BPh₄]₂

Scheme 66. Synthesis of RE Hydrocarbyls via Salt Elimination,^{635–638,640} Donor-Cleavage,⁶³¹ and Transmetalation Reactions^{12,503}

Scheme 67. Selected Examples of Protonolysis Reactivity of 226-RE with Brønsted Acids^{542,631,634}Scheme 68. Reactivity of Hexamethyl Complexes with [Et₃NH][BPh₄]⁶³⁹ and Synthesis of Terminal Yb Methyl Complex 243 from 230⁵⁰³Scheme 69. Synthesis of α -Silyl Alkyl complexes 235-RE,⁶⁴¹ 210-RE,⁵¹ and 236²⁹⁹

(Scheme 73).⁶⁷⁵ These starting materials can also be used to generate heteroleptic alkyl complexes with bidentate nitrogen donors, such as amidinates (257),⁶⁷⁵ guanidinates (258-RE; RE = Y, La, Dy, Lu),^{682,683} and ferrocenyl *bis*-amides (259-RE; RE = Nd, Gd, Ho, Er) (Scheme 73).³³⁹


Furthermore, RE benzyls can be employed to generate new organometallic complexes. Harder first showed that 245-RE can deprotonate HCp^{BIG} (Cp^{BIG} = {C₅Ar₅}⁻; Ar = C₆H₄R-4; R = Et, ⁱPr, ⁿBu) to afford the heteroleptic Cp derivatives [RE(Cp^{BIG})(Bn^{NMe2})₂] (260-RE; RE = Y, Nd, Dy, Tm).^{672,684} Interestingly, with 245-Sm and 245-Yb the metals are reduced to their divalent state and the homoleptic metallocenes [RE(Cp^{BIG})₂] (261-RE; RE = Sm, Yb) are formed, with concomitant formation of byproduct 1,2-di(2-Me₂N-phenyl)ethane as a result of the coupling of two benzyl radicals (Scheme 74).^{672,684} In addition to this, Liddle and co-workers used 246-RE and 249-RE to perform a double-deprotonation of H₂-BIPM and generate alkylidene complexes [RE(BIPM)(Bn)] (262-RE; RE = Y, Dy, Er),^{678,685} [RE(BIPM)(H-BIPM)] (263-RE; RE = Y, La, Ce, Pr, Nd, Sm, Gd, Tb, Dy),^{678,686,687} and [RE(BIPM)-I](THF)₂] (264-RE; RE = Y, Er) (Scheme 74).^{679,688} Finally, Schmidt and Behrle screened a series of protic substrates (*i.e.*, HODb, H₂NDipp, HN(SiMe₃)₂) with the DBA complexes 250-Y and 250-La and obtained clean conversion to [RE(ODb)₃] (72^{Db}-RE; RE = Y, La), RE(HNDipp)₃, and [RE{N(SiMe₃)₂}₃] (185-RE; RE = Y, La).⁵¹⁶

Applications of Ln(II) benzyl complexes are more scarce compared to their trivalent counterparts. Harder obtained Sm(II) metallocene 261-Sm from the direct deprotonation of HCp^{BIG} with 251-Sm (Scheme 75).⁶⁸⁹ Moreover, Anwender and co-workers reacted 252-Eu and 252-Yb with H₂NDipp (Scheme 75), obtaining a cubane cluster with bridging imido ligands, [Ln(η^3 -NDipp)(THF)]₄] (265-Ln; Ln = Eu, Yb).⁶⁸¹


9.4. Organoaluminates

Since the first report of [Y(AlMe₄)₃] (266-Y) and [Nd(AlMe₄)₃] (266-Nd) by Evans *et al.* in 1995,⁶⁹¹ RE tetraalkylaluminates have emerged as a very interesting class of synthetic precursors in RE and f-element chemistry.¹¹ One of the intriguing aspects of organoaluminates is their dual chemical behavior: on the one hand they can be deemed “masked-alkyl” complexes formed as adducts of AlR₃, which can be used for protonolysis reactivity; on the other hand they could be regarded as ionic complexes of the {AlR₄}⁻ ligand, which can

Scheme 70. Selected Examples of Protonolysis Reactivity of 235-RE with Ammonium Salts^{648,649} and Substituted Cyclopentadienes⁶⁵⁰ and Reactivity of 210-RE with $[\text{Re}(\text{Cp})_2\text{H}]$ ⁶⁵⁵





Scheme 71. Synthesis of RE(III) Benzyl Complexes



act as a *pseudo*-halide in salt elimination reactions. Their preparation and chemical properties were incorporated in the detailed account on RE alkyl chemistry by Anwander and Zimmermann in 2010;¹¹ therefore, this section aims to give a broad overview of their preparation and synthetic applications.




Evans *et al.* prepared $[\text{RE}(\text{AlMe}_4)_3]$ (**266-RE**; $\text{RE} = \text{Y, Nd}$) by treating amide precursor $\text{RE}(\text{NMe}_2)_3(\text{LiCl})_3$ with excess AlMe_3

(Scheme 76, A).⁶⁹¹ This synthetic approach has since been optimized by generating the amide precursors *in situ* from $\text{RECl}_3(\text{THF})_x$ and $\text{Li}(\text{NMe}_2)$, and the methodology has been extended to most of the REs.^{518,692,693} The overall reaction can be viewed as an amide-methyl metathesis generating $\text{Me}_2\text{NAlMe}_2$ and putative methyl complex “ $\text{RE}(\text{Me})_3$ ”, with the latter converted into **266-RE** upon adduct formation with

Scheme 72. Synthesis of Ln(II) Benzyl Complexes 251-Ln,⁶⁸⁰ 252-Ln, 253, and 254⁶⁸¹Scheme 73. Reactivity of RE(III) Benzyl Precursors with Bidentate N-Donors^{339,675,682,683}

excess AlMe_3 .¹¹ The Sc analogue **266-Sc** cannot be obtained *via* amide-methyl exchange (Scheme 76, B); nonetheless, treatment of the hexamethyl Sc complex $\text{Sc}(\mu\text{-Me})_6\text{Li}_3(\text{THF})_{1,2}$ ⁶³⁹ with more than six equivalents of AlMe_3 affords $[\text{Sc}(\text{AlMe}_4)_3] \cdot \text{Al}_2\text{Me}_6$ (**266-Sc**· Al_2Me_6). Organoaluminates obtained with these methods can cocrystallize with Al_2Me_6 , which can be removed upon recrystallization.^{691,692,694} Most of these species are also relatively thermally robust, especially if compared to alkyl derivatives, and can also be sublimed. However, an

important aspect of the chemistry of **266-RE** is the ability of polar solvents to trigger their degradation *via* donor-induced cleavage of the aluminate ligand (*vide infra*).^{11,695} It is possible to obtain **266-RE** also by exchanging other ligands with AlMe_3 , such as different amides and alkoxides, though the purification of the desired organoaluminates is complicated by the low volatility of the $\text{Me}_2\text{Al}(\text{L})$ byproducts ($\text{L} = \text{N}(\text{SiHMe}_2)_2, \text{OCH}_2^t\text{Bu}$).⁶⁹² Additionally, synthesis of **266-Y** *via* direct salt elimination reaction between YCl_3 and $\text{Li}(\text{AlMe}_4)$ has also been reported



Scheme 74. Reactivity of RE(III) Benzyl Precursors with H₂-BIPM^{678,679,685–688} and HCp^{BIG672,684}Scheme 75. Reactivity of Ln(II) Benzyls with HCp^{BIG} and H₂NDipp^{681,689}Scheme 76. Synthesis of RE(III) Aluminate^{691,692,694}

(Scheme 74, C).⁶⁹² Alternatively, homoleptic trimethyl precursors [Y(Me)₃]_n (**226-Y**) and [Lu(Me)₃]_n (**226-Lu**) can be converted into the corresponding organoaluminates by adduct formation with AlMe₃ (Scheme 74, D).^{631,692}

Amide-alkyl or alkoxide-alkyl exchange reactions can be used to access divalent Ln organoaluminates [Ln(AlMe₄)₂]_n (**267-Ln**; Ln = Sm, Yb) and [Ln(AlEt₄)₂]_n (**268-Ln**; Ln = Sm, Yb);^{556,696} these are achieved by reacting *bis*-silylamide precursors **185-Sm** and **185-Yb** (Scheme 77, A) or *bis*-alkoxides Ln(ODipp)₂(THF)_x with AlR₃ (R = Me, Et) (Scheme 77, B).^{556,601,696} Interestingly, **267-Ln** and **268-Ln** do not degrade


in the presence of donors or polar solvents, and the solid-state structures of THF adducts [Ln(AlEt₄)₂(THF)₂] (**269-Ln-2THF**; Ln = Sm, Yb) have also been reported.⁶⁹⁶ **267-Yb** can also be obtained *via* thermally induced self-reduction of **266-Yb**, with concomitant formation of Al₂Me₆ and C₂H₆ (Scheme 77, C).⁵¹⁸ Finally, Eu(III) *tris*-amide **186-Eu-2THF** can be converted into [Eu(AlR₄)₂]_n (**267-Eu** – R = Me; **268-Eu** – R = Et) upon treatment with an excess of AlR₃ (Scheme 77, D).⁵¹⁸

RE organoaluminates (Table 12) are excellent synthetic precursors in protonolysis reactivity with a range of substrates. Anwander demonstrated this with the preparation of Cp

Scheme 77. Synthesis of Divalent Alkylaluminates^{518,696}Scheme 78. Selected Examples of Protonolysis Reactivity of 266-RE and 267-Yb^{697–700}

derivatives $[\text{RE}(\text{Cp}^{\text{R}})(\text{AlMe}_4)_2]$ (**272-RE**; Cp^R = Cp^{*}, RE = Y, La, Nd, and Lu; **273-RE**; Cp^R = Cp^{*}; RE = Y, La, Nd, Sm, Gd, and Lu) *via* deprotonation of the corresponding cyclopentadiene in hexane or pentane under mild conditions (Scheme 78).^{697,698} Lappert and co-workers first showed that heteroleptic Cp-alkylaluminum complexes can undergo donor-induced cleavage of AlMe₃ upon treatment with a Lewis base.⁶⁹⁵ In a similar vein, **272-RE** can be converted to the “unmasked” alkyl analogues $[\text{RE}(\text{Cp}^*)(\text{Me})_2]_3$ (RE = Y, Lu) by addition of THF; the reaction is fully reversible, and **272-RE** is reobtained *via* donor addition reaction with AlMe₃ at −35 °C.⁶⁹⁸ Furthermore, the difference in the bonding character between divalent Ln and trivalent RE aluminates is exemplified by their divergent reactivity with cyclopentadienyls. When divalent **267-Yb** is reacted with HCp^{*}, the {AlMe₄}[−] ligand acts as a weakly coordinating anion and the separated ion pair complex $[\text{Yb}(\text{Cp}^*)(\text{THF})_4][\text{AlMe}_4]$ (**274**) is isolated from the reaction.⁵⁵⁶ Interestingly, complex **274** is obtained even when two equivalents of HCp^{*} are used in the reaction, and the authors detected unreacted pentamethylcyclopentadiene in the reaction mixture, together with AlMe₃(THF) and traces of Cp^{*}AlMe₂.⁵⁵⁶ Zimmermann *et al.* have also shown that proligand C₆NH₃[CH₂NH(Dipp)]₂-2,6 can be deprotonated by **266-RE** (RE = Y, La, Lu), yielding heteroleptic aluminate complex $[\text{RE}(\text{BDPPpyr})(\text{AlMe}_4)]$ (**275**; BDPPpyr = {C₆NH₃[CH₂N(Dipp)]₂-2,6}) (Scheme 78).⁶⁹⁹

Anwander and co-workers have also demonstrated that RE organoaluminates can be employed for salt elimination reactivity with various ligand transfer reagents.¹¹ Piano-stool derivative **272-Sc** is obtained from the 1:1 reaction between K(Cp^{*}) and **266-Sc**, while the metallocene complex $[\text{Sc}(\text{Cp}^*)_2(\text{AlMe}_4)]$ (**276**) is obtained when two equivalents of K(Cp^{*}) are used (Scheme 79).⁶³³ When K(Tp^{tBu,Me}) is reacted with **266-Y**, K(AlMe₃)₄ is eliminated together with one equivalent of AlMe₃ and the mixed alkylaluminum/alkyl complex $[\text{Y}(\text{Tp}^{\text{tBu,Me}})(\text{AlMe}_4)(\text{Me})]$ (**277**) is isolated (Scheme 79),⁷⁰¹ which can also be obtained *via* protonolysis reactivity between HTp^{tBu,Me} and **266-Y**.⁷⁰⁰ However, when **266-La** is used as a starting material, the reaction produces the “Tebbe-like” methylidene complex $[\text{La}(\text{Tp}^{\text{tBu,Me}})(\mu\text{-CH}_2)(\text{AlMe}_3)_2]$ (**278**) (Scheme 79).⁷⁰¹ Salt elimination protocols were also employed by Le Roux *et al.* to produce the piano-stool phospholyl complex $[\text{RE}\{\text{C}_4\text{PMe}_2(\text{SiMe}_3)_2\}(\text{AlMe}_4)_2]$ (**279-RE**; RE = La, Nd) (Scheme 79).⁷⁰² Another example of the interesting reactivity of RE organoaluminates has been shown recently by Barisic *et al.* with the salt metathesis reaction between **266-RE** (RE = Y, Lu) and two equivalents of K(2,4-dtbp) (2,4-dtbp = {CH₂C(^tBu)CHC(^tBu)CH₂}[−]). At room temperature the reaction affords the aluminabenzene complexes $[\text{RE}(2,4\text{-dtbp})(\text{C}_3\text{H}_3\text{AlMe-1-}^t\text{Bu}_2\text{-3,5})]$ (**280-RE**; RE = Y, Lu) (Scheme 75).⁷⁰³ However, when the same reaction is carried out at −40 °C, the *pseudo*-metallocenes $[\text{RE}(2,4\text{-dtbp})_2(\text{AlMe}_4)]$ (**281-RE**; RE = Y, Lu)

Scheme 79. Selected Examples of Salt Elimination Reactivity of 266-RE^{701–703}

are obtained instead, which can undergo thermal decomposition to give aluminabenzene derivatives **280-RE**.⁷⁰³

10. CONCLUSIONS AND FUTURE PERSPECTIVES

The landscape of RE and Ln synthetic chemistry has grown immensely since the first adventurous steps taken more than half a century ago by various pioneers of the discipline. This impressive growth has been supported by the enormous expansion of the RE synthetic toolbox and the opening of a myriad of synthetic possibilities, the potential of which is still far from being fulfilled. Nonetheless, there is still progress to be made in the development of synthetic precursors and methodologies applicable to divalent Lns outside the Sm, Eu, and Yb triad, which is exemplified by the relatively small number of synthetic applications compared to the trivalent counterparts.⁴² Similarly, the landscape of molecular Ln(IV) chemistry—historically limited to Ce(IV)—has been recently expanded with the stabilization of the first molecular Pr(IV) and Tb(IV) species,^{708–713} as a result, new advances will likely emerge for the preparation of specific starting materials and methodologies applicable to tetravalent Ln chemistry. Recently, Evans and Daly have demonstrated that mechanochemical synthesis can be used for the preparation of RE derivatives using salt elimination methodologies, thus offering a viable alternative to standard solution methods.^{714–716} Additionally, direct activation of metals is a synthetic technique that is still used by only a few groups around the world, largely because of the historic involvement of toxic Hg and Tl reagents; however, these methods have now been extended to more benign metals (Bi and Ag),¹⁷⁵ which should encourage research teams to incorporate these methods into their synthetic repertoire.

Many of the limitations that historically frustrated the progress of RE synthetic chemistry are gradually being challenged and overcome; an example of this is the successful stabilization and identification of the once elusive “RE(Me)₃” species by Anwander and co-workers,^{503,631,633} though this still has not been achieved with the larger Lns.⁷¹⁷ Finally, several of the synthetic precursors presented in this work have emerged over the past decade; therefore, because of these recent discoveries, there will certainly be many new and exciting synthetic avenues that will be opened-up in the years to come.

AUTHOR INFORMATION

Corresponding Author

Fabrizio Ortu – School of Chemistry, University of Leicester, LE1 7RH Leicester, U.K.; orcid.org/0000-0002-1743-8338; Email: fabrizio.ortu@leicester.ac.uk

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.chemrev.1c00842>

Notes

The author declares no competing financial interest.

Biography

Fabrizio Ortu is from Sassari (Sardinia, IT) and obtained his Ph.D. from the University of Nottingham in 2014, where he studied s-block organometallic chemistry under the supervision of Prof. Deborah Kays. After his Ph.D. studies he moved to the University of Manchester to work with Prof. David Mills as a postdoctoral researcher, covering a range of topics in rare earth and actinide synthetic chemistry. In June 2019 he was appointed to a lectureship at the University of Leicester.

His research interests lie in the coordination and organometallic chemistry of alkaline earth and rare earth metals.

ACKNOWLEDGMENTS

I acknowledge the School of Chemistry, University of Leicester for financial support and the Engineering and Physical Sciences Research Council for a New Investigator Award (EP/W00691X/1). Also, I thank Prof. David Mills and Dr. Conrad Goodwin for all the constructive and insightful discussions on RE synthetic chemistry we have had over the years, which provided the inspiration for this work.

ABBREVIATIONS

Ad	adamantyl
AE	alkaline Earth
An	actinide
Ar	aryl
BDI	β -diketiminato
BDPPpyr	$[\text{C}_6\text{NH}_3\{\text{CH}_2\text{N}(\text{Dipp})\}_2\text{-2,6}]^-$
BIPM-H ₂	bis(iminophosphorano)methane
BIPM-H	bis(iminophosphorano)methanide
BIPM-	bis(iminophosphorano)methanediide
bipy	2,2'-bipyridine
Bn	benzyl, $\{\text{CH}_2\text{Ph}\}^-$
Bn ^{Me}	$\{\text{CH}_2(\text{C}_6\text{H}_4\text{Me-4})\}^-$
Bn ^{tBu}	$\{\text{CH}_2(\text{C}_6\text{H}_4\text{tBu-4})\}^-$
Bn ^{NMe2}	$\{\text{CH}_2(\text{C}_6\text{H}_4(\text{NMe}_2)\text{-2})\}^-$
CAN	ceric ammonium nitrate
Carb	carbazole, $\{\text{C}_{12}\text{H}_8\text{N}\}^-$
COT	cyclooctatetraenyl, $\{\text{C}_8\text{H}_8\}^-$
Cp	cyclopentadienyl, $\{\text{C}_5\text{H}_5\}^-$
Cp ^{BIG}	$\{\text{C}_5\text{Ar}_5\}^-$ (Ar = C ₆ H ₄ R-4; R = Et, ⁱ Pr, ⁿ Bu)
Cp ^{iPr4}	$\{\text{C}_5\text{H}^i\text{Pr}_4\}^-$
Cp ^{iPr5}	$\{\text{C}_5\text{H}^i\text{Pr}_5\}^-$
Cp ^{Ph5}	$\{\text{C}_5\text{H}^i\text{Ph}_5\}^-$
Cp [']	$\{\text{C}_5\text{H}_4\text{SiMe}_3\}^-$
Cp ^{''}	$\{\text{C}_5\text{H}_3(\text{SiMe}_3)_2\}^-$
Cp ^{'''}	$\{\text{C}_5\text{H}_2(\text{SiMe}_3)_3\}^-$
Cp [*]	$\{\text{C}_5\text{Me}_5\}^-$
Cp ^{tet}	$\{\text{C}_5\text{HMe}_4\}^-$
Cp ^t	$\{\text{C}_5\text{H}_4\text{tBu}\}^-$
Cp ^{tt}	$\{\text{C}_5\text{H}_3\text{tBu}_2\}^-$
Cp ^{ttt}	$\{\text{C}_5\text{H}_2\text{tBu}_3\}^-$
DF	2,6-difluorophenyl, C ₆ H ₃ F ₂ -2,6
Db	2,6-di- <i>tert</i> -butylphenyl, C ₆ H ₃ ^t Bu ₂ -2,6
DBA	<i>N,N</i> -dimethylbenzylamine
Dbmp	2,6-di- <i>tert</i> -butyl-4-methylphenyl, C ₆ H ₂ ^t Bu ₂ -2,6-Me-4
Dipp	2,6-di- <i>iso</i> -propylphenyl, C ₆ H ₃ ⁱ Pr ₂ -2,6
Dipp-Bian	1,2-bis[(2,6-diisopropylphenyl)imino]-acenaphthene
Dmp	C ₆ H ₃ Mes ₂ -2,6
dpa	2,2'-dipyridylamide, $\{(\text{C}_6\text{H}_4\text{N})_2\text{N}\}^-$
Dpp	2,6-diphenylphenyl, C ₆ H ₃ Ph ₂ -2,6
2,4-dtbp	$\{\text{CH}_2\text{C}(\text{tBu})\text{CHC}(\text{tBu})\text{CH}_2\}^-$
EPR	electron paramagnetic resonance
Form	formamidate
HMPA	hexamethylphosphoramide
Htp	$\{\text{C}_4\text{H}_2\text{P}^t\text{Bu}_2\text{-2,5}\}^-$
IDipp	1,3-bis(Dipp) ₂ imidazolin-2-ylidene
IMe ₂	1,3-bis(Me) ₂ imidazolin-2-ylidene
IMes	1,3-bis(Mes) ₂ imidazolin-2-ylidene

Ln	lanthanoid (La–Lu)
Mes	2,4,6-trimethylphenyl
MVS	metal vapor synthesis
OTf	triflate, $\{\text{O}_3\text{SCF}_3\}^-$
Ph ₂ pz	$\{3,5\text{-Ph}_2\text{C}_3\text{HN}_2\}^-$
Phen	phenantroline
Phpip	<i>N</i> -phenylpiperazine
pyr	pyrrolide, $\{\text{C}_4\text{H}_4\text{N}\}^-$
pz	pyrazolate, $\{\text{C}_3\text{H}_3\text{N}_2\}^-$
RE	rare earth (Sc, Y, La–Lu)
RT	redox transmetalation
RTP	redox transmetalation protonolysis/protolysis
THF	tetrahydrofuran
TM	transition metal
TMG	tetramethylguanidinate
Tph	C ₆ H ₄ Tripp-2
Tripp	2,4,6-triisopropylphenyl

REFERENCES

- (1) Fischer, E. O. On the Way to Carbene and Carbyne Complexes. In *Adv. Organomet. Chem.*, Vol. 14; Stone, F. G. A., West, R., Eds.; Academic Press, 1976; pp 1–32.
- (2) Wilkinson, G.; Birmingham, J. M. Cyclopentadienyl Compounds of Sc, Y, La, Ce and Some Lanthanide Elements. *J. Am. Chem. Soc.* **1954**, *76*, 6210.
- (3) von Frankland, E. Notiz Über Eine Neue Reihe Organischer Körper, Welche Metalle, Phosphor u. s. w. Enthalten. *Justus Liebigs Ann. Chem.* **1849**, *71*, 213–216.
- (4) Cotton, F. A. Alkyls and Aryls of Transition Metals. *Chem. Rev.* **1955**, *55*, 551–594.
- (5) Cotton, S. A. Aspects of the Lanthanide-Carbon σ -Bond. *Coord. Chem. Rev.* **1997**, *160*, 93–127.
- (6) Edelmann, F.; Scandium, T. Yttrium, and the Lanthanide and Actinide Elements, Excluding Their Zero Oxidation State Complexes. *Compr. Organomet. Chem. II* **1995**, 11–212.
- (7) Ortu, F.; Mills, D. P. Low Coordinate Rare Earth and Actinide Complexes. In *Handbook of the Physics and Chemistry of Rare Earths*, Vol. 55; Bünzli, J.-C., Pecharsky, V. K., Ed.; Elsevier B.V., 2019; pp 1–87.
- (8) Anwander, R. Principles in Organolanthanide Chemistry. In *Lanthanides: Chemistry and Use in Organic Synthesis*; Kobayashi, S., Ed.; Springer: Berlin, Heidelberg, 1999; pp 1–61.
- (9) Edelmann, F. T. Lanthanides and Actinides. In *Synthetic Methods of Organometallic and Inorganic Chemistry*, Vol. 6; Thieme: Stuttgart, 1997; pp 1–143.
- (10) Mills, D. P.; Liddle, S. T. Ligand Design in Modern Lanthanide Chemistry. In *Ligand Design in Metal Chemistry: Reactivity and Catalysis*; Stradiotto, M., Lundgren, R. J., Eds.; John Wiley & Sons, Ltd, 2016; pp 330–363.
- (11) Zimmermann, M.; Anwander, R. Homoleptic Rare-Earth Metal Complexes Containing Ln-C σ -Bonds. *Chem. Rev.* **2010**, *110*, 6194–6259.
- (12) Guo, Z.; Huo, R.; Tan, Y. Q.; Blair, V.; Deacon, G. B.; Junk, P. C. Syntheses of Reactive Rare Earth Complexes by Redox Transmetalation/Protolysis Reactions—A Simple and Convenient Method. *Coord. Chem. Rev.* **2020**, *415*, 213232.
- (13) Cloke, F. G. N. Zero Oxidation State Compounds of Scandium, Yttrium, and the Lanthanides. *Chem. Soc. Rev.* **1993**, *22*, 17–24.
- (14) Taylor, M. D. Preparation of Anhydrous Lanthanide Halides. *Chem. Rev.* **1962**, *62*, 503–511.
- (15) Meyer, G. The Divalent State in Solid Rare Earth Metal Halides. In *Encyclopedia of Inorganic and Bioinorganic Chemistry*; John Wiley & Sons, Ltd., 2012; pp 1–13.
- (16) Marks, T. J.; Kolb, J. R. Covalent Transition Metal, Lanthanide, and Actinide Tetrahydroborate Complexes. *Chem. Rev.* **1977**, *77*, 263–293.

- (17) Ephritikhine, M. Synthesis, Structure, and Reactions of Hydride, Borohydride, and Aluminohydride Compounds of the f-Elements. *Chem. Rev.* **1997**, *97*, 2193–2242.
- (18) Makhaev, V. D. Structural and Dynamic Properties of Tetrahydroborate Complexes. *Russ. Chem. Rev.* **2000**, *69*, 727–746.
- (19) Visseaux, M.; Bonnet, F. Borohydride Complexes of Rare Earths, and Their Applications in Various Organic Transformations. *Coord. Chem. Rev.* **2011**, *255*, 374–420.
- (20) Guillaume, S. M.; Maron, L.; Roesky, P. W. Catalytic Behavior of Rare-Earth Borohydride Complexes in Polymerization of Polar Monomers. In *Handbook of the Physics and Chemistry of Rare Earths*, Vol. 44; Bünzli, J.-C., Pecharsky, V. K., Ed.; Elsevier B.V., 2014; pp 1–86.
- (21) Paskevicius, M.; Jepsen, L. H.; Schouwink, P.; Černý, R.; Ravnsbæk, D. B.; Filinchuk, Y.; Dornheim, M.; Besenbacher, F.; Jensen, T. R. Metal Borohydrides and Derivatives-Synthesis, Structure and Properties. *Chem. Soc. Rev.* **2017**, *46*, 1565–1634.
- (22) Anwander, R. Lanthanide Amides. In: *Organolanthanoid Chemistry: Synthesis, Structure, Catalysis. Topics in Current Chemistry*, Vol. 179; Springer: Berlin, 1996; pp 33–112.
- (23) Lappert, M. F.; Power, P. P.; Protchenko, A. V.; Sebert, A. Amides of the Group 3 and Lanthanide Metals. In *Metal Amide Chemistry*; John Wiley & Sons, Ltd., 2008; pp 79–120.
- (24) Goodwin, C. A. P.; Mills, D. P. Silylamides: Towards a Half-Century of Stabilising Remarkable f-Element Chemistry. In *SPR Organomet. Chem.*, Vol. 41; RSC Publishing: Cambridge, 2017; pp 123–156.
- (25) Bradley, D. C.; Mehrotra, R. C.; Gaur, D. P. *Metal Alkoxides*; Academic Press, Inc.: London, 1978; pp 1–411.
- (26) Mehrotra, R. C.; Singh, A.; Tripathi, U. M. Recent Advances in Alkoxo and Aryloxo Chemistry of Scandium, Yttrium, and Lanthanoids. *Chem. Rev.* **1991**, *91*, 1287–1303.
- (27) Bradley, D. C.; Mehrotra, R. C.; Rothwell, I. P.; Singh, A. *Alkoxo and Aryloxo Derivatives of Metals*; Elsevier Ltd: Amsterdam, 2001; pp 1–704.
- (28) Anwander, R. Routes to monomeric lanthanide alkoxides. In *Organolanthanoid Chemistry: Synthesis, Structure, Catalysis. Topics in Current Chemistry*, Vol. 179; Springer: Berlin, 1996; pp 149–245.
- (29) Boyle, T. J.; Ottley, L. A. M. Advances in Structurally Characterized Lanthanide Alkoxide, Aryloxo, and Silyloxo Compounds. *Chem. Rev.* **2008**, *108*, 1896–1917.
- (30) Parmar, V.; Gransbury, G. K.; Whitehead, G. F. S.; Mills, D. P.; Winpenny, R. E. P. Slow Magnetic Relaxation in Distorted Tetrahedral Dy(III) Aryloxo Complexes. *Chem. Commun.* **2021**, *57*, 9208–9211.
- (31) Lawrance, G. A. Coordinated Trifluoromethanesulfonate and Fluorosulfate. *Chem. Rev.* **1986**, *86*, 17–33.
- (32) Davidson, P. J.; Lappert, M. F.; Pearce, R. Metal σ -Hydrocarbyls, MR_n , Stoichiometry, Structures, Stabilities, and Thermal Decomposition Pathways. *Chem. Rev.* **1976**, *76*, 219–242.
- (33) Bochkarev, M. N.; Kalinina, G. S.; Bochkarev, L. N. Advances in the Chemistry of Organolanthanides. *Russ. Chem. Rev.* **1985**, *54*, 802–816.
- (34) Schumann, H.; Meese-Marktscheffel, J. A.; Esser, L. Synthesis, Structure, and Reactivity of Organometallic π -Complexes of the Rare Earths in the Oxidation State Ln^{3+} with Aromatic Ligands. *Chem. Rev.* **1995**, *95*, 865–986.
- (35) Deacon, G. B.; Forsyth, C. M.; Nickel, S. Bis(Pentafluorophenyl) Mercury - A Versatile Synthon in Organo-, Organooxo-, and Organoamido-Lanthanoid Chemistry. *J. Organomet. Chem.* **2002**, *647*, 50–60.
- (36) Edelman, F. T.; Freckmann, D. M. M.; Schumann, H. Synthesis and Structural Chemistry of Non-Cyclopentadienyl Organolanthanide Complexes. *Chem. Rev.* **2002**, *102*, 1851–1896.
- (37) Lyubov, D. M.; Trifonov, A. A. Ln(II) Alkyl Complexes: From Elusive Exotics to Catalytic Applications. *Inorg. Chem. Front.* **2021**, *8* (12), 2965–2986.
- (38) Izod, K. Alkyl, Carbonyl and Cyanide Complexes of the Group 3 Metals and Lanthanides. In *Compr. Organomet. Chem. IV*; Elsevier Ltd.: Amsterdam, 2021; pp 1–55.
- (39) Evans, W. J. The Organometallic Chemistry of the Lanthanide Elements in Low Oxidation States. *Polyhedron* **1987**, *6*, 803–835.
- (40) Anwander, R.; Herrmann, W. A. Features of Organolanthanide Complexes. In *Top. Current Chem.*, Vol. 179; Herrmann, W. A., Ed.; Springer-Verlag: Berlin, 2005; pp 1–32.
- (41) Liddle, S. Lanthanides: Organometallic Chemistry. In *Encyclopedia of Inorganic and Bioinorganic Chemistry*; John Wiley & Sons, Ltd., 2012; pp 1–21.
- (42) Nicholas, H. M.; Mills, D. P. Lanthanides: Divalent Organometallic Chemistry. In *Encyclopedia of Inorganic and Bioinorganic Chemistry*; John Wiley & Sons, Ltd., 2017; pp 1–10.
- (43) Layfield, R. A. Lanthanides. In *Compr. Organomet. Chem. III*; Elsevier Ltd.: Amsterdam, 2021; pp 418–470.
- (44) *The Rare Earth Elements: Fundamentals and Applications*; Atwood, D. A., Ed.; Wiley: New York, 2012; pp 1–624.
- (45) Kapustinskii, A. F. Lattice Energy of Ionic Crystals. *Q. Rev. Chem. Soc.* **1956**, *10*, 283–294.
- (46) Kaya, S. Relationships between Lattice Energies of Inorganic Ionic Solids. *Phys. B Condens. Matter* **2018**, *538*, 25–28.
- (47) Lovinger, G. J.; Aparece, M. D.; Morken, J. P. Pd-Catalyzed Consecutive Cross-Coupling between Grignard-Derived Boron “Ate” Complexes and $C(sp^2)$ Halides or Triflates: NaOTf as a Grignard Activator and Halide Scavenger. *J. Am. Chem. Soc.* **2017**, *139*, 3153–3160.
- (48) Bounioux, C.; Bar-Hen, A.; Yerushalmi-Rozen, R. Salting-in Effect in Organic Dispersions of Poly(3-Hexyl Thiophene)-Carbon-Nanotubes. *Chem. Commun.* **2015**, *51*, 6343–6345.
- (49) Bloor, E. G.; Kidd, R. G. Solvation of Sodium Ion Studied by ^{23}Na Nuclear Magnetic Resonance. *Can. J. Chem.* **1968**, *46*, 3425–3430.
- (50) Arkhipov, S. M.; Mikheeva, V. I.; Pruntsev, A. E. Joint Solubility of Lithium and Aluminium Bromides and Iodides in Ether and Toluene at 25° . *Bull. Acad. Sci. USSR Div. Chem. Sci.* **1975**, *24*, 1550–1552.
- (51) Hitchcock, P. B.; Lappert, M. F.; Smith, R. G.; Bartlett, R. A.; Power, P. P. Synthesis and Structural Characterisation of the First Neutral Homoleptic Lanthanide Metal(III) Alkyls: $[LnR_3]$ [$Ln = La$ or Sm , $R = CH(SiMe_3)_2$]. *J. Chem. Soc., Chem. Commun.* **1988**, 1007–1009.
- (52) *Determination of Organic Structures by Physical Methods*, 1st ed.; Nachod, F. C., Braude, E. A., Eds.; Academic Press, Inc.: New York, 1955; pp 1–824.
- (53) Joannis, A. Combinaisons Du Potassium et Du Sodium Avec Le Gaz Ammoniac. *Compt. Rend.* **1889**, *109*, 900–903.
- (54) Johnson, W. C.; Fernelius, W. C. Liquid Ammonia as a Solvent and the Ammonia System of Compounds. IV. Experimental Procedures Involved in the Manipulation of Liquid Ammonia Solutions. *J. Chem. Educ.* **1929**, *6*, 441–450.
- (55) Wayda, A. L.; Dye, J. L. A Versatile System for Vacuum-Line Manipulations. *J. Chem. Educ.* **1985**, *62*, 356–359.
- (56) Mentrel, B. Sure Le Baryum-Ammonium et l’amidure de Baryum. *Compt. Rend.* **1902**, *135*, 740–742.
- (57) Roederer, M. Sure Le Strontium Ammonium. *Compt. Rend.* **1905**, *140*, 1252–1253.
- (58) Moissan, H. Préparation Du Lithium-Ammonium, Du Calcium-Ammonium et Des Amidures de Lithium et de Calcium. *Compt. Rend.* **1898**, *127*, 685–693.
- (59) Moissan, H. Préparation et Propriétés d’un Ammonium Organique: Le Lithium-Monométhylammonium. *Compt. Rend.* **1899**, *128*, 26–30.
- (60) Moissan, H. Action de l’acetylene Sur Les Metaux-Ammoniums. *Compt. Rend.* **1898**, *127*, 911–917.
- (61) Kraus, C. A. Solutions of Metals in Non-Metallic Solvents; I. General Properties of Solutions of Metals in Liquid Ammonia. *J. Am. Chem. Soc.* **1907**, *29*, 1557–1571.
- (62) Kraus, C. A. Solutions of Metals in Non-Metallic Solvents; III. The Apparent Molecular Weight of Sodium Dissolved in Liquid Ammonia. *J. Am. Chem. Soc.* **1908**, *30*, 1197–1219.
- (63) Kraus, C. A. Solution of Metals in Non-Metallic Solvents; II. On the Formation of Compounds between Metals and Ammonia. *J. Am. Chem. Soc.* **1908**, *30*, 653–668.

- (64) Cottrell, F. G. Some Reactions of Liquid Anhydrous Ammonia and Acetylene. *J. Phys. Chem.* **1914**, *18*, 85–100.
- (65) Watt, G. W.; Jenkins, W. A.; McCuiston, J. M. Reactions of Some Thorium and Uranium Compounds in Liquid Ammonia. *J. Am. Chem. Soc.* **1950**, *72*, 2260–2262.
- (66) Warf, J. C.; Korst, W. L. Solutions of Europium and Ytterbium Metals in Liquid Ammonia. *J. Phys. Chem.* **1956**, *60* (11), 1590–1591.
- (67) Shannon, R. D. Revised Effective Ionic Radii and Systematic Studies of Interatomic Distances in Halides and Chalcogenides. *Acta Crystallogr. Sect. A Found. Crystallogr.* **1976**, *A32*, 751–767.
- (68) Harder, S. From Limestone to Catalysis: Application of Calcium Compounds as Homogeneous Catalysts. *Chem. Rev.* **2010**, *110*, 3852–3876.
- (69) Catterall, R.; Symons, M. C. R. Nature of the Ion Pair or Monomer Species in Metal-Ammonia Solutions. *J. Chem. Phys.* **1965**, *42*, 1466.
- (70) Symons, M. C. R. Solutions of Metals: Solvated Electrons. *Chem. Soc. Rev.* **1976**, *5*, 337–358.
- (71) Thompson, D. S.; Schaefer, D. W.; Waugh, J. S. Electronic Spectra of Solutions of Europium and Ytterbium in Liquid Ammonia. *Inorg. Chem.* **1966**, *5*, 325–326.
- (72) Peer, W. J.; Lagowski, J. J. Metal-Ammonia Solutions. 13. Solutions of Lanthanoid Metals. *J. Phys. Chem.* **1980**, *84*, 1110–1119.
- (73) Hadenfeldt, C.; Jacobs, H.; Juza, R. Über Die Amide Des Europiums Und Ytterbiums. *Z. Anorg. Allg. Chem.* **1970**, *379*, 144–156.
- (74) Stühr, A.; Jacobs, H.; Juza, R. Darstellung Und Kristallstruktur von $\text{Na}[\text{Yb}(\text{NH}_2)_4]$. *Z. Anorg. Allg. Chem.* **1973**, *398*, 1–14.
- (75) Rabenau, A. The Role of Hydrothermal Synthesis in Preparative Chemistry. *Angew. Chem., Int. Ed.* **1985**, *24*, 1026–1040.
- (76) Young, D. M.; Schimek, G. L.; Kolis, J. W. Synthesis and Characterization of $[\text{Yb}(\text{NH}_3)_8][\text{Cu}(\text{S}_4)_2] \cdot \text{NH}_3$, $[\text{Yb}(\text{NH}_3)_8][\text{Ag}(\text{S}_4)_2] \cdot 2\text{NH}_3$, and $[\text{La}(\text{NH}_3)_9][\text{Cu}(\text{S}_4)_2]$ in Supercritical Ammonia: Metal Sulfide Salts of the First Homoleptic Lanthanide Ammine Complexes. *Inorg. Chem.* **1996**, *35*, 7620–7625.
- (77) Quitmann, C. C.; Müller-Buschbaum, K. $[\text{Sm}(\text{NH}_3)_9][\text{Sm}(\text{Pyr})_6]$, Ein Komplexes Salz Mit Zwei Homoleptischen Ionen Aus Der Synthese in Flüssigem Ammoniak. *Z. Anorg. Allg. Chem.* **2005**, *631* (2–3), 564–568.
- (78) Müller, T. G.; Karau, F.; Schnick, W.; Kraus, F. A New Route to Metal Azides. *Angew. Chem., Int. Ed.* **2014**, *53*, 13695–13697.
- (79) Müller, T. G.; Mogk, J.; Conrad, M.; Kraus, F. Octaammine Eu^{II} and Yb^{II} Azides and Their Thermal Decompositions to the Nitrides. *Eur. J. Inorg. Chem.* **2016**, *2016*, 4162–4169.
- (80) Salot, S.; Warf, J. C. Chemical Generation of the Ammoniated Electron via Ytterbium(II). *J. Am. Chem. Soc.* **1968**, *90*, 1932–1933.
- (81) Howell, J. K.; Pytlewski, L. L. Synthesis of Divalent Europium and Ytterbium Halides in Liquid Ammonia. *J. Less-Common Met.* **1969**, *18*, 437–439.
- (82) Tilley, T. D.; Boncella, J. M.; Berg, D. J.; Burns, C. J.; Andersen, R. A. Bis[Bis(Trimethylsilyl)Amido[Bis(Diethylether)Ytterbium and (Diethyl Ether)Bis(η^5 -Pentamethylcyclopentadienyl)Ytterbium. *Inorg. Synth.* **1990**, *27*, 146–150.
- (83) Fischer, E. O.; Fischer, H. Über Dicyclopentadienyleuropium Und Dicyclopentadienylytterbium Und Tricyclopentadienyle Des Terbiums, Holmiums, Thuliums Und Lutetiums. *J. Organomet. Chem.* **1965**, *3*, 181.
- (84) Wayda, A. L.; Dye, J. L.; Rogers, R. D. Divalent Lanthanoid Synthesis in Liquid Ammonia. 1. The Synthesis and x-Ray Crystal Structure of $(\text{C}_5\text{Me}_5)_2\text{Yb}(\text{NH}_3)(\text{THF})$. *Organometallics* **1984**, *3* (11), 1605–1610.
- (85) Hayes, R. G.; Thomas, J. L. Synthesis of Cyclooctatetraenyleuropium and Cyclooctatetraenylytterbium. *J. Am. Chem. Soc.* **1969**, *91*, 6876.
- (86) Murphy, E.; Toogood, G. E. The Preparation of Europium(II) Propynide. *Inorg. Nucl. Chem. Lett.* **1971**, *7*, 755–759.
- (87) Wayda, A. L.; Mukerji, I.; Dye, I. J. L.; Rogers, R. D. Divalent Lanthanoid Synthesis In Liquid Ammonia: 2: The Synthesis and X-Ray Crystal Structure of $(\text{C}_8\text{H}_8)\text{Yb}(\text{C}_5\text{H}_5\text{N})_3 \cdot 1/2\text{C}_5\text{H}_5\text{N}$. *Organometallics* **1987**, *6* (6), 1328–1332.
- (88) Feistel, G. R.; Mathai, T. P. A New Type of Intramolecular Antiferromagnetism. *J. Am. Chem. Soc.* **1968**, *90*, 2988–2989.
- (89) Pappalardo, R. Absorption Spectra of Neutral Dipyrindyl Complexes. *Inorg. Chim. Acta* **1968**, *2*, 209–215.
- (90) Cetinkaya, B.; Hitchcock, P. B.; Lappert, M. F.; Smith, R. G. The First Neutral, Mononuclear 4f Metal Thiolates and New Methods for Corresponding Aryl Oxides and Bis(Trimethylsilyl)Amides. *J. Chem. Soc. Chem. Commun.* **1992**, 932–934.
- (91) Deacon, G. B.; Hitchcock, P. B.; Holmes, S. A.; Lappert, M. F.; Mackinnon, P.; Newnham, R. H. Four- and Five-Co-Ordinate Lanthanide(II) Aryloxides: X-Ray Structures of the Bis(2,6-Di-*t*-Butyl-4-Methylphenoxy)Ytterbium(II) Complexes $[\text{Yb}(\text{OAr})_2(\text{L})_2]$ and $[\text{Yb}(\text{OAr})_2(\text{L}')_3]$ [Ar = $\text{C}_6\text{H}_3\text{Bu}^t_2$ -2,6-Me-4, L = Tetrahydrofuran (thf) or OEt_2 , L' = thf]. *J. Chem. Soc. Chem. Commun.* **1989**, 935–937.
- (92) Evans, W. J.; Greci, M. A.; Ziller, J. W. Substituent Effects in the Formation of Aryloxy-Bridged Europium Complexes. *J. Chem. Soc., Dalton Trans.* **1997**, 3035–3039.
- (93) Quitmann, C. C.; Müller-Buschbaum, K. Low-Temperature Oxidation of Ytterbium with Pyrrole in Liquid Ammonia. Formation of the First Unsubstituted Pyrroles of a Rare Earth Element. *Z. Anorg. Allg. Chem.* **2004**, *630*, 2422–2430.
- (94) Müller-Buschbaum, K.; Zurawski, A. On the Mechanisms of Electride Induced Synthesis of Ytterbium Carbazoles, Formation of Coordination Polymers by Condensation and Polymer Degradation by Chemical Scissors. *Z. Anorg. Allg. Chem.* **2007**, *633*, 2300–2304.
- (95) Müller-Buschbaum, K.; Quitmann, C. C.; Zurawski, A. $2\infty[\text{Yb}_2(\text{NH}_2)_2(\text{Pz})_4][\text{Yb}(\text{NH}_3)_2(\text{Pz})_3\text{PzH}]$: Electride Induced Synthesis of a 2D-Ytterbium-Pyrazolate Network. *Monat. Chem.* **2007**, *138*, 813–817.
- (96) Müller-Buschbaum, K. Tieftemperatur-Oxidation in Flüssigem Ammoniak: $[\text{Eu}_2(\text{Ind})_4(\text{NH}_3)_6]$, Das Erste Indolat Eines Selten-Erd-Elementes. *Z. Anorg. Allg. Chem.* **2004**, *630*, 895–899.
- (97) Müller-Buschbaum, K. $[\text{Yb}(\text{NH}_3)_8][\text{Yb}(\text{Pyr})_6]$ - Elektrid-Induzierte Synthese Und Kristallisation Aus Flüssigem Ammoniak. *Z. Anorg. Allg. Chem.* **2007**, *633*, 1403–1406.
- (98) Müller-Buschbaum, K. The Utilization of Solid State Chemistry Reaction Routes as New Syntheses Strategies for the Coordination Chemistry of Rare Earth Amides. *Z. Anorg. Allg. Chem.* **2005**, *631*, 811–828.
- (99) Quitmann, C. C.; Müller-Buschbaum, K. Complete Nitrogen Coordination in Rare Earth Bipyridine Pyrrolate Complexes from Pyrrole under Solvothermal Conditions: $[\text{Ln}(\text{Pyr})_3(\text{Bipy})_2]$, Ln = Pr, Yb. *Z. Anorg. Allg. Chem.* **2005**, *631*, 2651–2654.
- (100) Quitmann, C. C.; Müller-Buschbaum, K. $[\text{Yb}_3\text{N}(\text{Dpa})_6][\text{Yb}(\text{Dpa})_3]$: A Molecular Nitride of a Rare-Earth Metal with a Yb_3N Unit. *Angew. Chem., Int. Ed.* **2004**, *43* (44), 5994–5996.
- (101) Beckmann, E. Einige Anwendungen von Metallischem Calcium. *Ber. Dtsch. Chem. Ges.* **1905**, *38*, 904.
- (102) Gilman, H.; Schulze, F. Organocalcium Iodides. *J. Am. Chem. Soc.* **1926**, *48*, 2463–2467.
- (103) Bryce-Smith, D.; Skinner, A. C. Organometallic Compounds of Group II. Part IV. Preparation and Reactions of Organocalcium Halides. *J. Chem. Soc.* **1963**, 577.
- (104) Westerhausen, M.; Gärtner, M.; Fischer, R.; Langer, J. Aryl Calcium Compounds: Syntheses, Structures, Physical Properties, and Chemical Behavior. *Angew. Chem., Int. Ed.* **2007**, *46*, 1950–1956.
- (105) Westerhausen, M. Heavy Grignard Reagents-Synthesis and Reactivity of Organocalcium Compounds. *Coord. Chem. Rev.* **2008**, *252*, 1516–1531.
- (106) De Bruin-Dickason, C. N.; Deacon, G. B.; Jones, C.; Junk, P. C.; Wiecko, M. Functionalised Alkaline Earth Iodides from Grignard Synthons “PhAeI(Thf)_n” (Ae = Mg-Ba). *Eur. J. Inorg. Chem.* **2019**, *2019*, 1030–1038.
- (107) Evans, D. F.; Fazakerley, G. V.; Phillips, R. F. Organometallic Compounds of Bivalent Ytterbium. *J. Chem. Soc. Chem. Commun.* **1970**, 244–244.
- (108) Evans, D. F.; Fazakerley, G. V.; Phillips, R. F. Organometallic Compounds of Bivalent Europium, Ytterbium, and Samarium. *J. Chem. Soc. A Inorg. Physical, Theor.* **1971**, 1931–1934.

- (109) Syutkina, O. P.; Rybakova, L. F.; Petrov, E. S.; Beletskaya, I. P. Thienyl and Perfluorophenyl Derivatives of Divalent Lanthanides. *J. Organomet. Chem.* **1985**, *280*, C67–C69.
- (110) Fieser, M. E.; Macdonald, M. R.; Krull, B. T.; Bates, J. E.; Ziller, J. W.; Furcher, F.; Evans, W. J. Structural, Spectroscopic, and Theoretical Comparison of Traditional vs Recently Discovered Ln²⁺ Ions in the [K(2.2.2-Cryptand)][(C₅H₄SiMe₃)₃Ln] Complexes: The Variable Nature of Dy²⁺ and Nd²⁺. *J. Am. Chem. Soc.* **2015**, *137*, 369–382.
- (111) Hitchcock, P. B.; Lappert, M. F.; Maron, L.; Protchenko, A. V. Lanthanum Does Form Stable Molecular Compounds in the +2 Oxidation State. *Angew. Chem., Int. Ed.* **2008**, *47*, 1488–1491.
- (112) Wiecko, M.; Deacon, G. B.; Junk, P. C. Organolanthanoid-Halide Synthons - A New General Route to Monofunctionalized Lanthanoid(II) Compounds? *Chem. Commun.* **2010**, *46*, 5076–5078.
- (113) Ali, S. H.; Deacon, G. B.; Junk, P. C.; Hamidi, S.; Wiecko, M.; Wang, J. Lanthanoid Pseudo-Grignard Reagents: A Major Untapped Resource. *Chem. - A Eur. J.* **2018**, *24*, 230–242.
- (114) Eaborn, C.; Hitchcock, P. B.; Izod, K.; Smith, J. D. A Monomeric Solvent-Free Bent Lanthanide Dialkyl and a Lanthanide Analog of a Grignard Reagent. Crystal Structures of Yb{C(SiMe₃)₃}₂ and [Yb{C(SiMe₃)₃}I-OEt₂]. *J. Am. Chem. Soc.* **1994**, *116*, 12071–12072.
- (115) Eaborn, C.; Hitchcock, P. B.; Izod, K.; Lu, Z.; Smith, J. D. Alkyl Derivatives of Europium(+2) and Ytterbium(+2). Crystal Structures of Eu{C(SiMe₃)₃}₂, Yb{C(SiMe₃)₂(SiMe₂CH = CH₂)}I-OEt₂ and Yb{C(SiMe₃)₂(SiMe₂OMe)}I-OEt₂. *Organometallics* **1996**, *15*, 4783–4790.
- (116) Heckmann, G.; Niemeyer, M. Synthesis and First Structural Characterization of Lanthanide(II) Aryls: Observation of a Schlenk Equilibrium in Europium(II) and Ytterbium(II) Chemistry. *J. Am. Chem. Soc.* **2000**, *122*, 4227–4228.
- (117) Niemeyer, M. Novel Synthesis of a Lanthanide Trialkyl ± Characterization and Crystal Structure of Yb(CH₂^tBu)₃(THF)₂. *Z. Anorg. Allg. Chem.* **2000**, *626*, 1027–1029.
- (118) Meyer, G. The Oxidation of Metals with Liebig Acids. *Z. Anorg. Allg. Chem.* **2008**, *634*, 201–222.
- (119) Evans, W. J.; Greci, M. A.; Ziller, J. W. Utility of 2-Methoxyethanol in the Synthesis of Poly Europium Complexes: {[Eu(OCH₂CH₂OMe)₂(OC₆H₃R₂-2,6)-][H⁺]}₄ (R = Me, ^tPr) and [EuAl₂(OCH₂CH₂OMe)₃Me₅]₂. *Inorg. Chem.* **1998**, *37*, 5221–5226.
- (120) Evans, W. J.; Greci, M. A.; Ziller, J. W. The Utility of N-Methylimidazole and Acetonitrile as Solvents for the Direct Reaction of Europium with Alcohols Including the First Example of Acetonitrile as a μ-η¹:η¹-Bridging Ligand. *Chem. Commun.* **1998**, 2367–2368.
- (121) Carretas, J. M.; Branco, J.; Marçalo, J.; Waerenborgh, J. C.; Marques, N.; Pires De Matos, A. The “dissolution” of Europium and Ytterbium in Alcohols. *J. Alloys Compd.* **1998**, *275–277*, 841–843.
- (122) Bradley, D. C.; Chudzynska, H.; Frigo, D. M.; Hammond, M. E.; Hursthouse, M. B.; Mazid, M. A. Pentanuclear Oxoalkoxide Clusters of Scandium, Yttrium, Indium and Ytterbium, X-Ray Crystal Structures of [M₅(μ₅-O)(μ₃-OPr)₄(μ₂-OPr)₄(OPr^t)₅] (M = In, Yb). *Polyhedron* **1990**, *9* (5), 719–726.
- (123) Deacon, G. B.; Feng, T.; Forsyth, C. M.; Gitlits, A.; Hockless, D. C. R.; Shen, Q.; Skelton, B. W.; White, A. H. A Simple Synthesis and a Structural Survey of Homoleptic Rare Earth(III) 2,6-Diphenylphenolates. *J. Chem. Soc., Dalton Trans.* **2000**, 961–966.
- (124) Mazdiyasi, K. S.; Lynch, C. T.; Smith, J. S. The Preparation and Some Properties of Yttrium, Dysprosium, and Ytterbium Alkoxides. *Inorg. Chem.* **1966**, *5*, 342–346.
- (125) Brown, L. M.; Mazdiyasi, K. S. Synthesis and Some Properties of Yttrium and Lanthanide Isopropoxides. *Inorg. Chem.* **1970**, *9*, 2783–2786.
- (126) Deacon, G. B.; Tuong, T. D.; Wilkinson, D. L.; Marks, T. J. Lanthanide Trichlorides by Reaction of Lanthanide Metals with Mercury(II) Chloride in Tetrahydrofuran. *Inorg. Synth.* **2007**, *42*, 286–291.
- (127) Poncelet, O.; Sartain, W. J.; Hubert-pfalzgraf, L. G.; Folting, K.; Caulton, K. G. Chemistry of Yttrium Triisopropoxide Revisited: Characterization and Crystal Structure of Y₅(μ₅-O)(μ₃-OPr)₄(μ₂-OPr)₄(OPr^t)₅. *Inorg. Chem.* **1989**, *28*, 263–267.
- (128) Hubert-Pfalzgraf, L. G.; Daniele, S.; Bennaceur, A.; Daran, J.-C.; Vaissermann, J. Praseodymium Alkoxide Chemistry: Synthesis and Molecular Structure of [Pr₄(μ₄-O)₂(μ₃, η²-OR)₂(μ, η²-OR)₄(μ, η¹-OR)(OR)(OPMe₃)₂] (R = C₂H₄OMe) and [Y₄Pr(μ₅-O)(μ₃-OR)₄(μ-OR)₄(OR)₅] (R = Pri). *Polyhedron* **1997**, *16* (7), 1223–1234.
- (129) Nief, F.; Mathey, F. Synthesis of Diphosphametalloocene Derivatives of Divalent Lanthanides. Phosphorus-Phosphorus Bond Cleavage by Metallic Ytterbium and Samarium. *Synlett.* **1991**, *1991*, 745–746.
- (130) Nief, F.; Ricard, L.; Mathey, F. Phospholyl (Phosphacyclopentadienyl) and Arsolyl (Arsacyclopentadienyl) Complexes of Ytterbium(II) and Samarium(II). Synthetic, Structural and Multi-nuclear (³¹P and ¹⁷¹Yb) NMR Studies. *Polyhedron* **1993**, *12* (1), 19–26.
- (131) Recknagel, A.; Edelman, F. T. One-Step Synthesis of Organolanthanide(II) Complexes from the Metal. *Angew. Chem., Int. Ed.* **1991**, *30*, 693–694.
- (132) Sanden, T.; Gamer, M. T.; Fagin, A. A.; Chudakova, V. A.; Konchenko, S. N.; Fedushkin, I. L.; Roesky, P. W. Synthesis of Unsupported Ln-Ga Bonds by Salt Metathesis and Ga-Ga Bond Reduction. *Organometallics* **2012**, *31*, 4331–4339.
- (133) Shriver, D. F.; Drezdson, M. A. *The Manipulation of Air-Sensitive Compounds*, 2nd ed.; Wiley, 1986; pp 1–336.
- (134) Deacon, G. B.; Forsyth, C. M.; Junk, P. C.; Skelton, B. W.; White, A. H. The Striking Influence of Intramolecular Lanthanoid-π-Arene Interactions on the Structural Architecture of the Homoleptic Aryloxolanthanoid(II) Complexes [Eu₂(Odp)₂(μ-Odp)₃] and [Yb₂(Odp)₂(μ-Odp)₂] and the Yb^{II}/Yb^{III} Trimetallic [Yb₂(μ-Odp)₃]⁺[Yb(Odp)₄]⁻ (Odp = 2,6-Diphenylphenolate). *Chem. - A Eur. J.* **1999**, *5*, 1452–1459.
- (135) Deacon, G. B.; Fanwick, P. E.; Gitlits, A.; Rothwell, I. P.; Skelton, B. W.; White, A. H. Rare Earth Complexes of Bulky 2,6-Diphenylphenolates Containing Additional, Potentially Buttressing 3,5-Substituents. *Eur. J. Inorg. Chem.* **2001**, *2001*, 1505–1514.
- (136) Cole, M. L.; Deacon, G. B.; Junk, P. C.; Proctor, K. M.; Scott, J. L.; Strauss, C. R. Direct Syntheses and Structural Novelty of Lanthanoid Aryloxides with Flexible Radial Arms. *Eur. J. Inorg. Chem.* **2005**, *2005*, 4138–4144.
- (137) Deacon, G. B.; Junk, P. C.; Moxey, G. J. Mono-, di-, tri- and tetranuclear rare earth complexes obtained using a moderately bulky aryloxide ligand. *Chem. - An Asian J.* **2009**, *4*, 1717–1728.
- (138) Deacon, G. B.; Forsyth, C. M.; Harika, R.; Junk, P. C.; Ziller, W.; Evans, W. J. Hydrocarbon-Soluble, Polymetallic, Lanthanoid Aryloxides Constructed Utilising Ligands with Distal But Groups. *J. Mater. Chem.* **2004**, *14*, 3144–3149.
- (139) Deacon, G. B.; Junk, P. C.; Moxey, G. J. Metal-π Interactions Dominate in the Solid-State Structures of Molecular Heterobimetallic Alkali-Metal-Europium(II). *Chem. - An Asian J.* **2009**, *4*, 1309–1317.
- (140) Deacon, G. B.; Junk, P. C.; Moxey, G. J.; Ruhlandt-Senge, K.; St. Prix, C.; Zuniga, M. F. Charge-Separated and Molecular Heterobimetallic Rare Earth-Rare Earth and Alkaline Earth-Rare Earth Aryloxo Complexes Featuring Intramolecular Metal-π-Arene Interactions. *Chem. - A Eur. J.* **2009**, *15*, 5503–5519.
- (141) Deacon, G. B.; Gitlits, A.; Roesky, P. W.; Bürgstein, M. R.; Lim, K. C.; Skelton, B. W.; White, A. H. Simple Syntheses, Structural Diversity, and Tishchenko Reaction Catalysis of Neutral Homoleptic Rare Earth(II or III) 3,5-Di-tert-Butylpyrazolates—The Structures of [Sc(^tBu₂pz)₃], [Ln₂(^tBu₂pz)₆] (Ln = La, Nd, Yb, Lu), and [Eu₄(^tBu₂pz)₈]. *Chem. - A Eur. J.* **2001**, *7*, 127–138.
- (142) Deacon, G. B.; Forsyth, C. M.; Gitlits, A.; Harika, R.; Junk, P. C.; Skelton, B. W.; White, A. H. Pyrazolate Coordination Continues to Amaze - The New μ-η²:η¹ Binding Mode and the First Case of Unidentate Coordination to a Rare Earth Metal. *Angew. Chem., Int. Ed.* **2002**, *41*, 3249–3251.
- (143) Deacon, G. B.; Gitlits, A.; Skelton, B. W.; White, A. H. The First Neutral Homoleptic Lanthanoid Pyrazolates, Including the Mixed Oxidation State Species [Yb₂(^tBu₂pz)₅] (^tBu₂pz = 3,5-di-tert-butylpyrazolate), from a Simple New Synthesis of Pyrazolate Complexes. *Chem. Commun.* **1999**, 1213–1214.

- (144) Deacon, G. B.; Forsyth, C. M.; Gitlits, A.; Skelton, B. W.; White, A. H. Reactions of Lanthanoid Metals with 3,5-Diphenylpyrazole at Elevated Temperatures: Synthesis and Structures of Both Homoleptic, $[\text{Ln}_3(\text{Ph}_2\text{pz})_3]$ (Ln = La, Nd), $[\text{Ln}_2(\text{Ph}_2\text{pz})_6]$ (Ln = Er, Lu), and Heteroleptic, $[\text{Ln}(\text{Ph}_2\text{pz})_3(\text{Ph}_2\text{pzH})_2]$ (Ln = La, Nd, Gd, Tb). *Er or Y. Dalton Trans.* **2004**, 1239–1247.
- (145) Qayyum, S.; Haberland, K.; Forsyth, C. M.; Junk, P. C.; Deacon, G. B.; Kempe, R. Small Steric Variations in Ligands with Large Synthetic and Structural Consequences. *Eur. J. Inorg. Chem.* **2008**, 2008, 557–562.
- (146) Müller-Buschbaum, K.; Mokaddem, Y.; Schappacher, F. M.; Pöttgen, R. $\infty 3[\text{Eu}(\text{Tzpy})_2]$: A Homoleptic Framework Containing $\{\text{Eu}^{\text{III}}\text{N}_{12}\}$ Icosahedra. *Angew. Chem., Int. Ed.* **2007**, 46, 4385–4387.
- (147) Müller-Buschbaum, K.; Quitmann, C. C. Two New Groups of Homoleptic Rare Earth Pyridylbenzimidazolates: $(\text{NC}_{12}\text{H}_8(\text{NH})_2)$ - $[\text{Ln}(\text{N}_3\text{C}_{12}\text{H}_8)_4]$ with Ln = Y, Tb. *Yb. Inorg. Chem.* **2003**, 42, 2742–2750.
- (148) Quitmann, C. C.; Müller-Buschbaum, K. The Unsubstituted Rare Earth Pyrazolates $[\text{Nd}(\text{Pz})_3(\text{PzH})_4]$ and $\infty 1[\text{Ho}(\text{Pz})_3(\text{PzH})_3]$: Structural Diversity from Monomers to a Coordination Polymer. *Z. Anorg. Allg. Chem.* **2005**, 631, 1191–1198.
- (149) Silverman, G. S.; Rakita, P. E. *Handbook of Grignard Reagents*, 1st ed.; CRC Press: Boca Raton, FL, 1996; pp 1–736.
- (150) Hamidi, S.; Deacon, G. B.; Junk, P. C.; Neumann, P. Direct Reaction of Iodine-Activated Lanthanoid Metals with 2,6-Diisopropylphenol. *Dalton Trans.* **2012**, 41, 3541–3552.
- (151) Deacon, G. B.; Junk, P. C.; Urbatsch, A. Lanthanoid and Alkaline Earth Complexes Involving New Substituted Pyrazolates. *Aust. J. Chem.* **2012**, 65, 802–810.
- (152) Deacon, G. B.; Junk, P. C.; Urbatsch, A. Trivalent Rare Earth Complexes of the Unsymmetrical 3-(2'-Thienyl)-5-(Trifluoromethyl)-Pyrazolate Ligand. *Eur. J. Inorg. Chem.* **2011**, 2011, 3592–3600.
- (153) Deacon, G. B.; Hamidi, S.; Junk, P. C.; Kelly, R. P.; Wang, J. Direct Reactions of Iodine-Activated Rare-Earth Metals with Phenols of Varying Steric Bulk. *Eur. J. Inorg. Chem.* **2014**, 2014, 460–468.
- (154) Guo, Z.; Blair, V. L.; Deacon, G. B.; Junk, P. C. Widely Contrasting Outcomes from the Use of Tris(Pentafluorophenyl)Bismuth or Pentafluorophenylsilver as Oxidants in the Reactions of Lanthanoid Metals with N,N'-Diarylformamidines. *Dalton Trans.* **2020**, 49, 13588–13600.
- (155) Guo, Z.; Blair, V. L.; Deacon, G. B.; Junk, P. C. A Simple One-Pot Route to Stable Formamidinatoiodidolanthanoid(III) Complexes from Lanthanoid Metals. *Chem. Commun.* **2021**, 57, 11513–11516.
- (156) Sazonova, A. V.; Bochkarev, L. N.; Malysheva, I. P.; Baranov, E. V. Facile Synthesis of Rare-Earth Pyrazolonates by the Reaction of Rare-Earth Metals with 1-Phenyl-3-Methyl-4-Isobutryryl-5-Pyrazolone. Crystal Structures of $[\text{Ln}(\text{PMIP})_3]_2$ (Ln = Y, Gd, Tb, Er, Tm). *Inorg. Chim. Acta* **2012**, 392, 454–458.
- (157) Mashima, K.; Nakayama, Y.; Nakamura, A.; Kanehisa, N.; Kai, Y.; Takaya, H. A New Convenient Preparation of Monocyclooctate-triaenyl-Lanthanide Complexes from Metallic Lanthanides and Oxidants. *J. Organomet. Chem.* **1994**, 473, 85–91.
- (158) Müller-Buschbaum, K.; Quitmann, C. C. First Homoleptic Rare Earth Carbazolates: Synthesis, Crystal Structures, Spectroscopic and Thermal Investigations of Divalent $\infty 1[\text{Ln}(\text{NC}_{12}\text{H}_8)_2]$ with Ln = Europium and Ytterbium. *Z. Anorg. Allg. Chem.* **2003**, 629, 1610–1616.
- (159) Werner, D.; Bayer, U.; Rad, N. E.; Junk, P. C.; Deacon, G. B.; Anwender, R. Unique and Contrasting Structures of Homoleptic Lanthanum(III) and Cerium(III) 3,5-Dimethylpyrazolates. *Dalton Trans.* **2018**, 47, 5952–5955.
- (160) Deacon, G. B.; Junk, P. C.; Werner, D. Enhancing the Value of Free Metals in the Synthesis of Lanthanoid Formamidinates: Is a Co-Oxidant Needed? *Chem. - A Eur. J.* **2016**, 22, 160–173.
- (161) Deacon, G. B.; Vince, D. G. Bis(Pentafluorophenyl)Ytterbium: A Transmetalation Synthesis of a Sigma-Bonded Lanthanide Organometallic Compound. *J. Organomet. Chem.* **1976**, 112, C1–C2.
- (162) Deacon, G. B.; Raverty, W. D.; Vince, D. G. Bis-(Polyfluorophenyl)Ytterbium Compounds. *J. Organomet. Chem.* **1977**, 135, 103–114.
- (163) Deacon, G. B.; Koplick, A. J. Bis(Phenylethynyl)Ytterbium - a Novel Organolanthanide. *J. Organomet. Chem.* **1978**, 146, C43–C45.
- (164) Deacon, G. B.; Koplick, A. J.; Raverty, W. D.; Vince, D. G. Preparation and Identification of Some Organolanthanoid Species in Solution. *J. Organomet. Chem.* **1979**, 182, 121–141.
- (165) Deacon, G. B.; Koplick, A. J.; Tuong, T. D. Preparations and Properties of Some Di(Alkynyl)Lanthanoids. *Aust. J. Chem.* **1982**, 35, 941–949.
- (166) Beaini, S.; Deacon, G. B.; Delbridge, E. E.; Junk, P. C.; Skelton, B. W.; White, A. H. Dimethyltin(IV) Bis(3,5-Diphenylpyrazolate) as a Synthetic Reagent in the Preparation of Rare-Earth Pyrazolate Complexes. *Eur. J. Inorg. Chem.* **2008**, 2008, 4586–4596.
- (167) Beaini, S.; Deacon, G. B.; Hilder, M.; Junk, P. C.; Turner, D. R. Organotin Compounds as Reagents for the Synthesis of Lanthanoid Complexes by Redox Transmetalation Reactions. *Eur. J. Inorg. Chem.* **2006**, 2006, 3434–3441.
- (168) Deacon, G. B.; Koplick, A. J.; Tuong, T. D. Reactions of Lanthanide Elements with Thallous Cyclopentadienide-A New Route to Cyclopentadienylanthanides. *Polyhedron* **1982**, 1, 423–424.
- (169) Deacon, G. B.; Forsyth, C. M.; Newnham, R. H.; Tuong, T. D. Organolanthanoids. X. Syntheses of Cyclopentadienylanthanoids by Transmetalation Reactions in Pyridine, Acetonitrile and Ethers. *Aust. J. Chem.* **1987**, 40, 895–906.
- (170) Deacon, G. B.; Feng, T.; Nickel, S.; Ogden, M. I.; White, A. H. Organoamido- and Aryloxo-Lanthanoids. IV. Synthesis and x-Ray Structures of Low-Coordinate $[\text{Yb}(\text{O}-2,4,6\text{-Bu}^t_3\text{C}_6\text{H}_2)_3(\text{Thf})]$ and $[\text{Yb}(\text{O}-2,4,6\text{-Bu}^t_3\text{C}_6\text{H}_2)_2(\mu\text{-OH})(\text{Thf})_2]$ (Thf = Tetrahydrofuran). *Aust. J. Chem.* **1992**, 45, 671–683.
- (171) Deacon, G. B.; Feng, T.; Nickel, S.; Skelton, B. W.; White, A. H. A Simple Synthesis of Tetrahydrofuran Complexes of Lanthanoid Trichlorides: Convenient Substitutes for Anhydrous Lanthanoid Chlorides. *J. Chem. Soc., Chem. Commun.* **1993**, 1328–1329.
- (172) Lin, G.; Wong, W. T. Synthesis and Crystal Structure of $[(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)\text{Ti}]_\infty$ and the Divalent Ytterbium Derivative $[(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)_2\text{Yb}(\text{DME})]$. *J. Organomet. Chem.* **1995**, 495, 203–208.
- (173) Lin, G.; Wong, W. Synthesis and Structural Characterization of Functionalized Organolanthanide Complexes $[(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)_2\text{Ln}(\text{DIME})]$ (Ln = Eu, Yb) (DIME = diethylene glycol dimethyl ether). *J. Organomet. Chem.* **1996**, 523, 93–98.
- (174) Deacon, G. B.; Delbridge, E. E.; Skelton, B. W.; White, A. H. Synthesis of the First Lanthanoid(II) Pyrazolate Complex by Redox Transmetalation from Thallium and Mercury Reagents. *Eur. J. Inorg. Chem.* **1998**, 1998, 543–545.
- (175) Guo, Z.; Blair, V.; Deacon, G. B.; Junk, P. C. Can Bismuth Replace Mercury in Redox Transmetalation/Protolysis Syntheses from Free Lanthanoid Metals? *Chem. - A Eur. J.* **2018**, 24, 17464–17474.
- (176) Bochkarev, L. N.; Stepantseva, T.; Zakharov, L. N.; Fukin, G. K.; Yanovsky, A. I.; Struchkov, Y. I. Synthesis and Crystal Structure of $\text{Ph}_3\text{Ln}(\text{THF})_3$ (Ln = Er, Tm). *Organometallics* **1995**, 14, 2127–2129.
- (177) Guo, Z.; Luu, J.; Blair, V.; Deacon, G. B.; Junk, P. C. Replacing Mercury: Syntheses of Lanthanoid Pyrazolates from Free Lanthanoid Metals, AgC_6F_5 , and Pyrazoles, Aided by a Facile Synthesis of Polyfluoroarylsilver Compounds. *Eur. J. Inorg. Chem.* **2019**, 2019, 1018–1029.
- (178) Gilman, H.; Jones, R. G. Organometallic Compounds of Titanium, Zirconium, and Lanthanum. *J. Org. Chem.* **1945**, 10, 505–515.
- (179) Deacon, G. B.; Pain, G. N.; Tuong, T. D. $(\eta^5\text{-Cyclopentadienyl})\text{Lanthanide}$ Complexes from the Metallic Elements: Cp_3Ln (Ln = Nd, Sm), $\text{Cp}_2\text{Y}(\text{DME})$. *Inorg. Synth.* **1990**, 27, 291–296.
- (180) Deacon, G. B.; Forsyth, C. M. A Half-Sandwich Perfluoroorganoytterbium(II) Complex from a Simple Redox Transmetalation/Ligand Exchange Synthesis. *Organometallics* **2003**, 22, 1349–1352.
- (181) Suleimanov, G. Z.; Bregadze, V. I.; Koval'chuk, N. A.; Beletskaya, I. P. Synthesis of Carboranyl Derivatives of Di- and Trivalent Lanthanides. *J. Organomet. Chem.* **1982**, 235, C17–C18.
- (182) De Bruin-Dickason, C. N.; Boutland, A. J.; Dange, D.; Deacon, G. B.; Jones, C. Redox Transmetalation Approaches to the Synthesis of

Extremely Bulky Amido-Lanthanoid(II) and -Calcium(II) Complexes. *Dalton Trans.* **2018**, 47, 9512–9520.

(183) Deacon, G. B.; Forsyth, C. M.; Newham, R. H. Synthesis of Divalent Aryloxy- and Diorganoamido-Lanthanides from Bis-(Pentafluorophenyl)Lanthanide(II) Complexes. *Polyhedron* **1987**, 6, 1143–1145.

(184) Deacon, G. B.; Forsyth, C. M. The Synthesis of a New Soluble Samarium(II) Diorganoamide. *Inorg. Chim. Acta* **1988**, 154, 121–122.

(185) Deacon, G. B.; Forsyth, C. M.; Gatehouse, B. M. Organoamido- and Aryloxo-Lanthanoids. I The Preparation and Characterization of Some Lanthanoid(II) Organoamides, and the X-Ray Crystal Structure of *Cis*-Bis(Carbazol-9-Yl)Tetrakis(Tetrahydrofuran)Europium(II). *Aust. J. Chem.* **1990**, 43, 795–806.

(186) Cole, M. L.; Deacon, G. B.; Forsyth, C. M.; Junk, P. C.; Konstas, K.; Wang, J. Steric Modulation of Coordination Number and Reactivity in the Synthesis of Lanthanoid(III) Formamidinates. *Chem. - A Eur. J.* **2007**, 13, 8092–8110.

(187) Deacon, G. B.; Junk, P. C.; Werner, D. Lanthanoid Induced C-F Activation of All Fluorine Atoms of One CF₃ Group. *Eur. J. Inorg. Chem.* **2015**, 2015, 1484–1489.

(188) Cole, M. L.; Deacon, G. B.; Forsyth, C. M.; Junk, P. C.; Konstas, K.; Wang, J.; Bittig, H.; Werner, D. Synthesis, Structures and Reactivity of Lanthanoid(II) Formamidinates of Varying Steric Bulk. *Chem. - A Eur. J.* **2013**, 19, 1410–1420.

(189) Deacon, G. B.; Junk, P. C.; Werner, D. The Synthesis and Structures of Rare Earth 2-Fluorophenyl- and 2,3,4,5-Tetrafluorophenyl-N,N'-Bis(Aryl)Formamidinate Complexes. *Polyhedron* **2016**, 103, 178–186.

(190) Deacon, G. B.; Junk, P. C.; Macreadie, L. K.; Werner, D. Structural and Reactivity Consequences of Reducing Steric Bulk of N,N-Diarylformamidinates Coordinated to Lanthanoid Ions. *Eur. J. Inorg. Chem.* **2014**, 2014 (30), 5240–5250.

(191) Cole, M. L.; Junk, P. C. The Synthesis of a Sterically Hindered Samarium(II) Bis(Amidinate) and Conversion to Its Homoleptic Trivalent Congener. *Chem. Commun.* **2005**, 2695–2697.

(192) Deacon, G. B.; Nickel, S.; Mackinnon, P.; Tiekink, E. R. T. Organoamido- and Aryloxo-Lanthanoids. II. Preparation of Tris(2,6-Diphenylphenoxy)-Lanthanoid(III) Complexes and the X-Ray Structures of Low-Coordinate [Yb(O-2,6-Ph₂C₆H₃)₃] Containing an Intramolecular Chelate Yb... π -Arene Interaction, and [Yb(O-2,6-Ph₂C₆H₃)₃(thf)₂].thf (thf = Tetrahydrofuran). *Aust. J. Chem.* **1990**, 43, 1245–1257.

(193) Melman, J. H.; Emge, T. J.; Brennan, J. G. Fluorinated Thiolates of Divalent and Trivalent Lanthanides. Ln-F Bonds and the Synthesis of LnF₃. *Inorg. Chem.* **2001**, 40, 1078–1081.

(194) Melman, J. H.; Rohde, C.; Emge, T. J.; Brennan, J. G. Trivalent Lanthanide Compounds with Fluorinated Thiolate Ligands: Ln-F Dative Interactions Vary with Ln and Solvent. *Inorg. Chem.* **2002**, 41, 28–33.

(195) Simler, T.; Feuerstein, T. J.; Yadav, R.; Gamer, M. T.; Roesky, P. W. Access to Divalent Lanthanide NHC Complexes by Redox-Transmetalation from Silver and CO₂ Insertion Reactions. *Chem. Commun.* **2019**, 55, 222–225.

(196) Schwarz, N.; Sun, X.; Yadav, R.; Koppe, R.; Simler, T.; Roesky, P. W. Application of the Redox-Transmetalation Procedure to Access Divalent Lanthanide and Alkaline-Earth NHC Complexes. *Chem. - A Eur. J.* **2021**, 27, 12857.

(197) Deacon, G.; Newham, R. Organolanthanoids.VIII. Some Ligand-Exchange Reactions of Pentafluorophenyllanthanoids and Bis(Phenylethynyl)Ytterbium(II). *Aust. J. Chem.* **1985**, 38, 1757.

(198) Deacon, G. B.; Feng, T.; Mackinnon, P.; Newham, R. H.; Nickel, S.; Skelton, B. W.; White, A. H. Organoamido- and Aryloxo-Lanthanoids. V. Preparations of Low-Coordinate Lanthanoid(II) Phenolates, Ln(OC₆H₂Bu^t₂-2,6-X-4)₂(thf)_n (Ln = Yb or Eu; X = H, Me or Bu^t; n = 2 or 3) and the X-Ray Crystal Structure of Five-Coordinate [Yb(OC₆H₂Bu^t₂-2,6-X-4)₂(thf)₃].thf. *Aust. J. Chem.* **1993**, 46, 387–399.

(199) Deacon, G. B.; Fallon, G. D.; Forsyth, C. M.; Harris, S. C.; Junk, P. C.; Skelton, W.; White, A. H. Manipulation of Reaction Pathways in

Redox Transmetalation – Ligand Exchange Syntheses of Lanthanoid-(II)/(III) Aryloxo Complexes. *Dalton Trans.* **2006**, 802–812.

(200) Deacon, G. B.; Feng, T.; Skelton, B. W.; White, A. H. Organoamido- and Aryloxo-Lanthanoids. XI. Syntheses and Crystal Structures of Nd(Odpp)₃, Nd(Odpp)₃(Thf) and [Nd(Odpp)₃(Thf)₂].2(Thf) (Odpp = 2,6-Diphenylphenolate): Variations in Intramolecular π -Ph-Nd Interactions. *Aust. J. Chem.* **1995**, 48, 741–756.

(201) Deacon, G. B.; Forsyth, C. M.; Jaroschik, F.; Junk, P. C.; Kay, D. L.; Maschmeyer, T.; Masters, A. F.; Wang, J.; Field, L. D. Accessing Decaphenylmetallocenes of Ytterbium, Calcium, and Barium by Desolvation of Solvent-Separated Ion Pairs: Overcoming Adverse Solubility Properties. *Organometallics* **2008**, 27, 4772–4778.

(202) Kelly, R. P.; Bell, T. D. M.; Cox, R. P.; Daniels, D. P.; Deacon, G. B.; Jaroschik, F.; Junk, P. C.; Le Goff, X. F.; Lemerrier, G.; Martinez, A.; et al. Divalent Tetra- and Penta-Phenylcyclopentadienyl Europium and Samarium Sandwich and Half-Sandwich Complexes: Synthesis, Characterization, and Remarkable Luminescence Properties. *Organometallics* **2015**, 34, 5624–5636.

(203) Deacon, G. B.; Jaroschik, F.; Junk, P. C.; Kelly, R. P. A Divalent Heteroleptic Lanthanoid Fluoride Complex Stabilised by the Tetraphenylcyclopentadienyl Ligand, Arising from C-F Activation of Pentafluorobenzene. *Chem. Commun.* **2014**, 50, 10655–10657.

(204) Forsyth, C. M.; Deacon, G. B.; Field, L. D.; Jones, C.; Junk, P. C.; Kay, D. L.; Masters, A. F.; Richards, A. F. Dinuclear Alkynyllanthanoid(II) Dications with Pentaphenylcyclopentadienyl or Tri-Tert-Butyldiphosphacyclopentadienyl Counter Ions. *Chem. Commun.* **2006**, 1003–1005.

(205) Deacon, G. B.; Jaroschik, F.; Junk, P. C.; Kelly, R. P. Bulky Group 2 Octaphenylmetallocenes and Direct Access to Calcium and Ytterbium Pseudo-Grignard Complexes. *Organometallics* **2015**, 34, 2369–2377.

(206) Deacon, G. B.; Delbridge, E. E.; Skelton, B. W.; White, A. H. Organoamido- and Aryloxo-Lanthanoids, 19. Synthesis and Structures of *Cisoid* and *Transoid* Bis(1,2-Dimethoxyethane)Bis(η^2 -Pyrazolato)-Lanthanoid(II) Complexes. *Eur. J. Inorg. Chem.* **1999**, 1999, 751–761.

(207) Pfeiffer, D.; Ximba, B. J.; Liable-Sands, L. M.; Rheingold, A. L.; Heeg, M. J.; Coleman, D. M.; Schlegel, H. B.; Kuech, T. F.; Winter, C. H. Synthesis, Structure, and Molecular Orbital Studies of Yttrium, Erbium, and Lutetium Complexes Bearing η^2 -Pyrazolato Ligands: Development of a New Class of Precursors for Doping Semiconductors. *Inorg. Chem.* **1999**, 38, 4539–4548.

(208) Deacon, G. B.; Harika, R.; Junk, P. C.; Skelton, B. W.; Werner, D.; White, A. H. The Synthesis, Structures and Polymorphism of the Dimeric Trivalent Rare-Earth 3,5-Dimethylpyrazolato Complexes [Ln(Me₂pz)₃(Thf)]₂. *Eur. J. Inorg. Chem.* **2014**, 2014, 2412–2419.

(209) Cosgriff, J. E.; Deacon, G. B.; Hemling, H.; Schumann, H. Monomeric Tris(η^2 -Pyrazolato)Lanthanoid Complexes with the Bulky 3,5-Di(*tert*-Butyl)Pyrazolato Ligand. *Angew. Chem., Int. Ed.* **1993**, 32, 874–875.

(210) Cosgriff, J. E.; Deacon, G. B.; Gatehouse, B. M.; Schumann, H.; Hemling, H. Organoamido- and Aryloxo-Lanthanoids. X. Tris(η^2 -3,5-Di-*t*-Butylpyrazolato) Lanthanoid(III) Complexes and the X-Ray Crystal Structure of Er(η^2 -Bu^t₂pz)₃(Thf)₂. *Aust. J. Chem.* **1994**, 47 (7), 1223–1235.

(211) Cole, M. L.; Deacon, G. B.; Junk, P. C.; Konstas, K. Steric Engineering of C-F Activation with Lanthanoid Formamidinates. *Chem. Commun.* **2005**, 1581–1583.

(212) Deacon, G. B.; Fallon, G. D.; Forsyth, C. M.; Schumann, H.; Weimann, R. Syntheses of Low Coordination Number Divalent Lanthanoid Organoamide Complexes, and the X-Ray Crystal Structures of Bis[(N-2,6-Diisopropylphenyl)(N-Trimethylsilyl)-Amido]Bis(Tetrahydrofuran)Samarium(II) and -Ytterbium(II). *Chem. Ber.* **1997**, 130, 409–415.

(213) Hauber, S. O.; Niemeyer, M. Stabilization of Unsolvated Europium and Ytterbium Pentafluorophenyls by π -Bonding Encapsulation through a Sterically Crowded Triazenido Ligand. *Inorg. Chem.* **2005**, 44, 8644–8646.

(214) Bard, A. J.; Parsons, R.; Jordan, J. *Standard Potentials in Aqueous Solutions*, 1st ed.; Taylor & Francis: Boca Raton, FL, 1985; pp 1–848.

- (215) Deacon, G. B.; Koplick, A. J.; Tuong, T. D. Organolanthanoids. VI Syntheses of Cyclopentadienyllanthanoids by Transmetalation Reactions Between Thallous Cyclopentadienides and Lanthanoid Elements. *Aust. J. Chem.* **1984**, *37*, 517–525.
- (216) Evans, W. J.; Bloom, I.; Hunter, W. E.; Atwood, J. L. Synthesis and X-Ray Crystal Structure of a Soluble Divalent Organosamarium Complex. *J. Am. Chem. Soc.* **1981**, *103*, 6507.
- (217) Cloke, F. G. N.; Green, M. L. H. Synthesis of Zerovalent Bis(η -Arene) Compounds of Zirconium, Hafnium, Niobium, Tantalum, and Tungsten Using the Metal Vapours. *J. Chem. Soc., Dalton Trans.* **1981**, 1938–1943.
- (218) Geoffrey, F.; Cloke, N.; Courtney, K. A.E.; Sameh, A. A.; Swain, A. C. Bis(η -Arene) Complexes Of The Early Transition Metals Derived From The 1,3,5-Tri-*t*-Butylbenzene Ligand. *Polyhedron* **1989**, *8*, 1641–1648.
- (219) Anderson, D. M.; Cloke, F. G. N.; Cox, P. A.; Edelstein, N.; Green, J. C.; Pang, T.; Sameh, A. A.; Shalimoff, G. On the Stability and Bonding in Bis(η -Arene)Lanthanide Complexes. *J. Chem. Soc. Chem. Commun.* **1989**, 53–55.
- (220) Cloke, F. G. N.; Khan, K.; Perutz, R. N. η -Arene Complexes of Scandium(0) and Scandium(II). *J. Chem. Soc., Chem. Commun.* **1991**, 1372–1373.
- (221) Arnold, P. L.; Cloke, F. G. N.; Hitchcock, P. B.; Nixon, J. F. The First Example of a Formal Scandium(I) Complex: Synthesis and Molecular Structure of the 22-Electron Scandium Triple Decker Incorporating the Novel 1,3,5-Triphosphabenzene Ring. *J. Am. Chem. Soc.* **1996**, *118*, 7630–7631.
- (222) DeKock, C. W.; Ely, S. R.; Hopkins, T. E.; Brault, M. A. Preparation, Crystal and Molecular Structure and Properties of an Asymmetric Lanthanide Cyclooctatetraene Complex $[\text{Ln}(\text{C}_8\text{H}_8)(\text{OC}_4\text{H}_8)_2][\text{Ln}(\text{C}_8\text{H}_8)_2]$, Where Ln = La, Ce, Nd and Er. *Inorg. Chem.* **1978**, *17* (3), 625–631.
- (223) Blackborow, J. R.; Eady, C. R.; Von Gustorf, E. A. K.; Scriver, A.; Wolfbeis, O. Chemical Syntheses with Metal Atoms. *J. Organomet. Chem.* **1976**, *108*, C32–C34.
- (224) Evans, W. J.; Engerer, S. C.; Piliero, P. A.; Wayda, A. L. Homogeneous Catalytic Activation of Molecular Hydrogen by Lanthanoid Metal Complexes. *J. Chem. Soc. Chem. Commun.* **1979**, 1007–1008.
- (225) Evans, W. J.; Engerer, S. C.; Coleson, K. M. Reactivity of Lanthanide Metals with Unsaturated Hydrocarbons: Terminal Alkyne Reactions. *J. Am. Chem. Soc.* **1981**, *103*, 6672–6677.
- (226) Ely, S. R.; Hopkins, T. E.; DeKock, C. W. Crystal and Molecular Structure of a Novel Asymmetrical Neodymium-Cyclooctatetraene Compound, Cyclooctatetraenylbis(Tetrahydrofuran)Neodymium(III) Bis(Cyclooctatetraenyl)Neodymate(III). *J. Am. Chem. Soc.* **1976**, *98*, 1624–1625.
- (227) Evans, W. J.; Bloom, I.; Hunter, W. E.; Atwood, J. L. Metal Vapor Synthesis of $(\text{C}_5\text{Me}_5)_2\text{Sm}(\text{THF})_2$ and $(\text{C}_5\text{Me}_4\text{Et})_2\text{Sm}(\text{THF})_2$ and Their Reactivity with Organomercurial Reagents. Synthesis and X-Ray Structural Analysis of $(\text{C}_5\text{Me}_5)_2\text{Sm}(\text{C}_6\text{H}_5)(\text{THF})$. *Organometallics* **1985**, *4*, 112–119.
- (228) Cloke, F. G. N.; de Lemos, H. C.; Sameh, A. a. Homoleptic Diazadiene Complexes of Titanium, Yttrium, and Some Lanthanoid Elements. *J. Chem. Soc., Chem. Commun.* **1986**, 1344.
- (229) Arnold, P. L.; Cloke, F. G. N.; Nixon, J. F. The First Stable Scandocene: Synthesis and Characterisation of Bis(η -2,4,5-Tri-Tert-Butyl-1,3-Diphosphacyclopentadienyl)Scandium(II). *Chem. Commun.* **1998**, 797–798.
- (230) Matignon, C.; Cazes, E. Le Chlorure Samareux. *Ann. Chem. Phys.* **1906**, *8*, 417–426.
- (231) McCollum, B. C.; Dudis, D. S.; Lachgar, A.; Corbett, J. D. The Layered, Metallic Scandium Iodide $\text{Sc}_{0.93}\text{I}_2$: Synthesis, Structure, and Properties. *Inorg. Chem.* **1990**, *29*, 2030–2032.
- (232) Jongen, L.; Mudring, A. V.; Meyer, G. The Molecular Solid $\text{Sc}_{24}\text{C}_{10}^{130}$: A Truncated, Hollow T4 Supertetrahedron of Iodine Filled with a T3 Supertetrahedron of Scandium That Encapsulates the Adamantoid Cluster Sc_4C_{10} . *Angew. Chem., Int. Ed.* **2006**, *45*, 1886–1889.
- (233) Corbett, J. D.; et al. Lanthanum Triiodide (and Other Rare-Earth Metal Triiodides). *Inorg. Synth.* **1983**, *30*, 11–16.
- (234) Druding, L. F.; Corbett, J. D. Lower Oxidation States of the Lanthanides. Neodymium(II) Chloride and Iodide. *J. Am. Chem. Soc.* **1961**, *83*, 2462–2467.
- (235) Corbett, J. D.; Druding, L. F.; Burkhard, W. J.; Lindahl, C. B. Metal + Metal Halide Systems for Lanthanum Cerium and Praseodymium Iodides. *Discuss. Faraday Soc.* **1961**, *32*, 79–83.
- (236) Corbett, J. D.; Pollard, D. L.; Mee, J. E. Rare Earth Metal-Metal Halide Systems. X. Phase Studies of the Yttrium and Erbium Chlorides and Iodides. A Correlation of the Reduction Characteristics of the Rare Earth Elements. *Inorg. Chem.* **1966**, *5*, 761–766.
- (237) Meyer, G. The Reduction of Rare-Earth Metal Halides with Unlike Metals - Wöhler's Metallothermic Reduction. *Z. Anorg. Allg. Chem.* **2007**, *633*, 2537–2552.
- (238) Johnson, K. E.; Mackenzie, J. R. Anhydrous Chlorides of Some Rare Earths. *J. Inorg. Nucl. Chem.* **1970**, *32*, 43–48.
- (239) Asprey, L. B.; Kruse, F. H. Divalent Thulium. Thulium Diiodide. *J. Inorg. Nucl. Chem.* **1960**, *13*, 32–35.
- (240) Haschke, J.; Preparation, M. Phase Equilibria, and Crystal Chemistry of Samarium Bromides. *Inorg. Chem.* **1976**, *15*, 298–303.
- (241) Hadenfeldt, C.; Held, W. Darstellung, Eigenschaften Und Kristallstruktur Der Europium(II)-Phosphidhalogenide Eu_2PCl , Eu_2PBr Und Eu_2PI . *J. Less-Common Met.* **1986**, *123*, 25–35.
- (242) Bochkarev, M. N.; Fagin, A. A. A New Route to Neodymium(II) and Dysprosium(II) Iodides. *Chem. - A Eur. J.* **1999**, *5*, 2990–2992.
- (243) Katkova, M. A.; Fukin, G. K.; Fagin, A. A.; Bochkarev, M. N. Reduction of Azobenzene by Neodymium(II), Dysprosium(II), and Thulium(II) Diiodides. *J. Organomet. Chem.* **2003**, *682*, 218–223.
- (244) Evans, W. J.; Allen, N. T.; Workman, P. S.; Meyer, J. C. Large Scale Synthesis of Dysprosium and Neodymium Diiodides. *Inorg. Chem.* **2003**, *42*, 3097–3099.
- (245) Girard, P.; Namy, J. L.; Kagan, B. Divalent Lanthanide Derivatives in Organic Synthesis. I. Mild Preparation of SmI_2 and YbI_2 and Their Use as Reducing or Coupling Agents. *J. Am. Chem. Soc.* **1980**, *102* (8), 2693–2698.
- (246) Williams, A. F.; Grandjean, F.; Long, G. J.; Ulibarri, T. A.; Evans, W. J. Europium-151 Mossbauer Effect Study of Several Organoeuropium(II) Complexes. *Inorg. Chem.* **1989**, *28*, 4584–4588.
- (247) Watson, P. L.; Tulip, T. H.; Williams, I. Defluorination of Perfluoroolefins by Divalent Lanthanoid Reagents: Activating C-F Bonds. *Organometallics* **1990**, *9*, 1999–2009.
- (248) Yoshihiro, O.; Toshiyuki, I. Characterizations of Divalent Lanthanoid Iodides in Tetrahydrofuran by UV-Vis, Fluorescence and ESR Spectroscopy. *Inorg. Chim. Acta* **1988**, *144*, 143–146.
- (249) Evans, W. J.; Gummersheimer, T. S.; Ziller, J. W. Coordination Chemistry of Samarium Diiodide with Ethers Including the Crystal Structure of Tetrahydrofuran-Solvated Samarium Diiodide, $\text{SmI}_2(\text{THF})_5$. *J. Am. Chem. Soc.* **1995**, *117*, 8999–9002.
- (250) Miller, R. S.; Sealy, J. M.; Shabangi, M.; Kuhlman, M. L.; Fuchs, J. R.; Flowers, R. A. Reactions of SmI_2 with Alkyl Halides and Ketones: Inner-Sphere vs Outer-Sphere Electron Transfer in Reactions of Sm(II) Reductants. *J. Am. Chem. Soc.* **2000**, *122*, 7718–7722.
- (251) Imamoto, T.; Ono, M. The Reaction of Samarium(III) Iodide with Samarium Metal in Tetrahydrofuran, A New Method for the Preparation of Samarium(II) Iodide. *Chem. Lett.* **1987**, *16*, 501–502.
- (252) Fedushkin, I. L.; Bochkarev, M. N.; Dechert, S.; Schumann, H. A Chemical Definition of the Effective Reducing Power of Thulium(II) Diiodide by Its Reactions with Cyclic Unsaturated Hydrocarbons. *Chem. - A Eur. J.* **2001**, *7*, 3558–3563.
- (253) Bochkarev, M. N.; Fedushkin, I. L.; Fagin, A. A.; Petrovskaya, T. V.; Ziller, J. W.; Broomhall-Dillard, R. N. R.; Evans, W. J. Synthesis and Structure of the First Molecular Thulium(II) Complex: $[\text{TmI}_2(\text{MeOCH}_2\text{CH}_2\text{OMe})_3]$. *Angew. Chem., Int. Ed.* **1997**, *36*, 133–135.
- (254) Bochkarev, M. N.; Fedushkin, I. L.; Dechert, S.; Fagin, A. A.; Schumann, H. $[\text{NdI}_2(\text{Thf})_5]$, the First Crystallographically Authenticated Neodymium(II) Complex. *Angew. Chem., Int. Ed.* **2001**, *40*, 3176–3178.

- (255) Maunder, G. H.; Sella, A. Simple Adducts of Samarium and Ytterbium Diodide: Synthesis and Molecular Structures of $\text{LnI}_2(3,5\text{-Lutidine})_4$ (Ln = Sm, Yb) and $\text{YbI}_2(4\text{-t-Butylpyridine})_4$. *Polyhedron* **1998**, *17*, 63–68.
- (256) Hou, Z.; Wakatsuki, Y. Isolation and X-Ray Structures of the Hexamethylphosphoramide (Hmpa)-Coordinated Lanthanide(II) Diodide Complexes $[\text{SmI}_2(\text{Hmpa})_4]$ and $[\text{Yb}(\text{Hmpa})_4(\text{Thf})_2]\text{I}_2$. *J. Chem. Soc. Chem. Commun.* **1994**, 1205–1206.
- (257) Xémard, M.; Cordier, M.; Molton, F.; Duboc, C.; Le Guennic, B.; Maury, O.; Cador, O.; Nocton, G. Divalent Thulium Crown Ether Complexes with Field-Induced Slow Magnetic Relaxation. *Inorg. Chem.* **2019**, *58*, 2872–2880.
- (258) Huh, D. N.; Ziller, J. W.; Evans, W. J. Facile Encapsulation of Ln(II) Ions into Cryptate Complexes from $\text{LnI}_2(\text{THF})_2$ Precursors (Ln = Sm, Eu, Yb). *Inorg. Chem.* **2019**, *58*, 9613–9617.
- (259) Spedding, F. H.; Daane, A. H. The Preparation of Rare Earth Metals. *J. Am. Chem. Soc.* **1952**, *74*, 2783–2785.
- (260) Selwood, P. W. Magnetochemical Properties of Samarium. *J. Am. Chem. Soc.* **1934**, *56*, 2392–2394.
- (261) Lebrun, A.; Namy, J. L.; Kagan, H. B. Samarium Dibromide an Efficient Reagent for the Pinacol Coupling Reactions. *Tetrahedron Lett.* **1993**, *34*, 2311–2314.
- (262) Jantsch, G.; Skalla, N. Über Samarium(II)Jodid Und Den Thermischen Abbau Des Samarium(III)Jodids. *Z. Anorg. Allg. Chem.* **1930**, *193*, 391–405.
- (263) Suta, M.; Lavoie-Cardinal, F.; Wickleder, C. Underestimated Color Centers: Defects as Useful Reducing Agents in Lanthanide-Activated Luminescent Materials. *Angew. Chem., Int. Ed.* **2020**, *59*, 10949–10954.
- (264) Huang, W.; Brosmer, J. L.; Diaconescu, P. L. In Situ Synthesis of Lanthanide Complexes Supported by a Ferrocene Diamide Ligand: Extension to Redox-Active Lanthanide Ions. *New J. Chem.* **2015**, *39*, 7696–7702.
- (265) Li, W.; Fujikawa, H.; Adachi, G.-ya; Shiokawa, J. Absorption and Emission Properties of Divalent Ytterbium Crown Ether Complexes. *Inorg. Chim. Acta* **1986**, *117*, 87–89.
- (266) Voos-Esquivel, C. A.; Eick, H. A. Synthesis of YbBr_2 and YbCl_2 and an X-Ray Diffraction Study of the System $\text{YbBr}_2\text{-YbCl}_2$. *J. Solid State Chem.* **1987**, *67*, 291–296.
- (267) Schilling, G.; Kunert, C.; Schleid, T.; Meyer, G. Metallothermische Reduktion Der Tribromide Und -Iodide von Thulium Und Ytterbium Mit Alkalimetallen. *Z. Anorg. Allg. Chem.* **1992**, *618*, 7–12.
- (268) Lasocha, W.; Voos-Esquivel, C. A.; Hodorowicz, S. A.; Kim, B. Y.; Eick, H. A. An X-Ray Powder Diffraction Study of the $\text{YbCl}_2\text{-YbI}_2$ System. *J. Solid State Chem.* **1988**, *74*, 67–73.
- (269) Mudring, A. V.; Babai, A.; Arenz, S.; Giernoth, R. The “Noncoordinating” Anion Tf_2N^- Coordinates to Yb^{2+} : A Structurally Characterized Tf_2N^- Complex from the Ionic Liquid $[\text{mppyr}][\text{Tf}_2\text{N}]$. *Angew. Chem., Int. Ed.* **2005**, *44*, 5485–5488.
- (270) Van Den Hende, J. R.; Hitchcock, P. B.; Holmes, S. A.; Lappert, M. F.; Leung, W. P.; Mak, T. C. W.; Prashar, S. Synthesis and Characterisation of Lanthanide(II) Aryloxides Including the First Structurally Characterised Europium(II) Compound $[\text{Eu}(\text{OC}_6\text{H}_2\text{Bu}^i_2\text{-2,6-Me-4})_2(\text{Thf})_3]\cdot\text{thf}$ (Thf = Tetrahydrofuran). *J. Chem. Soc., Dalton Trans.* **1995**, 1427–1433.
- (271) Berg, D. J.; Burns, C. J.; Andersen, R. A.; Zalkin, A. Electron-Transfer Reactions of Divalent Ytterbium Metallocenes. Synthesis of the Series $[(\text{Me}_5\text{C}_5)_2\text{Yb}]_2[\mu\text{-E}]$ (E = O, S, Se, or Te) and Crystal Structure of $[(\text{Me}_5\text{C}_5)_2\text{Yb}]_2[\mu\text{-Se}]$. *Organometallics* **1989**, *8*, 1865–1870.
- (272) Tilley, T. D.; Andersen, R. A.; Zalkin, A. Tertiary Phosphine Complexes of the F-Block Metals. Crystal Structure of $\text{Yb}[\text{N}(\text{SiMe}_3)_2]_2[\text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2]$: Evidence for a Ytterbium- γ -Carbon Interaction. *J. Am. Chem. Soc.* **1982**, *104*, 3725–3727.
- (273) Hitchcock, P. B.; Howard, J. A. K.; Lappert, M. F.; Prashar, S. The First Examples of Intermolecular Weak (Agostic) γ -Methyl-Metal Interactions in Organolanthanide Complexes: The Synthesis and X-Ray Structures of $[\{\text{Yb}(\eta\text{-Cp}^*)\}_2]_\infty$ and $[\{\text{Eu}(\eta\text{-Cp}^*)\}_2]_\infty$ [$\text{Cp}^* = \text{C}_5\text{H}_3(\text{SiMe}_3)_2\text{-1,3}$]. *J. Organomet. Chem.* **1992**, *437*, 177–189.
- (274) Plesniak, M. P.; Just-Baringo, X.; Ortu, F.; Mills, D. P.; Procter, D. J. SmCpR2-Mediated Cross-Coupling of Allyl and Propargyl Ethers with Ketoesters and a Telescoped Approach to Complex Cycloheptanols. *Chem. Commun.* **2016**, *52*, 13503–13506.
- (275) Palumbo, C. T.; Ziller, J. W.; Evans, W. J. Structural Characterization of the Bent Metallocenes, $[\text{C}_5\text{H}_3(\text{SiMe}_3)_2]_2\text{Sm}$ and $[\text{C}_5\text{H}_3(\text{CMe}_3)_2]_2\text{Ln}$ (Ln = Eu, Sm), and the Mono(Cyclopentadienyl) Tetraphenylborate Complex, $[\text{C}_5\text{H}_3(\text{CMe}_3)_2]\text{Eu}(\mu\text{-}\eta^5\text{-}\eta^1\text{-Ph})_2\text{BPh}_2$. *J. Organomet. Chem.* **2018**, *867*, 142–148.
- (276) Kilpatrick, A. F. R.; Cloke, F. G. N. A Base-Free Synthetic Route to Anti-Bimetallic Lanthanide Pentalene Complexes. *Dalton Trans.* **2017**, *46*, 5587–5597.
- (277) Evans, W. J.; Allen, N. T.; Ziller, J. W. Expanding Divalent Organolanthanide Chemistry: The First Organothulium(II) Complex and the in Situ Organodysprosium(II) Reduction of Dinitrogen. *Angew. Chem., Int. Ed.* **2002**, *41*, 359–361.
- (278) Yatabe, T.; Karasawa, M.; Isobe, K.; Ogo, S.; Nakai, H. A Naphthyl-Substituted Pentamethylcyclopentadienyl Ligand and Its Sm(II) Bent-Metallocene Complexes with Solvent-Induced Structure Change. *Dalton Trans.* **2012**, *41*, 354–356.
- (279) Sitzmann, H.; Dezember, T.; Schmitt, O.; Weber, F.; Wolmershäuser, G.; Ruck, M. Reactions of Free Cyclopentadienyl Radicals. 3 Metallocenes of Samarium, Europium, and Ytterbium with the Especially Bulky Cyclopentadienyl Ligands $\text{C}_5\text{H}(\text{CHMe}_2)_4$, $\text{C}_5\text{H}_2(\text{CMe}_3)_3$, and $\text{C}_5(\text{CHMe}_2)_5$. *Z. Anorg. Allg. Chem.* **2000**, *626*, 2241–2244.
- (280) Evans, W. J.; Allen, N. T.; Ziller, J. W. Facile Dinitrogen Reduction via Organometallic Tm(II) Chemistry. *J. Am. Chem. Soc.* **2001**, *123*, 7927–7928.
- (281) Khoroshen'kov, G. V.; Fagin, A. A.; Bochkarev, M. N.; Dechert, S.; Schumann, H. Reactions of Neodymium(II), Dysprosium(II), and Thulium(II) Diodides with Cyclopentadiene. Molecular Structures of Complexes $\text{CpTmI}_2(\text{THF})_3$ and $[\text{NdI}_2(\text{THF})_5]^+[\text{NdI}_4(\text{THF})_2]^-$. *Russ. Chem. Bull.* **2003**, *52*, 1715–1719.
- (282) Jaroschik, F.; Nief, F.; Ricard, L. Synthesis of a New Stable, Neutral Organothulium(II) Complex by Reduction of a Thulium(III) Precursor. *Chem. Commun.* **2006**, 426–428.
- (283) Nief, F.; De Borms, B. T.; Ricard, L.; Carmichael, D. New Complexes of Divalent Thulium with Substituted Phospholyl and Cyclopentadienyl Ligands. *Eur. J. Inorg. Chem.* **2005**, *2005*, 637–643.
- (284) Cantat, T.; Jaroschik, F.; Nief, F.; Ricard, L.; Mezailles, N.; Le Floch, P. New Mono- and Bis-Carbene Samarium Complexes: Synthesis, X-Ray Crystal Structures and Reactivity. *Chem. Commun.* **2005**, 5178–5180.
- (285) Nief, F.; Turcitu, D.; Ricard, L. Synthesis and Structure of Phospholyl- and Arsolylthulium(II) Complexes. *Chem. Commun.* **2002**, 1646–1647.
- (286) Turcitu, D.; Nief, F.; Ricard, L. Structure and Reactivity of Homoleptic Samarium(II) and Thulium(II) Phospholyl Complexes. *Chem. - A Eur. J.* **2003**, *9*, 4916–4923.
- (287) Tilley, T. D.; Andersen, R. A.; Zalkin, A. Divalent Lanthanide Chemistry. Preparation and Crystal Structures of Sodium Tris[Bis-(Trimethylsilyl)Amido]Europate(II) and Sodium Tris[Bis-(Trimethylsilyl)Amido]Ytterbate(II), $\text{NaM}[\text{N}(\text{SiMe}_3)_2]_3$. *Inorg. Chem.* **1984**, *23*, 2271–2276.
- (288) Evans, W. J.; Drummond, D. K.; Zhang, H.; Atwood, J. L. Synthesis and X-Ray Crystal Structure of the Divalent [Bis-(Trimethylsilyl)Amido] Samarium Complexes $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{Sm}(\text{THF})_2$ and $\{[(\text{Me}_3\text{Si})_2\text{N}]\text{Sm}(\mu\text{-I})(\text{DME})(\text{THF})\}_2$. *Inorg. Chem.* **1988**, *27* (3), 575–579.
- (289) Evans, W. J.; Zucchi, G.; Ziller, J. W. Dinitrogen Reduction by Tm(II), Dy(II), and Nd(II) with Simple Amide and Aryloxide Ligands. *J. Am. Chem. Soc.* **2003**, *125*, 10–11.
- (290) Evans, W. J.; Lee, D. S.; Rego, D. B.; Perotti, J. M.; Kozimor, S. A.; Moore, E. K.; Ziller, J. W. Expanding Dinitrogen Reduction Chemistry to Trivalent Lanthanides via the LnZ_3 /Alkali Metal Reduction System: Evaluation of the Generality of Forming $\text{Ln}_2(\mu\text{-}$

- $\eta^2:\eta^2\text{-N}_2$) Complexes via LnZ_3/K . *J. Am. Chem. Soc.* **2004**, *126*, 14574–14582.
- (291) Evans, W. J.; Fang, M.; Zucchi, G. L.; Furche, F.; Ziller, J. W.; Hoekstra, R. M.; Zink, J. I. Isolation of Dysprosium and Yttrium Complexes of a Three-Electron Reduction Product in the Activation of Dinitrogen, the $(\text{N}_2)^{3-}$ Radical. *J. Am. Chem. Soc.* **2009**, *131*, 11195–11202.
- (292) Goodwin, C. A. P.; Chilton, N. F.; Vettese, G. F.; Moreno Pineda, E.; Crowe, I. F.; Ziller, J. W.; Winpenny, R. E. P.; Evans, W. J.; Mills, D. P. Physicochemical Properties of Near-Linear Lanthanide(II) Bis(Silylamide) Complexes (Ln = Sm, Eu, Tm, Yb). *Inorg. Chem.* **2016**, *55*, 10057–10067.
- (293) Chilton, N. F.; Goodwin, C. A. P.; Mills, D. P.; Winpenny, R. E. P. The First Near-Linear Bis(Amide) f-Block Complex: A Blueprint for a High Temperature Single Molecule Magnet. *Chem. Commun.* **2015**, *51*, 101–103.
- (294) Goodwin, C. A. P.; Chilton, N. F.; Natrajan, L. S.; Boulon, M. E.; Ziller, J. W.; Evans, W. J.; Mills, D. P. Investigation into the Effects of a Trigonal-Planar Ligand Field on the Electronic Properties of Lanthanide(II) Tris(Silylamide) Complexes (Ln = Sm, Eu, Tm, Yb). *Inorg. Chem.* **2017**, *56*, 5959–5970.
- (295) Moutet, J.; Schleinitz, J.; La Droitte, L.; Tricoire, M.; Pointillart, F.; Gendron, F.; Simler, T.; Clavaguéra, C.; Le Guennic, B.; Cador, O.; et al. Bis-Cyclooctatetraenyl Thulium(II): Highly Reducing Lanthanide Sandwich Single-Molecule Magnets. *Angew. Chem., Int. Ed.* **2021**, *60*, 6042–6046.
- (296) Xémard, M.; Zimmer, S.; Cordier, M.; Goudy, V.; Ricard, L.; Clavaguéra, C.; Nocton, G. Lanthanidocenes: Synthesis, Structure, and Bonding of Linear Sandwich Complexes of Lanthanides. *J. Am. Chem. Soc.* **2018**, *140*, 14433–14439.
- (297) Clegg, W.; Eaborn, C.; Izod, K.; O'Shaughnessy, P.; Smith, J. D. The First Structurally Authenticated σ -Bonded Organosamarium(II) Derivative and Its Reaction with Benzophenone. *Angew. Chem., Int. Ed.* **1997**, *36*, 2815–2817.
- (298) Bowman, L. J.; Izod, K.; Clegg, W.; Harrington, R. W. Synthesis and Structures of Ln(II) and Ln(III) Dialkyls Derived from LnI_2 (Ln = Nd, Tm, Yb). *Organometallics* **2007**, *26*, 2646–2651.
- (299) Hitchcock, P. B.; Holmes, S. A.; Lappert, M. F.; Tian, S. Synthesis, Structures and Reactions of Ytterbium(II) Alkyls. *J. Chem. Soc., Chem. Commun.* **1994**, 2691–2692.
- (300) Yan, K. K.; Upton, B. M.; Ellern, A.; Sadow, A. D. Lewis Acid-Mediated β -Hydride Abstraction Reactions of Divalent $\text{M}(\text{C}(\text{SiHMe}_2)_3)_2\text{THF}_2$ (M = Ca, Yb). *J. Am. Chem. Soc.* **2009**, *131* (42), 15110–15111.
- (301) Yan, K.; Schoendorff, G.; Upton, B. M.; Ellern, A.; Windus, T. L.; Sadow, A. D. Intermolecular β -Hydrogen Abstraction in Ytterbium, Calcium, and Potassium Tris(Dimethylsilyl)Methyl Compounds. *Organometallics* **2013**, *32*, 1300–1316.
- (302) Pindwal, A.; Ellern, A.; Sadow, A. D. Homoleptic Divalent Dialkyl Lanthanide-Catalyzed Cross-Dehydrocoupling of Silanes and Amines. *Organometallics* **2016**, *35*, 1674–1683.
- (303) Hitchcock, P. B.; Khvostov, A. V.; Lappert, M. F. Synthesis and Structures of Crystalline Bis(Trimethylsilyl)Methanido Complexes of Potassium, Calcium and Ytterbium. *J. Organomet. Chem.* **2002**, *663*, 263–268.
- (304) Hill, M. S.; Hitchcock, P. B. Synthesis of a Homoleptic Sm(II) Bis(Phosphinimino)Methanide. *Dalton Trans.* **2003**, 608, 4570–4571.
- (305) Hitchcock, P. B.; Khvostov, A. V.; Lappert, M. F.; Protchenko, A. V. Heteroleptic Ytterbium(II) Complexes Supported by a Bulky β -Diketiminato Ligand. *Dalton Trans.* **2009**, 2383–2391.
- (306) Takats, J.; Zhang, W.; et al. Synthesis and Structure of Bis[Hydrotris(3,5-Dimethylpyrazolyl)Borato]Samarium(II), $\text{Sm}[\text{HB}(\text{3,5-Me}_2\text{pz})_3]_2$, and the Product of Its Reaction with Azobenzene. *Organometallics* **1993**, *12*, 4286–4288.
- (307) Hillier, A. C.; Zhang, X. W.; Maunder, G. H.; Liu, S. Y.; Eberspacher, T. A.; Metz, M. V.; McDonald, R.; Domingos, A.; Marques, N.; Day, V. W.; et al. Synthesis and Structural Comparison of a Series of Divalent $\text{Ln}(\text{Tp}^{\text{RR}})_2$ (Ln = Sm, Eu, Yb) and Trivalent $\text{Sm}(\text{Tp}^{\text{Me}_2})_2\text{X}$ (X = F, Cl, I, BPh_4) Complexes. *Inorg. Chem.* **2001**, *40*, 5106–5116.
- (308) Cheng, J.; Takats, J.; Ferguson, M. J.; McDonald, R. Heteroleptic Tm(II) Complexes: One More Success for Trofimenko's Scorpionates. *J. Am. Chem. Soc.* **2008**, *130*, 1544–1545.
- (309) Zhang, X. W.; Maunder, G. H.; Gießmann, S.; Macdonald, R.; Ferguson, M. J.; Bond, A. H.; Rogers, R. D.; Takats, J. Stable Heteroleptic Complexes of Divalent Lanthanides with Bulky Pyrazolylborate Ligands – Iodides, Hydrocarbyls and Triethylborohydrides. *Dalton Trans.* **2011**, *40*, 195–210.
- (310) Momin, A.; Carter, L.; Yang, Y.; McDonald, R.; Essafi née Labouille, S.; Nief, F.; Del Rosal, I.; Sella, A.; Maron, L.; Takats, J. To Bend or Not To Bend: Experimental and Computational Studies of Structural Preference in $\text{Ln}(\text{Tp}^{\text{IPr}_2})_2$ (Ln = Sm, Tm). *Inorg. Chem.* **2014**, *53*, 12066–12075.
- (311) Izod, K.; O'Shaughnessy, P.; Sheffield, J. M.; Clegg, W.; Liddle, S. T. Synthesis and Structural Characterization of Sm(II) and Yb(II) Complexes Containing Sterically Demanding, Chelating Secondary Phosphide Ligands. *Inorg. Chem.* **2000**, *39*, 4741–4748.
- (312) Zitz, R.; Hlina, J.; Gatterer, K.; Marschner, C.; Szilvási, T.; Baumgartner, J. Neutral “Cp-Free” Silyl-Lanthanide(II) Complexes: Synthesis, Structure, and Bonding Analysis. *Inorg. Chem.* **2015**, *54*, 7065–7072.
- (313) Réant, B. L. L.; Berryman, V. E. J.; Basford, A. R.; Nodaraki, L. E.; Wooles, A. J.; Tuna, F.; Kaltsayannis, N.; Mills, D. P.; Liddle, S. T. ^{29}Si NMR Spectroscopy as a Probe of s- And f-Block Metal(II)-Silanide Bond Covalency. *J. Am. Chem. Soc.* **2021**, *143*, 9813–9824.
- (314) Jones, C.; Stasch, A.; Woodul, W. D. Gallyl Lanthanide Complexes Containing Unsupported Ln-Ga (Ln = Sm, Eu, Yb or Tm) Bonds. *Chem. Commun.* **2009**, 113–115.
- (315) Hitchcock, P. B.; Khvostov, A. V.; Lappert, M. F.; Protchenko, A. V. Heteroleptic Ytterbium(II) Complexes Supported by a Bulky β -Diketiminato Ligand. *Dalton Trans.* **2009**, 2383–2391.
- (316) Jones, C.; Nembenna, S.; Stasch, A. Synthesis and Crystal Structure of a Bulky β -Diketiminato Ytterbium(II) Iodide Complex. *J. Chem. Crystallogr.* **2011**, *41*, 1490–1493.
- (317) Wen, Q.; Feng, B.; Xiang, L.; Leng, X.; Chen, Y. Divalent Ytterbium Hydrido Complex Supported by a β -Diketiminato-Based Tetradentate Ligand: Synthesis, Structure, and Reactivity. *Inorg. Chem.* **2021**, *60*, 13913–13919.
- (318) Liu, X.; Xiang, L.; Louyriac, E.; Maron, L.; Leng, X.; Chen, Y. Divalent Ytterbium Complex-Catalyzed Homo- and Cross-Coupling of Primary Arylsilanes. *J. Am. Chem. Soc.* **2019**, *141*, 138–142.
- (319) Liu, X.; Wen, Q.; Xiang, L.; Leng, X.; Chen, Y. Samarium(II) Monoalkyl Complex Supported by a β -Diketiminato-Based Tetradentate Ligand: Synthesis, Structure, and Catalytic Hydrosilylation of Internal Alkynes. *Chem. - A Eur. J.* **2020**, *26*, 5494–5499.
- (320) Xu, C.; Ye, Z.; Xiang, L.; Yang, S.; Peng, Q.; Leng, X.; Chen, Y. Insertion of Metal-Substituted Silylene into Naphthalene's Aromatic Ring and Subsequent Rearrangement for Silaspiro-Benzocycloheptenyl and Cyclobutenosilaindan Derivatives. *Angew. Chem., Int. Ed.* **2021**, *60*, 3189–3195.
- (321) Heitmann, D.; Jones, C.; Junk, P. C.; Lippert, K.; Stasch, A. Homoleptic Lanthanide(II)– Bis(Guanidinate) Complexes, $[\text{Ln}(\text{Giso})_2]$ (Giso = or Eu) vs Distorted Tetrahedral (Ln = Yb) Geometries. *Dalton Trans.* **2007**, 187–189.
- (322) Heitmann, D.; Jones, C.; Mills, D. P.; Stasch, A. Low Coordinate Lanthanide(II) Complexes Supported by Bulky Guanidinate and Amidinate Ligands. *Dalton Trans.* **2010**, 39, 1877.
- (323) Basalov, I. V.; Yurova, O. S.; Cherkasov, A. V.; Fukin, G. K.; Trifonov, A. A. Amido Ln(II) Complexes Coordinated by Bi- and Tridentate Amidinate Ligands: Nonconventional Coordination Modes of Amidinate Ligands and Catalytic Activity in Intermolecular Hydrophosphination of Styrenes and Toluene. *Inorg. Chem.* **2016**, *55*, 1236–1244.
- (324) Wiecko, M.; Roesky, P. W.; Burlakov, V. V.; Spannenberg, A. Bis(Phosphinimino)Methanides as Ligands in Divalent Samarium Chemistry: Synthesis, Structures and Catalysis. *Eur. J. Inorg. Chem.* **2007**, *2007*, 876–881.

- (325) Panda, T. K.; Zulus, A.; Gamer, M. T.; Roesky, P. W. Bis(Phosphinimino)Methanides as Ligands in Divalent Lanthanide and Alkaline Earth Chemistry - Synthesis, Structure, and Catalysis. *J. Organomet. Chem.* **2005**, *690*, 5078–5089.
- (326) Cheng, J.; Takats, J.; Ferguson, M. J.; McDonald, R. Heteroleptic Tm(II) Complexes: One More Success for Trofimenko's Scorpionates. *J. Am. Chem. Soc.* **2008**, *130*, 1544–1545.
- (327) Reid, A. F.; Wailes, P. C. The Reaction of Molten Magnesium Cyclopentadienide with Fluorides and Other Metal Halides. *Inorg. Chem.* **1966**, *5* (7), 1213–1216.
- (328) Bottomley, F.; Paez, D. E.; White, P. S. The Reaction of Scandium Trifluoride with Cyclopentadienyl Salts and the Crystal and Molecular Structure of the Trimer of Fluorodicyclopentadienylscandium. *J. Organomet. Chem.* **1985**, *291*, 35–41.
- (329) Ding, Y. S.; Yu, K. X.; Reta, D.; Ortu, F.; Winpenny, R. E. P.; Zheng, Y. Z.; Chilton, N. F. Field- and Temperature-Dependent Quantum Tunnelling of the Magnetisation in a Large Barrier Single-Molecule Magnet. *Nat. Commun.* **2018**, *9*, 3134.
- (330) Ortu, F.; Reta, D.; Ding, Y. S.; Goodwin, C. A. P.; Gregson, M. P.; McInnes, E. J. L.; Winpenny, R. E. P.; Zheng, Y. Z.; Liddle, S. T.; Mills, D. P.; et al. Studies of Hysteresis and Quantum Tunnelling of the Magnetisation in Dysprosium(III) Single Molecule Magnets. *Dalton Trans.* **2019**, *48* (24), 8541–8545.
- (331) Goodwin, C. A. P.; Schlimgen, A. W.; Albrecht-Schönzart, T. E.; Batista, E. R.; Gaunt, A. J.; Janicke, M. T.; Kozimor, S. A.; Scott, B. L.; Stevens, L. M.; White, F. D.; et al. Structural and Spectroscopic Comparison of Soft-Se vs. Hard-O Donor Bonding in Trivalent Americium/Neodymium Molecules. *Angew. Chem., Int. Ed.* **2021**, *60*, 9459–9466.
- (332) Windorff, C. J.; Sperling, J. M.; Albrecht-Schönzart, T. E.; Bai, Z.; Evans, W. J.; Gaiser, A. N.; Gaunt, A. J.; Goodwin, C. A. P.; Hobart, D. E.; Huffman, Z. K.; et al. A Single Small-Scale Plutonium Redox Reaction System Yields Three Crystallographically-Characterizable Organoplutonium Complexes. *Inorg. Chem.* **2020**, *59*, 13301–13314.
- (333) Goodwin, C. A. P.; Su, J.; Albrecht-Schmitt, T. E.; Blake, A. V.; Batista, E. R.; Daly, S. R.; Dehnen, S.; Evans, W. J.; Gaunt, A. J.; Kozimor, S. A.; et al. [Am(C₅Me₄H)₃]: An Organometallic Americium Complex. *Angew. Chem., Int. Ed.* **2019**, *58*, 11695–11699.
- (334) Wu, J.; Boyle, T. J.; Shreeve, J. L.; Ziller, J. W.; Evans, W. J. CP/MAS ⁸⁹Y NMR Spectroscopy: A Facile Method for Characterizing Yttrium-Containing Solids. *Inorg. Chem.* **1993**, *32*, 1130–1134.
- (335) Izod, K.; Liddle, S. T.; Clegg, W. A Convenient Route to Lanthanide Triiodide THF Solvates. Crystal Structures of LnI₃(THF)₄ [Ln = Pr] and LnI₃(THF)_{3.5} [Ln = Nd, Gd, Y]. *Inorg. Chem.* **2004**, *43*, 214–218.
- (336) Reed, J. B.; Hopkins, B. S.; Audrieth, L. F. Observations on the Rare Earths. XLIV. Preparation of Rare Earth Compounds by the Action of Fused and Solid "Onium" Salts on the Oxides. *J. Am. Chem. Soc.* **1935**, *57*, 1159–1160.
- (337) Meyer, G.; et al. The Ammonium Chloride Route To Anhydrous Rare Earth Chlorides-the Example of YCl₃. *Inorg. Synth.* **1973**, *25*, 146–150.
- (338) Young, R. C.; Hastings, J. L. Reaction of Lanthanum Oxide with Ammonium Iodide. *J. Am. Chem. Soc.* **1937**, *59*, 765–766.
- (339) Huang, W.; Upton, B. M.; Khan, S. I.; Diaconescu, P. L. Synthesis and Characterization of Paramagnetic Lanthanide Benzyl Complexes. *Organometallics* **2013**, *32*, 1379–1386.
- (340) Edleman, N. L.; Wang, A.; Belot, J. A.; Metz, A. W.; Babcock, J. R.; Kawaoka, A. M.; Ni, J.; Metz, M. V.; Flaschenriem, C. J.; Stern, C. L.; Liable-Sands, L. M.; et al. Synthesis and Characterization of Volatile, Fluorine-Free β-Ketoiminate Lanthanide MOCVD Precursors and Their Implementation in Low-Temperature Growth of Epitaxial CeO₂ Buffer Layers for Superconducting Electronics. *Inorg. Chem.* **2002**, *41*, 5005–5023.
- (341) Gompa, T. P.; Rice, N. T.; Russo, D. R.; Aguirre Quintana, L. M.; Yik, B. J.; Bacsa, J.; La Pierre, H. S. Diethyl Ether Adducts of Trivalent Lanthanide Iodides. *Dalton Trans.* **2019**, *48*, 8030–8033.
- (342) Windorff, C. J.; Dumas, M. T.; Ziller, J. W.; Gaunt, A. J.; Kozimor, S. A.; Evans, W. J. Small-Scale Metal-Based Syntheses of Lanthanide Iodide, Amide, and Cyclopentadienyl Complexes as Analogues for Transuranic Reactions. *Inorg. Chem.* **2017**, *56*, 11981–11989.
- (343) Deacon, G. B.; Feng, T.; Junk, P. C.; Meyer, G.; Scott, N. M.; Skelton, B. W.; White, A. H. Structural Variety in Solvated Lanthanoid(III) Halide Complexes. *Aust. J. Chem.* **2000**, *53*, 853–865.
- (344) Petriček, S. Syntheses of Lanthanide Bromide Complexes from Oxides and the Crystal Structures of [LnBr₃(DME)₂] (Ln = Pr, Nd, Sm, Eu), [LnBr₃(THF)₄] (Ln = Pr, Sm) and [EuBr₂(THF)₅]-[EuBr₄(THF)₂]. *Polyhedron* **2004**, *23*, 2293–2301.
- (345) Deacon, G. B.; Koplick, A. J. A Convenient Synthesis of Lanthanoid Trihalides in Tetrahydrofuran. *Inorg. Nucl. Chem. Lett.* **1979**, *15*, 263–265.
- (346) Carter, F. L.; Murray, J. F. Preparation of the Anhydrous Rare Earth Trichlorides, Tribromides, and Triiodides. *Mater. Res. Bull.* **1972**, *7*, 519–523.
- (347) Reed, J. B.; Hopkins, B. S.; Audrieth, L. F.; et al. Anhydrous Rare Earth Chlorides. *Inorg. Synth.* **1939**, *1*, 28–33.
- (348) Taylor, M. D.; Carter, C. P. Preparation of Anhydrous Lanthanide Halides, Especially Iodides. *J. Inorg. Nucl. Chem.* **1962**, *24*, 387–391.
- (349) Meyer, G.; Dötsch, S.; Staffel, T. The Ammonium-Bromide Route to Anhydrous Rare Earth Bromides MBr₃. *J. Less-Common Met.* **1987**, *127*, 155–160.
- (350) Carver, C. T.; Monreal, M. J.; Diaconescu, P. L. Scandium Alkyl Complexes Supported by a Ferrocene Diamide Ligand. *Organometallics* **2008**, *27*, 363–370.
- (351) Huang, W.; Brosmer, J. L.; Diaconescu, P. L. In Situ Synthesis of Lanthanide Complexes Supported by a Ferrocene Diamide Ligand: Extension to Redox-Active Lanthanide Ions. *New J. Chem.* **2015**, *39*, 7696–7702.
- (352) Freeman, J. H.; Smith, M. L. The Preparation of Anhydrous Inorganic Chlorides by Dehydration with Thionyl Chloride. *J. Inorg. Nucl. Chem.* **1958**, *7*, 224–227.
- (353) Wu, S. H.; Ding, Z. B.; Li, X. J. A Facile Method for Preparation of Tetrahydrofuran Complexes of Lanthanide Trichlorides. *Polyhedron* **1994**, *13*, 2679–2681.
- (354) Hazin, P. N.; Huffman, J. C.; Bruno, J. W. Synthetic and Structural Studies of Pentamethylcyclopentadienyl Complexes of Lanthanum and Cerium. *Organometallics* **1987**, *6*, 23–27.
- (355) Carver, C. T.; Monreal, M. J.; Diaconescu, P. L. Scandium Alkyl Complexes Supported by a Ferrocene Diamide Ligand. *Organometallics* **2008**, *27*, 363–370.
- (356) Atwood, J. L.; Smith, K. D. Crystal and Molecular Structure of Trichlorotris(Tetrahydrofuran)Scandium(III). *J. Chem. Soc., Dalton Trans.* **1974**, 921–923.
- (357) Boyle, T. J.; Ottley, L. A. M.; Alam, T. M.; Rodriguez, M. A.; Yang, P.; McIntyre, S. K. Structural Characterization of Methanol Substituted Lanthanum Halides. *Polyhedron* **2010**, *29*, 1784–1795.
- (358) Coles, M. P.; Hitchcock, P. B.; Lappert, M. F.; Protchenko, A. V. Syntheses and Structures of the Crystalline, Highly Crowded 1,3-Bis(Trimethylsilyl)Cyclopentadienyls [MCp^{''}₃] (M = Y, Er, Yb), [PbCp^{''}₂], [{YCp^{''}₂(μ-OH)}₂], [(ScCp^{''}₂)₂(μ-η²:η²-C₂H₄)], [YbCp^{''}₂Cl(μ-Cl)K(18-crown-6)], and [{KCp^{''}}_∞]. *Organometallics* **2012**, *31*, 2682–2690.
- (359) Sobota, P.; Utoko, J.; Szafert, S. Ionization of YCl₃ in Tetrahydrofuran. Crystal Structures of the [trans-YCl₂(THF)₃][trans-YCl₄(THF)₂] Salt and Polymeric [YCl₃·2THF]_∞ Compounds. *Inorg. Chem.* **1994**, *33*, 5203–5206.
- (360) Martinez-Arripe, E.; Jean-Baptiste-Dit-Dominique, F.; Auffrant, A.; Le Goff, X. F.; Thuilliez, J.; Nief, F. Synthesis and Characterization of Bidentate Rare-Earth Iminophosphorane o-Aryl Complexes and Their Behavior as Catalysts for the Polymerization of 1,3-Butadiene. *Organometallics* **2012**, *31*, 4854–4861.
- (361) Mou, Z.; Liu, B.; Liu, X.; Xie, H.; Rong, W.; Li, L.; Li, S.; Cui, D. Efficient and Heteroselective Heteroscorpionate Rare-Earth-Metal Zwitterionic Initiators for ROP of Rac-Lactide: Role of σ-Ligand. *Macromolecules* **2014**, *47*, 2233–2241.

- (362) Avent, A. G.; Cloke, F. G. N.; Elvidge, B. R.; Hitchcock, P. B. Yttrium Complexes Incorporating the Chelating Diamides $[\text{ArN}(\text{CH}_2)_x\text{NAr}]^{2-}$ (Ar = C_6H_3 -2,6- Pr_2 , $x = 2, 3$) and Their Unusual Reaction with Phenylsilane. *Dalton Trans.* **2004**, 1083–1096.
- (363) Kauzlarich, S. M.; Hughbanks, T.; Corbett, J. D.; Klavins, P.; Shelton, R. N. Two Extended Metal Chain Compounds, $\text{Y}_4\text{I}_3\text{C}$ and $\text{Y}_6\text{I}_7\text{C}_2$. Synthesis, Structure, Properties, and Bonding. *Inorg. Chem.* **1988**, *27*, 1791–1797.
- (364) Deacon, G. B.; Feng, T.; Junk, P. C.; Skelton, B. W.; Sobolev, A. N.; White, A. H. Preparation and X-Ray Crystal Structures of Tetrahydrofuran-Complexed Rare Earth Chlorides — a Structurally Rich Series. *Aust. J. Chem.* **1998**, *51*, 75–89.
- (365) Sánchez-Barba, L. F.; Hughes, D. L.; Humphrey, S. M.; Bochmann, M. New Bis(Allyl)(Diketiminato) and Tris(Allyl) Lanthanide Complexes and Their Reactivity in the Polymerization of Polar Monomers. *Organometallics* **2005**, *24*, 3792–3799.
- (366) Deacon, G. B.; Forsyth, C. M.; Junk, P. C.; Skelton, B. W.; White, A. H. Lithium and Lanthanoid(III) Complexes of the Chelating N,N-Dimethyl-N'-Trimethylsilyl-ethane-1,2-Diamine(1-) Ligand. *J. Chem. Soc., Dalton Trans.* **1998**, 1381–1388.
- (367) Carver, C. T.; Williams, B. N.; Ogilby, K. R.; Diaconescu, P. L. Coupling of Aromatic N-Heterocycles Mediated by Group 3 Complexes. *Organometallics* **2010**, *29*, 835–846.
- (368) Trifonov, A. A.; Van De Weghe, P.; Collin, J.; Domingos, A.; Santos, I. Synthesis of Lanthanide Complexes Coordinated by an Asymmetric Cyclopentadienyl Ligand. *J. Organomet. Chem.* **1997**, *527*, 225–237.
- (369) Evans, W. J.; Shreeve, J. L.; Ziller, J. W.; Doedens, R. J. Structural Diversity in Solvated Lanthanide Halide Complexes. *Inorg. Chem.* **1995**, *34*, 576–585.
- (370) Broderick, E. M.; Thuy-Boun, P. S.; Guo, N.; Vogel, C. S.; Sutter, J.; Miller, J. T.; Meyer, K.; Diaconescu, P. L. Synthesis and Characterization of Cerium and Yttrium Alkoxide Complexes Supported by Ferrocene-Based Chelating Ligands. *Inorg. Chem.* **2011**, *50*, 2870–2877.
- (371) Broderick, E. M.; Guo, N.; Wu, T.; Vogel, C. S.; Xu, C.; Sutter, J.; Miller, J. T.; Meyer, K.; Cantat, T.; Diaconescu, P. L. Redox Control of a Polymerization Catalyst by Changing the Oxidation State of the Metal Center. *Chem. Commun.* **2011**, *47*, 9897–9899.
- (372) Hitchcock, P. B.; Hulkes, A. G.; Lappert, M. F. Oxidation in Nonclassical Organolanthanide Chemistry: Synthesis, Characterization, and X-Ray Crystal Structures of Cerium(III) and -(IV) Amides. *Inorg. Chem.* **2004**, *43*, 1031–1038.
- (373) Liddle, S. T.; Arnold, P. L. Synthesis of Heteroleptic Cerium(III) Anionic Amido-Tethered N-Heterocyclic Carbene Complexes. *Organometallics* **2005**, *24*, 2597–2605.
- (374) Willey, G. R.; Woodman, T. J.; Drew, M. G. B. Lanthanide-(III) Chloride-Tetrahydrofuran Solvates: Structural Patterns within the Series $\text{LnCl}_3(\text{THF})_n$, Where $n = 2, 3, 3.5$ and 4: Crystal and Molecular Structures of $[\text{PrCl}(\mu\text{-Cl})_2(\text{THF})_2]_n$, $[\text{Nd}(\mu\text{-Cl})_3(\text{H}_2\text{O})(\text{THF})]_n$ and $\text{GdCl}_3(\text{THF})_4$. *Polyhedron* **1997**, *16*, 3385–3393.
- (375) Wang, Z.; Hu, N.; Sakata, K.; Hashimoto, M. Syntheses, Characterization and Crystal Structures of 5,14-Dihydro-6,8,15,17-Tetramethyldibenzo[*b,i*][1,4,8,11]Tetraazacyclotetradecine Rare Earth(III) Complexes Zenglin. *J. Chem. Soc., Dalton Trans.* **1999**, 1695–1700.
- (376) Wenqi, C.; Zhongsheng, J.; Yan, X.; Yuguo, F.; Guangdi, Y. Crystal Structure of $\text{NdCl}_3 \cdot 4\text{THF}$ and Its Catalytic Activity in Polymerization of Diene. *Inorg. Chim. Acta* **1987**, *130*, 125–129.
- (377) Balashova, T. V.; Kussyayev, D. M.; Kulikova, T. I.; Kuznetsova, O. N.; Edelmann, F. T.; Gießmann, S.; Blaurock, S.; Bochkarev, M. N. Use of Neodymium Diiodide in the Synthesis of Organosilicon, -Germanium and -Tin Compounds. *Z. Anorg. Allg. Chem.* **2007**, *633*, 256–260.
- (378) Clark, D. L.; Gordon, J. C.; Scott, B. L.; Watkin, J. G. Synthesis and Characterization of a Mixed-Ring Bis-Cyclopentadienyl Derivative of Neodymium. X-Ray Crystal Structures of $(\eta\text{-C}_5\text{Me}_5)\text{NdI}_2(\text{Py})_3$ and $(\eta\text{-C}_5\text{Me}_5)(\eta\text{-C}_3\text{H}_4\text{SiMe}_3)\text{NdI}(\text{Py})$. *Polyhedron* **1999**, *18* (10), 1389–1396.
- (379) Anfang, S.; Karl, M.; Faza, N.; Massa, W.; Dehnicke, K.; Magull, J. Synthese Und Kristallstrukturen Der Seltenerde-Komplexe $[\text{LaI}_2(\text{THF})_5]^+ \text{I}_3^-$, $[\text{SmCl}_3(\text{THF})_4]$, $[\text{ErCl}_2(\text{THF})_5]^+ [\text{ErCl}_4(\text{THF})_2]^-$, $[\text{ErCl}_3(\text{DME})_2]$ Und $[\text{Na}(18\text{-Krone-6})(\text{THF})_2]^+ [\text{YbBr}_4(\text{THF})_2]^-$. *Z. Anorg. Allg. Chem.* **1997**, *623*, 1425–1432.
- (380) Zamora, M. T.; Johnson, K. R. D.; Hänninen, M. M.; Hayes, P. G. Differences in the Cyclometalation Reactivity of Bisphosphinimine-Supported Organo-Rare Earth Complexes. *Dalton Trans.* **2014**, *43*, 10739–10750.
- (381) Collin, J.; Giuseppone, N.; Jaber, N.; Domingos, A.; Maria, L.; Santos, I. Iodo Bis Bistrimethylsilylamido Lanthanides. *J. Organomet. Chem.* **2001**, *628*, 271–274.
- (382) Bartenbach, D.; Wenzel, O.; Popescu, R.; Faden, L. P.; Reiß, A.; Kaiser, M.; Zimina, A.; Grunwaldt, J. D.; Gerthsen, D.; Feldmann, C. Liquid-Phase Synthesis of Highly Reactive Rare-Earth Metal Nanoparticles. *Angew. Chem., Int. Ed.* **2021**, *60*, 17373–17377.
- (383) Xie, Z.; Chiu, K. Y.; Wu, B.; Mak, T. C. W. Autoionization of SmI_3 in Tetrahydrofuran. X-Ray Crystal Structure of the Ionic Complex $[\text{SmI}_2(\text{THF})_5][\text{SmI}_4(\text{THF})_2]$. *Inorg. Chem.* **1996**, *35*, 5957–5958.
- (384) Hitchcock, P. B.; Lappert, M. F.; Tian, S. Synthesis, Characterisation and Reactions of 1,3-Bis(Trimethylsilyl)-1-Aza-Allyl-Lanthanide Complexes; X-Ray Structures of $[\text{Sm}(\text{LL}')_2](\text{Thf})$, $[\text{Yb}(\text{LL}')_2]$ and $[\text{RN}=\text{C}(\text{Bu}^i)\text{CH}(\text{R})_2]$ (Thf = Tetrahydrofuran, $\text{LL}' = \eta^3\text{-N}(\text{R})\text{C}(\text{Bu}^i)\text{CHR}$, R = SiMe₃). *J. Organomet. Chem.* **1997**, *549*, 1–12.
- (385) Lin, S. - H.; Dong, Z. - C.; Huang, J. - S.; Zhang, Q. - E.; Lu, J. - X. Structure of Trichlorotetrakis(Tetrahydrofuran)Europium(III). *Acta Crystallogr. Sect. C* **1991**, *47*, 426–427.
- (386) Joseph, S.; Radhakrishnan, P. K. Rare Earth Iodide Complexes of 4-Formyl-2, 3-Dimethyl-1-Phenyl-3-Pyrazolin-5-One. *Synth. React. Inorg. Met. Chem.* **1982**, *12*, 1219–1229.
- (387) Willey, G. R.; Meehan, P. R.; Rudd, M. D.; Clase, H. J.; Alcock, N. W. Synthesis Crystal Structure of $[\text{GdCl}_2\text{dibenzo-18-crown-6-MeCN}][\text{SbCl}_6] \cdot 2\text{MeCN}$ dibenzo-18-crown-6-2,3,11,12-dibenzo-1,4,7,10,13,16-hexaoxacyclooctadeca-2,11-diene. *Inorg. Chim. Acta* **1994**, *215*, 209–213.
- (388) Roitershtein, D. M.; Puntus, L. N.; Vinogradov, A. A.; Lyssenko, K. A.; Minyaev, M. E.; Dobrokhodov, M. D.; Taidakov, I. V.; Varaksina, E. A.; Churakov, A. V.; Nifant'Ev, I. E. Polyphenylcyclopentadienyl Ligands as an Effective Light-Harvesting μ -Bonded Antenna for Lanthanide +3 Ions. *Inorg. Chem.* **2018**, *57*, 10199–10213.
- (389) Willey, G. R.; Meehan, P. R.; Woodman, T. J.; Drew, M. G. B. Identification of the Dysprosium(III) Chloride Solvate $\text{DyCl}_3(\text{thf})_{3.5}$: Crystal of the Ion Pair $[\text{Trans-DyCl}_2(\text{thf})_5][\text{trans-DyCl}_4(\text{thf})_2]$. *Polyhedron* **1997**, *16*, 623–627.
- (390) Pan, Y.; Xu, T.; Yang, G. W.; Jin, K.; Lu, X. B. Bis(Oxazolonyl)-Phenyl-Ligated Rare-Earth-Metal Complexes: Highly Regioselective Catalysts for Cis -1,4-Polymerization of Isoprene. *Inorg. Chem.* **2013**, *52*, 2802–2808.
- (391) Raeder, J.; Reiners, M.; Baumgarten, R.; Münster, K.; Baabe, D.; Freytag, M.; Jones, P. G.; Walter, M. D. Synthesis and Molecular Structure of Pentadienyl Complexes of the Rare-Earth Metals. *Dalton Trans.* **2018**, *47*, 14468–14482.
- (392) Long, J.; Selikhov, A. N.; Mamontova, E.; Lyssenko, K. A.; Guari, Y.; Larionova, J.; Trifonov, A. A. Single-Molecule Magnet Behaviour in a Dy(III) Pentagonal Bipyramidal Complex with a Quasi-Linear Cl-Dy-Cl Sequence. *Dalton Trans.* **2019**, *48*, 35–39.
- (393) Petriček, S. Synthesis and Structural Similarities of Yttrium and Lanthanide Chloride Complexes with Diglyme and Tetrahydrofuran. *Acta Chim. Slov.* **2009**, *56*, 426–433.
- (394) Deacon, G. B.; Forsyth, C. M.; Scott, N. M. Solvent-Free Lanthanoid Complexes Derived From Chelation-Supported Organamide Ligands. *Eur. J. Inorg. Chem.* **2002**, *2002*, 1425–1438.
- (395) Johnson, K. R. D.; Côté, A. P.; Hayes, P. G. Four-Coordinate Erbium Organometallic and Coordination Complexes: Synthesis and Structure. *J. Organomet. Chem.* **2010**, *695*, 2747–2755.
- (396) Deacon, G. B.; Evans, D. J.; Junk, P. C. New Variations on the $\text{LnCl}_3(\text{L})_n$ (L = Tetrahydrofuran or 1, 2-Dimethoxyethane) Structural

Theme — NdCl₃(Dme)₂ and YbCl₃(Thf)_{3.5}. *Z. Anorg. Allg. Chem.* **2002**, *628*, 2033–2036.

(397) Emge, T. J.; Kornienko, A.; Brennan, J. G. Trans Influence in a Mer-Octahedral Triiodidolanthanide: Triiodidotris(Tetrahydrofuran-κO)Ytterbium(III). *Acta Crystallogr. Sect. C Cryst. Struct. Commun.* **2009**, *65*, 422–425.

(398) Niemeyer, M. Trans -Diiodopentakis(Tetrahydrofuran)-Ytterbium(III) Tetraiodo- Trans -Bis(Tetrahydrofuran)Ytterbium(III). *Acta Crystallogr. Sect. E Struct. Reports Online* **2001**, *57*, m363–m364.

(399) Caballo, J.; García-Castro, M.; Martín, A.; Mena, M.; Pérez-Redondo, A.; Yélamos, C. Molecular Nitrides with Titanium and Rare-Earth Metals. *Inorg. Chem.* **2011**, *50*, 6798–6808.

(400) Natrajan, L.; Pécaut, J.; Mazzanti, M.; LeBrun, C. Controlled Hydrolysis of Lanthanide Complexes of the N-Donor Tripod Tris(2-Pyridylmethyl)Amine versus Bisligand Complex Formation. *Inorg. Chem.* **2005**, *44*, 4756–4765.

(401) Birmingham, J. M.; Wilkinson, G. The Cyclopentadienides of Scandium, Yttrium and Some Rare Earth Elements. *J. Am. Chem. Soc.* **1956**, *78*, 42–44.

(402) MacDonald, M. R.; Fieser, M. E.; Bates, J. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Identification of the + 2 Oxidation State for Uranium in a Crystalline Molecular Complex, [K(2.2.2-Cryptand)]-[(C₅H₄SiMe₃)₃U]. *J. Am. Chem. Soc.* **2013**, *135*, 13310–13313.

(403) Stults, S. D.; Andersen, R. A.; Zalkin, A. Structural Studies on Cyclopentadienyl Compounds of Trivalent Cerium: Tetrameric (MeC₅H₄)₃Ce and Monomeric (Me₃SiC₅H₄)₃Ce and [(Me₃Si)₂C₅H₃]₃Ce and Their Coordination Chemistry. *Organometallics* **1990**, *9*, 115–122.

(404) Krinsky, J. L.; Minasian, S. G.; Arnold, J. Covalent Lanthanide Chemistry near the Limit of Weak Bonding: Observation of (CpSiMe₃)₃Ce-ECp* and a Comprehensive Density Functional Theory Analysis of Cp₃Ln-ECp (E = Al, Ga). *Inorg. Chem.* **2011**, *50*, 345–357.

(405) Peterson, J. K.; MacDonald, M. R.; Ziller, J. W.; Evans, W. J. Synthetic Aspects of (C₅H₄SiMe₃)₃Ln Rare-Earth Chemistry: Formation of (C₅H₄SiMe₃)₃Lu via [(C₅H₄SiMe₃)₂Ln]⁺ Metallocene Precursors. *Organometallics* **2013**, *32*, 2625–2631.

(406) MacDonald, M. R.; Bates, J. E.; Fieser, M. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Expanding Rare-Earth Oxidation State Chemistry to Molecular Complexes of Holmium(II) and Erbium(II). *J. Am. Chem. Soc.* **2012**, *134*, 8420–8423.

(407) MacDonald, M. R.; Ziller, J. W.; Evans, W. J. Synthesis of a Crystalline Molecular Complex of Y²⁺, [(18-Crown-6)K]-[(C₅H₄SiMe₃)₃Y]. *J. Am. Chem. Soc.* **2011**, *133*, 15914–15917.

(408) Hammel, A.; Schwarz, W.; Weidlein, J. Koordinationsverhältnisse in Cyclopentadienylverbindungen. I. Die Struktur Einer Monomeren Ln(Cp^R)₃-Verbindung Tris(Methylcyclopentadienyl)-Ytterbium. *J. Organomet. Chem.* **1989**, *363*, C29–C35.

(409) Xie, Z.; Hahn, F. E.; Qian, C. Synthesis and Molecular Structure of (MeCp)₃La (MeCp = CH₃C₅H₄): A Tetrameric Complex of the Type [(MeCp)₃La]₄. *J. Organomet. Chem.* **1991**, *414*, C12–C14.

(410) Burns, J. H.; Baldwin, W. H.; Fink, F. H. Crystal Structure of Neodymium Tris(Methylcyclopentadienide). *Inorg. Chem.* **1974**, *13*, 1916–1920.

(411) Angadol, M. A.; Woen, D. H.; Windorff, C. J.; Ziller, J. W.; Evans, W. J. Tert-Butyl(Cyclopentadienyl) Ligands Will Stabilize Nontraditional + 2 Rare-Earth Metal Ions. *Organometallics* **2019**, *38*, 1151–1158.

(412) Rodrigues, I.; Xue, T. Y.; Roussel, P.; Visseaux, M. (t-BuC₅H₄)₃Nd: A Triscyclopentadienyl Rare Earth Compound as Non-Classical Isoprene Polymerization Pre-Catalyst. *J. Organomet. Chem.* **2013**, *743*, 139–146.

(413) Sofield, C. D.; Andersen, R. A. A General Synthesis and Crystal Structure of [(Me₃C)₂C₅H₃]₃Ce. *J. Organomet. Chem.* **1995**, *501*, 271–276.

(414) Goodwin, C. A. P.; Ortu, F.; Reta, D.; Chilton, N. F.; Mills, D. P. Molecular Magnetic Hysteresis at 60 K in Dysprosocenium. *Nature* **2017**, *548*, 439–442.

(415) Goodwin, C. A. P.; Reta, D.; Ortu, F.; Chilton, N. F.; Mills, D. P. Synthesis and Electronic Structures of Heavy Lanthanide Metallophenium Cations. *J. Am. Chem. Soc.* **2017**, *139*, 18714–18724.

(416) Liu, J.; Reta, D.; Cleghorn, J. A.; Yeoh, Y. X.; Ortu, F.; Goodwin, C. A. P.; Chilton, N. F.; Mills, D. P. Light Lanthanide Metallocenium Cations Exhibiting Weak Equatorial Anion Interactions. *Chem. - A Eur. J.* **2019**, *25*, 7749–7758.

(417) Randall McClain, K.; Gould, C. A.; Chakarawet, K.; Teat, S. J.; Groshens, T. J.; Long, J. R.; Harvey, B. G. High-Temperature Magnetic Blocking and Magneto-Structural Correlations in a Series of Dysprosium(III) Metallocenium Single-Molecule Magnets. *Chem. Sci.* **2018**, *9*, 8492–8503.

(418) Jaroschik, F.; Nief, F.; Le Goff, X. F.; Ricard, L. Synthesis and Reactivity of Organometallic Complexes of Divalent Thulium with Cyclopentadienyl and Phospholyl Ligands. *Organometallics* **2007**, *26*, 3552–3558.

(419) Khoroshenkov, G. V.; Petrovskaya, T. V.; Fedushkin, I. L.; Bochkarev, M. N. On the Reactivity of Lanthanide Iodides LnI_x (x < 3) Formed in the Reactions of Lanthanide Metals with Iodine. *Z. Anorg. Allg. Chem.* **2002**, *628*, 699–702.

(420) Du, S.; Yin, J.; Chi, Y.; Xu, L.; Zhang, W. X. Dual Functionalization of White Phosphorus: Formation, Characterization, and Reactivity of Rare-Earth-Metal Cyclo-P₃ Complexes. *Angew. Chem., Int. Ed.* **2017**, *56*, 15886–15890.

(421) Lappert, M. F.; Singh, A.; Atwood, J. L.; Hunter, W. E. Use of the bis(trimethylsilyl)cyclopentadienyl ligand for stabilising early (f⁰–f³) lanthanocene chlorides; X-ray structure of [(Pr{η-[C₅H₃(SiMe₃)₂]}₂Cl)₂] and of isoleptic scandium and ytterbium complexes. *J. Chem. Soc. Chem. Commun.* **1981**, 1190–1191.

(422) Xie, Z.; Chui, K.; Liu, Z.; Xue, F.; Zhang, Z.; Mak, T. C. W.; Sun, J. Systematic Studies on the Reactions of Lanthanide Trichlorides with Na[1,3-bis(trimethylsilyl)cyclopentadienyl]. Crystal structures of [1,3-(Me₃Si)₂C₅H₃]₃Ln (Ln = La, Nd, Gd, Dy). *J. Organomet. Chem.* **1997**, *549*, 239–244.

(423) Thompson, M. E.; Baxter, S. M.; Bulls, A. R.; Burger, B. J.; Nolan, M. C.; Santarsiero, B. D.; Schaefer, W. P.; Bercaw, J. E. "σ Bond Metathesis" for C-H Bonds of Hydrocarbons and Sc-R (R = H, Alkyl, Aryl) Bonds of Permethylscandocene Derivatives. Evidence for Noninvolvement of the π System in Electrophilic Activation of Aromatic and Vinylic C-H Bonds. *J. Am. Chem. Soc.* **1987**, *109*, 203–219.

(424) Gong, L.; Streitwieser, A.; Zalkin, A. Structure of (η⁵-C₅Me₅)₂LuCl(C₄H₈O) and Exchange of Co-Ordinated Solvent. *J. Chem. Soc., Chem. Commun.* **1987**, 460–461.

(425) Evans, W. J.; Peterson, T. T.; Rausch, M. D.; Hunter, W. E.; Zhang, H.; Atwood, J. L. Synthesis and X-Ray Crystallographic Characterization of an Asymmetric Organoyttrium Halide Dimer: (C₅Me₅)₂Y(μ-Cl)YCl(C₅Me₅)₂. *Organometallics* **1985**, *4*, 554–559.

(426) Rausch, M. D.; Moriarty, K. J.; Atwood, J. L.; Weeks, J. A.; Hunter, W. E.; Brittain, H. G. Synthetic, X-Ray Structural, and Photoluminescence Studies on Pentamethylcyclopentadienyl Derivatives of Lanthanum, Cerium, and Praseodymium. *Organometallics* **1986**, *5*, 1281–1283.

(427) Evans, W. J.; Olofson, J. M.; Zhang, H.; Atwood, J. L. Synthesis and X-Ray Crystal Structure of an Unusual Oligomeric Bis-(Pentamethylcyclopentadienyl) Halide Complex of Cerium: [(C₅Me₅)₂CeCl₂K(THF)]_n. *Organometallics* **1988**, *7*, 629–633.

(428) Schumann, H.; Albrecht, I.; Loebel, J.; Hahn, E.; et al. Bis (Pentamethylcyclopentadienyl) Halide and Alkyl Derivatives of the Lanthanides. *Organometallics* **1986**, *5*, 1296–1304.

(429) Lin, H.-H.; Zeng, D.-S.; Shen, Q. CSD Entry FONGEU, Deposition Number 1159090.

(430) Junk, P. C.; Smith, M. K. Crystallographic Report: Bis(μ₂-Chloro)-{bis(Diethylether)-Lithium}-Bis(η⁵-Pentamethyl-Cyclopentadienyl)Samarium(III). *Appl. Organomet. Chem.* **2004**, *18*, 252–252.

(431) Meng, Y. S.; Zhang, Y. Q.; Wang, Z. M.; Wang, B. W.; Gao, S. Weak Ligand-Field Effect from Ancillary Ligands on Enhancing Single-Ion Magnet Performance. *Chem. - A Eur. J.* **2016**, *22*, 12724–12731.

- (432) Evans, W. J.; Grate, J. W.; Levan, K. R.; Bloom, I.; Peterson, T. T.; Doedens, R. J.; Zhang, H.; Atwood, J. L. Synthesis and X-Ray Crystal Structure of Bis (Pentamethylcyclopentadienyl) Lanthanide and Yttrium Halide Complexes. *Inorg. Chem.* **1986**, *25*, 3614–3619.
- (433) Watson, P. L.; Whitney, J. F.; Harlow, R. L. (Pentamethylcyclopentadienyl)Ytterbium and -Lutetium Complexes by Metal Oxidation and Metathesis. *Inorg. Chem.* **1981**, *20*, 3271–3278.
- (434) Beletskaya, I. P.; Voskoboinikov, A. Z.; Chuklanova, E. B.; Kirillova, N. I.; Shestakova, A. K.; Parshina, I. N.; Gusev, A. I.; Magomedov, G. K. I. Bimetallic Lanthanide Complexes with Lanthanide-Transition Metal Bonds. Molecular Structure of $(C_4H_8O)-(C_5H_5)_2Lu-Ru(CO)_2(C_5H_5)$. The Use Of ^{139}La NMR Spectroscopy. *J. Am. Chem. Soc.* **1993**, *115*, 3156–3166.
- (435) Ortu, F.; Packer, D.; Liu, J.; Burton, M.; Formanuk, A.; Mills, D. P. Synthesis and Structural Characterization of Lanthanum and Cerium Substituted Cyclopentadienyl Borohydride Complexes. *J. Organomet. Chem.* **2018**, *857*, 45–51.
- (436) Goodwin, C. A. P.; Reta, D.; Ortu, F.; Liu, J.; Chilton, N. F.; Mills, D. P. Terbenzenium: Completing a Heavy Lanthanide Metalloccenium Cation Family with an Alternative Anion Abstraction Strategy. *Chem. Commun.* **2018**, *54*, 9182–9185.
- (437) Liu, J.; Nodaraki, L. E.; Cobb, P. J.; Giansiracusa, M. J.; Ortu, F.; Tuna, F.; Mills, D. P. Synthesis and Characterisation of Light Lanthanide Bis-Phospholyl Borohydride Complexes. *Dalton Trans.* **2020**, *49*, 6504–6511.
- (438) Guo, F. S.; Day, B. M.; Chen, Y. C.; Tong, M. L.; Mansikkamäki, A.; Layfield, R. A. Magnetic Hysteresis up to 80 K in a Dysprosium Metallocene Single-Molecule Magnet. *Science* **2018**, *362*, 1400–1403.
- (439) White III, J. P.; Deng, H.; Shore, S. G. Borohydride Complexes of Europium(II) and Ytterbium(II) and Their Conversion to Metal Borides. Structure of $(L)_4Yb[BH_4]_2$ ($L = CH_3CN, C_5H_5N$). *Inorg. Chem.* **1991**, *30*, 2337–2342.
- (440) Makhaev, V. D.; Borisov, A. P. Tetrahydroborates of Divalent Samarium, Europium, and Ytterbium. *Russ. J. Inorg. Chem.* **1999**, *44*, 1411–1413.
- (441) Marks, S.; Heck, J. G.; Habicht, M. H.; Oña-Burgos, P.; Feldmann, C.; Roesky, P. W. $[Ln(BH_4)_2(THF)_2]$ ($Ln = Eu, Yb$)-A Highly Luminescent Material. Synthesis, Properties, Reactivity, and NMR Studies. *J. Am. Chem. Soc.* **2012**, *134*, 16983–16986.
- (442) Momin, A.; Bonnet, F.; Visseaux, M.; Maron, L.; Takats, J.; Ferguson, M. J.; Le Goff, X. F.; Nief, F. Synthesis and Structure of Divalent Thulium Borohydrides, and Their Application in ϵ -Caprolactone Polymerisation. *Chem. Commun.* **2011**, *47*, 12203–12205.
- (443) Schmid, M.; Guillaume, S. M.; Roesky, P. W. β -Diketiminato Rare Earth Borohydride Complexes: Synthesis, Structure, and Catalytic Activity in the Ring-Opening Polymerization of ϵ -Caprolactone and Trimethylene Carbonate. *Organometallics* **2014**, *33*, 5392–5401.
- (444) Schmid, M.; Oña-Burgos, P.; Guillaume, S. M.; Roesky, P. W. (Iminophosphoranyl)(Thiophosphoranyl)Methane Rare-Earth Borohydride Complexes: Synthesis, Structures and Polymerization Catalysis. *Dalton Trans.* **2015**, *44*, 12338–12348.
- (445) Schmid, M.; Guillaume, S. M.; Roesky, P. W. 2,5-Bis{N-(2,6-Diisopropylphenyl)iminomethyl}pyrrolyl Borohydride Complexes of the Divalent Lanthanides - Synthesis, Structures and Ring-Opening Polymerization of ϵ -Caprolactone. *J. Organomet. Chem.* **2013**, *744*, 68–73.
- (446) Jaroschik, F.; Bonnet, F.; Le Goff, X. F.; Ricard, L.; Nief, F.; Visseaux, M. Synthesis of Samarium(II) Borohydrides and Their Behaviour as Initiators in Styrene and ϵ -Caprolactone Polymerisation. *Dalton Trans.* **2010**, *39*, 6761–6766.
- (447) Zange, E. Entwicklung Eines Mikroverfahrens Zur Darstellung von Boranaten Der Schweren Lanthaniden. *Chem. Ber.* **1960**, *93*, 652–657.
- (448) Bernstein, E. R.; Chen, K. M. Spectroscopic Properties of Rare Earth Borohydrides: $Er(BH_4)_3 \cdot 3THF$ in Pure and Mixed Crystals. *Chem. Phys.* **1975**, *10*, 215–228.
- (449) Brukl, A.; Rossmannith, K. Über Die Umsetzung von Chloriden Der Seltenen Erden Mit Lithiumborhydrid. *Monat. Chem.* **1959**, *90*, 481–487.
- (450) Rossmannith, K.; Muckenhuber, E. Über Die Umsetzung von Chloriden Der Seltenen Erden Mit Lithiumborhydrid. *Monat. Chem.* **1961**, *92*, 600–604.
- (451) Andersen, R. A. Tris((Hexamethyldisilyl)Amido)Uranium-(III): Preparation and Coordination Chemistry. *Inorg. Chem.* **1979**, *18*, 1507–1509.
- (452) Cendrowski-Guillaume, S. M.; Le Gland, G.; Nierlich, M.; Ephritikhine, M. Lanthanide Borohydrides as Precursors to Organometallic Compounds. Mono(Cyclooctatetraenyl) Neodymium Complexes. *Organometallics* **2000**, *19*, S654–S660.
- (453) Yuan, F.; Li, T.; Li, L.; Zhou, Y. Polymerization of ϵ -Caprolactone by Lanthanide Trisborohydrides and Crystal Structure of $[Ce(BH_4)_2(THF)_5]$ $[Ce(BH_4)_4(THF)_2]$. *J. Rare Earths* **2012**, *30*, 753–756.
- (454) Guillaume, S. M.; Schappacher, M.; Soum, A. Polymerization of ϵ -Caprolactone Initiated by $Nd(BH_4)_3(THF)_3$: Synthesis of Hydroxytelechelic Poly(ϵ -Caprolactone). *Macromolecules* **2003**, *36*, 54–60.
- (455) Bonnet, F.; Visseaux, M.; Barbier-Baudry, D.; Hafid, A.; Vigier, E.; Kubicki, M. M. Organometallic Early Lanthanide Clusters: Syntheses and x-Ray Structures of New Monocyclopentadienyl Complexes. *Inorg. Chem.* **2004**, *43*, 3682–3690.
- (456) Visseaux, M.; Mainil, M.; Terrier, M.; Mortreux, A.; Roussel, P.; Mathivet, T.; Destarac, M. Cationic Borohydrido-Neodymium Complex: Synthesis, Characterization and Its Application as an Efficient Pre-Catalyst for Isoprene Polymerisation. *Dalton Trans.* **2008**, 4558–4561.
- (457) Morris, J. H.; Smith, W. E. Synthesis and Characterisation of a Tetrahydrofuran Derivative of Scandium Tetrahydroborate. *J. Chem. Soc., Chem. Commun.* **1970**, 245.
- (458) Olsen, J. E.; Frommen, C.; Sørby, M. H.; Hauback, B. C. Crystal Structures and Properties of Solvent-Free $LiYb(BH_4)_{4-x}Cl_x$, $Yb(BH_4)_3$ and $Yb(BH_4)_{2-x}Cl_x$. *RSC Adv.* **2013**, *3*, 10764–10774.
- (459) Olsen, J. E.; Frommen, C.; Jensen, T. R.; Riktor, M. D.; Sørby, M. H.; Hauback, B. C. Structure and Thermal Properties of Composites with RE-Borohydrides ($RE = La, Ce, Pr, Nd, Sm, Eu, Gd, Tb, Er, Yb$ or Lu) and $LiBH_4$. *RSC Adv.* **2014**, *4*, 1570–1582.
- (460) Jepsen, L. H.; Ley, M. B.; Černý, R.; Lee, Y. S.; Cho, Y. W.; Ravnsbæk, D.; Besenbacher, F.; Skibsted, J.; Jensen, T. R. Trends in Syntheses, Structures, and Properties for Three Series of Ammine Rare-Earth Metal Borohydrides, $M(BH_4)_3 \cdot nNH_3$ ($M = Y, Gd, \text{ and } Dy$). *Inorg. Chem.* **2015**, *54*, 7402–7414.
- (461) Grinderslev, J. B.; Möller, K. T.; Bremholm, M.; Jensen, T. R. Trends in Synthesis, Crystal Structure, and Thermal and Magnetic Properties of Rare-Earth Metal Borohydrides. *Inorg. Chem.* **2019**, *58*, 5503–5517.
- (462) Grinderslev, J. B.; Jensen, T. R. Trends in the Series of Ammine Rare-Earth-Metal Borohydrides: Relating Structural and Thermal Properties. *Inorg. Chem.* **2021**, *60*, 2573–2589.
- (463) Schumann, H.; Keitsch, M. R.; Demtschuk, J.; Mühle, S. Synthesis of Monomeric Bis(Cyclopentadienyl)Lanthanide Tetrahydroborates with Bulky Cyclopentadienyl Ligands. Single Crystal X-Ray Structures of $[(\eta^5-C_5Me_5)_2SmBH_4(THF)]$ and $[(\eta^5-C_5Me_4Et)_2Y(\mu-H)_2BH_2(THF)]$. *Z. Anorg. Allg. Chem.* **1998**, *624*, 1811–1818.
- (464) Visseaux, M.; Chenal, T.; Roussel, P.; Mortreux, A. Synthesis and X-Ray Structure of a Borohydrido Metallocene of Neodymium and Its Use as Pre-Catalyst in Nd/Mg Dual-Component Ethylene and Isoprene Polymerisations. *J. Organomet. Chem.* **2006**, *691*, 86–92.
- (465) Bonnet, F.; Da Costa Violante, C.; Roussel, P.; Mortreux, A.; Visseaux, M. Unprecedented Dual Behaviour of a Half-Sandwich Scandium-Based Initiator for Both Highly Selective Isoprene and Styrene Polymerisation. *Chem. Commun.* **2009**, 3380–3382.
- (466) Georges, S.; Bonnet, F.; Roussel, P.; Visseaux, M.; Zinck, P. β -Diketiminato-Supported Magnesium Alkyl: Synthesis, Structure and Application as Co-Catalyst for Polymerizations Mediated by a Lanthanum Half-Sandwich Complex. *Appl. Organomet. Chem.* **2016**, *30*, 26–31.

- (467) Bonnet, F.; Jones, C. E.; Semlali, S.; Bria, M.; Roussel, P.; Visseaux, M.; Arnold, P. L. Tuning the Catalytic Properties of Rare Earth Borohydrides for the Polymerisation of Isoprene. *Dalton Trans.* **2013**, 42, 790–801.
- (468) Barbier-Baudry, D.; Blacque, O.; Hafid, A.; Nyassi, A.; Sitzmann, H.; Visseaux, M. Synthesis and X-Ray Crystal Structures of $(C_5H^iPr_4)Ln(BH_4)_2(THF)$ ($Ln = Nd$ and Sm), Versatile Precursors for Polymerization Catalysts. *Eur. J. Inorg. Chem.* **2000**, 2000, 2333–2336.
- (469) Fadlallah, S.; Terrier, M.; Jones, C.; Roussel, P.; Bonnet, F.; Visseaux, M. Mixed Alkyl-Borohydride Lanthanide Complexes: Synthesis of $Ln(BH_4)_2(C_3H_5)(THF)_3$ ($Ln = Nd, Sm$), Characterization, and Reactivity toward Polymerization. *Organometallics* **2016**, 35, 456–461.
- (470) Arliguie, T.; Lance, M.; Nierlich, M.; Vigner, J.; Ephritikhine, M. Inverse Cycloheptatrienyl Sandwich Complexes. Crystal structure of $[U(BH_4)_2(OC_4H_8)_5][U(\mu-\eta^7, \eta^7-C_7H_7)U(BH_4)_3]$. *J. Chem. Soc., Chem. Commun.* **1994**, 847–848.
- (471) Cendrowski-Guillaume, S. M.; Nierlich, M.; Lance, M.; Ephritikhine, M. First Chemical Transformations of Lanthanide Borohydride Compounds: Synthesis and Crystal Structures of $[(\eta-C_8H_8)Nd(BH_4)(THF)]_2$ and $[(\eta-C_8H_8)Nd(THF)_4][BPh_4]$. *Organometallics* **1998**, 17, 786–788.
- (472) Robert, D.; Kondracka, M.; Okuda, J. Cationic Rare-Earth Metal Bis(Tetrahydridoborato) Complexes: Direct Synthesis, Structure and Ring-Opening Polymerisation Activity toward Cyclic Esters. *Dalton Trans.* **2008**, 2667–2669.
- (473) Arliguie, T.; Belkhir, L.; Bouaoud, S. E.; Thuéry, P.; Villiers, C.; Boucekine, A.; Ephritikhine, M. Lanthanide(III) and Actinide(III) Complexes $[M(BH_4)_2(THF)_5][BPh_4]$ and $[M(BH_4)_2(18-crown-6)][BPh_4]$ ($M = Nd, Ce, U$): Synthesis, Crystal Structure, and Density Functional Theory Investigation of the Covalent Contribution to Metal-Borohydride Bonding. *Inorg. Chem.* **2009**, 48, 221–230.
- (474) Long, J.; Selikhov, A. N.; Rad'kova, N. Y.; Cherkasov, A. V.; Guari, Y.; Larionova, J.; Trifonov, A. A. Synthesis, Structures and Magnetic Properties of Two Heteroleptic Dy^{3+} Borohydride Complexes. *Eur. J. Inorg. Chem.* **2021**, 2021, 3008–3012.
- (475) Yu, K. X.; Ding, Y. S.; Zhai, Y. Q.; Han, T.; Zheng, Y. Z. Equatorial Coordination Optimization for Enhanced Axiality of Mononuclear $Dy(III)$ Single-Molecule Magnets. *Dalton Trans.* **2020**, 49, 3222–3227.
- (476) He, M.; Guo, F. S.; Tang, J.; Mansikkamäki, A.; Layfield, R. A. Fulvalene as a Platform for the Synthesis of a Dimetallic Dysprosocenium Single-Molecule Magnet. *Chem. Sci.* **2020**, 11, 5745–5752.
- (477) He, M.; Guo, F. S.; Tang, J.; Mansikkamäki, A.; Layfield, R. A. Synthesis and Single-Molecule Magnet Properties of a Trimetallic Dysprosium Metallocene Cation. *Chem. Commun.* **2021**, 57, 6396–6399.
- (478) Nakamori, Y.; Li, H. W.; Kikuchi, K.; Aoki, M.; Miwa, K.; Towata, S.; Orimo, S. Thermodynamical Stabilities of Metal-Borohydrides. *J. Alloys Compd.* **2007**, 446–447, 296–300.
- (479) Makhav, V.; Borisov, A. P. Synthesis of Bis(Cyclopentadienyl)-Tetrahydroborato Complexes of Scandium by Reacting Scandium Tetrahydroborate with Sodium Cyclopentadienes. *Russ. J. Coord. Chem.* **1992**, 403–405.
- (480) Lobkovskii, E. B.; Kravchenko, S. E.; Semenenko, K. N. X-Ray Structure Study of Crystals of the Tetrahydrofuranate of Scandium Borohydride. *J. Struct. Chem.* **1977**, 18, 312–314.
- (481) Remhof, A.; Borgschulte, A.; Friedrichs, O.; Mauron, P.; Yan, Y.; Züttel, A. Solvent-Free Synthesis and Decomposition of $Y(BH_4)_3$. *Scr. Mater.* **2012**, 66, 280–283.
- (482) Sato, T.; Miwa, K.; Nakamori, Y.; Ohoyama, K.; Li, H. W.; Noritake, T.; Aoki, M.; Towata, S. I.; Orimo, S. I. Experimental and Computational Studies on Solvent-Free Rare-Earth Metal Borohydrides $R(BH_4)_3$ ($R = Y, Dy, \text{ and } Gd$). *Phys. Rev. B - Condens. Matter Mater. Phys.* **2008**, 77, 104114.
- (483) Segal, B. G.; Lippard, S. J. Transition Metal Hydroborate Complexes. 10. Crystal and Molecular Structure of Tris(Tetrahydroborato)Tris(Tetrahydrofuran)Yttrium(III). *Inorg. Chem.* **1978**, 17, 844–850.
- (484) Lobkovskii, E. B.; Kravchenko, S. E.; Kravchenko, O. V. Crystal and Molecular Structure of a 1:2 Complex of Tris(Tetrahydroborato)-Yttrium with Tetrahydrofuran. *J. Struct. Chem.* **1983**, 23, 582–586.
- (485) Volkov, V. V.; Khikmatov, M.; Mirsaidov, U.; Gabuda, S. P.; Kozlova, S. G. Internal Rotation and Phase Transitions in Lanthanum Tetrahydroborates. *J. Struct. Chem.* **1988**, 29, 58–61.
- (486) Li, J.; Ni, X.; Ling, J.; Shen, Z. Syntheses and Properties of Poly(Diethyl Vinylphosphonate) Initiated by Lanthanide Tris-(Borohydride) Complexes: Polymerization Controllability and Mechanism. *J. Polym. Sci. Part A Polym. Chem.* **2013**, 51, 2409–2415.
- (487) Gafurov, B. A.; Mirsaidov, I. U.; Nasrulloeva, D. K.; Badalov, A. Simulating the Synthesis and Thermodynamic Characteristics of the Desolvation of Lanthanide Borohydride Tris-Tetrahydrofuranates. *Russ. J. Phys. Chem. A* **2013**, 87, 1601–1606.
- (488) Grinderslev, J. B.; Jepsen, L. H.; Lee, Y. S.; Møller, K. T.; Cho, Y. W.; Černý, R.; Jensen, T. R. Structural Diversity and Trends in Properties of an Array of Hydrogen-Rich Ammonium Metal Borohydrides. *Inorg. Chem.* **2020**, 59, 12733–12747.
- (489) Payandeh Gharibdoust, S.; Ravnsbæk, D. B.; Černý, R.; Jensen, T. R. Synthesis, Structure and Properties of Bimetallic Sodium Rare-Earth (RE) Borohydrides, $NaRE(BH_4)_4$, $RE = Ce, Pr, Er$ or Gd . *Dalton Trans.* **2017**, 46, 13421–13431.
- (490) Heere, M.; Payandeh GharibDoust, S. H.; Frommen, C.; Humphries, T. D.; Ley, M. B.; Sørby, M. H.; Jensen, T. R.; Hauback, B. C. The Influence of LiH on the Rehydrogenation Behavior of Halide Free Rare Earth (RE) Borohydrides ($RE = Pr, Er$). *Phys. Chem. Chem. Phys.* **2016**, 18, 24387–24395.
- (491) Li, W.; Xue, M.; Zhang, Y.; Yao, Y.; Shen, Q. Direct Synthesis of Ion-Pair Lanthanide Borohydrides and Their High Activity for Polymerization of L-Lactide and ϵ -Caprolactone. *Z. Anorg. Allg. Chem.* **2014**, 640, 1455–1461.
- (492) Møller, K. T.; Jørgensen, M.; Fogh, A. S.; Jensen, T. R. Perovskite Alkali Metal Samarium Borohydrides: Crystal Structures and Thermal Decomposition. *Dalton Trans.* **2017**, 46, 11905–11912.
- (493) Humphries, T. D.; Ley, M. B.; Frommen, C.; Munroe, K. T.; Jensen, T. R.; Hauback, B. C. Crystal Structure and in Situ Decomposition of $Eu(BH_4)_2$ and $Sm(BH_4)_2$. *J. Mater. Chem. A* **2015**, 3, 691–698.
- (494) Gennari, F. C. Mechanochemical Synthesis of Erbium Borohydride: Polymorphism, Thermal Decomposition and Hydrogen Storage. *J. Alloys Compd.* **2013**, 581, 192–195.
- (495) Bradley, D. C.; Ghotra, J. S.; Hart, F. A. Three-Co-Ordination in Lanthanide Chemistry: Tris[Bis(Trimethylsilyl)Amido]Lanthanide(III) Compounds. *J. Chem. Soc., Chem. Commun.* **1972**, 349–350.
- (496) Bradley, D. C.; Ghotra, J. S.; Road, M. E.; Hart, A. Low Co-Ordination Numbers in Lanthanide and Actinide Compounds. Part I. The Preparation and Characterization of Tris{bis(Trimethylsilyl)-Amido}lanthanides. *J. Chem. Soc., Dalton Trans.* **1973**, 1021–1023.
- (497) Anwander, R.; Runte, O.; Eppinger, J.; Gerstberger, G.; Herdtweck, E.; Spiegler, M. Synthesis and Structural Characterisation of Rare-Earth Bis(Dimethylsilyl)Amides and Their Surface Organometallic Chemistry on Mesoporous MCM-41. *J. Chem. Soc., Dalton Trans.* **1998**, 847–858.
- (498) Boyle, T. J.; Bunge, S. D.; Clem, P. G.; Richardson, J.; Dawley, J. T.; Ottley, L. A. M.; Rodriguez, M. A.; Tuttle, B. A.; Avilucea, G. R.; Tissot, R. G. Synthesis and Characterization of a Family of Structurally Characterized Dysprosium Alkoxides for Improved Fatigue-Resistance Characteristics of PDyZT Thin Films. *Inorg. Chem.* **2005**, 44, 1588–1600.
- (499) Tilley, T. D.; Andersen, R. A.; Zalkin, A. Tertiary Phosphine Complexes of the F-Block Metals. Crystal Structure of $Yb[N(SiMe_3)_2]_2[Me_2PCH_2CH_2PMe_2]$: Evidence for a Ytterbium- γ -Carbon Interaction. *J. Am. Chem. Soc.* **1982**, 104, 3725–3727.
- (500) Tilley, T. D.; Andersen, R. A.; Zalkin, A. Divalent Lanthanide Chemistry. Preparation and Crystal Structures of Sodium Tris[Bis(Trimethylsilyl)Amido]Europate(II) and Sodium Tris[Bis-

- (Trimethylsilyl)Amido]Ytterbide(II), NaM[N(SiMe₃)₂]₃. *Inorg. Chem.* **1984**, *23*, 2271–2276.
- (501) Bienfait, A. M.; Schädle, C.; Maichle-Mössmer, C.; Törnroos, K. W.; Anwander, R. Europium Bis(Dimethylsilyl)Amides Including Mixed-Valent Eu₃[N(SiHMe₂)₂]₆[μ-N(SiHMe₂)₂]₂. *Dalton Trans.* **2014**, *43*, 17324–17332.
- (502) Tilley, T. D.; Zalkin, A.; Andersen, R. A.; Templeton, D. H. Divalent Lanthanide Chemistry. Preparation of Some Four- and Six-Coordinate Bis[(Trimethylsilyl)Amido] Complexes of Europium(II). Crystal Structure of Bis[Bis(Trimethylsilyl)Amido]Bis(1,2-Dimethoxyethane)Europium(II). *Inorg. Chem.* **1981**, *20*, 551–554.
- (503) Katzenmayer, M. M.; Wolf, B. M.; Mortis, A.; Maichle-Mössmer, C.; Anwander, R. Polymeric Dimethylytterbium and the Terminal Methyl Complex (Tp^t^{Bu},Me)Yb(CH₃)(Thf). *Chem. Commun.* **2021**, *57*, 243–246.
- (504) Van Den Hende, J. R.; Hitchcock, P. B.; Holmes, S. A.; Lappert, M. F. Synthesis and ¹⁷¹Yb-¹H nuclear magnetic resonance spectra of ytterbium(II) aryloxides [Yb(OR')₂(L)_n][(L)_n=(OEt)₂, (thf)₂, (thf)₃, (pyridine)₂ or Me₂PCH₂CH₂PMe₂] and [Yb(μ-OR')(X)]₂ (X = OR' or NR₂) (R' = C₆H₂Bu^t-2,6-Me-4, R = SiMe₃, thf = tetrahydrofuran). *J. Chem. Soc., Dalton Trans.* **1995**, 1435–1440.
- (505) Nagl, I.; Scherer, W.; Tafipolsky, M.; Anwander, R. The First Oligomeric Samarium(II) Silylamide: Coordinative Saturation through Agostic Sm...SiH Interactions. *Eur. J. Inorg. Chem.* **1999**, 1999, 1405–1407.
- (506) Rabe, G. W.; Rheingold, A. L.; Incarvito, C. D. Crystal Structure of Ytterbium-Bis[Bis(m-Dimethylsilylamido)Ytterbium-Dimethylsilylamide(Tetrahydrofuran)]—Tetrahydrofuran (1/1), C₃₂H₁₀₀N₆O₂Si₁₂Yb₃·C₄H₈O. *Z. Krist. NCS* **2000**, *215*, 560–562.
- (507) Richardson, G. M.; Douair, I.; Cameron, S. A.; Bracegirdle, J.; Keyzers, R. A.; Hill, M. S.; Maron, L.; Anker, M. D. Hydroarylation of Olefins Catalysed by a Dimeric Ytterbium(II) Alkyl. *Nat. Commun.* **2021**, *12*, 3147.
- (508) Shi, Y.; Li, J.; Cui, C. Synthesis of divalent ytterbium terphenylamide and catalytic application for regioselective hydro-silylation of alkenes. *Dalton Trans.* **2017**, *46*, 10957–10962.
- (509) van den Hende, J. R.; Hitchcock, P. B.; Lappert, M. F. Three-coordinate neutral ligand-free ytterbium(II) complexes [(YbX(μ-X))₂] (X = OAr 1 or OCBu^t₃ 3) or [(Yb(NR₂)(μ-X))₂] (X = OCBu^t₃ 2 or OAr 4) (Ar = C₆H₂Bu^t-2,6-Me-4, R = SiMe₃); the X-ray structures of 1 and 2. *J. Chem. Soc., Chem. Commun.* **1994**, 1413–1414.
- (510) Zhao, P.; Zhu, Q.; Fettinger, J. C.; Power, P. Characterization of a Monomeric, Homoleptic, Solvent-Free Samarium Bis(Aryloxyde). *Inorg. Chem.* **2018**, *57*, 14044–14046.
- (511) Deacon, G. B.; Forsyth, C. M.; Junk, P. C. η⁶:η⁶ Coordination of Tetraphenylborate to Ytterbium(II): A New Class of Lanthanoid Ansa-Metalloenes. *Eur. J. Inorg. Chem.* **2005**, 2005, 817–821.
- (512) Bienfait, A. M.; Wolf, B. M.; Törnroos, K. W.; Anwander, R. Ln(II)/Pb(II)-Ln(III)/Pb(0) Redox Approach toward Rare-Earth-Metal Half-Sandwich Complexes. *Organometallics* **2015**, *34*, 5734–5744.
- (513) Schuetz, S. A.; Day, V. W.; Sommer, R. D.; Rheingold, A. L.; Belot, J. A. Anhydrous Lanthanide Schiff Base Complexes and Their Preparation Using Lanthanide Triflate Derived Amides. *Inorg. Chem.* **2001**, *40*, 5292–5295.
- (514) Hitchcock, P. B.; Hulkes, A. G.; Lappert, M. F.; Li, Z. Cerium(III) dialkyl dithiocarbamates from [Ce{N(SiMe₃)₂}]₃ and tetraalkylthiuram disulfides, and [Ce(κ²-S₂CNET₂)₄] from the Ce^{III} precursor; Tb^{III} and Nd^{III} analogues. *Dalton Trans.* **2004**, 129–136.
- (515) Roger, M.; Barros, N.; Arliguie, T.; Thuéry, P.; Maron, L.; Ephritikhine, M. U(SMes*)_n (n = 3, 4) and Ln(SMes*)₃ (Ln = La, Ce, Pr, Nd): Lanthanide(III)/Actinide(III) Differentiation in Agostic Interactions and an Unprecedented η³ Ligation Mode of the Arylthiolate Ligand, from X-Ray Diffraction and DFT Analysis. *J. Am. Chem. Soc.* **2006**, *128*, 8790–8802.
- (516) Behrle, A. C.; Schmidt, J. A. R. Synthesis and Reactivity of Homoleptic -Metalated N,N-Dimethylbenzylamine Rare-Earth-Metal Complexes. *Organometallics* **2011**, *30*, 3915–3918.
- (517) Yuen, H. F.; Marks, T. J. Synthesis and Catalytic Properties of Phenylene-Bridged Binuclear Organolanthanide Complexes. *Organometallics* **2008**, *27*, 155–158.
- (518) Occhipinti, G.; Meermann, C.; Dietrich, H. M.; Litlabø, R.; Auras, F.; Törnroos, K. W.; Maichle-Mössmer, C.; Jensen, V. R.; Anwander, R. Synthesis and Stability of Homoleptic Metal(III) Tetramethylaluminates. *J. Am. Chem. Soc.* **2011**, *133*, 6323–6337.
- (519) Skår, H.; Seland, J. G.; Liang, Y.; Frøystein, N. Å.; Törnroos, K. W.; Anwander, R. Screening of the Relaxivity of Gadolinium-Loaded Periodic Mesoporous Silica Functionalized by Means of Soft Metal-organic Silylamide Grafting. *Eur. J. Inorg. Chem.* **2013**, 2013, 5969–5979.
- (520) Dietrich, H. M.; Meermann, C.; Törnroos, K. W.; Anwander, R. Sounding out the Reactivity of Trimethylyttrium. *Organometallics* **2006**, *25*, 4316–4321.
- (521) Hitchcock, P. B.; Lappert, M. F.; Singh, A. Three- and Four-Coordinate, Hydrocarbon-Soluble-Aryloxides of Scandium, Yttrium, and the Lanthanoids; X-Ray Crystal Structure of Tris(2,6-di-*t*-Butyl-4-Methylphenoxy)Scandium. *J. Chem. Soc., Chem. Commun.* **1983**, 1499–1501.
- (522) Barnhart, D. M.; Clark, D. L.; Gordon, J. C.; Huffman, J. C.; Vincent, R. L.; Watkin, J. G.; Zwick, B. D. Synthesis, Properties, and X-Ray Structures of the Lanthanide η⁶-Arene-Bridged Aryloxy Dimers Ln₂(O-2,6-*i*-Pr₂C₆H₃)₆ and Their Lewis Base Adducts Ln(O-2,6-*i*-Pr₂C₆H₃)(THF)₂ (Ln = Pr, Nd, Sm, Gd, Er, Yb, Lu). *Inorg. Chem.* **1994**, *33*, 3487–3497.
- (523) Fischbach, A.; Herdtweck, E.; Anwander, R.; Eickerling, G.; Scherer, W. Reactivity of Trimethylaluminum with Lanthanide Aryloxides: Adduct and Tetramethylaluminate Formation. *Organometallics* **2003**, *22*, 499–509.
- (524) Yin, H.; Carroll, P. J.; Manor, B. C.; Anna, J. M.; Schelter, E. J. Cerium Photosensitizers: Structure-Function Relationships and Applications in Photocatalytic Aryl Coupling Reactions. *J. Am. Chem. Soc.* **2016**, *138*, 5984–5993.
- (525) Cole, B. E.; Falcones, I. B.; Cheisson, T.; Manor, B. C.; Carroll, P. J.; Schelter, E. J. A Molecular Basis to Rare Earth Separations for Recycling: Tuning the TriNO_x Ligand Properties for Improved Performance. *Chem. Commun.* **2018**, *54*, 10276–10279.
- (526) Dong, X.; Robinson, J. R. The Role of Neutral Donor Ligands in the Ioselective Ring-Opening Polymerization of: Rac -β-Butyrolactone. *Chem. Sci.* **2020**, *11*, 8184–8195.
- (527) Evans, W. J.; Ansari, M. A.; Ziller, J. W.; Khan, S. I. Utility of Arylamido Ligands in Yttrium and Lanthanide Chemistry. *Inorg. Chem.* **1996**, *35*, 5435–5444.
- (528) Klimpel, M. G.; Görlitzer, H. W.; Tafipolsky, M.; Spiegler, M.; Scherer, W.; Anwander, R. β-SiH Agostic Bonding in Sterically Crowded Lanthanidocene Silylamide Complexes. *J. Organomet. Chem.* **2002**, *647*, 236–244.
- (529) Anwander, R.; Klimpel, M. G.; Dietrich, H. M.; Shorokhov, D. J.; Scherer, W. High Tetraalkylaluminate Fluxionality in Half-Sandwich Complexes of the Trivalent Rare-Earth Metals. *Chem. Commun.* **2003**, *3*, 1008–1009.
- (530) Booij, M.; Kiers, N. H.; Heeres, H. J.; Teuben, J. H. On the Synthesis of Monopentamethylcyclopentadienyl Derivatives of Yttrium, Lanthanum, and Cerium. *J. Organomet. Chem.* **1989**, *364*, 79–86.
- (531) Vitanova, D. V.; Hampel, F.; Hultsch, K. C. Rare Earth Metal Complexes Based on β-Diketiminato and Novel Linked Bis(β-Diketiminato) Ligands: Synthesis, Structural Characterization and Catalytic Application in Epoxide/CO₂-Copolymerization. *J. Organomet. Chem.* **2005**, *690*, 5182–5197.
- (532) Luo, Y.; Lei, Y.; Fan, S.; Wang, Y.; Chen, J. Synthesis of Mono-Amidinate-Ligated Rare-Earth-Metal Bis(Silylamide) Complexes and Their Reactivity with [Ph₃C][B(C₆F₅)₄], AlMe₃ and Isoprene. *Dalton Trans.* **2013**, *42*, 4040–4051.
- (533) Benndorf, P.; Jenter, J.; Zielke, L.; Roesky, P. W. Chiral Lutetium Benzamidinate Complexes. *Chem. Commun.* **2011**, *47*, 2574–2576.
- (534) Yin, H.; Carroll, P. J.; Anna, J. M.; Schelter, E. J. Luminescent Ce(III) Complexes as Stoichiometric and Catalytic Photoreductants

for Halogen Atom Abstraction Reactions. *J. Am. Chem. Soc.* **2015**, *137*, 9234–9237.

(535) Crozier, A. R.; Bienfait, A. M.; Maichle-Mössmer, C.; Törnroos, K. W.; Anwander, R. A Homoleptic Tetravalent Cerium Silylamide. *Chem. Commun.* **2013**, *49*, 87–89.

(536) Schneider, D.; Spallek, T.; Maichle-Mössmer, C.; Törnroos, K. W.; Anwander, R. Cerium Tetrakis(Diisopropylamide) - a Useful Precursor for Cerium(IV) Chemistry. *Chem. Commun.* **2014**, *50*, 14763–14766.

(537) Kim, J. E.; Bogart, J. A.; Carroll, P. J.; Schelter, E. J. Rare Earth Metal Complexes of Bidentate Nitroxide Ligands: Synthesis and Electrochemistry. *Inorg. Chem.* **2016**, *55*, 775–784.

(538) Levin, J. R.; Dorfner, W. L.; Dai, A. X.; Carroll, P. J.; Schelter, E. J. Density Functional Theory as a Predictive Tool for Cerium Redox Properties in Nonaqueous Solvents. *Inorg. Chem.* **2016**, *55*, 12651–12659.

(539) Woen, D. H.; Chen, G. P.; Ziller, J. W.; Boyle, T. J.; Furche, F.; Evans, W. J. Solution Synthesis, Structure, and CO₂ Reduction Reactivity of a Scandium(II) Complex, {Sc[N(SiMe₃)₂]₃}⁻. *Angew. Chem., Int. Ed.* **2017**, *56*, 2050–2053.

(540) Ghotra, J. S.; Hursthouse, M. B.; Welch, A. J. Three-Coordinate Scandium(III) and Europium(III); Crystal and Molecular Structures of Their Tris-hexamethylsilylamides. *J. Chem. Soc., Chem. Commun.* **1973**, 669–670.

(541) Westerhausen, M.; Hartmann, M.; Pfitzner, A.; Schwarz, W. Bis(Trimethylsilyl)Amide und -Methanide Des Yttriums - Molekülstrukturen von Tris(Diethylether)Lithium-(μ-Chloro)-Tris[Bis(Trimethylsilyl)Methyl]Yttriat, Solvensfreiem Yttrium-Tris[Bis(Trimethylsilyl)amid] Sowie Dem Bis(Benzonitril)-Komplex. *Z. Anorg. Allg. Chem.* **1995**, *621*, 837–850.

(542) Dietrich, H. M.; Meermann, C.; Törnroos, K. W.; Anwander, R. Sounding out the Reactivity of Trimethylyttrium. *Organometallics* **2006**, *25*, 4316–4321.

(543) Herrmann, W. A. Reaktivitätsbestimmender Einfluß Der Ligandenkonstitution Bei Seltenerd-Amiden: Herstellung Und Struktur Sterisch Überladener Alkoxid-Komplexe. *Zeitschrift für Naturforsch. - Sect. B J. Chem. Sci.* **1994**, *49b*, 1789.

(544) Bradley, D. C.; Ghotra, J. S.; Hart, F. A.; Hursthouse, M. B.; Raithby, P. R. Syntheses, Properties, and Crystal and Molecular Structures of Triphenylphosphine Oxide and Peroxo-Derivatives of [Bis(Trimethylsilyl)Amido]Lanthanoids. *J. Chem. Soc., Dalton Trans.* **1977**, 1166–1172.

(545) Rees, W. S.; Just, O.; Van Derveer, D. S. Molecular Design of Dopant Precursors for Atomic Layer Epitaxy of SrS:Ce. *J. Mater. Chem.* **1999**, *9*, 249–252.

(546) Bubrin, D.; Niemeyer, M. Formation of Novel T-Shaped NNN Ligands via Rare-Earth Metal-Mediated Si-H Activation. *Inorg. Chem.* **2014**, *53* (3), 1269–1271.

(547) Andersen, R. A.; Templeton, D. H.; Zalkin, A. Structure of Tris(Bis(Trimethylsilyl)Amido)Neodymium(III), Nd[N(Si(CH₃)₂)₃]₃. *Inorg. Chem.* **1978**, *17*, 2317–2319.

(548) Brady, E. D.; Clark, D. L.; Gordon, J. C.; Hay, P. J.; Keogh, D. W.; Poli, R.; Scott, B. L.; Watkin, J. G. Tris(Bis(Trimethylsilyl)Amido)Samarium: X-Ray Structure and DFT Study. *Inorg. Chem.* **2003**, *42*, 6682–6690.

(549) Sundermeyer, J.; Khvorost, A.; Harms, K. Tris[Bis(Trimethylsilyl)Amido]Samarium Tetrahydrofuran Solvate. *Acta Crystallogr. Sect. E Struct. Reports Online* **2004**, *60*, m1117–m1119.

(550) Rabe, G. W.; Yap, G. P. A. Crystal Structure of Samarium Tris[Bis(Dimethylsilyl)Amide]Bis(Tetrahydrofuran), Sm[N(SiHMe₂)₂]₃(THF)₂. *Z. Krist. NCS* **2000**, *215*, 457–458.

(551) Bienfait, A. M.; Wolf, B. M.; Törnroos, K. W.; Anwander, R. Trivalent Rare-Earth-Metal Bis(Trimethylsilyl)Amide Halide Complexes by Targeted Oxidations. *Inorg. Chem.* **2018**, *57*, 5204–5212.

(552) Herrmann, W. A.; Anwander, R.; Kleine, M.; Scherer, W. Lanthanoiden-Komplexe, I Solvensfreie Alkoxid-Komplexe Des Neodyms Und Dysprosiums. Kristall- Und Molekülstruktur von Trans-Bis(Acetonitril)Tris(Tri-tert-butylmethoxy)Neodym. *Chem. Ber.* **1992**, *125*, 1971–1979.

(553) Rastätter, M.; Zulus, A.; Roesky, P. W. Bis(Phosphinimino)-Methanide Rare Earth Amides: Synthesis, Structure, and Catalysis of Hydroamination/Cyclization, Hydrosilylation, and Sequential Hydroamination/Hydrosilylation. *Chem. - A Eur. J.* **2007**, *13*, 3606–3616.

(554) Hitchcock, P. B.; Khvostov, A. V.; Lappert, M. F.; Protchenko, A. V. Ytterbium(II) Amides and Crown Ethers: Addition versus Amide Substitution. *J. Organomet. Chem.* **2002**, *647*, 198–204.

(555) Niemeyer, M. Reactions of Hypersilyl Potassium with Rare-Earth Metal Bis(Trimethylsilylamides): Addition versus Peripheral Deprotonation. *Inorg. Chem.* **2006**, *45*, 9085–9095.

(556) Sommerfeldt, H. M.; Meermann, C.; Schrems, M. G.; Törnroos, K. W.; Fröystein, N. A.; Miller, R. J.; Scheidt, E. W.; Scherer, W.; Anwander, R. Characterization and Reactivity of Peralkylated Ln^{II}Al^{III} Heterobimetallic Complexes. *Dalton Trans.* **2008**, 1899–1907.

(557) Niemeyer, M. Synthesis and Structural Characterization of Several Ytterbium Bis(Trimethylsilyl)Amides Including Base-Free[Yb-{N(SiMe₃)₂]₂(μ-Cl)₂] — A Coordinatively Unsaturated Complex with Additional Agostic Yb⋯(H3C—Si) Interactions. *Z. Anorg. Allg. Chem.* **2002**, *628*, 647–657.

(558) Scarel, G.; Wiemer, C.; Fanciulli, M.; Fedushkin, I. L.; Fukin, G. K.; Domrachev, G. A.; Lebedinskii, Y.; Zenkevich, A.; Pavia, G. [(Me₃Si)₂N]₃Lu: Molecular Structure and Use as Lu and Si Source for Atomic Layer Deposition of Lu Silicate Films. *Z. Anorg. Allg. Chem.* **2007**, *633*, 2097.

(559) So, Y. M.; Leung, W. H. Recent Advances in the Coordination Chemistry of Cerium(IV) Complexes. *Coord. Chem. Rev.* **2017**, *340*, 172–197.

(560) Kim, J. E.; Bogart, J. A.; Carroll, P. J.; Schelter, E. J. Rare Earth Metal Complexes of Bidentate Nitroxide Ligands: Synthesis and Electrochemistry. *Inorg. Chem.* **2016**, *55*, 775–784.

(561) Gulino, A.; Casarin, M.; Conticello, V. P.; Gaudiello, J. G.; Mauermann, H.; Fragalá, I.; Marks, T. J. Efficient Synthesis, Redox Characteristics, and Electronic Structure of a Tetravalent Tris-(Cyclopentadienyl)Cerium Alkoxide Complex. *Organometallics* **1988**, *7*, 2360–2364.

(562) Evans, W. J.; Deming, T. J.; Ziller, J. W. The Utility of Ceric Ammonium Nitrate Derived Alkoxide Complexes in the Synthesis of Organometallic Cerium(IV) Complexes: Synthesis and First X-Ray Crystallographic Determination of a Tetravalent Cerium Cyclopentadienide Complex, (C₅H₅)₃Ce(OCMe₃). *Organometallics* **1989**, *8*, 1581–1583.

(563) Moehring, S. A.; Miehlich, M.; Hoerger, C. J.; Meyer, K.; Ziller, J. W.; Evans, W. J. A Room-Temperature Stable Y(II) Aryloxide: Using Steric Saturation to Kinetically Stabilize Y(II) Complexes. *Inorg. Chem.* **2020**, *59*, 3207–3214.

(564) Xu, X.; Yao, Y.; Zhang, Y.; Shen, Q. Crystal Structure of Tris(2,6-Di-Tert-Butyl-4-Methylphenolato-O) (Tetrahydrofuran-O) Erbium Toluene Solvate. *Appl. Organomet. Chem.* **2004**, *18*, 382–383.

(565) Butcher, R. J.; Clark, D. L.; Grumbine, S. K.; Vincent-Hollis, R. L.; Scott, B. L.; Watkin, J. G. Ammoniacal Synthesis of Lanthanum Aryloxide Complexes and Observation of a Unique Interconversion between Oxygen- and η-Arene-Bridged Dimeric Species. X-Ray Crystal Structures of La₂(OAr)₆(NH₃)_n (n = 0, 2), La(OAr)₃(NH₃)₄, and La(OAr)₃(THF)₂ (Ar = 2,6-i-Pr₂C₆H₃). *Inorg. Chem.* **1995**, *34*, 5468–5476.

(566) Evans, W. J.; Johnston, M. A.; Greci, M. A.; Ziller, J. W. Facile Formation of Luminescent Terbium(III) Aryloxide Complexes Directly from Terbium Metal Including the X-Ray Crystal Structures of Tb(OC₆H₃Me₂-2,6)₃(THF)₃ and Tb(OC₆H₃i-Pr₂-2,6)₃(THF)₂. *Polyhedron* **2001**, *20*, 277–280.

(567) Xia, Q.; Cui, Y.; Yuan, D.; Wang, Y.; Yao, Y. Synthesis and Characterization of Lanthanide Complexes Stabilized by N-Aryl Substituted β-Ketoimino Ligands and Their Application in the Polymerization of Rac-Lactide. *J. Organomet. Chem.* **2017**, *846*, 161–168.

(568) Hong, Y.; Zheng, Y.; Xue, M.; Yao, Y.; Zhang, Y.; Shen, Q. Phenylene-Bridged β-Ketoimino Dilanthanide Aryloxides: Synthesis, Structure, and Catalytic Activity for Addition of Amines to Carbodiimides. *Z. Anorg. Allg. Chem.* **2015**, *641*, 1230–1237.

- (569) Peng, H.; Zhang, Z.; Qi, R.; Yao, Y.; Zhang, Y.; Shen, Q.; Cheng, Y. Synthesis, Reactivity, and Characterization of Sodium and Rare-Earth Metal Complexes Bearing a Dianionic N-Aryloxo-Functionalized β -Ketoiminate Ligand. *Inorg. Chem.* **2008**, *47*, 9828–9835.
- (570) Gu, W.; Xu, P.; Wang, Y.; Yao, Y.; Yuan, D.; Shen, Q. Synthesis and Characterization of Yttrium and Ytterbium Complexes Supported by Salen Ligands and Their Catalytic Properties for Rac-Lactide Polymerization. *Organometallics* **2015**, *34*, 2907–2916.
- (571) Heeres, H. J.; Meetsma, A.; Teuben, J. H.; Rogers, R. D. Mono(Pentamethylcyclopentadienyl) Complexes of Cerium(III). Synthesis, Molecular Structure Thermal Stability, and Reactivity of (C_5Me_5) CeX_2 ($X = 2,6$ -Di-Tert-Butylphenoxy, $CH(SiMe_3)_2$, and $N(SiMe_3)_2$) Complexes. *Organometallics* **1989**, *8*, 2637–2646.
- (572) Butcher, R. J.; Clark, D. L.; Gordon, J. C.; Watkin, J. G. Synthesis and X-Ray Crystal Structures of the Samarium Mono-(Pentamethylcyclopentadienyl) Aryloxo Complexes (η - C_5Me_5)Sm(O-2,6-*t*-Bu $_2$ C $_6$ H $_3$) $_2$ (THF) and [$(\eta$ - C_5Me_5)Sm(O-2,6-*i*-Pr $_2$ C $_6$ H $_3$) $_3$ Li(THF)]. Differences in metathesis chemistry of 2,6-di-*iso*-propylphenoxy and 2,6-di-*tert*-butylphenoxy ligands. *J. Organomet. Chem.* **1999**, *577*, 228–237.
- (573) Conley, M. P.; Lapadula, G.; Sanders, K.; Gajan, D.; Lesage, A.; Del Rosal, I.; Maron, L.; Lukens, W. W.; Copéret, C.; Andersen, R. A. The Nature of Secondary Interactions at Electrophilic Metal Sites of Molecular and Silica-Supported Organolanthanum Complexes from Solid-State NMR Spectroscopy. *J. Am. Chem. Soc.* **2016**, *138*, 3831–3843.
- (574) Avent, A. G.; Caro, C. F.; Hitchcock, P. B.; Lappert, M. F.; Li, Z.; Wei, X. H. Synthetic and Structural Experiments on Yttrium, Cerium and Magnesium Trimethylsilylmethyls and Their Reaction Products with Nitriles; With a Note on Two Cerium β -Diketiminates. *J. Chem. Soc., Dalton Trans.* **2004**, 1567–1577.
- (575) Sridharan, V.; Menéndez, J. C. Cerium(IV) Ammonium Nitrate as a Catalyst in Organic Synthesis. *Chem. Rev.* **2010**, *110*, 3805–3849.
- (576) Bradley, D. C.; Chatterjee, A. K.; Wardlaw, W. Structural Chemistry of the Alkoxides. Part VI. Primary Alkoxides of Quadrivalent Cerium and Thorium. *J. Chem. Soc.* **1956**, 2260–2264.
- (577) Bradley, D. C.; Chatterjee, A. K.; Wardlaw, W. Structural Chemistry of the Alkoxides. Part VII. Secondary Alkoxides of Quadrivalent Cerium and Thorium. *J. Chem. Soc.* **1956**, 3469–3472.
- (578) Bradley, D. C.; Chatterjee, A. K.; Wardlaw, W. Structural Chemistry of the Alkoxides. Part IX. Tert-Alkoxides of Quadrivalent Cerium. *J. Chem. Soc.* **1957**, 2600–2604.
- (579) Gradeff, P. S.; Schreiber, F. G.; Brooks, K. C.; Sievers, R. E. Simplified Method for the Synthesis of Ceric Alkoxides from Ceric Ammonium Nitrate. *Inorg. Chem.* **1985**, *24*, 1110–1111.
- (580) Gradeff, P. S.; Schreiber, F. G.; Mauermann, H. Preparation of Ceric Alkoxides in Glycol Ethers. *J. Less-Common Met.* **1986**, *126*, 335–338.
- (581) Evans, W. J.; Deming, T. J.; Olofson, J. M.; Ziller, J. W. Synthetic and Structural Studies of a Series of Soluble Cerium(IV) Alkoxide and Alkoxide Nitrate Complexes. *Inorg. Chem.* **1989**, *28*, 4027–4034.
- (582) Friedrich, J.; Schneider, D.; Bock, L.; Maichle-Mössner, C.; Anwender, R. Cerium(IV) Neopentoxide Complexes. *Inorg. Chem.* **2017**, *56*, 8114–8127.
- (583) Williams, U. J.; Schneider, D.; Dorfner, W. L.; Maichle-Mössner, C.; Carroll, P. J.; Anwender, R.; Schelter, E. J. Variation of Electronic Transitions and Reduction Potentials of Cerium(IV) Complexes. *Dalton Trans.* **2014**, *43*, 16197–16206.
- (584) Sutton, A. D.; Clark, D. L.; Scott, B. L.; Gordon, J. C. Synthesis and Characterization of Cerium(IV) Metallocenes. *Inorganics* **2015**, *3*, 589–596.
- (585) Haddow, M. F.; Newland, R. J.; Tegner, B. E.; Mansell, S. M. Reversible Temperature-Induced Polymorphic Phase Transitions of [Y(OAr) $_3$] and [Ce(OAr) $_3$] (Ar = 2,6-*t*-Bu $_2$ -4-MeC $_6$ H $_2$): Interconversions between Pyramidal and Planar Geometries. *CrystEngComm* **2019**, *21*, 2884–2892.
- (586) Skår, H.; Bienfait, A. M.; Schnitzlbaumer, M.; Törnroos, K. W.; Anwender, R. High 1H Relaxivity Gadolinium-Grafted Mesoporous Silica Materials. *Z. Anorg. Allg. Chem.* **2014**, *640*, 604–615.
- (587) Stecher, H. A.; Sen, A.; Rheingold, A. L. Synthesis, Structure, and Reactivity of Tricoordinate Cerium(III) Aryloxides. The First Structurally Characterized Monomeric Ln(OR) $_3$ Complexes. *Inorg. Chem.* **1988**, *27*, 1130–1132.
- (588) Steele, L. A. M.; Boyle, T. J.; Kemp, R. A.; Moore, C. The Selective Insertion of Carbon Dioxide into a Lanthanide(III) 2,6-Di-*t*-Butyl-Phenoxy Bond. *Polyhedron* **2012**, *42*, 258–264.
- (589) Schläfer, J.; Tyrra, W.; Mathur, S. Octakis(Tert-Butoxy)-Dicerium(IV) [Ce $_2$ (O t Bu) $_8$]: Synthesis, Characterization, Decomposition, and Reactivity. *Inorg. Chem.* **2014**, *53*, 2751–2753.
- (590) Schläfer, J.; Stucky, S.; Tyrra, W.; Mathur, S. Heterobi- and Trimetallic Cerium(IV) Tert -Butoxides with Mono-, Di-, and Trivalent Metals (M = K(I), Ge(II), Sn(II), Pb(II), Al(III), Fe(III)). *Inorg. Chem.* **2013**, *52*, 4002–4010.
- (591) Amberger, H. D.; Reddmann, H.; Guttenberger, C.; Unrecht, B.; Zhang, L.; Apostolidis, C.; Walter, O.; Kanellakopoulos, B. Zur Elektronenstruktur Hochsymmetrischer Verbindungen Der F-Elemente. Spektroskopische Und Strukturelle Charakterisierung von Tris(2,6-Di-*t*-Butyl-Phenolato)Lanthanid(III) (Ln(OAr') $_3$; Ln = Pr, Nd) Sowie Parametrische Analyse Des Kristallfeld-Aufspal. *Z. Anorg. Allg. Chem.* **2003**, *629*, 1522–1534.
- (592) Qi, G. Z.; Shen, Q.; Lin, Y. H. A Bis(2,6-Di-Tert-Butyl-4-Methylphenolato)Samarium(II) Complex, [Sm(OAr) $_2$ (Thf) $_3$] \cdot Thf. *Acta Crystallogr. Sect. C Cryst. Struct. Commun.* **1994**, *50*, 1456–1458.
- (593) Evans, W. J.; Anwender, R.; Ansari, M. A.; Ziller, J. W. Samarium(II) Surrounded by Only Oxygen Donor Ligands: [KSm(μ -OC $_6$ H $_3$ Bu $_2$ -2,6-Me-4) $_3$ (THF)] $_n$. *Inorg. Chem.* **1995**, *34*, 5–6.
- (594) Yao, Y. M.; Shen, Q.; Zhang, Y.; Xue, M. Q.; Sun, J. Syntheses and Reactivities of Bisaryloxo Lanthanide Chlorides and Crystal Structures of [Ln(ArO) $_2$ Cl(THF) $_2$] \cdot (MePh) (Ln = Er, Yb) and [Sm(ArO) $_2$ (THF) $_3$] \cdot (MePh) (ArO = 2,6-di-*tert*-butyl-4-methylphenoxy). *Polyhedron* **2001**, *20*, 3201–3208.
- (595) Hou, Z.; Fujita, A.; Yoshimura, T.; Jesorka, A.; Zhang, Y.; Yamazaki, H.; Wakatsuki, Y. Heteroleptic Lanthanide Complexes with Aryloxo Ligands. Synthesis and Structural Characterization of Divalent and Trivalent Samarium Aryloxo/Halide and Aryloxo/Cyclopentadienide Complexes. *Inorg. Chem.* **1996**, *35*, 7190–7195.
- (596) Qi, G.; Lin, Y.; Hu, J.; Shen, Q. I. Synthesis and Crystal Structure of a Four-Coordinate Aryloxo Samarium Complex [Sm(OAr) $_3$ (THF)](THF) (OAr = 2,6-Di-*tert*-butyl-4-Methylphenoxy). *Polyhedron* **1995**, *14*, 413–415.
- (597) Carretas, J.; Branco, J.; Marçalo, J.; Domingos, Â.; Pires de Matos, A. P. Europium(II) and Ytterbium(II) Aryloxo Chemistry: Synthesis and Crystal Structure of [Eu(OC $_6$ H $_3$ Bu $_2$ -2,6) $_2$ (THF) $_3$] \cdot 0.75C $_7$ H $_8$ and [Yb(OC $_6$ H $_3$ Bu $_2$ -2,6) $_2$ (NCMe) $_4$]. *Polyhedron* **2003**, *22* (11), 1425–1429.
- (598) Zhang, H.; Nakanishi, R.; Katoh, K.; Breedlove, B. K.; Kitagawa, Y.; Yamashita, M. Low Coordinated Mononuclear Erbium(III) Single-Molecule Magnets with C $_3v$ symmetry: A Method for Altering Single-Molecule Magnet Properties by Incorporating Hard and Soft Donors. *Dalton Trans.* **2018**, *47*, 302–305.
- (599) Bochkarev, M. N.; Fagin, A. A.; Fedushkin, I. L.; Petrovskaya, T. V.; Evans, W. J.; Greci, M. A.; Ziller, J. W. Alkyl(Aryl)Oxy Derivatives of Thulium(III). *Russ. Chem. Bull.* **1999**, *48*, 1782–1785.
- (600) Deacon, G. B.; Meyer, G.; Stellfeldt, D.; Zelesny, G.; Skelton, B. W.; White, A. H. Syntheses of Bis(Aryloxo)Halogenoytterbium(III) Complexes: [Yb(OAr) $_2$ X(THF) $_2$] (X = Cl, Br, I; OAr = OC $_6$ H $_2$ -2,6-*t*-Bu $_2$ -4-R; R = H, Me, t Bu) and the X-Ray Crystal Structures of [Yb(OC $_6$ H $_2$ -2,6- t Bu $_2$ -4-Me) $_2$ I(THF) $_2$] and [Yb(OC $_6$ H $_3$ -2,6- t Bu $_2$)OH(THF) $_2$]. *Z. Anorg. Allg. Chem.* **2001**, *627*, 1652–1658.
- (601) Fischbach, A.; Herdtweck, E.; Anwender, R.; Eickerling, G.; Scherer, W. Reactivity of Trimethylaluminum with Lanthanide Aryloxides: Adduct and Tetramethylaluminate Formation. *Organometallics* **2003**, *22*, 499–509.
- (602) Cao, Y.; Du, Z.; Li, W.; Li, J.; Zhang, Y.; Xu, F.; Shen, Q. Activation of Carbodiimide and Transformation with Amine to Guanidinate Group by Ln(OAr) $_3$ (THF) $_2$ (Ln: Lanthanide and Yttrium) and Ln(OAr) $_3$ (THF) $_2$ as a Novel Precatalyst for Addition

- of Amines to Carbodiimides: Influence of Aryloxy Group. *Inorg. Chem.* **2011**, *50*, 3729–3737.
- (603) Fukuzumi, S.; Fujii, Y.; Suenobu, T. Metal Ion-Catalyzed Cycloaddition vs Hydride Transfer Reactions of NADH Analogues with p-Benzoquinones. *J. Am. Chem. Soc.* **2001**, *123* (42), 10191–10199.
- (604) Ekkehardt Hahn, F.; Mohr, J. Synthese Und Strukturen von Bis(Trietanolamin)Lanthanoid-Komplexen. *Chem. Ber.* **1990**, *123*, 481–484.
- (605) Massaux, J.; Duyckaerts, G. Methode de Preparation de Sels Anhydres de Lanthanides Pour La Polarographie En Solvants Non Aqueux Aprotiques. *Anal. Chim. Acta* **1974**, *73*, 416–419.
- (606) Caravan, P.; Tóth, E.; Rockenbauer, A.; Merbach, A. E. Nuclear and Electronic Relaxation of Eu^{2+} (aq): An Extremely Labile Aqua Ion. *J. Am. Chem. Soc.* **1999**, *121*, 10403–10409.
- (607) Teprovich, J. A.; Antharjanam, P. K. S.; Prasad, E.; Pesciotta, E. N.; Flowers, R. A. Generation of SmII Reductants Using High Intensity Ultrasound. *Eur. J. Inorg. Chem.* **2008**, *2008*, 5015–5019.
- (608) Maisano, T.; Tempest, K. E.; Sadasivam, D. V.; Flowers, R. A. A Convenient Pathway to Sm(II)-Mediated Chemistry in Acetonitrile. *Org. Biomol. Chem.* **2011**, *9*, 1714–1716.
- (609) Collin, J.; Giuseppone, N.; Machrouhi, F.; Namy, J. L.; Nief, F. New Synthesis and Reactions of $[\text{Sm}(\text{OTf})_2(\text{DME})_2]$, a Salt-Free Samarium(II) Triflate. *Tetrahedron Lett.* **1999**, *40*, 3161–3164.
- (610) Hanamoto, T.; Sugimoto, Y.; Sugino, A.; Inanaga, J. Preparation and Reaction of Lanthanide(II) Trifluoromethanesulfonates. *Synlett.* **1994**, *1994* (5), 377–378.
- (611) van den Hende, J. R.; Hitchcock, P. B.; Lappert, M. F. Reactions of Ytterbium(II) Amides with Various Brønsted Acids, CS_2 or LiNR_2 ; Crystal Structures of $[\{\text{Yb}(\text{NR}_2)(\mu\text{-OCBu}_3^t)\}_2]$ and $[\text{Yb}(\text{OCBu}_3^t)_2(\text{thf})_2]$ (R = SiMe_3 , thf = tetrahydrofuran). *J. Chem. Soc., Dalton Trans.* **1995**, 2251–2258.
- (612) Bodizs, G.; Raabe, I.; Scopelliti, R.; Krossing, I.; Helm, L. Synthesis, Structures and Characterisations of Truly Homoleptic Acetonitrile Ln^{3+} Complexes in Solid State and in Solution. *Dalton Trans.* **2009**, 5137–5147.
- (613) Xémard, M.; Jaoul, A.; Cordier, M.; Molton, F.; Cador, O.; Le Guennic, B.; Duboc, C.; Maury, O.; Clavaguéra, C.; Nocton, G. Divalent Thulium Triflate: A Structural and Spectroscopic Study. *Angew. Chem., Int. Ed.* **2017**, *56*, 4266–4271.
- (614) Imamoto, T.; Koide, Y.; Hiyama, S. Cerium(IV) Trifluoromethanesulfonate as a Strong Oxidizing Agent. *Chem. Lett.* **1990**, *19*, 1445–1446.
- (615) Berthet, J. C.; Lance, M.; Nierlich, M.; Ephritikhine, M. Simple Preparations of the Anhydrous and Solvent-Free Uranyl and Cerium(IV) Triflates $\text{UO}_2(\text{OTf})_2$ and $\text{Ce}(\text{OTf})_4$ - Crystal Structures of $\text{UO}_2(\text{OTf})_2(\text{Py})_3$ and $[\{\text{UO}_2(\text{Py})_4\}_2(\mu\text{-O})][\text{OTf}]_2$. *Eur. J. Inorg. Chem.* **2000**, *2000* (9), 1969–1973.
- (616) Hickson, J. R.; Horsewill, S. J.; Bamforth, C.; McGuire, J.; Wilson, C.; Sproules, S.; Farnaby, J. H. The Modular Synthesis of Rare Earth-Transition Metal Heterobimetallic Complexes Utilizing a Redox-Active Ligand. *Dalton Trans.* **2018**, *47*, 10692–10701.
- (617) Mahoney, B. D.; Piro, N. A.; Carroll, P. J.; Schelter, E. J. Synthesis, Electrochemistry, and Reactivity of Cerium(III/IV) Methylene-Bis-Phenolate Complexes. *Inorg. Chem.* **2013**, *52*, 5970–5977.
- (618) Rabe, G. W.; Riede, J.; Schier, A. Synthesis, X-Ray Crystal Structure Determination, and NMR Spectroscopic Investigation of Two Homoleptic Four-Coordinate Lanthanide Complexes: Trivalent $(\text{Bu}_2\text{P})_2\text{La}[(\mu\text{-PtBu}_2)_2\text{Li}(\text{thf})]$ and Divalent $\text{Yb}[(\mu\text{-PtBu}_2)_2\text{Li}(\text{thf})_2]$. *Inorg. Chem.* **1996**, *35*, 40–45.
- (619) Moehring, S. A.; Ziller, J. W.; Evans, W. J. Rare-Earth Complexes of the Asymmetric Amide Ligands, $\text{N}(\text{SiMe}_3)\text{Ph}$ and $\text{N}(\text{SiMe}_3)\text{Cy}$. *Polyhedron* **2019**, *168*, 72–79.
- (620) Wooten, A. J.; Carroll, P. J.; Walsh, P. J. Insight into Substrate Binding in Shibasaki's $\text{Li}_3(\text{THF})_n(\text{BINOLate})_3\text{Ln}$ Complexes and Implications in Catalysis. *J. Am. Chem. Soc.* **2008**, *130* (23), 7407–7419.
- (621) Smith, P. H.; Reyes, Z. E.; Lee, C. W.; Raymond, K. N. Characterization of a Series of Lanthanide Amine Cage Complexes. *Inorg. Chem.* **1988**, *27*, 4154–4165.
- (622) Cassani, M. C.; Gun'ko, Y. K.; Hitchcock, P. B.; Lappert, M. F.; Laschi, F. Synthesis and Characterization of Organolanthanidocene-(III) (Ln) La, Ce, Pr, Nd) Complexes Containing the 1,4-Cyclohexa-2,5-Dienyl Ligand (Benzene 1,4-Dianion): Structures of $[\text{K}(\{[18\text{-crown-6}]\})[\text{Ln}\{\eta^5\text{-C}_5\text{H}_3(\text{SiMe}_3)_2\text{-1,3}\}_2(\text{C}_6\text{H}_6)] [\text{Cp}'' = \eta^5\text{-C}_5\text{H}_3(\text{SiMe}_3)_2\text{-1,3}; \text{Ln} = \text{La, Ce, Nd}]$. *Organometallics* **1999**, *18* (26), 5539–5547.
- (623) Hitchcock, P. B.; Hulkes, A. G.; Lappert, M. F.; Protchenko, A. V. Diversity of Triflate Coordination Modes in Neodymocene Complexes Containing Bulky Bis(Trimethylsilyl)Cyclopentadienyl Ligands. *Inorg. Chim. Acta* **2006**, *359*, 2998–3006.
- (624) Kilimann, U.; Edelmann, F. T. Cyclooctatetraenyl-Komplexe Der Fruhen Ubergangsmetalle Und Lanthanoid II. Neue Cyclooctatetraenyl-Halbsandwich-Komplexe Des Yttrium. *J. Organomet. Chem.* **1994**, *469*, C5–C9.
- (625) Simpson, C. K.; White, R. E.; Carlson, C. N.; Wroblewski, D. A.; Kuehl, C. J.; Croce, T. A.; Steele, I. M.; Scott, B. L.; Young, V. G.; Hanusa, T. P.; et al. The Role of Alkali Metal Cations in MMA Polymerization Initiated by Neutral and Anionic Allyl Lanthanide Complexes. *Organometallics* **2005**, *24*, 3685–3691.
- (626) White, R. E.; Carlson, C. N.; Veauthier, J. M.; Simpson, C. K.; Thompson, J. D.; Scott, B. L.; Hanusa, T. P.; John, K. D. Observation of Internal Electron Transfer in Bulky Allyl Ytterbium Complexes with Substituted Terpyridine Ligands. *Inorg. Chem.* **2006**, *45*, 7004–7009.
- (627) Lauenstein, Å.; Tegenfeldt, J. Ion Coordination and Chain Mobility in $\text{La}(\text{CF}_3\text{SO}_3)_3\text{PEO}_n$ Studied with NMR Spectroscopy. *J. Phys. Chem. B* **1997**, *101*, 3311–3317.
- (628) Fuming, M.; Guangxing, L.; Jin, N.; Huibi, X. A Novel Catalyst for Transesterification of Dimethyl Carbonate with Phenol to Diphenyl Carbonate: Samarium Trifluoromethanesulfonate. *J. Mol. Catal. A Chem.* **2002**, *184*, 465–468.
- (629) Egashira, K.; Yoshimura, Y.; Kanno, H.; Suzuki, Y. TG-DTA Study on the Lanthanoid Trifluoromethanesulfonate Complexes. *J. Therm. Anal. Calorim.* **2003**, *71*, 501–508.
- (630) Sella, A.; Brown, S. E.; Steed, J. W.; Tocher, D. A. Synthesis and Solid-State Structures of Pyrazolymethane Complexes of the Rare Earths. *Inorg. Chem.* **2007**, *46*, 1856–1864.
- (631) Dietrich, H. M.; Raudaschl-Sieber, G.; Anwender, R. Trimethylyttrium and Trimethyltutetium. *Angew. Chem., Int. Ed.* **2005**, *44*, 5303–5306.
- (632) Hart, F. A.; Massey, A. G.; Saran, M. S. Phenyls and Alkyls of the Group IIIA Metals. *J. Organomet. Chem.* **1970**, *21*, 147–154.
- (633) Barisic, D.; Diether, D.; Maichle-Mössner, C.; Anwender, R. Trimethylscandium. *J. Am. Chem. Soc.* **2019**, *141*, 13931–13940.
- (634) Zimmermann, M.; Rauschmaier, D.; Eichele, K.; Törnroos, K. W.; Anwender, R. Amido-Stabilized Rare-Earth Metal Mixed Methyl Methylidene Complexes. *Chem. Commun.* **2010**, *46*, 5346–5348.
- (635) Schumann, H.; Müller, J. Tris[(N,N,N',N'-Tetramethylethylenediamine) Lithium]Hexamethylerbate(III) and -Lutetate(III). *Angew. Chem., Int. Ed.* **1978**, *17*, 276.
- (636) Schumann, H.; Pickardt, J.; Bruncks, N. Crystal and Molecular Structure of $[\text{Li}(\text{Tmen})]_3[\text{Er}(\text{CH}_3)_6]$. *Angew. Chem., Int. Ed.* **1981**, *20*, 120–121.
- (637) Schumann, H.; Lauke, H.; Hahn, E.; Pickardt, J. Metalorganische Verbindungen Der Lanthanoiden. XVIII. Synthese Und Struktur von Tris[(1,2-Dimethoxyethan)Lithium]Hexamethyltutetate(III). *J. Organomet. Chem.* **1984**, *263*, 29–35.
- (638) Schumann, H.; Müller, J.; Bruncks, N.; Lauke, H.; Pickardt, J.; Schwarz, H.; Eckart, K. Organometallic Compounds of the Lanthanides. Tris[(Tetramethylethylenediamine)Lithium] Hexamethyl Derivatives of the Rare Earths. *Organometallics* **1984**, *3*, 69–74.
- (639) Kramer, M. U.; Robert, D.; Arndt, S.; Zeimentz, P. M.; Spaniol, T. P.; Yahia, A.; Maron, L.; Eisenstein, O.; Okuda, J. Cationic Methyl Complexes of the Rare-Earth Metals: An Experimental and Computational Study on Synthesis, Structure, and Reactivity. *Inorg. Chem.* **2008**, *47*, 9265–9278.

- (640) Berger, T.; Lebon, J.; Maichle-Mössmer, C.; Anwander, R. $\text{CeCl}_3/n\text{-BuLi}$: Unraveling Imamoto's Organocerium Reagent. *Angew. Chem., Int. Ed.* **2021**, *60*, 15622–15631.
- (641) Lappert, M. F.; Pearce, R. Stable Silylmethyl and Neopentyl Complexes of Scandium(III) and Yttrium(III). *J. Chem. Soc., Chem. Commun.* **1973**, 126.
- (642) Arndt, S.; Voth, P.; Spaniol, T. P.; Okuda, J. Dimeric Hydrido Complexes of Rare-Earth Metals Containing a Linked Amido-Cyclopentadienyl Ligand: Synthesis, Characterization, and Monomer-Dimer Equilibrium. *Organometallics* **2000**, *19*, 4690–4700.
- (643) Schumann, H.; Freckmann, D. M. M.; Dechert, S. The Molecular Structure of Tris (Trimethylsilylmethyl) Samarium, -Erbium, -Ytterbium, and -Lutetium. *Z. Anorg. Allg. Chem.* **2002**, *628*, 2422–2426.
- (644) Atwood, J. L.; Hunter, W. E.; Rogers, R. D.; Holton, J.; McMeeking, J.; Pearce, R.; Lappert, M. F. Neutral and Anionic Silylmethyl Complexes of the Group 3a and Lanthanoid Metals; the X-ray crystal and molecular structure of $[\text{Li}(\text{thf})_4][\text{Yb}\{\text{CH}(\text{SiMe}_3)_2\}_3\text{Cl}]$ (thf = tetrahydrofuran). *J. Chem. Soc., Chem. Commun.* **1978**, 140–142.
- (645) Evans, W. J.; Brady, J. C.; Ziller, J. W. Double Deprotonation of a Cyclopentadienyl Alkene to Form a Polydentate Trianionic Cyclopentadienyl Alkyl Ligand System. *J. Am. Chem. Soc.* **2001**, *123* (31), 7711–7712.
- (646) Barker, G. K.; Lappert, M. F. Stabilisation of Transition Metals in a Low Coordinative Environment Using the Bis(Trimethylsilyl)-Methyl Ligand; a Monomeric Cr^{III} Alkyl, $\text{Cr}[\text{CH}(\text{SiMe}_3)_2]_3$, and Related Complexes. *J. Organomet. Chem.* **1974**, *76*, C45–C46.
- (647) Mortis, A.; Barisic, D.; Eichele, K.; Maichle-Mössmer, C.; Anwander, R. Scandium Bis(Trimethylsilyl)methyl Complexes Revisited: Extending the ^{45}Sc NMR Chemical Shift Range and a New Structural Motif of $[\text{Li}\{\text{CH}(\text{SiMe}_3)_2\}]$. *Dalton Trans.* **2020**, *49*, 7829–7841.
- (648) Arndt, S.; Spaniol, T. P.; Okuda, J. Homogeneous Ethylene-Polymerization Catalysts Based on Alkyl Cations of the Rare-Earth Metals: Are Dicationic Mono(Alkyl) Complexes the Active Species? *Angew. Chem., Int. Ed.* **2003**, *42*, 5075–5079.
- (649) Elvidge, B. R.; Arndt, S.; Zeimentz, P. M.; Spaniol, T. P.; Okuda, J. Cationic Rare-Earth Metal Trimethylsilylmethyl Complexes Supported by THF and 12-Crown-4 Ligands: Synthesis and Structural Characterization. *Inorg. Chem.* **2005**, *44*, 6777–6788.
- (650) Li, X.; Nishiura, M.; Hu, L.; Mori, K.; Hou, Z. Alternating and Random Copolymerization of Isoprene and Ethylene Catalyzed by Cationic Half-Sandwich Scandium Alkyls. *J. Am. Chem. Soc.* **2009**, *131*, 13870–13882.
- (651) Nishiura, M.; Baldamus, J.; Shima, T.; Mori, K.; Hou, Z. Synthesis and Structures of the $\text{C}_5\text{Me}_4\text{SiMe}_3$ -Supported Polyhydride Complexes over the Full Size Range of the Rare Earth Series. *Chem. - A Eur. J.* **2011**, *17*, 5033–5044.
- (652) Butovskii, M. V.; Kempe, R. Rare Earth-Metal Bonding in Molecular Compounds: Recent Advances, Challenges, and Perspectives. *New J. Chem.* **2015**, *39*, 7544–7558.
- (653) Butovskii, M. V.; Tok, O. L.; Wagner, F. R.; Kempe, R. Bismetallocenes: Lanthanoid-Transition-Metal Bonds through Alkane Elimination. *Angew. Chem., Int. Ed.* **2008**, *47*, 6469–6472.
- (654) Butovskii, M. V.; Oelkers, B.; Bauer, T.; Bakker, J. M.; Bezugly, V.; Wagner, F. R.; Kempe, R. Lanthanoid-Transition-Metal Bonding in Bismetallocenes. *Chem. - A Eur. J.* **2014**, *20*, 2804–2811.
- (655) Butovskii, M. V.; Döring, C.; Bezugly, V.; Wagner, F. R.; Grin, Y.; Kempe, R. Molecules Containing Rare-Earth Atoms Solely Bonded by Transition Metals. *Nat. Chem.* **2010**, *2*, 741–744.
- (656) Estler, F.; Eickerling, G.; Herdtweck, E.; Anwander, R. Organo-Rare-Earth Complexes Supported by Chelating Diamide Ligands. *Organometallics* **2003**, *22*, 1212–1222.
- (657) Aillerie, A.; Rodriguez-Ruiz, V.; Carlino, R.; Bourdreux, F.; Guillot, R.; Bezzenine-Lafollée, S.; Gil, R.; Prim, D.; Hannedouche, J. Asymmetric Assisted Tandem Catalysis: Hydroamination Followed by Asymmetric Friedel–Crafts Reaction from a Single Chiral N,N,N',N'-Tetradentate Pyridylmethylamine-Based Ligand. *Chem. Catal. Chem.* **2016**, *8*, 2455–2460.
- (658) Emslie, D. J. H.; Piers, W. E.; Parvez, M.; McDonald, R. Organometallic Complexes of Scandium and Yttrium Supported by a Bulky Salicylaldehyde Ligand. *Organometallics* **2002**, *21*, 4226–4240.
- (659) Niemeyer, M. A Trigonal-Bipyramidal Coordinated Ytterbium(III) Alkyl: Tris(trimethylsilylmethyl)Bis(tetrahydrofuran-O)-Ytterbium(III). *Acta Crystallogr. Sect. E Struct. Reports Online* **2001**, *57*, m553–m555.
- (660) Chen, S. M.; Zhang, Y. Q.; Xiong, J.; Wang, B. W.; Gao, S. Adducts of Tris(Alkyl) Holmium(III) Showing Magnetic Relaxation. *Inorg. Chem.* **2020**, *59*, 5835–5844.
- (661) Yan, C.; Liu, Z.-X.; Xu, T.-Q. Regioselective, Stereoselective, and Living Polymerization of Divinyl Pyridine Monomers Using Rare Earth Catalyst. *Polym. Chem.* **2020**, *11*, 2044–2052.
- (662) Avent, A. G.; Caro, C. F.; Hitchcock, P. B.; Lappert, M. F.; Li, Z.; Wei, X.-H. Synthetic and Structural Experiments on Yttrium, Cerium and Magnesium Trimethylsilylmethyls and Their Reaction Products with Nitriles; with a Note on Two Cerium β -Diketiminates. *Dalton Trans.* **2004**, 1567–1577.
- (663) Qi, G.; Nitto, Y.; Saiki, A.; Tomohiro, T.; Nakayama, Y.; Yasuda, H. Isospecific Polymerizations of Alkyl Methacrylates with a Bis(Alkyl) Yb Complex and Formation of Stereocomplexes with Syndiotactic Poly(Alkyl Methacrylate)s. *Tetrahedron* **2003**, *59*, 10409–10418.
- (664) Zhu, X.; Guo, D.; Yao, F.; Huang, Z.; Li, Y.; Xie, Z.; Wang, S. Synthesis, Characterization, and Catalytic Activity of Dinuclear Rare-Earth Metal Alkyl Complexes Bearing 2,6-Diisopropylphenylamido-methyl Functionalized Pyrrolyl Ligand. *Organometallics* **2021**, *40*, 1633–1641.
- (665) Cui, D.; Nishiura, M.; Hou, Z. Alternating Copolymerization of Cyclohexene Oxide and Carbon Dioxide Catalyzed by Organo Rare Earth Metal Complexes. *Macromolecules* **2005**, *38*, 4089–4095.
- (666) Lukešová, L.; Ward, B. D.; Bellemin-Lapponaz, S.; Wadepohl, H.; Gade, L. H. High Tacticity Control in Organolanthanide Polymerization Catalysis: Formation of Isotactic Poly(α -Alkenes) with a Chiral C_3 -Symmetric Thulium Complex. *Dalton Trans.* **2007**, 920–922.
- (667) Hong, D.; Zhu, X.; Wang, S.; Wei, Y.; Zhou, S.; Huang, Z.; Zhu, S.; Wang, R.; Yue, W.; Mu, X. Synthesis, Characterization, and Reactivity of Dinuclear Organo-Rare-Earth-Metal Alkyl Complexes Supported by 2-Amidate-Functionalized Indolyl Ligands: Substituent Effects on Coordination and Reactivity. *Dalton Trans.* **2019**, *48*, 5230–5242.
- (668) Yan, C.; Xu, T. Q.; Lu, X. B. From Stereochemically Tunable Homopolymers to Stereomultiblock Copolymers: Lewis Base Regulates Stereochemistry in the Coordination Polymerization of 2-Vinylpyridine. *Macromolecules* **2018**, *51*, 2240–2246.
- (669) Rufanov, K. A.; Freckmann, D. M. M.; et al. Studies on the Thermolysis of Ether-Stabilized $\text{Lu}(\text{CH}_2\text{SiMe}_3)_3$. Molecular Structure of $\text{Lu}(\text{CH}_2\text{SiMe}_3)_3(\text{THF})(\text{Diglyme})$. *Z. fur Naturforsch. - Sect. B J. Chem. Sci.* **2005**, *3*, 533–537.
- (670) Manzer, L. E. Paramagnetic Organometallic Compounds of the Early Transition Metals Stabilized by Chelating Benzyl and Phenyl Ligands. *J. Am. Chem. Soc.* **1978**, *100*, 8068–8073.
- (671) Harder, S. Syntheses and Structures of Homoleptic Lanthanide Complexes with Chelating O-Dimethylaminobenzyl Ligands: Key Precursors in Lanthanide Chemistry. *Organometallics* **2005**, *24*, 373–379.
- (672) Ruspic, C.; Moss, J. R.; Schürmann, M.; Harder, S. Remarkable Stability of Metallocenes with Superbulky Ligands: Spontaneous Reduction of Sm^{III} to Sm^{II} . *Angew. Chem., Int. Ed.* **2008**, *47*, 2121–2126.
- (673) Harder, S.; Ruspic, C.; Bhriain, N. N.; Berkermann, F.; Schürmann, M. Benzyl Complexes of Lanthanide(II) and Lanthanide(III) Metals: Trends and Comparisons. *Z. Naturforsch. - Sect. B J. Chem. Sci.* **2008**, *63*, 267–274.
- (674) Zhang, W.; Nishiura, M.; Mashiko, T.; Hou, Z. Range of Group 3 and Lanthanide Metals. Synthesis, Structural Characterization, and Catalysis of Phosphine P-H Bond Addition to Carbodiimides. *Chem. - A Eur. J.* **2008**, *14*, 2167–2179.

- (675) Bambirra, S.; Meetsma, A.; Hessen, B. Lanthanum Tribenzyl Complexes as Convenient Starting Materials for Organolanthanum Chemistry. *Organometallics* **2006**, *25*, 3454–3462.
- (676) Harder, S.; Ruspig, C.; Bhriain, N.; Berkermann, F.; Schürmann, M. Benzyl Complexes of Lanthanide(II) and Lanthanide(III) Metals: Trends and Comparisons. *Z. Naturforsch.* **2008**, *63*, 267–274.
- (677) Meyer, N.; Roesky, P. W.; Bambirra, S.; Meetsma, A.; Hessen, B.; Saliu, K.; Takats, J. Synthesis and Structures of Scandium and Lutetium Benzyl Complexes. *Organometallics* **2008**, *27*, 1501–1505.
- (678) Wooles, A. J.; Mills, D. P.; Lewis, W.; Blake, A. J.; Liddle, S. T. Lanthanide Tri-Benzyl Complexes: Structural Variations and Useful Precursors to Phosphorus-Stabilised Lanthanide Carbenes. *Dalton Trans.* **2010**, *39*, 500–510.
- (679) Mills, D. P.; Wooles, A. J.; McMaster, J.; Lewis, W.; Blake, A. J.; Liddle, S. T. Heteroleptic $[M(CH_2C_6H_5)_2(I)(THF)_3]$ Complexes ($M = Y$ or Er): Remarkably Stable Precursors to Yttrium and Erbium T-Shaped Carbenes. *Organometallics* **2009**, *28*, 6771–6776.
- (680) Harder, S. The Chemistry of Ca^{II} and Yb^{II} : Astoundingly Similar but Not Equal! *Angew. Chem., Int. Ed.* **2004**, *43*, 2714–2718.
- (681) Wolf, B. M.; Stuhl, C.; Anwander, R. Synthesis of Homometallic Divalent Lanthanide Organonimides from Benzyl Complexes. *Chem. Commun.* **2018**, *54*, 8826–8829.
- (682) Kong, F.; Li, M.; Zhou, X.; Zhang, L. Synthesis, Structure and Reactivity of Guanidinate Rare Earth Metal Bis(O-Aminobenzyl) Complexes. *RSC Adv.* **2017**, *7*, 29752–29761.
- (683) Jiang, W.; Zhang, L. J.; Zhang, L. X. Synthesis, Structure, and Reactivity of Monoguanidinate Rare-Earth Metal Aminobenzyl Enolate Complexes. *Eur. J. Inorg. Chem.* **2020**, *2020*, 2153–2164.
- (684) Harder, S.; Naglav, D.; Ruspig, C.; Wickleder, C.; Adlung, M.; Hermes, W.; Eul, M.; Pöttgen, R.; Rego, D. B.; Poineau, F.; et al. Physical Properties of Superbulky Lanthanide Metallocenes: Synthesis and Extraordinary Luminescence of $[Eu^{II}(Cp^{BIG})_2]$ ($Cp^{BIG} = (4-nBu-C_6H_4)_5-Cyclopentadienyl$). *Chem. - A Eur. J.* **2013**, *19*, 12272–12280.
- (685) Mills, D. P.; Cooper, O. J.; McMaster, J.; Lewis, W.; Liddle, S. T. Synthesis and Reactivity of the Yttrium-Alkyl-Carbene Complex $[Y(BIPM)(CH_2C_6H_5)(THF)]$ ($BIPM = \{C(PPh_2NSiMe_3)_2\}$). *Dalton Trans.* **2009**, 4547–4555.
- (686) Gregson, M.; Chilton, N. F.; Ariciu, A.-M.; Tuna, F.; Crowe, I. F.; Lewis, W.; Blake, A. J.; Collison, D.; McInnes, E. J. L.; Winpenny, R. E. P.; et al. A Monometallic Lanthanide Bis(Methanediide) Single Molecule Magnet with a Large Energy Barrier and Complex Spin Relaxation Behaviour. *Chem. Sci.* **2016**, *7*, 155–165.
- (687) Gregson, M.; Lu, E.; Mills, D. P.; Tuna, F.; McInnes, E. J. L.; Hennig, C.; Scheinost, A. C.; McMaster, J.; Lewis, W.; Blake, A. J.; et al. The Inverse-Trans-Influence in Tetravalent Lanthanide and Actinide Bis(Carbene) Complexes. *Nat. Commun.* **2017**, *8*, 14137.
- (688) Liddle, S. T.; Mills, D. P.; Gardner, B. M.; McMaster, J.; Jones, C.; Woodul, W. D. A Heterobimetallic Gallyl Complex Containing an Unsupported Ga-Y Bond. *Inorg. Chem.* **2009**, *48*, 3520–3522.
- (689) Van Velzen, N. J. C.; Harder, S. Deca-Arylsamarocene: An Unusually Inert Sm(II) Sandwich Complex. *Organometallics* **2018**, *37*, 2263–2271.
- (690) Xie, H.; Wu, C.; Cui, D.; Wang, Y. Ligand-Free Scandium Alkyl and Alkoxide Complexes for Immortal Ring-Opening Polymerization of Lactide. *J. Organomet. Chem.* **2018**, *875*, 5–10.
- (691) Evans, W. J.; Anwander, R.; Ziller, J. W. Inclusion of Al_2Me_6 in the Crystalline Lattice of the Organometallic Complexes $LnAl_3Me_{12}$. *Organometallics* **1995**, *14*, 1107–1109.
- (692) Zimmermann, M.; Frøystein, N. Å.; Fischbach, A.; Sirsch, P.; Dietrich, H. M.; Törnroos, K. W.; Herdtweck, E.; Anwander, R. Homoleptic Rare-Earth Metal(III) Tetramethylaluminates: Structural Chemistry, Reactivity, and Performance in Isoprene Polymerization. *Chem. - A Eur. J.* **2007**, *13*, 8784–8800.
- (693) Nieland, A.; Mix, A.; Neumann, B.; Stämmler, H. G.; Mitzel, N. W. Lanthanoid Tetramethylaluminates and Their Paramagnetic NMR Parameters. *Eur. J. Inorg. Chem.* **2014**, *2014*, 51–57.
- (694) Klooster, W. T.; Lu, R. S.; Anwander, R.; Evans, W. J.; Koetzle, T. F.; Bau, R. Neutron Diffraction Study of $[Nd(AlMe_4)_3] \cdot 0.5 Al_2Me_6$ at 100 K: The First Detailed Look at a Bridging Methyl Group with a Trigonal-Bipyramidal Carbon Atom. *Angew. Chem., Int. Ed.* **1998**, *37*, 1268–1270.
- (695) Holton, J.; Lappert, M. F.; Ballard, D. G. H.; Pearce, R.; Atwood, J. L.; Hunter, W. E. Alkyl-Bridged Complexes of the d- and f-Block Elements. Part 2. Bis[Bis(η -Cyclopentadienyl)Methylmetal(III)] Complexes, and the Crystal and Molecular Structure of the Yttrium and Ytterbium Species. *J. Chem. Soc., Dalton Trans.* **1979**, 54–61.
- (696) Schrems, M. G.; Dietrich, H. M.; Törnroos, K. W.; Anwander, R. $[Ln^{II}Al^{III}_2(Alkyl)_8]_x$: Donor Addition Instead of Donor-Induced Cleavage. *Chem. Commun.* **2005**, *2* (47), 5922–5924.
- (697) Robert, D.; Spaniol, T. P.; Okuda, J. Neutral and Monocationic Half-Sandwich Methyl Rare-Earth Metal Complexes: Synthesis, Structure, and 1,3-Butadiene Polymerization Catalysis. *Eur. J. Inorg. Chem.* **2008**, *2008*, 2801–2809.
- (698) Dietrich, H. M.; Zapilko, C.; Herdtweck, E.; Anwander, R. $Ln(AlMe_4)_3$ as New Synthetic Precursors in Organolanthanide Chemistry: Efficient Access to Half-Sandwich Hydrocarbyl Complexes. *Organometallics* **2005**, *24*, 5767–5771.
- (699) Zimmermann, M.; Estler, F.; Herdtweck, E.; Törnroos, K. W.; Anwander, R. Distinct C-H Bond Activation Pathways in Diamidopyridine-Supported Rare-Earth Metal Hydrocarbyl Complexes. *Organometallics* **2007**, *26*, 6029–6041.
- (700) Zimmermann, M.; Takats, J.; Kiel, G.; Törnroos, K. W.; Anwander, R. Ln(III) Methyl and Methylidene Complexes Stabilized by a Bulky Hydrotris(Pyrazolyl)Borate Ligand. *Chem. Commun.* **2008**, 612–614.
- (701) Litlabø, R.; Zimmermann, M.; Saliu, K.; Takats, J.; Törnroos, K. W.; Anwander, R. A Rare-Earth Metal Variant of the Tebbe Reagent. *Angew. Chem., Int. Ed.* **2008**, *47*, 9560–9564.
- (702) Le Roux, E.; Nief, F.; Jaroschik, F.; Törnroos, K. W.; Anwander, R. Mono-Phosphacyclopentadienyl Bis(Tetramethylaluminate) Lanthanide Complexes. *Dalton Trans.* **2007**, 4866–4870.
- (703) Barisic, D.; Schneider, D.; Maichle-Mössmer, C.; Anwander, R. Formation and Reactivity of an Aluminabenzene Ligand at Pentadienyl-Supported Rare-Earth Metals. *Angew. Chem., Int. Ed.* **2019**, *58*, 1515–1518.
- (704) Dietrich, H. M.; Törnroos, K. W.; Anwander, R. $LaAl_3Et_2$: A Homoleptic Ethyllanthanum Complex. *Angew. Chem., Int. Ed.* **2011**, *50*, 12089–12093.
- (705) Klimpel, M. G.; Anwander, R.; Tafipolsky, M.; Scherer, W. Peralkylated Ytterbium(II) Aluminate Complexes $YbAl_2R_8$. New Insights into the Nature of Aluminate Coordination. *Organometallics* **2001**, *20*, 3983–3992.
- (706) Hollfelder, C. O.; Jende, L. N.; Diether, D.; Zelger, T.; Stauder, R.; Maichle-Mössmer, C.; Anwander, R. 1,3-Diene Polymerization Mediated By Homoleptic Tetramethylaluminates of the Rare-Earth Metals. *Catalysts* **2018**, *8*, 61.
- (707) König, S. N.; Chilton, N. F.; Maichle-Mössmer, C.; Pineda, E. M.; Pugh, T.; Anwander, R.; Layfield, R. A. Fast Magnetic Relaxation in an Octahedral Dysprosium Tetramethyl-Aluminate Complex. *Dalton Trans.* **2014**, *43*, 3035–3038.
- (708) Gomba, T. P.; Ramanathan, A.; Rice, N. T.; La Pierre, H. S. The Chemical and Physical Properties of Tetravalent Lanthanides: Pr, Nd, Tb, and Dy. *Dalton Trans.* **2020**, *49*, 15945–15987.
- (709) Rice, N. T.; Popov, I. A.; Russo, D. R.; Bacsa, J.; Batista, E. R.; Yang, P.; Telsler, J.; La Pierre, H. S. Design, Isolation, and Spectroscopic Analysis of a Tetravalent Terbium Complex. *J. Am. Chem. Soc.* **2019**, *141*, 13222–13233.
- (710) Rice, N. T.; Popov, I. A.; Russo, D. R.; Gomba, T. P.; Ramanathan, A.; Bacsa, J.; Batista, E. R.; Yang, P.; La Pierre, H. S. Comparison of Tetravalent Cerium and Terbium Ions in a Conserved, Homoleptic Imidophosphorane Ligand Field. *Chem. Sci.* **2020**, *11*, 6149–6159.
- (711) Willauer, A. R.; Palumbo, C. T.; Fadaei-Tirani, F.; Zivkovic, I.; Douair, I.; Maron, L.; Mazzanti, M. Accessing the + IV Oxidation State in Molecular Complexes of Praseodymium. *J. Am. Chem. Soc.* **2020**, *142*, 5538–5542.
- (712) Willauer, A. R.; Palumbo, C. T.; Scopelliti, R.; Zivkovic, I.; Douair, I.; Maron, L.; Mazzanti, M. Stabilization of the Oxidation State

+ IV in Siloxide-Supported Terbium Compounds. *Angew. Chem., Int. Ed.* **2020**, *59*, 3549–3553.

(713) Palumbo, C. T.; Zivkovic, I.; Scopelliti, R.; Mazzanti, M. Molecular Complex of Tb in the + 4 Oxidation State. *J. Am. Chem. Soc.* **2019**, *141*, 9827–9831.

(714) Woen, D. H.; Kotyk, C. M.; Mueller, T. J.; Ziller, J. W.; Evans, W. J. Tris(Pentamethylcyclopentadienyl) Complexes of Late Lanthanides Tb, Dy, Ho, and Er: Solution and Mechanochemical Syntheses and Structural Comparisons. *Organometallics* **2017**, *36*, 4558–4563.

(715) Woen, D. H.; White, J. R. K.; Ziller, J. W.; Evans, W. J. Mechanochemical C–H Bond Activation: Synthesis of the Tuckover Hydrides, $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1:\eta^5-CH_2C_5Me_4)Ln(C_5Me_5)$ from Solvent-Free Reactions of $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$ with KC_5Me_5 . *J. Organomet. Chem.* **2019**, *899*, 120885.

(716) Fetrov, T. V.; Daly, S. R. Mechanochemical Synthesis and Structural Analysis of Trivalent Lanthanide and Uranium Diphenylphosphinodiboranates. *Dalton Trans.* **2021**, *50*, 11472–11484.

(717) Gerber, L. C. H.; Le Roux, E.; Törnroos, K. W.; Anwander, R. Elusive Trimethylanthanum: Snapshots of Extensive Methyl Group Degradation in La–Al Heterobimetallic Complexes. *Chem. - A Eur. J.* **2008**, *14*, 9555–9564.

NOTE ADDED AFTER ASAP PUBLICATION

This paper was published ASAP on January 31, 2022, with errors in Schemes 21 and 79. The corrected version was reposted on February 2, 2022.