Re(CO)₅Br-Catalyzed Coupling of Epoxides with CO₂ Affording Cyclic Carbonates under Solvent-Free Conditions

Jia-Li Jiang, Gao Feixue, Ruimao Hua*, Xianqing Qiu

*Department of Chemistry, Tsinghua University, Key Laboratory of Organic Optoelectronics & Molecular Engineering of Ministry of Education, Beijing 100084, China
ruimao@mail.tsinghua.edu.cn*

Supporting Information:

- 1. Experimental Section (p S1 ~ S2)**
- 2. The charts of ¹H and ¹³C NMR of 2a ~ 2e (p S3 ~ S12)**

1. Experimental Section

Re(CO)₅Br, Re(CO)₅Cl, CpRe(CO)₃ and Re₂(CO)₁₀ were commercial complexes from Strem. All products are known compounds, which were identified by ¹H, ¹³C NMR spectra and GCMS. ¹H and ¹³C NMR spectra were recorded on a JOEL JNM-ECA 300 spectrometer at 300 MHz and 75 MHz, respectively. ¹H and ¹³C NMR chemical shifts (δ) were referenced to internal solvent resonances.

A typical experiment for the synthesis of 4-chloromethyl-[1, 3]dioxolan-2-one 2a (Table 1, entry 8): Chloromethyloxirane **1a** (10.0 mmol) and Re(CO)₅Br (0.01 mmol) were charged in a 25 mL-autoclave, and then CO₂ was introduced at an initial pressure of 5.5

MPa at room temperature. The autoclave was heated at 110 °C with stirring for 24 h. After the autoclave was cooled to room temperature, CO₂ was released slowly. To the reaction mixture, 3.0 mmol of mesitylene as an internal standard material for GC analysis was added with stirring. Analysis of the resulting mixture by GC and GC-MS revealed that the conversion of **1a** was 100 %, **2a** was formed in 97 % GC yield. **2a** was obtained in 93 % as an isolated yield by Kugelrohr distillation.

2a ~ e were known compounds and identified by ¹H, ¹³C NMR and GC-MS.

2. The charts of ¹H and ¹³C NMR of 2a ~ e

jiang031121.8
single_pulse

☑ X : parts per Million : 1H

jiang-040427-362-H-single_pulse.2
single_pulse

2b

jiang-040427-362-C-single_pulse_dec.2
single pulse decoupled gated NOE

jiang-040427-377-H-single_pulse.2
single_pulse

jiang-040427-377-C-single_pulse_dec.2
single pulse decoupled gated NOE

2c

jiang-040427-393-H-single_pulse.2
single_pulse

jiang-040427-393-H-single_pulse_dec.2
single pulse decoupled gated NOE

jiang-040427-383-H-single_pulse.2
single_pulse

jiang-04027-383-C-single_pulse_dec.2
single pulse decoupled gated NOE

Δ X : parts per Million : 13C