Recyclable and Reusable Pd(OAc)₂/DABCO/PEG-400 System for Suzuki-Miyaura Cross-Coupling Reaction

Jin-Heng Li,* Wen-Jie Liu, and Ye-Xiang Xie

*Key Laboratory of Chemical Biology & Traditional Chinese Medicine Research, College of
Chemistry and Chemical Engineering, Hunan Normal University, Changsha 410081, China.*

E-mail address: jhli@hunnu.edu.cn

Supporting Information

List of Contents

(A) Typical experimental procedure for the palladium-catalyzed Suzuki-Miyaura cross-coupling reaction in PEG	S2
(B) Analytical data for 3-14	S2-5
(C) References	S5
(D) NMR spectra	S6-29

(A) Typical Experimental Procedure for the Palladium-Catalyzed Suzuki-Miyaura cross-coupling reaction in PEG-400.

A mixture of aryl halide **1** (0.5 mmol), arylboronic acid **2** (0.7 mmol), Pd(OAc)₂/DABCO (1 : 2; the identical amount in Table 1 and 2), K₂CO₃ (3 equiv), and PEG-400 (2 g) was added to a sealed tube, and stirred at 110 °C for desired time until complete consumption of starting material as judged by TLC. After the mixture was extracted with dry ethyl ether (3×10 mL) and evaporated, the residue was purified by flash column chromatography (hexane or hexane/ethyl acetate) to afford the desired coupled products **3—14**.

After extracting with diethyl ether, the mixture of Pd(OAc)₂, DABCO, and PEG-400 were solidified (cooled, then evaporated under vacuo) and subjected to a second run of the Suzuki-Miyaura cross-coupling reaction by charging with the same substrates (aryl halide **1**, arylboronic acid **2**, and K₂CO₃).

(B) Analytical data for 3-14

4-Nitro-biphenyl (3)

¹H NMR (400 MHz, CDCl₃) δ: 8.31 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 8.8 Hz, 2H), 7.64 (d, *J* = 7.2 Hz, 2H), 7.53—7.46 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 147.6, 147.1, 138.8, 129.2, 128.91, 127.8, 127.4, 124.1.

2-Nitro-biphenyl (4)

¹H NMR (400 MHz, CDCl₃): δ 7.86 (d, *J* = 8 Hz, 1H), 7.62 (t, *J* = 7.2 Hz, 1H),

7.51—7.40 (m, 5H), 7.34—7.31 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.7, 140.3, 137.3, 128.8, 128.7, 128.6, 127.3, 127.0, 115.7, 115.5.

Biphenyl (5)

^1H NMR (300 MHz, CDCl_3) δ : 7.59 (d, $J = 8.4$ Hz, 4H), 7.43 (t, $J = 7.2$ Hz, 4H), 7.36 (t, $J = 7.8$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ : 141.6, 129.1, 127.6, 127.5.

4-Fluoro-biphenyl (6)

^1H NMR (400 MHz, CDCl_3) δ : 7.57—7.53 (m, 4H), 7.44 (t, $J = 7.6$ Hz, 2H), 7.35 (t, $J = 7.2$ Hz, 1H), 7.13 (t, $J = 8.8$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.7, 161.2, 140.2, 137.3, 128.8, 127.2, 115.8, 115.6.

4-Trifluoromethyl-biphenyl (7)

^1H NMR (400 MHz, CDCl_3) δ : 7.69 (s, 4H), 7.60 (d, $J = 8$ Hz, 2H), 7.48 (t, $J = 7.2$ Hz, 2H), 7.40 (t, $J = 7.2$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ : 144.7, 139.8, 129.0, 128.2, 127.6, 127.4, 127.3, 125.9, 125.7.

4-Chloro-biphenyl (8)

^1H NMR (400 MHz, CDCl_3): δ 7.57—7.49 (m, 4H), 7.46—7.34 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3): δ 140.0, 139.6, 133.3, 129.0, 128.9, 128.4, 127.6, 127.0.

4-Methoxy-biphenyl (9)

^1H NMR (300 MHz, CDCl_3): δ 7.54 (t, $J = 8.4$ Hz, 4H), 7.41 (t, $J = 7.2$ Hz, 2H), 7.31 (t, $J = 7.2$, 1H), 6.98 (d, $J = 6.7$, 2H), 3.85 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 159.1, 140.8, 133.7, 128.7, 128.1, 126.7, 126.6, 114.2, 55.3.

1-Biphenyl-4-yl-ethanone (10)

^1H NMR (400 MHz, CDCl_3): δ 8.04 (d, $J = 8.4$ Hz, 2H), 7.69 (d, $J = 8.4$ Hz, 2H), 7.64 (d, $J = 7.6$ Hz, 2H), 7.48 (t, $J = 7.6$ Hz, 2H), 7.41 (t, $J = 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ 197.8, 145.8, 139.9, 135.8, 128.9, 128.9, 128.2, 127.3, 127.2, 26.7.

1-(4'-Fluoro-biphenyl-4-yl)-ethanone (11)

^1H NMR (400 MHz, CDCl_3) δ : 8.03 (d, $J = 8$ Hz, 2H), 7.64 (d, $J = 8$ Hz, 2H), 7.60—7.58 (m, 2H), 7.17 (t, $J = 8.4$ Hz, 2H), 2.65 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.7, 164.2, 161.7, 144.7, 135.9, 135.8, 129.0, 128.9, 127.1, 116.0, 115.8, 26.7.

2-methyl-4'-methoxy-biphenyl (12)

^1H NMR (400 MHz, CDCl_3) δ : 7.26-7.22 (m, 6H), 6.95 (d, $J = 8.4$ Hz, 2H), 3.85 (s, 3H), 2.28 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 158.5, 141.5, 135.5, 134.3, 130.3, 130.2, 129.9, 127.0, 125.8, 113.5, 55.3, 20.6.

4, 4''-Difluoro-[1,1';4',1'']terphenyl (13)

^1H NMR (400 MHz, CDCl_3) δ : 7.51—7.48 (m, 6H), 7.14—7.10 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.6, 161.2, 136.4, 131.9, 128.6, 128.5, 115.8, 115.6.

4-Methyl-biphenyl (14)

^1H NMR (400 MHz, CDCl_3) δ : 7.58 (t, $J = 7.6$ Hz, 2H), 7.49 (d, $J = 8.0$ Hz, 2H), 7.42 (t, $J = 7.6$ Hz, 2H), 7.31 (t, $J = 7.6$ Hz, 1H), 7.24 (d, $J = 8.0$ Hz, 2H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 141.1, 138.3, 137.0, 129.5, 128.7, 127.3, 127.2, 127.0, 21.1.

(C) References

- (1) (a) Tao, B.; Boykin, D. W. *Tetrahedron Lett.* **2002**, *43*, 4955.
(b) Alonso, D. A.; Nájera, C.; Pacheco, M. C. *Org. Lett.* **2000**, *2*, 1823.
(c) Botella, L.; Nájera, C. *Angew. Chem. Int. Ed.* **2002**, *41*, 179.
(d) Bedford, R. B.; Cazin, C. S. *Chem. Commun.* **2001**, 1540.
- (2) (a) Tao, B.; Boykin, D. W. *Tetrahedron Lett.* **2003**, *44*, 7993.
(b) Tao, B.; Boykin, D. W. *J. Org. Chem.* **2004**, ASAP.
- (3) Grasa, G. A.; Hillier, A. C.; Nolan, S. P. *Org. Lett.* **2001**, *3*, 1077.
- (4) Dai, M.; Liang, B.; Wang, C.; Chen, J.; Yang, Z. *Org. Lett.* **2004**, *6*, 221.

(D) NMR spectra

4-Nitro-biphenyl (3)

4-Nitro-biphenyl (3)

2-Nitro-biphenyl (4)

2-Nitro-biphenyl (4)

Biphenyl (5)

Biphenyl (5)

4-Fluoro-biphenyl (6)

4-Fluoro-biphenyl (6)

4-Trifluoromethyl-biphenyl (7)

4-Trifluoromethyl-biphenyl (7)

$$DHI + p\text{-ClC}_6\text{H}_4\text{SO}_3\text{H}, 2$$

4-Chloro-biphenyl (8)

4-Methoxy-biphenyl (9)

4-Methoxy-biphenyl (9)

$$p\text{-BrPhCOCH}_3 + \text{PhCOCH}_3$$

1-Biphenyl-4-yl-ethanone (10)

1-(4'-Fluoro-biphenyl-4-yl)-ethanone (11)

1-(4'-Fluoro-biphenyl-4-yl)-ethanone (11)

2-methyl-4'-methoxy-biphenyl (12)

2-methyl-4'-methoxy-biphenyl (12)

(4,4')-Difluoro-[1,1';4',1'']Terphenyl (13)

(4,4')-Difluoro-[1,1';4',1'']Terphenyl (13)

4-Methyl-biphenyl (14)

4-Methyl-biphenyl (14)