2,3-Dihydroisoxazoles as Precursors to β -Amino Ketones and β -Amino Alcohols

Patrick Aschwanden, Lisbet Kværnø, Roger Geisser, Florian Kleinbeck and Erick M. Carreira*

Laboratorium für Organische Chemie, ETH Hönggerberg,
CH-8093 Zürich, Switzerland

carreira@org.chem.ethz.ch

Supplementary Material

IR-Spectra were recorded on a *Perkin Elmer Spectrum RX I FT-IR spectrophotometer*. The samples were prepared as either KBr pellets or thin films on NaCl salt plates and are reported as absorption maxima in cm^{-1} .

^1H -NMR-Spectra were recorded on a *Varian Mercury 300* operating at 300 MHz or a *Gemini 200* operating at 200 MHz. Chemical shifts δ (ppm) were referenced to the internal solvent signals. Multiplicities are reported as follows: s (singlet), d (doublet), t (triplet), q (quartet) and m (multiplet). Coupling constants J are given in Hz.

^{13}C -NMR-Spectra were recorded on a *Varian Mercury 300* operating at 75 MHz. Chemical shifts δ (ppm) were referenced to the internal solvent signals.

Melting points were measured using a *Büchi 510* melting point apparatus and are uncorrected.

Elemental analysis was performed by the Mikroelementaranalytisches Laboratorium at the ETH, Zürich. Calculated and observed data are given in mass percent.

High resolution MALDI (HRMALDI) was performed by the Massenspektroskopie-Service at the ETH, Zürich. Detected ionmasses are given as m/z with intensities related to the most intensive signal (%).

TLC: Thin layer chromatography was performed using *Merck Silica Gel 60 F₂₅₄ TLC plates* and visualized either with ultraviolet light at 254 nm, stained iodine or by a solution of 5 g NaHCO_3 , 1.5 g KMnO_4 and 400 mL H_2O or 25 g phosphomolybdic acid, 10 g $\text{Ce}(\text{SO}_4)_2 \cdot \text{H}_2\text{O}$, 60 mL conc. H_2SO_4 and 940 mL H_2O .

Chromatographic purification of products was accomplished using forced flow chromatography on *Fluka Silica Gel 60* (40 - 63 μm) according to the method of Still.¹

Solvents were all purchased from either Aldrich or Fluka chemical companies and used without prior purification. For chromatographic purification, solvents were distilled prior to use.

Reactions: Reactions were all performed using oven dried glassware under an atmosphere of dry nitrogen or argon. Solvents and chemicals were transferred by syringe. Reactions were monitored by TLC. All chiral compounds were obtained in racemic form. ZnI_2 (>99.998%) was used unless otherwise noted.

(1) Still, W. C.; Ammon, H. L.; DeShong, P. J. *Am. Chem. Soc.* **1995**, *117*, 5166.

Preparation of Benzaldehyde Oxime²

To a solution of benzaldehyde (31.8 g, 0.300 mol) and hydroxylamine hydrochloride (64.5 g, 1.00 mol) in 1.00 L 90% EtOH was added powdered NaOH (108 g, 2.70 mol) in small portions. The mixture was allowed to stir at 25 °C for 30 min and then refluxed for another 30 min. The reaction mixture was then cooled to 25°C, poured into a mixture of concentrated HCl (120 mL) and water (460 mL), carefully concentrated to one third of the original volume and finally extracted with CH₂Cl₂. The organic layer was washed with brine, dried over MgSO₄ and concentrated in vacuo. The crude yellow oil was purified by micro distillation (0.5 torr, bp.: 71°C) to give 32.6 g (90%) benzaldehyde oxime as a colorless oil which slowly crystallized below 20 °C.

All spectroscopic data were in agreement with the literature.

Preparation of *N*-Benzyl-Hydroxylamine³

To a solution of benzaldehyde oxime (24.2 g, 200 mmol) and NaBH₃CN (21.4 g, 340 mmol) in 300 mL methanol at 0 °C, 12 N HCl (33.6 mL, 400 mmol) was added drop wise. After addition the reaction mixture was allowed to stir at 25 °C for 4 h before adding 6 N NaOH until pH ~ 9. The reaction mixture was concentrated in *vacuo* and the product was extracted with CH₂Cl₂, washed with brine, dried over anhydrous MgSO₄ and concentrated in *vacuo*. The crude *N*-benzyl-hydroxylamine was recrystallized from hexane/methanol = 4:1 to afford 21.6 g (87 %) as colorless crystals.

All spectroscopic data were in agreement with the literature.

Preparation of *N*-Benzyl-Nitrones⁴

(*Z*)-*N*-Ethylidene-benzylamine-*N*-oxide

N-benzyl-hydroxylamine (4.92 g, 40.0 mmol, 1.00 equiv), freshly distilled acetaldehyde (8.81 g, 200 mmol, 5.00 equiv), Na₂SO₄ (5.68 g, 40.0 mmol, 1.00 equiv) and a trace of NaHCO₃ were stirred in 100 mL dichloromethane at 0 °C under an argon atmosphere for 1 h. The mixture was filtered and the filtrate concentrated in *vacuo* to afford 5.47 g (92%) (*Z*)-*N*-ethylidene-benzylamine-*N*-oxide as a colorless solid: 30-35 °C.

¹H NMR (300 MHz, CDCl₃) δ 7.37-7.26 (m, 5H), 6.70 (q, *J* = 5.9 Hz, 1H), 4.84 (s, 2H), 1.95 (d, *J* = 7.8 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 134.7, 132.9, 129.4, 128.9, 69.0, 12.7;

FTIR (thin film) ν 3401, 3089, 3032, 1612, 1497, 1456, 1360, 1171, 1101, 1040, 760, 704;

Anal. Calcd. for C₉H₁₁NO (149.19): C, 72.46; H, 7.43; N, 9.39. Found: C, 72.53; H, 7.60; N, 9.21.

(2) Ohemeng, K.A.; Podlogar, B.L.; Nguyen, V. N. et al., *J. Med. Chem.*, **1997**, Vol. 40, No. 20, 3292.

(3) Kawase, M.; Kikugawa, Y. *J. Chem. Soc. Perkin 1*, **1979**, 643.

(4) Dondoni, A.; Franco, S.; Junquera, F.; Merchan, F.; Merino, P.; Tejero, T. *Synthetic Communications* **1994**, 24(18), 2537.

(Z)-N-(2-Methylpropylidene)benzylamine-N-oxide

N-benzyl-hydroxylamine (4.92 g, 40.0 mmol, 1.00 equiv), 2-methyl-propionaldehyde (2.88 g, 40.0 mmol, 1.00 equiv) and MgSO₄ (4.81 g, 40.0 mmol, 1.00 equiv) were stirred in 200 mL ether for 15 h. Recrystallization from ethyl acetate/hexane = 1:4 afforded 5.91 g (84%) (Z)-N-(2-methylpropylidene)benzylamine-*N*-oxide.

All spectroscopic data were in agreement with the literature.

(Z)-N-(1-Cyclohexylmethylidene)benzylamine-N-oxide

N-benzyl-hydroxylamine (504 mg, 4.1 mmol), cyclohexane carboxaldehyde (0.5 mL, 4.1 mmol) and MgSO₄ (497 mg, 4.1 mmol) were stirred in 35 mL ether for 18 h. The reaction mixture was filtered and concentrated in vacuo. Purification using flash chromatography on silica gel (EtOAc/hexane, 19:1) afforded 787 mg (88 %) (Z)-N-(1-cyclohexylmethylidene)benzylamine-*N*-oxide as a white solid: 101-103 °C.

¹H NMR (400 MHz, CDCl₃) δ 7.41-7.33 (m, 5H), 6.47 (d, *J* = 7.4 Hz, 1H), 4.85 (s, 2H), 3.03-2.94 (m, 1H), 1.89-1.83 (m, 2H), 1.71-1.61 (m, 3H), 1.36 (m, 2H), 1.17 (m, 3H);

¹³C NMR (100 MHz, CDCl₃) δ 143.1, 133.2, 129.0, 128.8, 128.7, 69.3, 35.0, 28.7, 25.9, 25.2;

FTIR (KBr): ν 3064, 2919, 2849, 2360, 2341, 1592, 1496, 1446, 1173, 1122, 935, 701 cm⁻¹;

MS (EI): 217.2 ([M]⁺, 1), 220.2 (6), 172.2 (3), 149.2 (3), 146.2 (6), 133.1 (36), 91.1 (100), 65.1 (8);

Anal. Calcd. for C₁₄H₁₉NO: C, 77.38%; H, 8.81%; N, 6.45%. Found: C, 77.47%; H, 8.82%; N, 6.44%.

(Z)-N-(2,2-Dimethylpropylidene)benzylamine-N-oxide

N-benzyl-hydroxylamine (4.92 g, 40.0 mmol, 1.00 equiv), 2,2-dimethyl-propionaldehyde (3.45 g, 40.0 mmol, 1.00 equiv) and MgSO₄ (4.81 g, 40.0 mmol, 1.00 equiv) were stirred in 200 mL ether for 15 h. Recrystallization from heptane/ethanol = 1:1 afforded 6.50 g (85%) (Z)-N-(2,2-dimethylpropylidene)-benzylamine-*N*-oxide.

All spectroscopic data were in agreement with the literature.

(Z)-N-Benzylidene-benzylamine-N-oxide

N-benzyl-hydroxylamine (4.92 g, 40.0 mmol, 1.00 equiv), benzaldehyde (4.25 g, 40.0 mmol, 1.00 equiv) and MgSO₄ (4.82 g, 40.0 mmol, 1.00 equiv) were stirred in 200 mL CH₂Cl₂ for 15 h. Recrystallization from ether afforded 9.53 g (90%) (Z)-N-benzylidene-benzylamine-*N*-oxid.

All spectroscopic data were in agreement with the literature.

Preparation of Propargyl *N*-Hydroxylamines⁵

General Procedure 1 (GP 1)

A flask was charged with $\text{Zn}(\text{OTf})_2$ and purged with argon for 15 min. To the flask was added dichloromethane, alkyne and *Hünig's* base. The resulting mixture was stirred at 23 °C for 20 minutes. The nitron was dissolved in dichloromethane and added to the flask via syringe in one portion. Upon completion, the reaction was quenched by the addition of a saturated aqueous ammonium chloride solution. The reaction mixture was poured into a separatory funnel containing dichloromethane. The layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with brine, dried over sodium sulfate and concentrated in vacuo. Purification of the crude material by chromatography on silica gel afforded the analytically pure propargyl *N*-hydroxylamines.

N-Benzyl-*N*-(1-methyl-hept-2-ynyl)-hydroxylamine

According to GP 1 a solution of 1-hexyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-ethylidene-benzylamine-*N*-oxide (1.49 g, 10.0 mmol, 1.0 equiv), $\text{Zn}(\text{OTf})_2$ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH_2Cl_2 for 1 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.16 g (93%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl_3) δ 7.39-7.24 (m, 5H), 5.32 (s, OH), 4.02 and 3.80 (2d, AB, $J_{\text{AB}} = 12.8$ Hz, 2H), 3.66-3.59 (dq, $J = 6.8$ Hz, 1.9 Hz, 1H), 2.28 (dt, $J = 7.0$, 1.9 Hz, 2H), 1.61-1.42 (m, 4H), 1.37 (d, $J = 6.8$ Hz, 3H), 0.94 (t, $J = 7.3$ Hz, 3H);

¹³C NMR (75 MHz, CDCl_3) δ 137.1, 129.9, 128.7, 128.4, 127.5, 86.6, 61.5, 53.6, 31.0, 22.0, 19.7, 18.5, 13.7;

FTIR (thin film) ν 3238, 3031, 2957, 2934, 2872, 1496, 1455, 1310, 913, 742, 698;

Anal. Calcd. for $\text{C}_{15}\text{H}_{21}\text{NO}$ (231.34): C, 77.88; H, 9.15; N, 6.05. Found: C, 77.90; H, 9.14; N, 6.04.

N-Benzyl-*N*-(1,4,4-trimethyl-pent-2-ynyl)-hydroxylamine

According to GP 1 a solution of 3,3-dimethyl-1-butyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-ethylidene-benzylamine-*N*-oxide (1.49 g, 10.0 mmol, 1.0 equiv), $\text{Zn}(\text{OTf})_2$ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH_2Cl_2 for 2 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.14 g (93%) of the title compound as a colorless solid: mp 94 °C.

¹H NMR (300 MHz, CDCl_3) δ 7.39-7.25 (m, 5H), 5.03 (s, OH), 4.03 and 3.78 (2d, AB, $J_{\text{AB}} = 12.8$ Hz, 2H), 3.61 (q, $J = 6.9$ Hz, 1H), 1.36 (d, $J = 6.9$ Hz, 3H), 1.28 (s, 9H);

¹³C NMR (75 MHz, CDCl_3) δ 137.3, 129.7, 128.4, 127.4, 83.6, 61.8, 53.8, 31.3, 27.5, 19.7;

(5) Frantz, D. E.; Fässler, R.; Carreira, E. M. *J. Am. Chem. Soc.* **1999**, *121*, 11245.

FTIR (thin film) ν 3240, 3032, 2969, 2868, 1496, 1455, 1362, 1309, 1263, 739, 698;

Anal. Calcd. for $C_{15}H_{21}NO$ (231.34): C, 77.88; H, 9.15; N, 6.05. Found: C, 77.75; H, 9.15; N, 6.10.

***N*-benzyl-*N*-(4-phenylbut-3-yn-2-yl)-hydroxylamine**

According to *GP 1* a solution of phenylacetylene (1.53 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-ethylidenebenzylamine-*N*-oxide (1.49 g, 10.0 mmol, 1.0 equiv), $Zn(OTf)_2$ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH_2Cl_2 for 31 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.13 g (85%) of the title compound as a colorless solid: 80-81 °C;

1H NMR (300 MHz, $CDCl_3$) δ 7.54-7.47 (m, 2H), 7.45-7.39 (m, 2H), 5.81-4.89 (br s, 1H), 4.12 (d, J = 12.8 Hz, 1H), 3.92 (d, J = 12.8 Hz, 1H), 3.88 (q, J = 6.9 Hz, 1H), 1.51 (d, J = 7.2 Hz, 3H);

^{13}C NMR (75 MHz, $CDCl_3$) δ 137.1, 131.9, 129.7, 128.4, 128.3, 128.2, 127.5, 122.9, 86.9, 86.2, 61.7, 54.3, 19.4;

FTIR ($CHCl_3$) ν 3446, 3026, 2867, 1636, 1596, 1489, 1456, 1306, 1226, 1108, 1063, 755, 690 cm^{-1} .

MS (EI) 251.2 ($[M]^+$, <1), 233.2 ($[M-H_2O]^+$, 37), 91.0 (100);

Anal. Calcd. for $C_{17}H_{17}NO$: C, 81.24%; H, 6.82%; N, 5.57%. Found: C, 80.99%; H, 6.95%; N, 5.49%.

***N*-Benzyl-*N*-(1-isopropyl-hept-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 1-hexyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-(2-methylpropylidene)benzylamine-*N*-oxide (1.77 g, 10.0 mmol, 1.0 equiv), $Zn(OTf)_2$ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH_2Cl_2 for 2 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.42 g (93%) of the title compound as a colorless solid: mp 48-50 °C.

1H NMR (300 MHz, $CDCl_3$) δ 7.41-7.24 (m, 5H), 4.42 (s, OH), 4.12 and 3.78 (2d, AB, J_{AB} = 13.0 Hz, 2H), 3.14-3.11 (m, 1H), 2.32 (dt, J = 6.9, 2.2 Hz, 2H), 2.04-1.92 (m, 1H), 1.62-1.42 (m, 4H), 1.03 (d, J = 6.9, 3H), 1.01 (d, J = 6.9, 3H), 0.95 (t, J = 7.2 Hz, 3H);

^{13}C NMR (75 MHz, $CDCl_3$) δ 137.7, 129.4, 128.3, 127.3, 88.3, 75.2, 66.0, 62.2, 31.2, 31.0, 22.0, 20.0, 19.9, 18.5, 13.7;

FTIR (thin film) ν 3466, 3031, 2959, 2872, 1496, 1455, 1383, 1328, 1074, 1030, 773, 699;

Anal. Calcd. for $C_{17}H_{25}NO$ (259.39): C, 78.72; H, 9.71; N, 5.40. Found: C, 78.60; H, 9.71; N, 5.40.

***N*-Benzyl-*N*-(1-isopropyl-4,4-dimethyl-pent-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 3,3-dimethyl-1-butyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-(2-methylpropylidene)benzylamine-*N*-oxide (1.77 g, 10.0 mmol, 1.0 equiv), Zn(OTf)₂ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH₂Cl₂ for 2 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.44 g (94%) of the title compound as a colorless solid.

All spectroscopic data were in agreement with the literature.

***N*-Benzyl-*N*-(1-isopropyl-3-phenyl-prop-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of phenyl acetylene (0.77 g, 7.5 mmol, 1.5 equiv), (*Z*)-*N*-(2-methylpropylidene)benzylamine-*N*-oxide (0.89 g, 5.0 mmol, 1.0 equiv), Zn(OTf)₂ (0.18 g, 0.50 mmol, 0.10 equiv) and *Hünig's* base (0.16 g, 1.2 mmol, 0.25 equiv) was stirred in 5.0 mL CH₂Cl₂ for 1 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 1.2 g (90%) of the title compound as a yellowish solid.

All spectroscopic data were in agreement with the literature.

***N*-Benzyl-*N*-(1-*tert*-butyl-hept-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 1-hexyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-(2,2-dimethylpropylidene)benzylamine-*N*-oxide (1.91 g, 10.0 mmol, 1.0 equiv), Zn(OTf)₂ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH₂Cl₂ for 3 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.50 g (91%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.40-7.24 (m, 5H), 4.17 (s, OH), 4.07 and 3.76 (2d, AB, *J*_{AB} = 12.8 Hz, 2H), 3.20 (t, *J* = 2.2 Hz, 1H), 2.34 (dt, *J* = 6.9, 2.2 Hz, 2H), 1.64-1.45 (m, 4H), 0.99 (s, 9H), 0.96 (t, *J* = 7.2 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 137.5, 129.3, 128.3, 127.4, 90.0, 73.0, 68.3, 64.7, 35.3, 31.2, 27.8, 22.0, 18.5, 13.6;

FTIR (thin film) ν 3467, 3122, 2956, 2867, 1456, 1393, 1364, 1298, 1191, 913;

Anal. Calcd. for C₁₈H₂₇NO (273.42): C, 79.07; H, 9.95; N, 5.12. Found: C, 79.02; H, 10.03; N, 5.12.

***N*-Benzyl-*N*-(1-*tert*-butyl-4,4-dimethyl-pent-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 3,3-dimethyl-1-butyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-(2,2-dimethylpropylidene)benzylamine-*N*-oxide (1.91 g, 10.0 mmol, 1.0 equiv), Zn(OTf)₂ (0.364 g, 1.00 mmol, 0.1 equiv) and *Hünig's* base (0.323 g, 2.50 mmol, 0.25 equiv) was stirred in 10 mL CH₂Cl₂ for 3 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.61 g (95%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.40-7.26 (m, 5H), 4.10 (s, OH), 4.06 and 3.74 (2d, AB, *J*_{AB} = 12.8 Hz, 2H), 3.18 (s, 1H), 1.32 (s, 9H), 0.98 (s, 9H);

¹³C NMR (75 MHz, CDCl₃) δ 137.4, 129.3, 128.4, 127.4, 99.4, 70.9, 68.0, 64.8, 35.2, 31.4, 27.8, 27.5;

FTIR (thin film) ν 3467, 3064, 3031, 2967, 2903, 2868, 1477, 1455, 1393, 1363, 1263, 913, 748, 698;

Anal. Calcd. for C₁₈H₂₇NO (273.42): C, 79.07; H, 9.95; N, 5.12. Found: C, 78.88; H, 9.90; N, 5.19.

***N*-Benzyl-*N*-(1-cyclohexyl-3-phenylprop-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of phenylacetylene (0.58 ml, 5.2 mmol, 1.5 equiv), (*Z*)-*N*-(1-cyclohexylmethylidene)benzylamine-*N*-oxide (0.75 g, 3.5 mmol, 1.0 equiv), Zn(OTf)₂ (0.13 g, 1.0 mmol, 0.1 equiv) and *Hünig's* base (0.15 ml, 0.87 mmol, 0.25 equiv) was stirred in 8 mL CH₂Cl₂ for 6 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 3:1) afforded 1.1 g (95%) of the title compound as a colorless solid: mp 126-128 °C.

¹H NMR (300 MHz, CDCl₃) δ 7.26-7.53 (m, 10H), 4.45 (s, 1H), 4.21 (d, 1H, *J* = 13.4), 3.90 (d, 1H, *J* = 13.1), 3.47 (d, 1H, *J* = 9.0), 0.92-2.18 (m, 11H);

¹³C NMR (75 MHz, CDCl₃) δ 137.4, 131.8, 129.2, 128.2, 128.2, 128.2, 128.1, 127.2, 122.9, 88.0, 85.0, 65.1, 62.3, 40.1, 30.6, 26.7, 26.2, 26.0;

FTIR (thin film) ν 3536, 3061, 3030, 2925, 2850, 1950, 1876, 1803, 1598, 1490, 1450, 756, 691;

Anal. Calcd. for C₂₂H₂₅NO (319.44): C, 82.72; H, 7.89; N, 4.38. Found: C, 82.75; H, 7.67; N, 4.31.

***N*-Benzyl-*N*-(1-phenyl-hept-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 1-hexyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-benzylidene-benzylamine-*N*-oxide (2.11 g, 10.0 mmol, 1.0 equiv), Zn(OTf)₂ (3.64 g, 10.0 mmol, 1.0 equiv) and *Hünig's* base (1.94 g, 15.0 mmol, 1.5 equiv) was stirred in 10 mL CH₂Cl₂ for 4 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.56 g (87%) of the title compound as a colorless solid: mp 80-81 °C.

¹H NMR (300 MHz, CDCl₃) δ 7.57-7.23 (m, 10H), 5.25 (s, OH), 4.71 (s, 1H), 3.89 (s, 2H), 2.40 (dt, *J* = 2.2, 7.2 Hz, 2H), 1.68-1.45 (m, 4H), 0.97 (t, *J* = 7.2 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 138.0, 137.3, 129.7, 128.9, 128.4, 128.2, 127.9, 127.5, 89.5, 74.7, 62.5, 31.1, 22.1, 18.7, 13.7;

FTIR (thin film) ν 3234, 3031, 2931, 1493, 1451, 1332, 1291, 1073, 1026, 912, 735, 640;

Anal. Calcd. for C₂₀H₂₃NO (293.41): C, 81.87; H, 7.90; N, 4.77. Found: C, 81.75; H, 7.85; N, 4.75.

***N*-Benzyl-*N*-(4,4-dimethyl-1-phenyl-pent-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of 3,3-dimethyl-1-butyne (1.23 g, 15.0 mmol, 1.5 equiv), (*Z*)-*N*-benzylidene-benzylamine-*N*-oxide (2.11 g, 10.0 mmol, 1.0 equiv), Zn(OTf)₂ (3.64 g, 10.0 mmol, 1.0 equiv) and *Hünig's* base (1.94 g, 15.0 mmol, 1.5 equiv) was stirred in 10 mL CH₂Cl₂ for 5 h at 23 °C. Purification using chromatography on silica gel (hexane/ethyl acetate, 4:1) afforded 2.49 g (85%) of the title compound as a colorless solid: mp 130-132 °C.

¹H NMR (300 MHz, CDCl₃) δ 7.58-7.25 (m, 10H), 4.78 (s, 1H), 4.73 (s, OH), 3.98 and 3.88 (2d, AB, *J*_{AB} = 13.1 Hz, 2H), 1.37 (s, 9H);

¹³C NMR (75 MHz, CDCl₃) δ 138.2, 137.5, 129.5, 128.7, 128.4, 128.2, 127.8, 127.4, 98.4, 76.6, 62.7, 31.3, 27.8;

FTIR (thin film) ν 3240, 3064, 3031, 2968, 2867, 1495, 1454, 1362, 1270, 1030, 913, 743, 698;

Anal. Calcd. for C₂₀H₂₃NO (293.41): C, 81.78; H, 7.90; N, 4.77. Found: C, 81.92; H, 7.77; N, 4.85.

***N*-Benzyl-*N*-(1,3-diphenyl-prop-2-ynyl)-hydroxylamine**

According to *GP 1* a solution of phenylacetylene (0.77 g, 7.5 mmol, 1.5 equiv), (*Z*)-*N*-benzylidene-benzylamine-*N*-oxide (1.06 g, 5.00 mmol, 1.0 equiv), Zn(OTf)₂ (1.82 g, 5.00 mmol, 1.0 equiv) and *Hünig's* base (0.65 g, 5.0 mmol, 1.0 equiv) was stirred in 5.0 mL CH₂Cl₂ for 18 h at 23 °C. Purification using chromatography on silica gel (hexane/methylene chloride, 1:1) afforded 1.4 g (92%) of the title compound as a white solid: mp 134-136°C.

¹H NMR (300 MHz, CDCl₃) δ 7.64-7.26 (m, 15H), 5.80 (s, OH), 4.88 (s, 1H), 4.01 and 3.90 (2d, AB, *J*_{AB} = 12.9 Hz, 2H);

¹³C NMR (75 MHz, CDCl₃) δ 137.6, 137.3, 132.0, 129.7, 129.0, 128.5, 128.4, 128.4, 128.1, 127.5, 122.9, 88.8, 84.7, 63.2, 60.6;

FTIR (thin film) ν 3529, 3234, 3063, 3031, 2907, 1953, 1883, 1811, 1598, 1490, 1454, 1336, 1219, 1028, 756, 698;

Anal. Calcd. for C₂₂H₁₉NO (313.40): C, 84.32; H, 6.11; N, 4.47. Found: C, 84.35; H, 6.16; N, 4.53.

Preparation of 2,3-Dihydroisoxazoles 1-8⁶

General Procedure 2 (GP 2)

A flask was charged with ZnI₂ (0.10 equiv) and purged with nitrogen for 15 min. To the flask was added a solution of DMAP (0.10 equiv) in dichloromethane. The resulting mixture was stirred at 23 °C for 10 minutes. The Propargyl *N*-hydroxylamine (1.0 equiv) was dissolved in dichloromethane and added to the flask in one portion. Upon completion, the reaction was quenched by the addition of saturated aqueous ammonium chloride solution. The reaction mixture was poured into a separatory funnel containing CH₂Cl₂. The layers were separated and the aqueous layer was extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over sodium sulfate and concentrated in vacuo. Purification of the crude material by chromatography on silica gel afforded the analytically pure 2,3-dihydroisoxazoles.

2-Benzyl-5-butyl-3-methyl-2,3-dihydroisoxazole (1)

According to GP 2 a solution of *N*-benzyl-*N*-(1-methyl-hept-2-ynyl)-hydroxylamine (463 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 3 h at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 379 mg (82%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.42-7.25 (m, 5H), 4.52 (d, *J* = 2.2 Hz, 1H), 4.18 and 3.83 (2d, AB, *J*_{AB} = 12.9 Hz, 2H), 3.86 (dq, *J* = 6.2, 2.2 Hz, 1H), 2.15 (t, *J* = 7.2 Hz, 2H), 1.56-1.42 (m, 3H), 1.19 (d, *J* = 6.2 Hz, 3H), 0.93 (t, *J* = 7.2 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 154.6, 137.0, 129.3, 128.5, 128.3, 127.3, 96.2, 65.4, 62.9, 29.0, 25.7, 22.3, 19.8;

FTIR (thin film) ν 3064, 3030, 2960, 2930, 2872, 1712, 1674, 1454, 1367, 1075, 912, 736;

Anal. Calcd. for C₁₅H₂₁NO (231.34): C, 77.88; H, 9.15; N, 6.05. Found: C, 77.92; H, 9.02; N, 6.05.

2-Benzyl-5-*tert*-butyl-3-methyl-2,3-dihydroisoxazole (2)

According to GP 2 a solution of *N*-benzyl-*N*-(1,4,4-trimethyl-pent-2-ynyl)-hydroxylamine (463 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 11 h at 50 °C in a pressure tube. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 391 mg (84%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.42-7.25 (m, 5H), 4.49 (d, *J* = 2.5 Hz, 1H), 4.15 and 3.78 (2d, AB, *J*_{AB} = 12.6 Hz, 2H), 3.81 (dq, *J* = 6.2, 2.5 Hz, 1H), 1.16 (s, 9H), 1.11 (d, *J* = 6.2 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 162.5, 137.1, 129.5, 128.3, 127.3, 93.6, 65.1, 62.6, 31.1, 28.3, 22.7;

FTIR (thin film) ν 3064, 3031, 2967, 2924, 2870, 1660, 1496, 1456, 1364, 1264, 1201, 1109, 912, 734;

(6) Aschwanden, P.; Frantz, D. E.; Carreira, E. M. *Org. Lett.* **2000**, 2, 2331

Anal. Calcd. for C₁₅H₂₁NO (231.34): C, 77.88; H, 9.15; N, 6.05. Found: C, 77.79; H, 9.27; N, 6.09.

2-Benzyl-5-butyl-3-isopropyl-2,3-dihydroisoxazole (3)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-isopropyl-hept-2-ynyl)-hydroxylamine (519 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 2 h at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 485 mg (93%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.41-7.24 (m, 5H), 4.50 (d, *J* = 2.5 Hz, 1H), 4.13 and 3.76 (2d, AB, *J*_{AB} = 12.7 Hz, 2H), 3.48-3.44 (m, 1H), 2.15 (t, *J* = 7.2 Hz, 2H), 1.59-1.30 (m, 5H), 0.91 (t, *J* = 7.3 Hz, 3H), 0.79 (d, *J* = 7.2 Hz, 6H);

¹³C NMR (75 MHz, CDCl₃) δ 155.1, 137.2, 129.7, 128.2, 127.3, 92.3, 76.1, 64.0, 33.8, 29.1, 25.7, 22.3, 18.6, 18.2, 13.8;

FTIR (thin film) ν 3031, 2957, 2871, 1676, 1496, 1455, 1382, 1306, 1159, 1081, 946, 732, 697;

Anal. Calcd. for C₁₇H₂₅NO (259.39): C, 78.72; H, 9.71; N, 5.40. Found: C, 78.77; H, 9.78; N, 5.39.

2-Benzyl-5-*tert*-butyl-3-isopropyl-2,3-dihydroisoxazole (4)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-isopropyl-4,4-dimethyl-pent-2-ynyl)-hydroxylamine (519 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 14 h at 50 °C in a pressure tube. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 449 mg (87%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.41-7.24 (m, 5H), 4.47 (d, *J* = 2.5 Hz, 1H), 4.11 and 3.71 (2d, AB, *J*_{AB} = 12.4 Hz, 2H), 3.41 (dd, *J* = 6.5, 2.5 Hz, 1H), 1.58-1.49 (m, 1H), 1.15 (s, 9H), 0.80 (d, *J* = 6.5 Hz, 3H), 0.78 (d, *J* = 6.5 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 163.0, 137.1, 129.9, 128.2, 127.3, 89.7, 75.7, 63.7, 33.7, 31.2, 28.4, 18.5, 18.2;

FTIR (thin film) ν 3033, 2968, 2906, 2870, 1664, 1460, 1365, 1202, 1104, 921, 755, 651;

Anal. Calcd. for C₁₇H₂₅NO (259.39): C, 78.72; H, 9.71; N, 5.40. Found: C, 78.73; H, 9.80; N, 5.40.

2-Benzyl-5-butyl-3-*tert*-butyl-2,3-dihydroisoxazole (5)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-*tert*-butyl-hept-2-ynyl)-hydroxylamine (547 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 2 h at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 521 mg (95%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.42-7.24 (m, 5H), 4.46 (d, *J* = 2.5 Hz, 1H), 4.09 and 3.74 (2d, AB, *J*_{AB} = 12.8 Hz, 2H), 3.43 (d, *J* = 2.5 Hz, 1H), 2.17-2.12 (m, 2H), 1.55-1.33 (m, 4H), 0.91 (t, *J* = 7.2 Hz, 3H), 0.77 (s, 9H);

¹³C NMR (75 MHz, CDCl₃) δ 155.4, 137.4, 129.7, 128.2, 127.3, 90.9, 79.5, 64.8, 35.0, 29.2, 25.7, 25.5, 22.3, 13.8;

FTIR (thin film) ν 3031, 2955, 2865, 1676, 1456, 1390, 1361, 1260, 1182, 1071, 913;

Anal. Calcd. for C₁₈H₂₇NO (273.42): C, 79.07; H, 9.95; N, 5.12. Found: C, 79.02; H, 9.87; N, 5.06.

2-Benzyl-3,5-di-*tert*-butyl-2,3-dihydroisoxazole (6)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-*tert*-butyl-4,4-dimethyl-pent-2-ynyl)-hydroxylamine (547 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 13 h at 50 °C in a pressure tube. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 490 mg (90%) of the title compound as a colorless solid:

mp 44-45 °C.

¹H NMR (300 MHz, CDCl₃) δ 7.41-7.26 (m, 5H), 4.43 (d, *J* = 2.5 Hz, 1H), 4.08 and 3.68 (2d, AB, *J*_{AB} = 12.5 Hz, 2H), 3.38 (d, *J* = 2.5 Hz, 1H), 1.15 (s, 9H), 0.75 (s, 9H);

¹³C NMR (75 MHz, CDCl₃) δ 163.3, 137.1, 130.1, 128.1, 127.3, 88.2, 79.0, 64.4, 34.9, 31.3, 28.4, 25.4;

FTIR (thin film) ν 3031, 2959, 2904, 2867, 1663, 1477, 1459, 1361, 1204, 1109, 923, 722, 697;

Anal. Calcd. for C₁₈H₂₇NO (273.42): C, 79.07; H, 9.95; N, 5.12. Found: C, 79.02; H, 9.94; N, 5.28.

2-Benzyl-5-butyl-3-phenyl-2,3-dihydroisoxazole (7)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-phenyl-hept-2-ynyl)-hydroxylamine (587 mg, 2.00 mmol, 1.0 equiv), ZnI₂ (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 1 h at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded

561 mg (96%) of the title compound as a yellowish oil.

¹H NMR (300 MHz, CDCl₃) δ 7.43-7.18 (m, 10H), 4.87 (d, *J* = 1.1 Hz, 1H), 4.69 (d, *J* = 1.1 Hz, 1H), 4.33 and 4.04 (2d, AB, *J*_{AB} = 12.8 Hz, 2H), 2.27-2.22 (m, 2H), 1.62-1.35 (m, 6H), 0.95 (t, *J* = 7.2, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 155.7, 142.9, 136.7, 129.5, 128.4, 128.3, 127.5, 127.3, 127.0, 94.8, 73.1, 63.5, 29.0, 25.7, 22.4, 13.9;

FTIR (thin film) ν 3063, 3029, 2957, 2931, 2872, 1674, 1603, 1495, 1454, 1309, 1157, 1074, 913, 733;

Anal. Calcd. for C₂₀H₂₃NO (293.41): C, 81.87; H, 7.90; N, 4.77. Found: C, 81.73; H, 8.03; N, 4.72.

2-Benzyl-5-*tert*-butyl-3-phenyl-2,3-dihydroisoxazole (8)

According to *GP 2* a solution of *N*-benzyl-*N*-(1,1-dimethyl-1-phenyl-pent-2-ynyl)-hydroxylamine (587 mg, 2.00 mmol, 1.0 equiv), ZnI_2 (63.8 mg, 0.200 mmol, 0.1 equiv), and DMAP (24.4 mg, 0.200 mmol, 0.1 equiv) in dichloromethane (4.00 mL) was stirred for 3 h at 50 °C in a pressure tube. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 95:5) afforded 542 mg (92%) of the title compound as a yellowish solid: mp 52-54 °C.

^1H NMR (300 MHz, CDCl_3) δ 7.43-7.15 (m, 10H), 4.81 (d, $J = 2.5$ Hz, 1H), 4.66 (d, $J = 2.5$ Hz, 1H), 4.32 and 3.99 (2d, AB, $J_{\text{AB}} = 12.5$ Hz, 2H), 1.23 (s, 9H);

^{13}C NMR (75 MHz, CDCl_3) δ 163.5, 143.0, 136.7, 129.7, 128.4, 128.3, 127.5, 127.3, 127.0, 92.2, 72.8, 63.4, 31.4, 28.4;

FTIR (thin film) ν 3063, 3029, 2966, 2931, 2870, 1661, 1494, 1455, 1363, 1201, 1102, 1029, 758, 697;

Anal. Calcd. for $\text{C}_{20}\text{H}_{23}\text{NO}$ (293.41): C, 81.87; H, 7.90; N, 4.77. Found: C, 81.98; H, 7.95; N, 4.84.

2-benzyl-3-methyl-5-phenyl-2,3-dihydroisoxazole (17a)

According to *GP 2* a solution of *N*-benzyl-*N*-(4-phenylbut-3-yn-2-yl)-hydroxylamine (0.55 g, 2.2 mmol, 1.0 equiv), ZnI_2 (0.21 g, 0.67 mmol, 0.30 equiv), and DMAP (82 mg, 0.67 mmol, 0.30 equiv) in dichloromethane (3.5 mL) was stirred for 2 h at 23 °C. Workup and purification using chromatography on silica gel (dichloromethane/hexane, 1:1 to 9:1) afforded 0.33 g (60%) of the title compound as a yellowish oil.

^1H NMR (300 MHz, CDCl_3) δ 7.27-7.54 (m, 10H), 5.29 (d, 1H, $J = 2.8$), 4.29 (d, 1H, $J = 12.8$), 4.07 (dq, 1H, $J = 2.5$, $J = 6.2$), 3.91 (d, 1H, $J = 12.8$), 1.23 (d, 3H, $J = 6.2$);

^{13}C NMR (75 MHz, CDCl_3) δ 152.3, 137.1, 129.7, 129.3, 129.0, 128.6, 128.6, 127.7, 125.8, 97.6, 66.3, 63.1, 22.5;

FTIR (thin film) ν 3028, 2969, 2922, 2861, 2346, 1945, 1885, 1811, 1652, 1600, 1578, 1494, 1448, 736, 696;

Anal. Calcd. for $\text{C}_{17}\text{H}_{17}\text{NO}$ (251.13): C, 81.24; H, 6.82; N, 5.57. Found: C, 81.25; H, 6.90; N, 5.51;

2-Benzyl-3-isopropyl-5-phenyl-2,3-dihydroisoxazole (17b)

According to *GP 2* a solution of *N*-benzyl-*N*-(1-isopropyl-3-phenyl-prop-2-ynyl)-hydroxylamine (140 mg, 0.500 mmol, 1.0 equiv), ZnI_2 (16 mg, 0.050 mmol, 10 mol%), and DMAP (6 mg, 0.05 mmol, 10 mol%) in dichloromethane (1.00 mL) was stirred for 3 h at 23 °C. Workup and purification using chromatography on silica gel (methylene chloride/hexane, 6:4) afforded 132 mg (94%) of the title compound as a colorless oil.

^1H NMR (300 MHz, CDCl_3) δ 7.57-7.26 (m, 10H), 5.31 (d, 1 Vinyl-H, $J = 2.8$), 4.28 and 3.87 (2d, AB, $J_{\text{AB}} = 12.5$ Hz, 2H), 3.69 (dd, $J = 2.8$, 6.5 Hz, 1H), 1.76-1.64 (m, 1H), 3.69 (dd, $J = 1.4$, 6.7 Hz, 6H).

^{13}C NMR (75 MHz, CDCl_3) δ 152.5, 136.8, 129.9, 129.2, 128.8, 128.4, 128.3, 127.5, 125.6, 93.7, 76.6, 64.0, 33.8, 18.7, 18.4;

FTIR (thin film) ν 3028, 2956, 2868, 1947, 1875, 1806, 1676, 1604, 1496, 1454, 1156, 1074, 1030, 941, 733, 698;

Anal. Calcd. for $C_{19}H_{21}NO$ (279.38): C, 81.68; H, 7.58; N, 5.01. Found: C, 81.60; H, 7.65; N, 4.95.

2-Benzyl-3-cyclohexyl-5-phenyl-2,3-dihydroisoxazole (17c)

According to *GP 2* a solution of *N*-Benzyl-*N*-(1-cyclohexyl-3-phenylprop-2-ynyl)-hydroxylamine (935 mg, 2.90 mmol, 1.0 equiv), ZnI_2 (280 mg, 0.877 mmol, 0.30 equiv), and DMAP (108 mg, 0.884 mmol, 0.30 equiv) in dichloromethane (10.0 mL) was stirred for 2 h at 23 °C in a pressure tube. Workup and purification using chromatography on silica gel (dichloromethane/hexane, 1:1 to 1:0) afforded 813 mg (87%) of the title compound as a yellowish solid: mp 88-91 °C.

1H NMR (300 MHz, $CDCl_3$) δ 7.29-7.56 (m, 10H), 5.32 (d, 1H, $J = 3.0$), 4.25 (d, 1H, $J = 12.6$), 3.84 (d, 1H, $J = 12.6$), 3.67 (dd, 1H, $J = 2.8$, $J = 7.1$), 1.89 (d, 1H, $J = 12.9$), 1.66 (d, 4H, $J = 9.6$), 1.41 (m, 1H), 0.74-1.28 (m, 5H);

^{13}C NMR (75 MHz, $CDCl_3$) δ 152.2, 136.8, 129.8, 129.2, 128.7, 128.3, 128.2, 127.4, 125.6, 93.9, 75.6, 63.8, 43.5, 29.3, 29.0, 26.5, 26.1, 25.9;

FTIR (thin film) ν 3050, 3028, 2923, 2850, 1950, 1881, 1806, 1650, 1600, 1574, 1494, 1448, 1021, 726, 695;

Anal. Calcd. for $C_{22}H_{25}NO$ (319.44): C, 82.72; H, 7.89; N, 4.38. Found: C, 82.57; H, 7.90; N, 4.34.

2-Benzyl-3,5-diphenyl-2,3-dihydroisoxazole (17d)

According to *GP 2* a solution of *N*-benzyl-*N*-(1,3-diphenyl-prop-2-ynyl)-hydroxylamine (157 mg, 0.500 mmol, 1.0 equiv), ZnI_2 (16 mg, 0.050 mmol, 10 mol%), and DMAP (6 mg, 0.05 mmol, 10 mol%) in dichloromethane (1.00 mL) was stirred for 4 h at 23 °C. Workup and purification using chromatography on silica gel (methylene chloride/hexane, 6:4) afforded 143 mg (91%) of the title compound as a white solid: mp 102-104 °C.

1H NMR (300 MHz, $CDCl_3$) δ 7.60-7.23 (m, 15H), 5.44 (d, $J = 2.8$, 1 Vinyl-H), 5.07 (d, $J = 2.8$ Hz, 1H), 4.46 and 4.13 (2d, AB, $J_{AB} = 12.8$ Hz, 2H);

^{13}C NMR (75 MHz, $CDCl_3$) δ 153.2, 142.4, 136.7, 129.9, 129.3, 129.0, 128.8, 128.7, 128.6, 127.8, 127.4, 126.0, 95.9, 73.8, 63.6;

FTIR (thin film) ν 3062, 3028, 2888, 1953, 1884, 1807, 1651, 1600, 1494, 1454, 1320, 1271, 1220, 1022, 915, 770, 727, 698;

Anal. Calcd. for $C_{22}H_{19}NO$ (313.40): C, 84.32; H, 6.11; N, 4.47. Found: C, 84.48; H, 6.29; N, 4.43.

Preparation of β -Amino Ketones 9-16

General Procedure 3 (GP 3)

To a vigorously stirred mixture of activated zinc dust⁷ (5.0 equiv) in acetic acid/water (~3:1) was added a solution of 2,3-dihydro-isoxazole (1.0 equiv) in dichloromethane (0.50 mL). After 5 minutes of stirring another portion of zinc dust (5.0 equiv) was added to the reaction mixture. Upon completion, the reaction mixture was diluted with dichloromethane and poured into a separatory funnel containing saturated NaHCO₃ solution (20 mL). The layers were separated and the aqueous layer was extracted with dichloromethane (3 x 10 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated in *vacuo*. Purification of the crude material by flash chromatography on silica gel afforded the analytically pure β -Amino ketones.

2-Benzylamino-octan-4-one (9)

According to GP 3 a solution of 2-benzyl-5-butyl-3-methyl-2,3-dihydro-isoxazole (**1**) (231 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ L CH₂Cl₂ was stirred for 15 min at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 1:9) afforded 173-190 mg (74-81%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.33-7.19 (m, 5H), 3.82 and 3.70 (2d, AB, J_{AB} = 13.1 Hz, 2H), 3.21-3.13 (m, 1H), 2.57 (dd, J = 16.5, 7.2 Hz, 1H), 2.45 (dd, J = 16.5, 5.6 Hz, 1H), 2.37 (t, J = 7.5 Hz, 2H), 1.79 (s, NH), 1.58-1.48 (m, 2H), 1.35-1.23 (m, 2H), 1.11 (d, J = 6.2 Hz, 3H), 0.89 (t, J = 7.5 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 210.9, 140.4, 128.4, 128.1, 126.9, 51.3, 49.9, 48.9, 43.3, 25.8, 22.3, 20.4, 13.9;

FTIR (thin film) ν 3326, 3063, 3028, 2959, 2932, 2873, 1709, 1631, 1453, 1376, 1135, 1029, 734;

Anal. Calcd. for C₁₅H₂₃NO (233.35): C, 77.21; H, 9.93; N, 6.00. Found: C, 77.15; H, 10.01; N, 6.05.

5-Benzylamino-2,2-dimethyl-hexan-3-one (10)

According to GP 3 a solution of 2-benzyl-5-*tert*-butyl-3-methyl-2,3-dihydro-isoxazole (**2**) (231 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ L CH₂Cl₂ was stirred for 15 min at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 1:9) afforded 180-192 mg (77-82%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.33-7.21 (m, 5H), 3.83 and 3.71 (2d, AB, J_{AB} = 13.1 Hz, 2H), 3.24-3.17 (m, 1H), 2.66 (dd, J = 17.7, 6.9 Hz, 1H), 2.58 (dd, J = 17.7, 5.9 Hz, 1H), 1.92 (s, NH), 1.12 (s, 9H), 1.11 (t, J = 7.2 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 215.7, 140.3, 128.5, 128.2, 126.9, 51.5, 48.9, 44.3, 43.8, 26.3, 20.3;

FTIR (thin film) ν 3331, 3028, 2968, 2871, 1702, 1495, 1478, 1367, 1153, 1060, 1004, 734, 698;

(7) Activation was achieved by stirring zinc powder in 5% HCl for 5 minutes. Subsequent washing with water, ethanol, acetone and diethylether afforded the activated zinc dust as a pale grey powder which was dried in *vacuo*. See also: Yamamura, S.; Hirata, Y. *J. Chem. Soc. C* **1968**, 2887.

Anal. Calcd. for C₁₅H₂₃NO (233.35): C, 77.21; H, 9.93; N, 6.00. Found: C, 77.18; H, 9.86; N, 5.98;

3-Benzylamino-2-methyl-nonan-5-one (11)

According to *GP 3* a solution of 2-benzyl-5-butyl-3-isopropyl-2,3-dihydro-isoxazole (**3**) (259 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH₂Cl₂ was stirred for 30 min at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 224-238 mg (86-91%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.33-7.21 (m, 5H), 3.77 and 3.69 (2d, AB, J_{AB} = 12.8 Hz, 2H), 3.00-2.94 (m, 1H), 2.48 (dd, J = 15.9, 4.4 Hz, 1H), 2.40 (t, 2H, J = 7.1 Hz), 2.38 (dd, J = 15.9, 8.7 Hz, 1H), 1.96-1.87 (m, 1H), 1.59-1.51 (m, 2H), 1.49 (s, NH), 1.36-1.23 (m, 2H), 0.92-0.87 (m, 9H);

¹³C NMR (75 MHz, CDCl₃) δ 211.5, 140.8, 128.3, 128.2, 126.9, 58.7, 51.6, 43.9, 43.2, 29.7, 25.8, 22.4, 19.0, 17.2, 13.9;

FTIR (thin film) ν 3335, 3028, 2958, 2872, 1710, 1465, 1367, 1267, 1108, 1029, 982, 735, 699;

Anal. Calcd. for C₁₇H₂₇NO (261.41): C, 78.11; H, 10.41; N, 5.36. Found: C, 77.99; H, 10.32; N, 5.32.

5-Benzylamino-2,2,6-trimethyl-heptan-3-one (12)

According to *GP 3* a solution of 2-benzyl-5-*tert*-butyl-3-isopropyl-2,3-dihydro-isoxazole (**4**) (259 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH₂Cl₂ was stirred for 30 min at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 229-236 mg (88-90%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.34-7.20 (m, 5H), 3.76 and 3.70 (2d, AB, J_{AB} = 12.8 Hz, 2H), 3.03-2.97 (m, 1H), 2.60 (dd, J = 17.2, 4.0 Hz, 1H), 2.47 (dd, J = 17.2, 8.1 Hz, 1H), 1.94-1.88 (m, 1H), 1.50 (s, NH), 1.14 (s, 9H), 0.91 (d, J = 6.8 Hz, 3H), 0.89 (d, J = 6.8 Hz, 3H);

¹³C NMR (75 MHz, CDCl₃) δ 216.0, 140.9, 128.3, 128.2, 126.8, 58.4, 52.0, 44.6, 37.6, 29.8, 26.5, 18.8, 17.4;

FTIR (thin film) ν 3335, 3063, 3028, 2964, 2871, 1703, 1624, 1478, 1366, 1301, 1075, 984, 736, 699;

Anal. Calcd. for C₁₇H₂₇NO (261.41): C, 78.11; H, 10.41; N, 5.36. Found: C, 78.24; H, 10.32; N, 5.43.

3-Benzylamino-2,2-dimethyl-nonan-5-one (13)

According to *GP 3* a solution of 2-benzyl-5-butyl-3-*tert*-butyl-2,3-dihydro-isoxazole (**5**) (273 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH_2Cl_2 was stirred for 30 min at 23 $^\circ\text{C}$. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 250-264 mg (91-96%) of the title compound as a colorless oil.

^1H NMR (300 MHz, CDCl_3) δ 7.33-7.21 (m, 5H), 3.77 and 3.70 (2d, AB, J_{AB} = 12.8 Hz, 2H), 2.89 (dd, J = 7.8, 3.7 Hz, 1H), 2.63 (dd, J = 15.6, 3.7 Hz, 1H), 2.45 (t, 2H, J = 7.5), 2.37 (dd, J = 15.6, 7.8 Hz, 1H), 1.92-1.51 (m, 2H), 1.38-1.26 (m, 2H + NH), 0.91 (t, J = 7.5 Hz, 3H), 0.91 (s, 9H);

^{13}C NMR (75 MHz, CDCl_3) δ 211.3, 141.1, 128.3, 128.2, 126.9, 62.7, 54.3, 45.6, 43.5, 35.3, 26.7, 25.9, 22.4, 13.9;

FTIR (thin film) ν 3349, 3032, 2958, 2871, 1711, 1604, 1495, 1467, 1365, 1286, 1204, 1116, 1029, 913;

Anal. Calcd. for $\text{C}_{18}\text{H}_{29}\text{NO}$ (275.43): C, 78.49; H, 10.61; N, 5.09. Found: C, 78.41; H, 10.55; N, 4.96.

5-Benzylamino-2,2,6,6-tetramethyl-heptan-3-one (14)

According to *GP 3* a solution of 2-benzyl-3,5-di-*tert*-butyl-2,3-dihydro-isoxazole (**6**) (273 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH_2Cl_2 was stirred for 30 min at 23 $^\circ\text{C}$. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 242-253 mg (88-92%) of the title compound as a colorless oil.

^1H NMR (300 MHz, CDCl_3) δ 7.33-7.21 (m, 5H), 3.77 and 3.68 (2d, AB, J_{AB} = 12.8 Hz, 2H), 2.97 (dd, J = 7.8, 3.7 Hz, 1H), 2.69 (dd, J = 17.1, 3.7 Hz, 1H), 2.50 (dd, J = 17.1, 7.8 Hz, 1H), 1.17 (s, 9H), 0.92 (s, 9H);

^{13}C NMR (75 MHz, CDCl_3) δ 215.4, 141.3, 128.4, 128.2, 126.8, 61.9, 54.6, 44.5, 39.4, 35.3, 26.8, 26.6;

FTIR (thin film) ν 3351, 3063, 3029, 2962, 2907, 2869, 1705, 1624, 1478, 1365, 1296, 1220, 1114, 1076, 736, 699;

Anal. Calcd. for $\text{C}_{18}\text{H}_{29}\text{NO}$ (275.43): C, 78.49; H, 10.61; N, 5.09. Found: C, 78.60; H, 10.72; N, 5.00.

1-Benzylamino-1-phenyl-heptan-3-one (15)

According to *GP 3* a solution of 2-Benzyl-5-butyl-3-phenyl-2,3-dihydro-isoxazole (**7**) (293 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH_2Cl_2 was stirred for 30 min at 23 $^\circ\text{C}$. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 175-215 mg (59-73%) of the title compound as a colorless oil.

^1H NMR (300 MHz, CDCl_3) δ 7.38-7.23 (m, 10H), 4.15 (dd, J = 8.6, 5.0 Hz, 1H), 3.62 and 3.53 (2d, AB, J_{AB} = 13.1 Hz, 2H), 2.81 (dd, J = 16.7, 8.6 Hz, 1H), 2.72 (dd, J = 16.7, 5.0 Hz, 1H), 2.35-2.29 (m, 2H), 2.11 (s, NH), 1.55-1.45 (m, 2H), 1.29-1.22 (m, 1H), 0.87 (t, J = 7.5, 3H);

^{13}C NMR (75 MHz, CDCl_3) δ 210.17, 143.02, 140.36, 128.65, 128.37, 128.23, 127.40, 127.24, 126.90, 58.3, 51.5,

51.2, 43.1, 25.7, 22.3, 13.9;

FTIR (thin film) ν 3335, 3062, 2970, 2905, 1702, 1605, 1477, 1394, 1201, 1119, 1028, 992, 745;

Anal. Calcd. for $C_{20}H_{25}NO$ (295.42): C, 81.31; H, 8.53; N, 4.74. Found: C, 81.35; H, 8.56; N, 4.65.

1-Benzylamino-4,4-dimethyl-1-phenyl-pentan-3-one (16)

According to *GP 3* a solution of 2-Benzyl-5-*tert*-butyl-3-phenyl-2,3-dihydro-isoxazole (**8**) (293 mg, 1.00 mmol, 1.0 equiv), zinc dust (654 mg, 10.0 mmol, 10 equiv), acetic acid (1.20 g, 20.0 mmol, 20 equiv), water (360 mg, 20.0 mmol, 20 equiv) and 500 μ l CH_2Cl_2 was stirred for 30 min at 23 °C. Workup and purification using chromatography on silica gel (hexane/ethyl acetate, 7:3) afforded 191-231 mg (65-78%) of the title compound as a colorless solid: mp 61-63 °C.

1H NMR (300 MHz, $CDCl_3$) δ 7.43-7.22 (m, 10H), 4.18 (t, J = 6.4 Hz, 1H), 3.63 and 3.56 (2d, AB, J_{AB} = 13.1 Hz, 2H), 2.86 (d, J = 6.4 Hz, 2H), 2.22 (s, NH), 1.07 (s, 9H);

^{13}C NMR (75 MHz, $CDCl_3$) δ 214.9, 143.4, 140.5, 128.6, 128.4, 128.2, 127.4, 127.3, 126.9, 58.5, 51.8, 45.5, 44.3, 26.2;

FTIR (thin film) ν 3334, 3028, 2968, 2870, 1702, 1603, 1454, 1366, 1076, 992, 744, 701;

Anal. Calcd. for $C_{20}H_{25}NO$ (295.42): C, 81.31; H, 8.53; N, 4.74. Found: C, 81.22; H, 8.58; N, 4.66.

Preparation of *syn*- β -Amino Alcohols 19

The relative *syn*-stereochemistry for the NaBH₄ reduction of 2,3-dihydroisoxazoles to the corresponding isoxazolidines was confirmed by NOE-measurements. NOE's were observed between H^a and both H^b and H^c.

(±)-*syn*-2-Benzyl-3-methyl-5-phenylisoxazolidine (**18a**) (Table 2, Entry 1)

To 2-benzyl-3-methyl-5-phenyl-2,3-dihydroisoxazole (**17a**) (138.8 mg, 0.55 mmol) in AcOH (1.7 mL) was added NaBH₄ (63 mg, 1.67 mmol), and the reaction mixture was stirred at 23 °C for 20 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (30 mL). After extraction with dichloromethane (3 x 20 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (20 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 1:19) afforded the title compound (82.1 mg, 59 %) as a clear, colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 1.24 (d, *J* = 6.2, 3H), 2.01 (m, 1H), 2.90 (m, 1H), 3.22 (m, 1H), 4.05 (d, *J* = 13.7, 1H), 4.11 (d, *J* = 13.7, 1H), 5.20 (dd, *J* = 7.8, *J* = 7.8, 1H), 7.26-7.49 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) δ 19.2, 46.1, 60.8, 61.7, 77.8, 125.9, 127.0, 127.1, 128.1, 128.2, 128.9, 137.4, 142.1;

FTIR (thin film) ν 2968, 2918, 1494, 1454, 1373, 734, 697;

HRMS-MALDI Calcd. for C₁₇H₂₀NO [M+H]⁺, 254.1545. Found, 254.1541.

(±)-*syn*-3-(Benzylamino)-1-phenylbutan-1-ol (**19a**) (Table 2, Entry 1)

To (±)-*syn*-2-benzyl-3-methyl-5-phenylisoxazolidine (**18a**) (67.9 mg, 0.26 mmol) in AcOH (1.5 mL) was added activated zinc dust (171 mg, 2.62 mmol) in two portions (5 min interval), and the reaction mixture was stirred at 23 °C for 40 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (15 mL). After extraction with dichloromethane (3 x 10 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (10 mL) and dried over MgSO₄. Concentration by rotary evaporation afforded the title compound (64 mg, 96 %) as a clear, colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 1.22 (d, *J* = 6.3, 3H), 1.53-1.65 (m, 1H), 1.75 (td, *J* = 2.5, *J* = 14.6, 1H), 3.06-3.17 (m, 1H), 3.79 (d, *J* = 12.6, 1H), 4.01 (d, *J* = 12.4, 1H), 4.25 (bs, 1H), 4.92 (dd, *J* = 2.2, *J* = 10.7, 1H), 7.27-7.40 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) δ 20.9, 45.9, 50.7, 54.2, 75.3, 125.5, 126.9, 127.3, 128.2, 128.4, 128.6, 139.1, 145.2;

FTIR (thin film) ν 3016, 2925, 2837, 2362, 1602, 1490, 1452, 748, 698;

HRMS-MALDI Calcd. for C₁₉H₂₆NO [M+H]⁺, 284.2014. Found, 284.2004;

Anal. Calcd. for C₁₇H₂₂NO [M+H]⁺, 256.1701; found, 256.1692.

2-Benzyl-3-isopropyl-5-phenylisoxazolidine (**18b**) (Table 2, Entry 2)

To 2-benzyl-3-isopropyl-5-phenyl-2,3-dihydroisoxazole (**17b**) (591 mg, 2.1 mmol) in AcOH (7 mL) was added NaBH₄ (134 mg, 3.54 mmol), and the reaction mixture was stirred at 23 °C for 15 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (100 mL). After extraction with dichloromethane (3 x 70 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (70 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 1:49 → 1:19) afforded the title compound (528.6 mg, 89 %) as a clear, colorless oil.

¹H NMR (300 MHz, CDCl₃) δ 0.95 (d, *J* = 6.5, 3H), 1.04 (d, *J* = 6.9, 3H), 1.86 (m, 1H), 2.11 (m, 1H), 2.81 (ddd, *J* = 6.9, *J* = 7.8, *J* = 14.3, 1H), 2.99 (ddd, *J* = 7.5, *J* = 7.5, *J* = 7.5, 1H), 4.02 (d, *J* = 13.4, 1H), 4.25 (d, *J* = 13.4, 1H), 5.31 (dd, *J* = 7.7, *J* = 15.9, 1H), 7.30-7.54 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) for *syn*-**18b** δ 18.7, 20.6, 32.5, 41.8, 62.4, 72.7, 78.2, 126.4, 127.2, 127.6, 128.2, 128.3, 129.3, 137.8, 140.3;

FTIR (thin film) ν 3062, 3029, 2956, 2870, 1951, 1877, 1807, 1603, 1495, 1454, 1368, 1029, 756, 732, 697;

HRMS-MALDI Calcd. for C₁₉H₂₄NO [M+H]⁺, 282.1858. Found, 282.1847;

Anal. Calcd. for C₁₉H₂₃NO (281.39): C, 81.10; H, 8.24; N, 4.98. Found: C, 81.03; H, 8.23; N, 4.92.

(±)-*syn*-3-(Benzylamino)-4-methyl-1-phenylpentan-1-ol (**19b**) (Table 2, Entry 2)

To 2-benzyl-3-isopropyl-5-phenylisoxazolidine (**18b**) (166 mg, 0.59 mmol) in AcOH (3 mL) was added activated zinc dust (390 mg, 6 mmol) in two portions (10 min interval), and the reaction mixture was stirred at 23 °C for 45 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (20 mL). After extraction with dichloromethane (3 x 10 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (10 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 2:3 → 19:1) afforded title compound (120.6 mg, 72 %) and the *anti*-diastereoisomer (7.4 mg, 4 %) as a yellowish oil.

¹H NMR (300 MHz, CDCl₃) δ 0.84 (d, *J* = 6.9, 3H), 0.96 (d, *J* = 7.1, 3H), 1.39 (ddd, *J* = 11.3, *J* = 11.3, *J* = 22.0, 1H), 1.71 (ddd, *J* = 2.2, *J* = 2.2, *J* = 14.3, 1H), 2.18 (m, 1H), 2.91 (ddd, *J* = 1.9, *J* = 3.6, *J* = 11.3, 1H), 3.76 (d, *J* = 12.1, 1H), 3.98 (d, *J* = 12.4, 1H), 4.87 (dd, *J* = 1.9, *J* = 10.7, 1H), 7.27-7.40 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) δ 15.5, 19.9, 28.2, 37.4, 50.7, 63.7, 75.7, 125.5, 126.9, 127.3, 128.2, 128.4, 128.6, 139.2, 145.5;

FTIR (thin film) ν 3027, 2955, 2359, 2341, 1495, 1452, 1095, 1065, 750, 699, 668;

HRMS-MALDI Calcd. for C₁₉H₂₆NO [M+H]⁺, 284.2014. Found, 284.2004;

Anal. Calcd. for C₁₉H₂₅NO (283.41): C, 80.52; H, 8.89; N, 4.94. Found: C, 80.50; H, 9.03; N, 4.88.

2-Benzyl-3-cyclohexyl-5-phenylisoxazolidine (**18c**) (Table 2, Entry 3)

To 2-benzyl-3-cyclohexyl-5-phenyl-2,3-dihydroisoxazole (**17c**) (406 mg, 1.27 mmol) in AcOH (5.5 mL) was added NaBH₄ (146 mg, 3.86 mmol), and the reaction mixture was stirred at 23 °C. After 2 h the reaction mixture was treated with an additional quantity of NaBH₄ (97 mg, 2.56 mmol), and stirring was continued for another 30 min at 23 °C. The reaction was quenched by addition of saturated aqueous NaHCO₃ (90 mL). After extraction with dichloromethane (3 x 60 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (60 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 1:19) afforded the title compound (348 mg, 85 %) as a clear, colorless oil.

¹H NMR (300 MHz, CDCl₃) for *syn*-**18c** δ 0.76-0.98 (m, 2H), 1.02-1.30 (m, 3H), 1.63-1.71 (bm, 4H), 1.99-2.09 (m, 2H), 2.69-2.78 (m, 1H), 2.93 (q, *J* = 7.7, 1H), 3.91 (d, *J* = 13.5, 1H), 4.17 (d, *J* = 13.5, 1H), 5.22 (dd, *J* = 6.5, *J* = 9.5, 1H), 7.22-7.45 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) δ 25.9, 26.0, 26.3, 29.3, 31.0, 41.9, 42.4, 62.4, 71.6, 77.9, 126.3, 127.0, 127.4, 128.0, 128.1, 129.1, 137.8, 140.0;

FTIR (thin film) ν 3015, 2922, 2849, 1952, 1811, 1602, 1494, 1450, 750, 732, 697;

HRMS-MALDI Calcd. for C₂₂H₂₈NO [M+H]⁺, 322.2171. Found, 322.2166.

(±)-*syn*-3-(Benzylamino)-3-cyclohexyl-1-phenylpropan-1-ol (**19c**) (Table 2, Entry 3)

To 2-benzyl-3-cyclohexyl-5-phenylisoxazolidine (**18c**) (322.6 mg, 1 mmol) in AcOH (6 mL) was added activated zinc dust (655 mg, 10.02 mmol) in two portions (5 min interval), and the reaction mixture was stirred at 23 °C for 40 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (60 mL). After extraction with dichloromethane (3 x 40 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (40 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 2:3 → 2:1) afforded the title compound (256.3 mg, 79 %) as a clear, colorless oil, which solidified upon standing at 4 °C for several hours (mp 101-103 °C), and the *anti*-diastereoisomer (24.2 mg, 7 %).

¹H NMR (300 MHz, CDCl₃) δ 0.90-1.52 (m, 6H), 1.62-1.85 (m, 7H), 2.91 (d, *J* = 12.1, 1H), 3.78 (d, *J* = 12.4, 1H), 4.00 (d, *J* = 12.4, 1H), 4.89 (dd, *J* = 1.7, *J* = 10.4, 1H), 7.27-7.43 (m, 10H);

¹³C NMR (75 MHz, CDCl₃) δ 15.5, 19.9, 28.2, 37.4, 50.7, 63.7, 75.7, 125.5, 126.9, 127.3, 128.2, 128.4, 128.6, 139.2, 145.5;

FTIR (thin film) ν 3021, 2922, 2849, 2344, 1947, 1879, 1598, 1495, 1450, 750, 699;

HRMS-MALDI Calcd. for C₂₂H₃₀NO [M+H]⁺, 324.2327. Found, 324.2312.

Anal. Calcd. for C₂₂H₂₉NO (323.47): C, 81.69; H, 9.04; N, 4.33. Found: C, 81.84; H, 9.10; N, 4.34.

2-Benzyl-3,5-diphenylisoxazolidine (**18d**) (Table 2, Entry 4)

To 2-benzyl-3,5-diphenyl-2,3-dihydroisoxazole (**17d**) (77 mg, 0.25 mmol) in AcOH (1 mL) was added NaBH₄ (29 mg, 0.77 mmol), and the reaction mixture was stirred at 23 °C. After 150 min the reaction mixture was treated with an additional quantity of NaBH₄ (29 mg, 0.77 mmol), and stirring was continued for another 18.5 h at 23 °C. The reaction was quenched by addition of saturated aqueous NaHCO₃ (30 mL). After extraction with dichloromethane (3 x 20 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (20 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 1:9) afforded the title compound (62.5 mg, 79 %) as a clear, colourless oil.

¹H NMR (300 MHz, CDCl₃) for *syn*-**18d** δ 2.44 (ddd, *J* = 7.2, *J* = 9.2, *J* = 12.4, 1H), 3.17 (ddd, *J* = 7.7, *J* = 7.7, *J* = 12.4, 1H), 3.93 (d, *J* = 14.0, 1H), 4.12 (d, *J* = 14.0, 1H), 4.07-4.12 (dd, 1H), 5.28 (dd, *J* = 7.6, *J* = 7.6, 1H), 7.27-7.47 (m, 15H).

All spectroscopic data were in agreement with the literature⁸

(±)-*syn*-3-(Benzylamino)-1,3-diphenylpropan-1-ol (**19d**) (Table 2, Entry 4)

To 2-benzyl-3,5-diphenylisoxazolidine (**18d**) (61.9 mg, 0.2 mmol) in AcOH (1.4 mL) was added activated zinc dust (132 mg, 2.02 mmol) in two portions (5 min interval), and the reaction mixture was stirred at 23 °C for 45 min. The reaction was quenched by addition of saturated aqueous NaHCO₃ (15 mL). After extraction with dichloromethane (3 x 10 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (10 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 2:3 → 1:1) afforded the title compound (47 mg, 75 %) and the *anti*-diastereoisomer (5.4 mg, 9 %) as a clear, colourless oil.

¹H NMR (300 MHz, CDCl₃) δ 1.92-2.08 (m, 2H), 3.63 (d, *J* = 12.5, 1H), 3.73 (d, *J* = 12.8, 1H), 4.00 (dd, *J* = 3.6, *J* = 10.4, 1H), 5.03 (dd, *J* = 3.0, *J* = 9.8, 1H), 7.27-7.43 (m, 15H);

¹³C NMR (75 MHz, CDCl₃) δ 46.21, 50.94, 63.08, 75.17, 125.46, 126.35, 127.03, 127.33, 127.54, 128.20, 128.42, 128.57, 128.83, 138.95, 142.78, 144.82.

FTIR (thin film) ν 3295, 3060, 3026, 2909, 2850, 2360, 1948, 1876, 1809, 1601, 1493, 1452, 750, 699;

HRMS-MALDI Calcd. for C₂₂H₃₀NO [M+H]⁺, 318.1858. Found, 318.1850;

Anal. Calcd. for C₂₂H₂₉NO (317.42): C, 83.24; H, 7.30; N, 4.41. Found: C, 83.12; H, 7.44; N, 4.24.

(8) Shimizu, T.; Ishizaki, M.; Nitada, N. *Chem. Pharm. Bull.* **2002**, *50*, 908-921.

One-Pot Procedure for the Preparation of *syn*- β -Amino Alcohols 19

($\pm$)-*syn*-3-(benzylamino)-4-methyl-1-phenylpentan-1-ol (19b) (Table 2, Entry 3)

To 2-benzyl-3-isopropyl-5-phenyl-2,3-dihydroisoxazole (**17b**) (294.6 mg, 1.05 mmol) in AcOH (5 mL) was added NaBH₄ (120 mg, 3.17 mmol), and the reaction mixture was stirred at 23 °C. After 30 min the reaction mixture was treated with an additional quantity of NaBH₄ (40 mg, 1.06 mmol), and stirring was continued for another 30 min at 23 °C until complete consumption of starting material was observed. After addition of activated zinc dust (689 mg, 10.54 mmol) in two portions (5 min interval), the reaction mixture was stirred for 30 min at 23 °C. The reaction was quenched by addition of saturated aqueous NaHCO₃ (80 mL). After extraction with dichloromethane (3 x 50 mL), the combined organic layers were washed with saturated aqueous NaHCO₃ (50 mL), dried over MgSO₄ and concentrated by rotary evaporation. Chromatography on silica gel (EtOAc/hexane, 2:3 $\rightarrow$ 4:1) afforded the title compound (186 mg, 63 %) and the *anti*-diastereoisomer (34.6 mg, 12 %) as a yellowish oil.

Preparation of *anti*- β -Amino Alcohols 19

($\pm$)-*anti*-3-(benzylamino)-4-methyl-1-phenylpentan-1-ol

To a solution of 2-benzyl-3-isopropyl-5-phenyl-2,3-dihydroisoxazole (**17b**) (0.14 g, 0.50 mmol, 1.0 equiv) in AcOH (2.0 mL) was added activated zinc dust (0.16 g, 2.5 mmol, 5.0 equiv.) in two portions (5 min interval). The reaction mixture was stirred vigorously for 30 min at 23 °C. Upon completion, the reaction mixture was diluted with dichloromethane (5.0 mL) and poured into a separatory funnel containing saturated NaHCO₃ solution (10 mL) and dichloromethane (10 mL). The layers were separated and the aqueous layer was extracted with dichloromethane (3 x 10 mL). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄ and concentrated in *vacuo*. Purification of the crude material by flash chromatography on silica gel (0.5% NEt₃ in EtOAc/hexane = 3:7) afforded the title compound (96 mg, 68 %) as a yellowish oil.

¹H NMR (300 MHz, CDCl₃) δ 7.39-7.20 (m, 10H), 5.12 (dd, J = 6.4, 3.8 Hz, 1H), 3.77 and 3.53 (2d, AB, J_{AB} = 12.6 Hz, 2H), 2.53 (ddd, J = 6.9, 6.8, 3.2 Hz, 1H), 2.10-2.00 (m, 1H), 1.95-1.79 (m, 2H), 0.89 (dd, J = 19.0, 6.8, 6H);

¹³C NMR (75 MHz, CDCl₃) δ 145.5, 139.4, 128.7, 128.5, 128.2, 128.1, 127.4, 126.6, 125.4, 72.2, 60.7, 51.2, 35.2, 28.4, 19.9, 17.5.

HRMS-MALDI Calcd. for C₁₉H₂₆NO [M+H]⁺, 284.2104. Found 284.2007;

Anal. Calcd for C₁₉H₂₅NO (283.41): C, 80.52; H, 8.89; N, 4.94; O, 5.65. Found: C, 80.62; H, 8.76; N, 4.90; O, 5.73.