A Regioselective Entry to Bromo- γ -hydroxybutenolides: Useful Building Blocks for Assembling Natural Product-*like* Libraries

M. Aquino, I. Bruno, R. Riccio* and L. Gomez-Paloma[†]

Dipartimento di Scienze Farmaceutiche, Università degli Studi di Salerno, via Ponte Don Melillo, 84084, Fisciano (SA), Italy

riccio@unisa.it

SUPPORTING INFORMATION

General Experimental Data

All water and air sensitive reactions were carried out under an inert atmosphere (N₂) in oven- or flame-dried glassware. Methylene chloride, toluene and diethyl ether were distilled from CaH₂ immediately prior to use. Water was degassed under vacuum (10 mbar). All reagents were used from commercial sources without any further purification. Organic extracts were dried over anhydrous Na₂SO₄.

The photo-oxidation reaction was performed using a 300W tungsten lamp.

Reactions were monitored on silica gel 60 F254 (Merck) plates and visualized with potassium permanganate or cerium sulphate and under UV (λ = 254, 365 nm). Flash column chromatography was performed using Merck 60/230-400 mesh silica gel.

Analytical and semipreparative reverse phase HPLC purifications were performed on a Jupiter C-18 column (250 x 4.60mm, 5 μ , 300 Å; 250 x 10.00mm, 10 μ , 300 Å, respectively).

Microwave reactions were performed on a CEM Discover[®] single mode platform using 10 ml pressurized vials.

Reaction yields refer to chromatographically and spectroscopically pure products.

Proton detected (¹H, HMBC, HSQC) and carbon detected NMR spectra were recorded on Bruker instruments of Avance series operating at 300, 400, 600 MHz and 75, 100 and 150 MHz, respectively.

Chemical shifts are expressed in parts per million (ppm) on the delta (δ) scale. The solvent peak was used as internal reference: for ¹H NMR CDCl₃ = 7.26 ppm, MeOD = 3.34 ppm; for ¹³C NMR: CDCl₃ = 77.0 ppm, MeOD = 47.7 ppm. Multiplicities are reported as follows: (s = singlet; d = doublet; t = triplet; q = quartet; dd = doublet of doublets; dt = doublet of triplets; b = broad). High resolution mass spectra were recorded on a Q-TOF Premier Waters (Milford, Massachusetts, USA) mass spectrometer using an electrospray ion source (ESI-MS) and on a ThermoQuest GC-MS equipped with a Zebron[™] Z5-ms column (capillary in cage column 30m x 0.25mm x 0.25mm) using an electron impact ion source (EI-MS).

General procedure for photo-oxidation of 1.

A two neck flask with 150 ml of dry dichloromethane was cooled to -78 °C. Upon stirring, 3-bromofuran (**1**) (612 μ l, 6.8 mmol) and finely powdered polystyrene-bound rose bengal catalyst (150 mg) were added. Then the base (2 equiv.) was introduced and the oxygen bubbled for 15 min. After the 300 W lamp was turned on, the solution was stirred at -78 °C in a continuous flow of oxygen until the reaction appeared complete by TLC (4 to 6 h depending on the base, see Table 1). The mixture was

[†] Deceased

allowed to warm to 0 °C, the catalyst was filtered off and the solution rapidly washed with an aqueous solution of HCl 1M (100 ml). The aqueous layer was extracted with dichloromethane (2 x 100 ml). The organics were finally washed with distilled water, dried, filtered and concentrated *in vacuo*. The residue was purified on a semipreparative C-18 RP-HPLC column, leaving the desired product (**3** or **4**) as an oil.

4-Bromo-5-hydroxy-5H-furan-2-one (3).

¹H-NMR (300 MHz, CDCl₃): δ 6.42 (1H, s, *H*-3), 6.04 (1H, s, *H*-5), 3.6 (1H, bs, OH).

¹³C-NMR (75 MHz, CDCl₃): δ 169.4 (*C*-2), 148.1 (*C*-4), 124.0 (*C*-3), 99.6 (*C*-5).

HRMS calc. for C₄H₂BrO₃: [M - H]⁻ 176.9187, 178.9167 (ratio 1:1); found 176.9178, 178.9163 (ratio 1:1).

3-Bromo-5-hydroxy-5H-furan-2-one (4).

¹H-NMR (300 MHz, CDCl₃): δ 7.34 (1H, s, *H*-4), 6.24 (1H, s, *H*-5).

¹³C-NMR (75 MHz, CDCl₃): δ 167.7 (*C*-2), 149.3 (*C*-4), 117.8 (*C*-3), 98.7 (*C*-5).

HRMS calc. for C₄H₂BrO₃: [M - H]⁻ 176.9187, 178.9167 (ratio 1:1); found 176.9175, 178.9161 (ratio 1:1).

3,4-Dibromo-5-(2-methoxy-ethoxymethoxy)-5H-furan-2-one (6).

To a solution of mucobromic acid (**5**) (100 mg, 0.387 mmol) in 10 ml of dry dichloromethane, MEM-Cl (66 µl, 0.581 mmol) was added. Diisopropylethylamine (101 µl, 0.581 mmol) was dropwise added over a period of 15 min. After 3.30 h, the reaction was quenched with 20 ml of HCl 1M. The aqueous layer was extracted with dichloromethane (2 x 30 ml) and the organics were dried, filtered and concentrated *in vacuo* to leave dark oil. The crude oil was purified by flash chromatography (10% diethyl ether/hexane) to give **6** (115 mg, 86% yield).

¹H-NMR δ (300 MHz; CDCl₃): δ 6.07 (1H, s, *H*-5), 5.11 (1H, d, *J*=7 Hz, *H*-1'^A), 4.93 (1H, d, *J*=7 Hz, *H*-1'^B), 3.89-3.85 (1H, m, *H*-2'^A), 3.82-3.76 (1H, m, *H*-2'^B), 3.60 (2H, t, *J*=4.5 Hz, *H*-3'), 3.40 (3H, s, *H*-4').

¹³C-NMR δ (75 MHz; CDCl₃): δ 167.4 (*C*-2), 144.2 (*C*-4), 119.1 (*C*-3), 100.4 (*C*-5), 95.2 (*C*-1'), 72.1 (*C*-2'), 69.0 (*C*-3'), 59.8 (*C*-4').

HRMS calc. for C₈H₁₁Br₂O₅: [M + H]⁺ 344.8973, 346.8953, 348.8932 (ratio 1:2:1); found 344.8942, 346.8925, 348.8918 (ratio 1:2:1).

Protection of 3 and 4 as MEM ethers.

To a solution of the desired isomer (**3** or **4**) (100 mg, 0.562 mmol) in 10 ml of dry dichloromethane MEM-Cl (95.8 µl, 0.843 mmol) was added. Diisopropylethylamine (146.8 µl, 0.843 mmol) was dropwise added over a period of 15 min. After 5 h, the reaction was quenched with 20 ml of HCl 1M. The aqueous layer was extracted with dichloromethane (2 x 30 ml) and the organics were dried, filtered and concentrated *in vacuo* to leave dark oil. The crude oil was purified by flash chromatography (10% diethyl ether/hexane) to give, depending on the starting material, **7** or **8** (123 mg, 82% yield).

3-Bromo-4-oxo-but-2-enoic acid 2-methoxy-ethoxymethyl ester (7).

$^1\text{H-NMR}$ δ (300 MHz; CDCl_3): δ 10.23 (1H, s, H -4), 7.29 (1H, s, H -2), 5.46 (2H, s, H_2 -1'), 3.85-3.82 (2H, m, H_2 -2'), 3.57-3.56 (2H, m, H_2 -3'), 3.38 (3H, s, H_3 -4').

HRMS calc. for $\text{C}_8\text{H}_{12}\text{BrO}_5$: $[\text{M} + \text{H}]^+$ 266.9868, 268.9848 (ratio 1:1); found 266.9875, 268.9857 (ratio 1:1).

2-Bromo-4-oxo-but-2-enoic acid 2-methoxy-ethoxymethyl ester (8).

$^1\text{H-NMR}$ δ (300 MHz; CDCl_3): δ 10.08 (1H, d, J = 6.6 Hz, H -4), 6.88 (1H, d, J = 6.6 Hz, H -3), 5.53 (2H, s, H_2 -1'), 3.87-3.86 (2H, m, H_2 -2'), 3.58-3.56 (2H, m, H_2 -3'), 3.37 (3H, s, H_3 -4').

$^{13}\text{C-NMR}$ δ (75 MHz; CDCl_3): δ 188.1 (C -4), 160.8 (C -1), 140.5 (C -3), 131.5 (C -2), 92.0 (C -1'), 71.0 (C -2'), 70.2 (C -3'), 58.8 (C -4').

HRMS calc. for $\text{C}_8\text{H}_{12}\text{BrO}_5$: $[\text{M} + \text{H}]^+$ 266.9868, 268.9848 (ratio 1:1); found 266.9880, 268.9859 (ratio 1:1).

Protection of 3 and 4 as THP ethers.

A solution of the hydroxyfuranone (40 mg, 0.22 mmol) and DHP (51 μl , 0.56 mmol) in 4 ml of dry dichloromethane was cooled to 0 $^\circ\text{C}$ in ice-bath. Upon stirring *p*-toluenesulfonic acid monohydrate (192 mg, 1.12 mmol) was slowly added keeping the temperature at 0 $^\circ\text{C}$ for 10 min. The mixture was allowed to warm to room temperature and the reaction was stopped after 2 h. The solution was diluted with 20 ml of dichloromethane and then washed with a 1:1:2 mixture of saturated brine, saturated sodium bicarbonate and water (20 ml). The organic layer was dried, filtered and concentrated in *vacuo* to leave brown oil. The crude oil was purified by flash chromatography (10% diethyl ether/hexane) to give, depending on the starting material, **11** (55.9 mg, 95% yield) or **12** (51.2 mg, 87% yield).

4-Bromo-5-(tetrahydropyran-2-yloxy)-5H-furan-2-one (11).

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 6.38 (1H, s, H -3), 6.10 (1H, s, H -5), 5.16 (1H, s, H -2'), 3.96-3.90 (1H, m, H -6'A), 3.81-3.75 (1H, m, H -6'B), 1.78-1.54 (6H, m, H_2 -3', H_2 -4', H_2 -5').

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 168.7 (C -2), 148.6 (C -4), 124.3 (C -3), 98.8 (C -5), 98.4 (C -2'), 64.3 (C -6'), 30.2 (C -3'), 25.4 (C -5'), 19.0 (C -4').

EI-MS calc. for $\text{C}_9\text{H}_{11}\text{BrO}_4$: $[\text{M}]^+$ 261.9, 263.9 (ratio 1:1), found 261.9, 263.9 (ratio 1:1).

3-Bromo-5-(tetrahydropyran-2-yloxy)-5H-furan-2-one (12).

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.32 (1 H, d, J =1.5 Hz, H -4), 6.21 (1 H, d, J =1.5 Hz, H -5), 5.12-5.10 (1H, m, H -2'), 3.87-3.80 (1H, m, H -6'A), 3.66-3.61 (1H, m, H -6'B), 1.80-1.64 (6H, m, H_2 -3', H_2 -4', H_2 -5').

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 166.5 (C -2), 147.8 (C -4), 118.1 (C -3), 98.2 (C -5), 95.0 (C -2'), 63.2 (C -6'), 30.1 (C -3'), 25.4 (C -5'), 19.0 (C -4').

EI-MS calc. for $\text{C}_9\text{H}_{11}\text{BrO}_4$: $[\text{M}]^+$ 261.9, 263.9 (ratio 1:1), found 261.9, 263.9 (ratio 1:1).

Suzuki coupling of 11 and 12.

In a 10 ml pressurized vessel the desired isomer **11** or **12** (40 mg, 0.153 mmol), the desired boronic acid (0.229 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (33 μg , 0.0459 $\times 10^{-3}$ mmol) and TBAB (49.2 mg, 0.153 mmol) were

placed. 500 μ l of degassed THF and CsF (93 mg, 0.612 mmol) in 500 μ l of degassed water were added under a N_2 atmosphere. The mixture was then irradiated with microwaves in a CEM Discover[®] apparatus for 6 min. at 200 W, setting the temperature to 120 $^{\circ}$ C and switching the *power-max* on. The solution was then diluted with 10 ml of dichloromethane and washed with HCl 1N aqueous solution (10 ml). The aqueous layer was re-extracted (2 x 15 ml) and the organics were dried, filtered and concentrated *in vacuo*. The brown oil was purified by flash chromatography (20% diethyl ether/hexane) to give, depending on the starting material (see table 3), the desired coupled adduct.

General procedure for THP cleavage.

To a solution of the THP-derivatives (**13a-22a**) in 8 ml of THF (volumes are based on starting amount of **13a**), was added HCl 1N aqueous solution (8 ml) at room temperature and upon stirring. After 3 h, the reaction was quenched with sat. $NaHCO_3$ (15 ml), concentrated *in vacuo* and extracted with AcOEt (3 x 20 ml). The organics were dried, filtered and concentrated *in vacuo*, to leave white precipitate that was purified by flash chromatography (30% diethyl ether/hexane). All the products (**13-22**) were obtained as white solids.

5-Hydroxy-4-phenyl-5H-furan-2-one (**13**).

80.2 mg (0.308 mmol) of compound **13a** were used, obtaining 48.3 mg of compound **13** (89% yield).

1H -NMR (600 MHz, $CDCl_3$): δ 7.73-7.71 (2H, dd, *H*-2'A, *H*-2'B), 7.52-7.48 (3H, m, *H*-3'A, *H*-3'B, *H*-4'), 6.55 (1H, s, *H*-5), 6.43 (1H, s, *H*-3).

^{13}C -NMR (75 MHz, $CDCl_3$): δ 171.1 (*C*-2), 162.3 (*C*-4), 131.9 (*C*-1'), 129.5 (*C*-2'A, *C*-2'B), 129.4 (*C*-4'), 128.2 (*C*-3'A and *C*-3'B), 116.0 (*C*-3), 98.6 (*C*-5).

HRMS calc. for $C_{10}H_7O_3$: [$M - H$]⁻ 175.0395, found 175.0388.

4-Benzo[*b*]tiophen-2-yl-5-hydroxy-5H-furan-2-one (**14**).

19.6 mg (0.062 mmol) of compound **14a** were used, obtaining 12.9 mg of compound **14** (90% yield).

1H -NMR (600 MHz, $CDCl_3$): δ 7.87 (2H, d, $J=7.7$ Hz, *H*-5', *H*-8'), 7.83 (1H, s, *H*-3'), 7.48-7.42 (2H, m, *H*-6', *H*-7'), 6.52 (1H, s, *H*-5), 6.29 (1H, s, *H*-3).

^{13}C -NMR (150 MHz, $CDCl_3$): δ 169.6 (*C*-2), 156.4 (*C*-4), 141.1 (*C*-9'), 139.3 (*C*-4'), 131.8 (*C*-2'), 128.0 (*C*-3'), 127.6 (*C*-7'), 125.2 (*C*-6'), 125.2 (*C*-5'), 122.3 (*C*-8'), 114.9 (*C*-3), 97.3 (*C*-5).

HRMS calc. for $C_{12}H_7O_3S$: [$M - H$]⁻ 231.0116, found 231.0105.

4-(2-Hydroxy-5-oxo-2,5-dihydro-furan-3-yl)-benzaldehyde (**15**).

23.2 mg (0.080 mmol) of compound **15a** were used, obtaining 15.3 mg of compound **15** (93% yield).

1H -NMR (600 MHz, $CDCl_3$): δ 10.08 (1H, s, CHO), 7.99 (2H, d, $J=8.3$ Hz, *H*-3'A, *H*-3'B), 7.88 (2H, d, $J=8.3$ Hz, *H*-2'A, *H*-2'B), 6.61 (1H, s, *H*-5), 6.54 (1H, s, *H*-3).

^{13}C -NMR (150 MHz, $CDCl_3$): δ 191.6 (CHO), 170.4 (*C*-2), 162.2 (*C*-4), 137.3 (*C*-1'), 135.9 (*C*-4'), 129.4 (*C*-3'A, *C*-3'B), 128.5 (*C*-2'A, *C*-2'B), 114.5 (*C*-3), 97.8 (*C*-5).

HRMS calc. for $C_{11}H_7O_4$: [$M - H$]⁻ 203.0344, found 203.0328.

5-Hydroxy-4-(4-methoxy-phenyl)-5H-furan-2-one (16).

22.5 mg (0.077 mmol) of compound **16a** were used, obtaining 14.5 mg of compound **16** (91% yield).

¹H-NMR (600 MHz, MeOD): δ 7.75 (2H, d, *J*=9 Hz, *H*-2'A, *H*-2'B), 7.02 (2H, d, *J*=9 Hz, *H*-3'A, *H*-3'B), 6.52 (1H, s, *H*-5), 6.37 (1H, s, *H*-3), 3.86 (3H, s, OCH₃).

¹³C-NMR (150 MHz, MeOD): δ 172.2 (*C*-2), 163.8 (*C*-4'), 162.4 (*C*-4), 129.3 (*C*-2'A and *C*-2'B), 121.9 (*C*-1'), 113.5 (*C*-3'A, *C*-3'B), 111.0 (*C*-3), 98.0 (*C*-5), 54.4 (OCH₃).

HRMS calc. for C₁₁H₉O₄: [M - H]⁻ 205.0501, found 205.0515.

5-Hydroxy-4-naphthalen-1-yl-5H-furan-2-one (17).

22.4 mg (0.072 mmol) of compound **17a** were used, obtaining 14.4 mg of compound **17** (88% yield).

¹H-NMR (300 MHz, MeOD): δ 8.18 (1H, d, *H*-2'), 8.00-7.95 (2H, m, *H*-6', *H*-9'), 7.70 (1H, d, *H*-4'), 7.63-7.54 (3H, m, *H*-3', *H*-7', *H*-8'), 6.75 (1H, s, *H*-5), 6.48 (1H, s, *H*-3).

¹³C-NMR (150 MHz, MeOD): δ 169.8 (*C*-2), 163.8 (*C*-4), 133.5 (*C*-1'), 130.8 (*C*-5'), 130.4 (*C*-10'), 128.9 (*C*-6'), 128.2 (*C*-4'), 127.6 (*C*-8'), 127.2 (*C*-7'), 126.9 (*C*-3'), 126.5 (*C*-9'), 124.4 (*C*-2'), 119.8 (*C*-3), 99.8 (*C*-5).

HRMS calc. for C₁₄H₉O₃: [M - H]⁻ 225.0552, found 225.0542.

5-Hydroxy-4-phenyl-5H-furan-2-one (18).

40.1 mg (0.154 mmol) of compound **18a** were used, obtaining 24.1 mg of compound **18** (89% yield).

¹H-NMR (600 MHz, MeOD): δ 7.71-7.68 (2H, m, *H*-2'A, *H*-2'B), 7.48-7.44 (3H, m, *H*-3'A, *H*-3'B, *H*-4'), 7.35 (1H, s, *H*-4), 6.72 (1H, s, *H*-5).

¹³C-NMR (150 MHz, MeOD): δ 168.6 (*C*-2), 139.5 (*C*-4), 131.5 (*C*-3), 131.3 (*C*-1'), 129.3 (*C*-3'A, *C*-3'B), 128.1 (*C*-2'A, *C*-2'B), 128.0 (*C*-4'), 97.0 (*C*-5).

HRMS calc. for C₁₀H₇O₃: [M - H]⁻ 175.0395, found 175.0388.

3-Benzo[b]thiophen-2-yl-5-hydroxy-5H-furan-2-one (19).

23.8 mg (0.075 mmol) of compound **19a** were used, obtaining 16.4 mg of compound **19** (94% yield).

¹H-NMR (600 MHz, CDCl₃): δ 8.02 (1H, s, *H*-3'), 7.86-7.81 (2H, m, *H*-5', *H*-8'), 7.41-7.38 (2H, m, *H*-6', *H*-7'), 7.24 (1H, s, *H*-4), 6.28 (1H, s, *H*-5).

¹³C-NMR (150 MHz, CDCl₃): δ 168.4 (*C*-2), 142.5 (*C*-4), 142.3 (*C*-9'), 139.9 (*C*-4'), 132.9 (*C*-3), 130.1 (*C*-2'), 127.6 (*C*-3'), 127.8 (*C*-7'), 126.4 (*C*-6'), 126.6 (*C*-5'), 122.7 (*C*-8'), 96.7 (*C*-5).

HRMS calc. for C₁₂H₇O₃S: [M - H]⁻ 231.0116, found 231.0112.

3-(5-Hydroxy-2-oxo-2,5-dihydro-furan-3-yl)-benzaldehyde (20).

20.2 mg (0.070 mmol) of compound **20a** were used, obtaining 13.0 mg of compound **20** (91% yield).

¹H-NMR (600 MHz, MeOD): δ 10.02 (1H, s, CHO), 8.12 (2H, d, *J*=8.3 Hz, *H*-3'A, *H*-3'B), 7.97 (2H, d, *J*=8.3 Hz, *H*-2'A, *H*-2'B), 7.76 (1H, s, *H*-4), 6.22 (1H, s, *H*-5).

¹³C-NMR (150 MHz, MeOD): δ 191.9 (CHO), 169.9 (*C*-2), 142.7 (*C*-4), 136.8 (*C*-1'), 135.1 (*C*-4'), 132.4 (*C*-3), 129.3 (*C*-3'A, *C*-3'B), 127.7 (*C*-2'A, *C*-2'B), 96.7 (*C*-5).

HRMS calc. for C₁₁H₇O₄: [M - H]⁻ 203.0344, found 203.0333.

5-Hydroxy-3-(4-methoxy-phenyl)-5H-furan-2-one (21).

24.6 mg (0.085 mmol) of compound **21a** were used, obtaining 15.7 mg of compound **21** (90% yield).

¹H-NMR (600 MHz, MeOD): δ 7.86 (2H, d, *J*=9.0 Hz, *H*-2'A, *H*-2'B), 7.42 (1H, s, *H*-4), 6.97 (2H, d, *J*=9.0 Hz, *H*-3'A, *H*-3'B), 6.15 (1H, s, *H*-5), 3.82 (1H, s, OCH₃).

¹³C-NMR (150 MHz, MeOD): δ 170.5 (*C*-2), 160.6 (*C*-4'), 142.0 (*C*-4), 132.5 (*C*-3), 128.3 (*C*-2'A, *C*-2'B), 121.2 (*C*-1'), 113.5 (*C*-3'A, *C*-3'B), 96.3 (*C*-5), 54.1 (OCH₃).

HRMS calc. for C₁₁H₉O₄: [M - H]⁻ 205.0501, found 205.0512.

5-Hydroxy-3-(1-naphthalen-1-yl)-5H-furan-2-one (22).

23.0 mg (0.074 mmol) of compound **22a** were used, obtaining 15.1 mg of compound **22** (90% yield).

¹H-NMR (600 MHz, MeOD): δ 7.98-7.91 (3H, m, *H*-2', *H*-6', *H*-9'), 7.59-7.52 (4H, m, *H*-3', *H*-4', *H*-7', *H*-8'), 7.49 (1H, s, *H*-4), 6.36 (1H, s, *H*-5).

¹³C-NMR (150 MHz, MeOD): δ 171.2 (*C*-2), 149.2 (*C*-4), 132.4 (*C*-3), 134.2 (*C*-1'), 133.3 (*C*-5'), 131.1 (*C*-10'), 127.7 (*C*-6'), 127.2 (*C*-4'), 126.3 (*C*-8'), 125.8 (*C*-7'), 125.4 (*C*-3'), 124.7 (*C*-9'), 124.4 (*C*-2'), 97.3 (*C*-5).

HRMS calc. for C₁₄H₉O₃: [M - H]⁻ 225.0552, found 225.0542.

Method Set

Column: Jupiter C-18 semipreparative, **p/n:** 00G-4055-N0

Eluent: CH₃CN (0.1% TFA)/H₂O (0.1% TFA)

Gradient: 5/95 to 100/0, **time:** 45 minutes

Flow Rate: 4 mL/min

Detection: UV at 280 nm, 1.0 AUFS

Injection: 100 µl (0.1 mg/µL)