Regioselective Lewis Acid-Mediated [1,3] Rearrangement of Allylvinyl Ethers

Christopher G. Nasveschuk, and Tomislav Rovis*

*Department of Chemistry Colorado State University, Fort Collins, CO, 80523*E-mail: rovis@lamar.colostate.edu

General Methods: All reactions were performed under an inter atmosphere of argon in flame-dried glassware with magnetic stirring. Dichloromethane and toluene were degassed with argon and passed through two columns of neutral alumina. Column chromatography was performed on EM Science silica gel 60 (230-400 mesh). Thin layer chromatography was performed on EM Science 0.25 mm silica gel 60-F plates. Visualization was accomplished with UV light, KMnO₄, and aqueous ceric ammonium molybdate followed by heating.

Et₂AlCl was purchased as a 1.8M solution in toluene. EtAlCl₂ was purchased as a 1.0M solution in hexane. CuOTf₂ was purchased from Strem Chemical Co. All other Lewis acids used were purchased from Aldrich Chemical Co. and used without further purification.

Infrared spectra were obtained on a Nicolet Avatar 320 FT-IR spectrometer. ¹H NMR spectra were recorded on a Varian 400 MHz spectrometer at ambient temperature. Data are reported as follows: chemical shift in parts per million (δ, ppm) from an internal standard [deuterated chloroform (CDCl₃)], multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet), integration, and coupling constant (Hz). ¹³C NMR were recorded on a Varian 100 MHz spectrometer at ambient temperature. Chemical shifts are reported in ppm from (CDCl₃) taken as 77.23 ppm. Mass spectra were obtained on Fisons VG Autospec. Analytical high performance liquid chromatography (HPLC) was performed on a Dynamax model SD-200 HPLC equipped with a Dynamax model UV-1 variable wavelength UV detector using Chiracel chiral columns as indicated.

General Procedure for the Lewis acid-mediated [1,3] rearrangement of allylvinyl ethers. A flame-dried round bottom flask was charged with Lewis acid (1.05 eq.) under an inert atmosphere of argon and subsequently diluted with CH₂Cl₂ or toluene (1-2 mL) and mixed for 10 min. at ambient temperature. The mixture was then cooled to the appropriate temperature, allylvinyl ether (0.25 mmol) was then added neat, drop wise via syringe and the reaction was stirred for 1-5 h. The reaction mixture was quenched with 5 mL sat. aq. NH₄Cl and allowed to warm to ambient temperature. The biphasic mixture was poured onto 10 mL of H₂O, the organic layer was separated and the aqueous layer was extracted with CH₂Cl₂ (3 X 10 mL). The organic layers were combined, dried over Na₂SO₄, filtered, then concentrated *in vacuo*, and purified via column chromatography. Yamamoto-type Lewis acids were formed in situ by dilution of the appropriate phenol (2eq.) in toluene, subsequently cooling the mixture to 0 °C, followed by drop wise addition of Me₃Al (1eq.) and mixing for 1h.

(3-Vinyloxy-but-1-enyl)-benzene (1).¹ ¹H NMR (400 MHz CDCl₃) δ 7.49-7.22 (5H, m), 6.56 (1H, d, *J* = 16.0 Hz), 6.38 (1H, dd, *J* = 14.1, 6.6 Hz), 6.17 (1H, dd, *J* = 16.0, 6.6 Hz), 4.51 (1H, dq, *J* = 6.5, 6.5 Hz), 4.36 (1H, dd, *J* = 14.3, 1.4 Hz), 4.03 (1H, dd, *J* = 6.6, 1.5 Hz), 1.41 (3H, d, *J* = 6.6 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.7, 136.6, 131.3, 130.6, 128.8, 128.0, 126.7, 89.2, 76.6, 21.4; IR (NaCl dep from CH₂Cl₂) 2979, 1633, 1322, 1191, 967, 693 cm⁻¹.

1-Methyl-4(3-vinyloxy-but-1-enyl)-benzene (4). ¹H NMR (400 MHz CDCl₃) δ 7.30-7.11 (4H, m), 6.52 (1H, d, *J* = 16.0 Hz), 6.37 (1H, dd, *J* = 14.1, 6.6 Hz), 6.12 (1H, dd, *J* = 16.0, 6.8 Hz), 4.49 (1H, dq, *J* = 6.6, 6.6 Hz), 4.35 (1H, d, *J* = 14.1 Hz), 4.02 (1H, d, *J* = 6.6 Hz), 2.34 (3H, s), 1.41 (3H, d, *J* = 6.4 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.7, 137.9, 133.8, 131.3, 129.6, 129.5, 126.6, 89.2, 76.8, 21.4, 21.4; IR (NaCl dep from CH₂Cl₂) 2978, 1633, 1514, 1191, 1065, 800 cm⁻¹.

1-Methoxy-4(3-vinyloxy-but-1-enyl)-benzene (5). ¹H NMR (400 MHz CDCl₃) δ 7.35-6.83 (4H, m), 6.50 (1H, d, *J* = 16.0 Hz), 6.38 (1H, dd, *J* = 14.1, 6.6 Hz), 6.03 (1H, dd, *J* = 16.0, 6.8 Hz), 4.48 (1H, dq, *J* = 6.6, 6.6 Hz), 4.35 (1H, dd, *J* = 14.1, 1.3 Hz), 4.02 (1H, dd, *J* = 6.4, 1.1 Hz), 3.81 (3H, s), 1.40 (3H, d, *J* = 6.4 Hz); ¹³C NMR (100 MHz CDCl₃) δ 159.6, 150.7, 130.9, 129.3, 128.4, 127.9, 114.2, 89.1, 76.9, 55.5, 21.5; IR (NaCl dep from CH₂Cl₂) 2977, 1634, 1512, 1250, 1035, 820 cm⁻¹.

1-Trifluoromethyl-4(3-vinyloxy-but-1-enyl)-benzene (6). ¹H NMR (400 MHz CDCl₃) δ 7.59-7.45 (4H, m), 6.59 (1H, d, *J* = 16.2 Hz), 6.38 (1H, dd, *J* = 14.1, 6.6 Hz), 6.28 (1H, dd, *J* = 16.2, 6.4 Hz), 4.55 (1H, dq, *J* = 6.4, 6.4 Hz), 4.36 (1H, dd, *J* = 14.1, 1.5 Hz), 4.07 (1H, dd, *J* = 6.6, 1.5 Hz), 1.43 (3H, d, *J* = 6.4 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.5, 140.1, 133.3, 129.6, 126.9, 126.9, 125.8, 125.7, 89.4, 76.0, 21.2; IR (NaCl dep from CH₂Cl₂) 2982, 1616, 1415, 1326, 1124, 819 cm⁻¹.

(1-Vinyloxy-but-2-enyl)-benzene (7). ¹H NMR (400 MHz CDCl₃) δ 7.41-7.27 (5H, m), 6.39 (1H, dd, *J* = 14.1, 6.6 Hz), 5.79-5.59 (2H, m), 5.17 (1H, d, *J* = 6.8 Hz), 4.34 (1H, d, *J* = 14.1 Hz), 4.03 (1H, d, *J* = 6.6 Hz), 1.72 (3H, d, *J* = 6.2 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.5, 140.8, 131.2, 129.3, 128.7, 127.9, 126.7, 89.8, 82.4, 18.0; IR (NaCl dep from CH₂Cl₂) 3030, 1635, 1452, 1318, 1175, 965, 698 cm⁻¹.

(1-Methyl-3-vinyloxy-but-1-enyl)-benzene (8).¹ ¹H NMR (400 MHz CDCl₃) δ 7.44-7.23 (5H, m), 6.38 (1H, dd, *J* = 14.3, 6.6 Hz), 5.76 (1H, dd, *J* = 8.4, 1.1 Hz), 4.79 (1H, dq, *J* = 6.6, 6.6 Hz), 4.32 (1H, dd, *J* = 13.9, 1.1 Hz), 4.03 (1H, dd, *J* = 6.6, 1.1 Hz), 2.10 (3H, d, *J* = 1.1 Hz), 1.39 (3H, d, *J* = 6.6

¹ Cresson, P. *Bull. Soc. Chim. Fr.* **1964**, 10, 2629-2635.

Hz); ^{13}C NMR (100 MHz CDCl_3) δ 150.6, 142.8, 137.0, 129.8, 128.5, 127.5, 126.0, 88.8, 72.9, 21.1, 16.4; IR (NaCl dep from CH_2Cl_2) 2978, 1633, 1444, 1192, 821, 696 cm^{-1} .

3-Methyl-5-phenyl-hex-4-enal (9). ^1H NMR (400 MHz CDCl_3) δ 9.74 (1H, t, $J = 2.13\text{ Hz}$), 7.36-7.19 (5H, m), 5.55 (1H, d, $J = 9.6\text{ Hz}$), 3.14 (1H, m), 2.46 (2H, dd, $J = 7.0, 2.1\text{ Hz}$), 2.07 (3H, s), 1.11 (3H, d, $J = 6.6\text{ Hz}$); ^{13}C NMR (100 MHz CDCl_3) δ 202.5, 143.6, 134.9, 132.3, 128.4, 127.1, 125.9, 51.2, 28.6, 21.2, 16.3; IR (NaCl dep from CH_2Cl_2) 2961, 1724, 1445, 1026, 759, 697 cm^{-1} ; HRMS (FAB+) calcd for $\text{C}_{13}\text{H}_{16}\text{O}$, 188.1201. Found 188.1193.

(1-Methyl-3-vinyloxy-pent-1-enyl)-benzene (11sm). ^1H NMR (400 MHz CDCl_3) δ 7.46-7.24 (5H, m), 6.40 (1H, dd, $J = 13.9, 6.6\text{ Hz}$), 5.73 (1H, d, $J = 8.4\text{ Hz}$), 4.56 (1H, dt, $J = 6.6, 6.6\text{ Hz}$), 4.30 (1H, dd, $J = 14.3, 1.5\text{ Hz}$), 4.01 (1H, dd, $J = 6.6, 1.1\text{ Hz}$), 2.12 (3H, s), 1.91-1.58 (2H, m), 0.99 (3H, t, $J = 7.3\text{ Hz}$); ^{13}C NMR (100 MHz CDCl_3) δ 150.9, 143.0, 137.9, 128.7, 128.5, 127.5, 126.0, 88.6, 78.2, 28.5, 16.7, 9.8; IR (NaCl dep from CH_2Cl_2) 2966, 1632, 1446, 1190, 757, 696 cm^{-1} .

3-Ethyl-5-phenyl-hex-4-enal (11). ^1H NMR (400 MHz CDCl_3) δ 9.73 (1H, t, $J = 2.3\text{ Hz}$), 7.37-7.19 (5H, m), 5.49 (1H, dd, $J = 10.0, 1.3\text{ Hz}$), 2.93 (1H, m), 2.51 (1H, ddd, $J = 16.0, 5.8, 2.3\text{ Hz}$), 2.41 (1H, ddd, $J = 16.0, 8.1, 2.3\text{ Hz}$), 2.06 (3H, s), 1.55 (1H, m), 1.37 (1H, m), 0.91 (3H, t, $J = 7.5\text{ Hz}$); ^{13}C NMR (100 MHz CDCl_3) δ 202.7, 143.8, 136.3, 131.1, 128.4, 127.1, 126.0, 49.7, 35.5, 28.9, 16.7, 11.9; IR (NaCl dep from CH_2Cl_2) 2961, 2721, 1724, 1494, 1027, 697 cm^{-1} ; HRMS (FAB+) calcd for $\text{C}_{14}\text{H}_{18}\text{O}$, 202.1358. Found 202.1355.

(1,4-Dimethyl-3-vinyloxy-pent-1-enyl)-benzene (12sm). ^1H NMR (400 MHz CDCl_3) δ 7.43-7.22 (5H, m), 6.36 (1H, dd, $J = 14.1, 6.4\text{ Hz}$), 5.7 (1H, dd, $J = 9.2, 1.3\text{ Hz}$), 4.32 (1H, dd, $J = 9.0, 6.4\text{ Hz}$), 4.25 (1H, dd, $J = 14.1, 1.5\text{ Hz}$), 4.95 (1H, dd, $J = 6.6, 1.3\text{ Hz}$), 2.09 (3H, s), 1.99-1.90 (2H, m), 1.00 (3H, d, $J = 6.8\text{ Hz}$), 0.94 (3H, d, $J = 7.0\text{ Hz}$); ^{13}C NMR (100 MHz CDCl_3) δ 151.2, 143.1, 138.5, 128.4, 127.5, 127.3, 126.1, 88.4, 81.7, 33.5, 18.6, 18.2, 17.0; IR (NaCl dep from CH_2Cl_2) 2960, 1632, 1446, 1194, 1041, 696 cm^{-1} .

3-Isopropyl-5-phenyl-hex-4-enal (12). ^1H NMR (400 MHz CDCl_3) δ 9.72 (1H, t, $J = 2.4\text{ Hz}$), 7.38-7.20 (5H, m), 5.57 (1H, d, $J = 10.3\text{ Hz}$), 2.87 (1H, ddt, $J = 15.8, 10.3, 5.9\text{ Hz}$), 2.63-2.37 (2H, m), 2.08 (3H, s), 1.74 (1H, dh, $J = 13.6, 6.6\text{ Hz}$), 0.97 (3H, d, $J = 7.0\text{ Hz}$), 0.94 (3H, d, $J = 7.0\text{ Hz}$); ^{13}C NMR (100 MHz CDCl_3) δ 203.0, 144.0, 136.8, 129.5, 128.4, 127.1, 126.1, 47.5, 40.0, 33.0, 20.7, 19.4, 16.9; IR (NaCl dep from CH_2Cl_2) 2959, 2720, 1724, 1494, 1027, 698 cm^{-1} ; HRMS (FAB+) calcd for $\text{C}_{15}\text{H}_{19}\text{O}$, 216.1514. Found 216.1513.

(1,4,4-Trimethyl-3-vinyloxy-pent-1-enyl)-benzene (13sm). ^1H NMR (400 MHz CDCl_3) δ 7.43-7.24 (5H, m), 6.36 (1H, dd, $J = 14.3, 6.6\text{ Hz}$), 5.72 (1H, dq, $J = 9.6, 1.3\text{ Hz}$), 4.26-4.21 (2H, m), 3.93 (1H, dd, $J = 6.6, 1.3\text{ Hz}$), 2.11 (3H, s), 0.98 (9H, s); ^{13}C NMR (100 MHz CDCl_3) δ

151.7, 143.3, 138.9, 128.5, 127.5, 126.5, 126.1, 87.9, 84.1, 36.4, 26.1, 17.1; IR (NaCl dep from CH₂Cl₂) 2961, 1724, 1494, 1365, 1028, 698 cm⁻¹.

3-tert-Butyl-5-phenyl-hex-4-enal (13). ¹H NMR (400 MHz CDCl₃) δ 9.65 (1H, t, *J* = 3.0), 7.35-7.29 (5H, m), 5.57 (1H, d, *J* = 11.1 Hz), 2.82 (1H, dt, 10.9, 3.4 Hz), 2.60 (1H, m), 2.35 (1H, m), 2.06 (3H, s), 0.94 (9H, s); ¹³C NMR (100 MHz CDCl₃) δ 203.3, 144.1, 137.1, 129.0, 128.4, 127.1, 126.1, 45.2, 43.5, 34.4, 27.6, 17.0; IR (NaCl dep from CH₂Cl₂) 2961, 1724, 1494, 1365, 1028, 698 cm⁻¹; HRMS (FAB+) calcd for C₁₆H₂₂O, 230.1671. Found 2230.1679.

(1-Ethyl-3-vinyloxy-but-1-enyl)-benzene (14sm). ¹H NMR (400 MHz CDCl₃) δ 7.39-7.23 (5H, m), 6.37 (1H, dd, *J* = 14.3, 6.6 Hz), 5.6 (1H, d, *J* = 8.5 Hz), 4.78 (1H, m), 4.32 (1H, dd, *J* = 14.1, 1.3 Hz), 4.02 (1H, dd, *J* = 6.8, 1.5 Hz), 2.61-2.51 (2H, m), 1.39 (3H, d, *J* = 6.4 Hz), 0.98 (3H, t, *J* = 7.5 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.6, 144.0, 141.8, 128.5, 128.3, 128.1, 126.7, 89.0, 72.8, 23.5, 21.7, 13.8; IR (NaCl dep from CH₂Cl₂) 2973, 1633, 1191, 1063, 762, 698 cm⁻¹.

3-Methyl-5-phenyl-hept-4-enal (14). ¹H NMR (400 MHz CDCl₃) δ 9.74 (1H, t, *J* = 2.4 Hz), 7.32-7.19 (5H, m), 5.39 (1H, d, *J* = 9.8 Hz), 3.14 (1H, m), 2.53 (2H, ddd, *J* = 21.1, 14.3, 7.3 Hz), 2.44 (2H, dd, *J* = 7.0, 2.4 Hz), 1.11 (3H, d, *J* = 6.6 Hz), 0.96 (3H, t, *J* = 7.5 Hz); ¹³C NMR (100 MHz CDCl₃) δ 202.5, 142.7, 141.8, 132.0, 128.4, 127.1, 126.7, 51.4, 28.4, 23.4, 21.7, 14.0; IR (NaCl dep from CH₂Cl₂) 2966, 1725, 1454, 1075, 763, 699 cm⁻¹; HRMS (FAB+) calcd for C₁₄H₁₈O, 202.1358. Found 202.1358.

4-Methyl-2-vinyloxy-oct-3-ene (15sm). ¹H NMR (400 MHz CDCl₃) δ 6.31 (1H, dd, *J* = 14.1, 6.6 Hz), 5.14 (1H, d, *J* = 8.5 Hz), 4.60 (1H, dq, *J* = 13.4, 6.6 Hz), 4.25 (1H, d, *J* = 14.3 Hz), 3.96 (1H, d, *J* = 6.6 Hz), 2.00 (2H, d, *J* = 7.2 Hz), 1.66 (3H, s), 1.44-1.28 (4H, m), 1.26 (3H, d, *J* = 6.2 Hz), 0.89 (3H, t, *J* = 7.3 Hz); ¹³C NMR (100 MHz CDCl₃) δ 150.7, 138.9, 126.3, 88.4, 72.8, 39.3, 30.0, 22.5, 21.3, 16.7, 14.2; IR (NaCl dep from CH₂Cl₂) 2930, 1633, 1456, 1194, 1065, 811 cm⁻¹.

3,5-Dimethyl-non-4-enal (15). ¹H NMR (400 MHz CDCl₃) δ 9.68 (1H, t, *J* = 2.6 Hz), 4.94 (1H, d, *J* = 9.4 Hz), 3.01-2.89 (1H, m), 2.37-2.29 (2H, m), 1.94 (3H, t, *J* = 7.0 Hz), 1.62 (3H, s), 1.39-1.18 (4H, m), 1.01 (3H, d, *J* = 6.8 Hz), 0.88 (3H, t, *J* = 7.2 Hz); ¹³C NMR (100 MHz CDCl₃) δ 203.3, 135.7, 128.7, 51.5, 39.5, 30.3, 26.3, 23.0, 21.5, 16.4, 14.3; IR (NaCl dep from CH₂Cl₂) 2958, 2872, 2716, 1727, 1456, 1380 cm⁻¹; HRMS (FAB+) calcd for C₁₁H₂₀O, 168.1592. Found 168.1591.

(5-Methyl-3-vinyloxy-non-4-enyl)-benzene (16sm). ¹H NMR (400 MHz CDCl₃) δ 7.31-7.16 (5H, m), 6.31 (1H, dd, *J* = 14.3, 6.6 Hz), 5.14 (1H, dd, *J* = 8.7, 1.1 Hz), 4.41 (1H, dt, *J* = 8.7, 6.6

Hz), 4.24 (1H, dd, $J = 14.1, 1.3$ Hz), 3.96 (1H, dd, $J = 6.6, 1.3$ Hz), 2.71-2.64 (2H, m), 2.08-1.98 (3H, m), 1.83-1.72 (1H, m), 1.61 (3H, s), 1.44-1.24 (4H, m), 0.90 (3H, t, $J = 7.3$ Hz); ^{13}C NMR (100 MHz CDCl_3) δ 150.8, 142.0, 140.1, 128.6, 128.6, 126.0, 125.1, 88.5, 75.9, 39.4, 37.1, 31.6, 30.1, 22.5, 16.9, 14.2; IR (NaCl dep from CH_2Cl_2) 2929, 1632, 1455, 1193, 822, 699 cm^{-1} .

5-Methyl-3-phenethyl-non-4-enal (16). ^1H NMR (400 MHz CDCl_3) δ 9.65 (1H, t, $J = 2.3$ Hz), 7.30-7.13 (5H, m), 4.94 (1H, d, $J = 10.0$ Hz), 2.90-2.79 (1H, m), 2.72-2.60 (2H, m), 2.43-2.28 (2H, m), 2.01 (2H, t, $J = 7.0$ Hz), 1.60 (3H, s), 1.45-1.1 (6H, m), 0.91 (3H, t, $J = 7.3$ Hz); ^{13}C NMR (100 MHz CDCl_3) δ 203.1, 142.4, 138.1, 128.7, 128.6, 128.5, 126.0, 50.1, 39.7, 37.9, 33.7, 33.0, 30.4, 22.5, 16.8, 14.3; IR (NaCl dep from CH_2Cl_2) 2929, 2857, 1724, 1454, 1031, 699 cm^{-1} ; HRMS (FAB+) calcd for $\text{C}_{18}\text{H}_{26}\text{O}$, 258.1984. Found 258.1971.

(3-Isopropenyloxy-1-methyl-but-1-enyl)-benzene (17). ^1H NMR (400 MHz CDCl_3) δ 7.45-7.21 (5H, m), 5.77 (1H, dd, $J = 7.7, 1.1$ Hz), 4.88 (1H, dq, $J = 12.8, 6.6$ Hz), 3.87 (2H, d, $J = 13.9$ Hz), 2.09 (3H, s), 1.83 (3H, s), 1.39 (3H, d, $J = 6.6$ Hz); ^{13}C NMR (100 MHz CDCl_3) δ 158.5, 142.9, 135.5, 130.9, 128.4, 127.3, 125.9, 82.8, 70.8, 21.7, 20.9, 16.3; IR (NaCl dep from CH_2Cl_2) 2978, 1655, 1445, 1275, 1079, 696 cm^{-1} .

4-Methyl-6-phenyl-hept-5-en-2-one (18). ^1H NMR (400 MHz CDCl_3) δ 7.38-7.18 (5H, m), 5.54 (1H, dd, $J = 9.5, 1.1$ Hz), 3.19-3.03 (1H, m), 2.48 (2H, dd, $J = 6.6, 3.3$ Hz), 2.14 (3H, s), 2.07 (3H, s), 1.07 (3H, d, $J = 6.6$ Hz); ^{13}C NMR (100 MHz CDCl_3) δ 208.5, 143.8, 134.5, 132.9, 128.4, 127.0, 125.9, 51.3, 30.8, 29.8, 21.1, 16.2; IR (NaCl dep from CH_2Cl_2) 2960, 1716, 1445, 1358, 1165, 698; HRMS (FAB+) calcd for $\text{C}_{14}\text{H}_{18}\text{O}$, 202.1358. Found 202.1357.

Deuterium Crossover Experiment: Conducted according to the general procedure. The products were analyzed by GC-MS. Integration of the individual mass ions showed the following results:

Spectra: