Supporting Information

to accompany

Remote Asymmetric Induction with Vinylketene Silyl *N,O*-acetal

**Shin-ichi Shirokawa, Masato Kamiyama, Tomoaki Nakamura,
Masakazu Okada, Atsuo Nakazaki, Seiji Hosokawa,
Susumu Kobayashi***

*Faculty of Pharmaceutical Sciences, Tokyo University of Science (RIKADAI),
2641 Yamazaki, Nodashi, Chiba 278-8510, Japan*

Experimental Section

General.

IR spectra were recorded on a JASCO FT/IR-410. ^1H and ^{13}C NMR spectra were recorded on a JEOL JNM-LD400 or JNM-LD500 spectrometer in CDCl_3 or C_6D_6 as a solvent. Tetramethylsilane (TMS) served as internal standard (δ) for ^1H NMR. CDCl_3 (δ 7.0) or C_6D_6 (δ 28.0) were used as internal references for ^{13}C NMR. HPLC was carried out using a GL Science PU614, UV620, CO630N-10 and DG660-2. Optical rotations were recorded on a JASCO P-1030. Flash column chromatography was performed on PSQ 100B (Fuji Silysia Co., Ltd., Japan). Analytical thin-layer chromatography was performed on Silica gel 60 F_{254} plates (Merck). Preparative thin layer chromatography was performed on Wakogel B-5F. Mass Spectra were recorded on a Applied Biosystems mass spectrometer (API QSTAR pulsar i) under conditions as high resolution, using poly(ethylene glycol) as internal standard. Melting points were recorded on Yanaco MP-3S. All air and water sensitive reactions were performed in flame-dried glassware.

General Procedure A for preparation of vinylketene silyl *N,O*-acetal.

To a solution of imide (14.2 mmol) in THF (150 ml) was added NaHMDS (21.1 ml of a 1.01 M solution in THF, 21.3 mmol) at 478 °C. After the mixture had been stirred for 90 min at 478 °C, a solution of TBSCl (6.42 g, 42.6 mmol) in THF (30 ml) was added at 478 °C. Stirring was continued for 30 min at 478 °C, then the reaction was quenched with a saturated aq. NH_4Cl , and extracted with ethyl acetate. The extract was washed with water and brine, dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (hexane / ethyl acetate = 5 / 1) to give the vinylketene silyl *N,O*-acetal.

(4*S*)-3-[(1*E*)-1-(*tert*-Butyldimethylsilyloxy)-2-methylbuta-1,3-dienyl]-4-isopropylloxazolidin-2-one **4**.

The title compound was prepared according to the General Procedure A and isolated by column chromatography as a colorless solid (85%): m.p. 48.7 °C, IR (neat) 3509, 2957, 2859, 1763, 1648, 1472, 1255, 784 cm^{-1} ; ^1H NMR (400 MHz) δ 0.14-0.18 (3H, br s), 0.20 (3H, s), 0.93 (6H, d, $J=6.8\text{Hz}$), 0.99 (9H, s), 1.80 (3H, s), 1.95 (1H, sept d, $J=6.8, 2.2\text{Hz}$), 3.97-4.07 (1H, m), 4.12 (1H, dd, $J=8.3, 8.8\text{Hz}$), 4.32 (1H, t, $J=8.8\text{Hz}$), 5.04 (1H, d, $J=10.7\text{Hz}$), 5.15 (1H, d, $J=17.1\text{Hz}$), 6.54 (1H, dd, $J=10.7, 17.1\text{Hz}$); ^{13}C NMR (100 MHz) δ 44.7, 44.2, 11.7,

16.4, 18.2, 18.4, 25.8, 29.5, 59.5, 64.6, 112.5, 115.3, 134.0, 136.7, 155.9; $[\alpha]_{\text{D}}^{25} = 465.7^\circ$ (c 0.87, CHCl_3); HRMS calcd for $\text{C}_{17}\text{H}_{31}\text{NO}_3\text{Na}$ ($[\text{M}+\text{Na}]^+$). 348.1970. found 348.1966.

(4*S*)-3-[(1*Z*)-1-(*tert*-Butyldimethylsilyloxy) buta-1,3-dienyl]-4-isopropylloxazolidin-2-one **7**.

The title compound was prepared according to the General Procedure A and isolated by column chromatography as a colorless oil (63%): IR (neat) 3506, 2959, 2932, 2896, 2860, 1761, 1657, 1472, 1256, 784 cm^{-1} ; ^1H NMR (400 MHz) δ 0.17 (3H, s), 0.21 (3H, s), 0.90 (6H, d, $J=7.1\text{Hz}$), 0.99 (9H, s), 2.07 (1H, sept d, $J=7.1, 3.7\text{Hz}$), 4.00 (1H, ddd, $J=3.7, 4.9, 8.8\text{Hz}$), 4.11 (1H, dd, $J=4.9, 9.0\text{Hz}$), 4.25 (1H, dd, $J=8.8, 9.0\text{Hz}$), 4.98 (1H, dd, $J=1.5, 10.7\text{Hz}$), 5.16 (1H, dd, $J=1.5, 17.3\text{Hz}$), 5.65 (1H, d, $J=10.5\text{Hz}$), 6.49 (1H, ddd,

$J=10.5, 10.7, 17.3\text{Hz}$); ^{13}C NMR (100 MHz) δ 44.7, 43.6, 15.1, 17.6, 18.1, 25.5, 29.0, 59.1, 63.1, 108.8, 114.9, 130.4, 139.1, 155.1; $[\zeta]_{\text{D}}^{25} = 484.5\#(c\ 1.17, \text{CHCl}_3)$; HRMS calcd for $\text{C}_{16}\text{H}_{29}\text{NO}_3$ ($[\text{M}]^+$). 311.1917. found 311.1921.

(4*S*)-3-[(1*E*,3*E*)-1-(*tert*-Butyldimethylsilyoxy)-2-methylpenta-1,3-dienyl]-4-isopropylloxazolidin-2-one **10**.

10

The title compound was prepared according to the General Procedure A and isolated by column chromatography as a colorless oil (90%): IR (neat) 3503, 2958, 2931, 2858, 1762, 1652, 1631, 1472, 1286, 784 cm^{-1} ; ^1H NMR (400 MHz) δ 0.14 (3H, s), 0.19 (3H, s), 0.93 (6H, d, $J=7.1\text{Hz}$), 0.98 (9H, s), 1.75-1.82 (3H, m), 1.78 (3H, d, $J=6.6\text{Hz}$), 1.88-2.04 (1H, m), 3.95-4.04 (1H, m), 4.13 (1H, dd, $J=8.3, 8.6\text{Hz}$), 4.32 (1H, dd, $J=8.6, 8.8\text{Hz}$), 5.63 (1H, qd, $J=6.6, 15.4\text{Hz}$), 6.21 (1H, d, $J=15.4\text{Hz}$); ^{13}C NMR (100 MHz) δ 44.9, 44.3, 12.3, 16.3, 18.0, 18.3, 18.8, 25.6, 29.3, 59.4, 64.4, 115.0, 124.4, 128.1, 134.6, 155.9; $[\zeta]_{\text{D}}^{25} = 450.4\#(c\ 0.84, \text{CHCl}_3)$; HRMS calcd for $\text{C}_{18}\text{H}_{33}\text{NO}_3$ Na ($[\text{M}+\text{Na}]^+$). 362.2127. found 362.2129.

General Procedure B for Vinylogous Mukaiyama aldol reaction of with vinylketene silyl *N,O*-acetal **4** with aldehyde.

To a solution of aldehyde (2.46 mmol) in CH_2Cl_2 (6.0 ml) was added TiCl_4 (1.23 ml of a 1.0 M solution in CH_2Cl_2 , 1.23 mmol) and a solution of **4** (400 mg, 1.23 mmol) in CH_2Cl_2 (6.0 ml) at 478 $^\circ\text{C}$. After stirring for 6 hr at 478 $^\circ\text{C}$, the reaction was quenched with pyridine. After a saturated aq. Rochelle Salt and a saturated aq. NaHCO_3 were added, the mixture was warmed to room temperature and stirred vigorously until the resulting white slurry was completely dissolved, and extracted with ethyl acetate. The extract was washed with water and brine, dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (hexane / ethyl acetate = 5 / 1) to give the aldol adducts.

(4*S*)-3-[(2*E*,5*R*)-5-Hydroxy-2-methyldec-2-enoyl]-4-isopropylloxazolidin-2-one **5a**.

5a

The title compound was prepared according to the General Procedure B and isolated by column chromatography as a colorless oil (97 %): IR (neat) 3524, 2961, 2930, 2860, 1779, 1685, 1466, 1296, 1209, 1119, 755 cm^{-1} ; ^1H NMR (400 MHz) δ 0.89 (3H, t, $J=6.6\text{Hz}$), 0.92 (3H, d, $J=6.8\text{Hz}$), 0.94 (3H, d, $J=6.8\text{Hz}$), 1.24-1.56 (8H, m), 1.95 (3H, d, $J=1.5\text{Hz}$), 2.23-2.44 (3H, m), 2.85 (1H, d, $J=3.4\text{Hz}$), 3.66-3.77 (1H, m), 4.19 (1H, dd, $J=5.6, 9.0\text{Hz}$), 4.34 (1H, t, $J=9.0\text{Hz}$), 4.56 (1H, ddd, $J=4.4, 5.6, 9.0\text{Hz}$), 6.03 (1H, qt, $J=1.5, 7.8\text{Hz}$); ^{13}C NMR (100 MHz) δ 3.7, 14.0, 15.1, 17.8, 22.6, 25.6, 28.4, 31.8, 36.8, 36.9, 58.1, 63.4, 70.6, 133.0, 135.3, 154.6, 171.6; $[\zeta]_{\text{D}}^{24} = 233.3\#(c\ 0.97, \text{CHCl}_3)$; HRMS calcd for $\text{C}_{17}\text{H}_{29}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 334.1994. found 334.1990. Diastereoselectivity was determined by HPLC analysis with Mightysil Si-60 (hexane / isopropanol = 60 / 1, 0.3 ml/min, 151 MPa, 254 nm), major diastereomer = 64.5 min, minor diastereomer = 73.4 min; d.s. = 42 : 1.

(4*S*)-3-[(2*E*,5*R*)-5-Hydroxy-2-methylhexadec-2-enoyl]-4-isopropylloxazolidin-2-one **5b**.

The title compound was prepared according to the General Procedure

B and isolated by column chromatography as a colorless oil (92%).;

IR (neat) 3526, 2925, 2854, 1780, 1685, 1466, 1297, 1210, 1119, 756 cm^{-1} ; ^1H NMR (400 MHz) δ 0.88 (3H, t, $J=6.6\text{Hz}$), 0.93 (3H, d, $J=6.8\text{Hz}$), 0.94 (3H, d, $J=6.6\text{Hz}$), 1.24-1.35 (18H, m), 1.40-1.50 (2H,

$J=6.8\text{Hz}$), 1.95 (3H, d, $J=1.5\text{Hz}$), 2.26-2.42 (3H, m), 2.85 (1H, br s), 3.67-3.76 (1H, m), 4.19 (1H, dd, $J=5.4, 9.0\text{Hz}$), 4.34 (1H, dd, $J=8.8,$

9.0Hz), 4.54-4.59 (1H, m), 6.03 (1H, qt, $J=1.5, 8.0\text{Hz}$); ^{13}C NMR (100 MHz, C_6D_6) δ 3.9, 14.3, 14.9, 17.4, 23.1, 26.4, 28.5, 29.8, 30.1 (3 carbons), 30.2 (2 carbons), 32.2, 37.3, 37.5, 57.9, 63.0, 70.6, 133.2, 135.7, 154.3, 171.3; $[\zeta]_{\text{D}}^{23} = 211.9$ (c 1.02, CHCl_3);

HRMS calcd for $\text{C}_{23}\text{H}_{41}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 418.2936. found 418.2933. Diastereoselectivity was determined by HPLC analysis with

Mightysil Si-60 (hexane / isopropanol = 60 / 1, 0.3 ml/min, 162 MPa, 254 nm), major diastereomer = 63.6 min, minor diastereomer = 73.1 min; d.s. = 94 : 1.

(4*S*)-3-[(2*E*,5*S*)-5-Hydroxy-2,6-dimethylhept-2-enoyl]-4-isopropylloxazolidin-2-one **5c**.

The title compound was prepared according to the General Procedure B and isolated by column

chromatography as a colorless oil (78%).; IR (neat) 3524, 2963, 2931, 2876, 1778, 1685, 1468, 1297,

1210, 1119, 754 cm^{-1} ; ^1H NMR (400 MHz) δ 0.93 (3H, t, $J=6.6\text{Hz}$), 0.95 (3H, d, $J=7.1\text{Hz}$), 0.96 (3H, d, $J=6.8\text{Hz}$), 1.00 (3H, d, $J=6.8\text{Hz}$), 1.74 (1H, qqd, $J=6.6, 6.8, 7.1\text{Hz}$), 1.95 (3H, d, $J=1.4\text{Hz}$),

2.26-2.43 (3H, m), 2.88 (1H, br s), 3.41-3.51 (1H, m), 4.19 (1H, dd, $J=5.4, 9.0\text{Hz}$), 4.34 (1H, t,

$J=9.0\text{Hz}$), 4.50-4.59 (1H, m), 6.04 (1H, qt, $J=1.4, 7.6\text{Hz}$); ^{13}C NMR (100 MHz) δ 3.7, 15.1, 17.8, 18.3, 18.8, 28.4, 33.5, 33.7, 58.1, 63.4, 75.4, 132.9, 135.8, 154.3, 171.6; $[\zeta]_{\text{D}}^{24} = 237.4$ (c 1.01, CHCl_3); HRMS calcd for $\text{C}_{15}\text{H}_{25}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 306.1681. found

306.1677. Diastereoselectivity was determined by HPLC analysis with Mightysil Si-60 (hexane / isopropanol = 60 / 1, 1.0 ml/min, 17 MPa, 254 nm), major diastereomer = 21.6 min, minor diastereomer = 25.1 min; d.s. = 40 : 1.

(4*S*)-3-[(2*E*,5*S*,6*E*)-5-Hydroxy-2-methylocta-2,6-dienoyl]-4-isopropylloxazolidin-2-one **5d**.

The title compound was prepared according to the General Procedure B and isolated by column

chromatography as a colorless oil (54% (conversion yield 87%)).; IR (neat) 3423, 2965, 1781,

1682, 1298, 1210, 1118, 774 cm^{-1} ; ^1H NMR (400 MHz) δ 0.93 (6H, t, $J=6.8\text{Hz}$), 1.70 (3H, dd, $J=1.2, 6.3\text{Hz}$), 1.95 (3H, d, $J=1.4\text{Hz}$), 2.32-2.54 (3H, m), 2.70 (1H, d, $J=3.9\text{Hz}$), 4.16-4.26 (1H,

m) 4.20 (1H, dd, $J=5.4, 9.0\text{Hz}$), 4.34 (1H, t, $J=9.0\text{Hz}$), 4.52-4.58 (1H, m), 5.55 (1H, qdd, $J=6.3,$

1.7, 15.3Hz), 5.72 (1H, qd, $J=1.2, 15.3\text{Hz}$), 6.02 (1H, qdd, $J=1.4, 6.6, 9.3\text{Hz}$); ^{13}C NMR (100 MHz) δ 3.9, 15.1, 17.7, 17.9, 28.4, 36.8, 58.2, 63.5, 71.3, 126.8, 132.8, 133.4, 134.3, 154.1, 171.6; $[\zeta]_{\text{D}}^{25} = 221.2$ (c 1.01, CHCl_3); HRMS calcd for $\text{C}_{15}\text{H}_{23}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 304.1514. found 304.1524. Diastereoselectivity was determined by HPLC analysis with Mightysil Si-60 (hexane /

isopropanol = 60 / 1, 0.3 ml/min, 161 MPa, 254 nm), major diastereomer = 170.6 min, minor diastereomer = 190.3 min; d.s. = 20 : 1.

(4*S*)-3-[(2*E*,5*S*,6*E*)-5-Hydroxy-2,6-dimethylnona-2,6-dienoyl]-4-isopropylloxazolidin-2-one **5e**.

5f

The title compound was prepared according to the General Procedure B and isolated by column chromatography as a colorless oil (55% (conversion yield 65%)).; IR (neat) 3511, 2965, 2931, 2874, 1772, 1686, 1301, 1211, 1119, 755 cm^{-1} ; ^1H NMR (400 MHz) δ 0.91 (3H, t, $J=6.8\text{Hz}$), 0.93 (3H, t, $J=6.8\text{Hz}$), 0.96 (3H, t, $J=7.6\text{Hz}$), 1.65 (3H, s), 1.95 (3H, d, $J=1.4\text{Hz}$), 2.04 (2H, qd, $J=7.6, 7.1\text{Hz}$), 2.30-2.41 (2H, m), 2.54 (1H, m), 2.66 (1H, d, $J=4.6\text{Hz}$), 4.13

(1H, dd, $J=4.6, 8.5\text{Hz}$), 4.19 (1H, dd, $J=5.4, 9.0\text{Hz}$), 4.33 (1H, dd, $J=8.8, 9.0\text{Hz}$), 4.51-4.59 (1H, m), 5.45 (1H, t, $J=7.1\text{Hz}$), 6.00 (1H, qdd, $J=1.4, 6.3, 8.8\text{Hz}$); ^{13}C NMR (100 MHz) δ 1.9, 13.8, 14.1, 15.0, 17.8, 20.8, 28.4, 34.9, 58.1, 63.4, 75.6, 128.1, 133.0, 134.9, 135.3, 154.0, 171.6; $[\zeta]_{\text{D}}^{25} = 214.3$ (c 0.34, CHCl_3); HRMS calcd for $\text{C}_{17}\text{H}_{27}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 332.1837. found 332.1838. Diastereoselectivity was determined by HPLC analysis with Mightysil Si-60 (hexane / isopropanol = 60 / 1, 0.3 ml/min, 161 hMPa, 254 nm), major diastereomer = 170.6 min, minor diastereomer = 190.3 min; d.s. = 20 : 1.

(4*S*)-3-[(2*E*,5*S*)-5-Hydroxy-2-methyl-5-phenylpent-2-enoyl]-4-isopropylloxazolidin-2-one **5f**.

5f

The title compound was prepared according to the General Procedure B and isolated by column chromatography as a colorless oil (94%); IR (neat) 3505, 2964, 2926, 2875, 1778, 1684, 1298, 1211, 1119, 756 cm^{-1} ; ^1H NMR (400 MHz) δ 0.93 (3H, d, $J=6.8\text{Hz}$), 0.94 (3H, d, $J=6.8\text{Hz}$), 1.98 (3H, d, $J=1.4\text{Hz}$), 2.37 (1H, sept d, $J=6.8, 4.4\text{Hz}$), 2.54-2.68 (2H, m), 3.27 (1H, br s), 4.21 (1H, dd, $J=5.1, 9.0\text{Hz}$), 4.35 (1H, dd, $J=8.8, 9.0\text{Hz}$), 4.52-4.59 (1H, m), 4.85 (1H, dd, $J=3.7, 8.8\text{Hz}$),

6.04 (1H, qdd, $J=1.4, 6.3, 9.5\text{Hz}$), 7.24-7.43 (5H, m); ^{13}C NMR (100 MHz) δ 3.8, 15.1, 17.8, 28.4, 38.9, 58.3, 63.5, 72.6, 125.6, 127.4, 128.4, 133.2, 133.9, 143.5, 154.3, 171.4; $[\zeta]_{\text{D}}^{23} = 425.7$ (c 1.08, CHCl_3); HRMS calcd for $\text{C}_{18}\text{H}_{23}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 340.1548. found 340.1541. Diastereoselectivity was determined by HPLC analysis with Mightysil Si-60 (hexane / isopropanol = 60 / 1, 1.0 ml/min, 17 MPa, 254 nm), major diastereomer = 32.7min, minor diastereomer = 35.0 min; d.s. = 86 : 1.

(4*S*)-3-[(2*E*,5*S*)-5-Hydroxydec-2-enoyl]-4-isopropylloxazolidin-2-one **8**.

8

To a solution of hexanal 0.15 ml (1.28 mmol) in CH_2Cl_2 (3.0 ml) was added TiCl_4 (0.64 ml of a 1.0 M solution in CH_2Cl_2 , 0.64 mmol) and a solution of **7** (200 mg, 0.64 mmol) in CH_2Cl_2 (3.0 ml) at 478 $^\circ\text{C}$. After stirring for 6.0 hr at 478 $^\circ\text{C}$, the reaction was quenched with pyridine. After a saturated aq. Rochelle Salt and saturated aq. NaHCO_3 were added, the mixture was warmed to room temperature and stirred vigorously until the resulting

white slurry was completely dissolved, and extracted with ethyl acetate. The extract was washed with water and brine, dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (hexane / ethyl acetate = 5 / 1) to give 73 mg (38%) of **8** as a colorless oil. IR (neat) 3486, 3093, 2959, 2930, 1779, 1685, 1634, 1206, 1121, 755 cm^{-1} ; ^1H NMR (400 MHz) δ 0.87 (3H, t, $J=6.8\text{Hz}$), 0.89 (3H, d, $J=6.8\text{Hz}$), 0.91 (3H, d, $J=6.8\text{Hz}$), 1.22-1.36 (6H, m), 1.40-1.54 (3H, m), 2.34-2.52 (3H, m), 3.73-3.83 (1H, m), 4.11 (1H, dd, $J=3.6, 8.4\text{Hz}$), 4.26 (1H, dd, $J=8.4, 9.2\text{Hz}$), 4.50-4.54 (1H,

m), 7.09-7.18 (1H, m), 7.31 (1H, ddd, $J=1.2, 15.2\text{Hz}$); ^{13}C NMR (100 MHz) δ 4.0, 14.7, 18.0, 22.5, 25.2, 28.5, 31.7, 37.2, 40.5, 58.5, 63.4, 70.7, 122.8, 147.2, 154.0, 164.7; HRMS calcd for $\text{C}_{16}\text{H}_{27}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 320.1837. found 320.1834.

General Procedure C for Vinylogous Mukaiyama aldol reaction of with vinylketene silyl *N,O*-acetal **10** with aldehyde.

To a solution of aldehyde (1.17 mmol) in CH_2Cl_2 (3.0 ml) was added TiCl_4 (0.59 ml of a 1.0 M solution in CH_2Cl_2 , 0.59 mmol) and a solution of **10** (200 mg, 0.59 mmol) in CH_2Cl_2 (3.0 ml) at 478°C . After stirring for 19 hr, the reaction was quenched with pyridine. After a saturated aq. Rochelle Salt and a saturated aq. NaHCO_3 were added, the mixture was warmed to room temperature and stirred vigorously until the resulting white slurry was completely dissolved, and extracted with ethyl acetate. The extract was washed with water and brine, dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (hexane / ethyl acetate = 5 / 1) to give the aldol adducts.

(4*S*)-3-[(2*E*,4*S*,5*R*)-5-Hydroxy-2,4-dimethyldec-2-enoyl]-4-isopropylloxazolidin-2-one **11a**.

The title compound was prepared according to the General Procedure C, the reaction mixture was stirred at 478°C , and isolated by column chromatography as a colorless oil (92%); IR (neat) 3521, 2961, 2930, 2872, 1771, 1685, 1300, 1211, 755 cm^{-1} ; ^1H NMR (500 MHz) δ 0.89 (3H, t, $J=6.8\text{Hz}$), 0.93 (3H, d, $J=6.1\text{Hz}$), 0.98 (3H, d, $J=6.8\text{Hz}$), 1.22-1.46 (8H, m), 1.61 (3H, d, $J=7.4\text{Hz}$), 1.94 (3H, s), 2.34 (1H, qd, $J=6.8, 11.6\text{Hz}$), 2.56 (1H, m), 3.02-3.17 (1H, br s), 3.34 (1H, m), 4.19 (1H, dd, $J=5.8, 8.9\text{Hz}$), 4.34 (1H, dd, $J=8.9, 9.2\text{Hz}$), 4.56-4.62 (1H, m), 5.82 (1H, qd, $J=1.3, 10.4\text{Hz}$); ^{13}C NMR (100 MHz) δ 3.9, 14.1, 15.2, 16.1, 17.8, 22.7, 25.3, 28.4, 32.0, 33.8, 40.1, 58.1, 63.4, 75.3, 131.0, 142.4, 154.5, 171.6; $[\zeta]_{\text{D}}^{24} = 221.6$ (c 0.86, CHCl_3); HRMS calcd for $\text{C}_{18}\text{H}_{31}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 348.2150. found 348.2135.

(4*S*)-3-[(2*E*,4*S*,5*R*)-5-Hydroxy-2,4,6-trimethylhept-2-enoyl]-4-isopropylloxazolidin-2-one **11b**.

The title compound was prepared according to the General Procedure C, the reaction mixture was stirred at 478°C , and isolated by column chromatography as a colorless oil (99%); IR (neat) 3528, 2965, 2933, 2875, 1772, 1686, 1300, 1210, 1119, 755 cm^{-1} ; ^1H NMR (400 MHz) δ 0.90-0.97 (12H, m), 1.04 (3H, t, $J=6.8\text{Hz}$), 1.82-1.91 (1H, m), 1.95 (3H, d, $J=1.5\text{Hz}$), 2.35 (1H, sept d, $J=6.8, 4.4\text{Hz}$), 2.64-2.76 (1H, m), 2.96 (1H, d, $J=2.7\text{Hz}$), 3.18 (1H, td, $J=2.7, 8.3\text{Hz}$), 4.19 (1H, dd, $J=5.9, 9.0\text{Hz}$), 4.34 (1H, t, $J=9.0\text{Hz}$), 4.54-4.59 (1H, m), 5.82 (1H, qd, $J=1.5, 10.4\text{Hz}$); ^{13}C NMR (100 MHz) δ 3.9, 14.7, 15.2, 15.8, 17.8, 120.6, 28.4, 29.2, 37.4, 58.1, 63.4, 79.1, 131.0, 142.4, 154.5, 171.7; $[\zeta]_{\text{D}}^{23} = 26.7$ (c 2.00, CHCl_3); HRMS calcd for $\text{C}_{16}\text{H}_{27}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$). 320.1837. found 320.1823.

(4*S*)-3-[(2*E*,4*S*,5*S*,6*E*)-5-Hydroxy-2,4,6-trimethylnona-2,6-dienoyl]-4-isopropylloxazolidin-2-one **11c**.

The title compound was prepared according to the General Procedure C, the reaction mixture was stirred at 440 °C, and isolated by column chromatography as a colorless oil (67% (conversion yield 81%)).; IR (neat) 3510, 2965, 2931, 2874, 1772, 1686, 1301, 1211, 1119, 756 cm⁻¹; ¹H NMR (400 MHz) δ 0.83 (3H, d, *J*=6.8Hz), 0.90-1.00 (9H, m), 1.67 (3H, s), 1.98 (3H, d, *J*=1.6Hz), 2.07 (2H, quint, *J*=6.8Hz), 2.35 (1H, sept d, *J*=6.8, 4.4Hz), 2.69-2.81 (1H, m), 3.36

(1H, br s), 3.65 (1H, d, *J*=9.2Hz), 4.20 (1H, dd, *J*=5.6, 9.2Hz), 4.35 (1H, t, *J*=9.2Hz), 4.55-4.62 (1H, m), 5.39 (1H, t, *J*=6.8Hz), 5.79 (1H, qd, *J*=1.6, 10.4Hz); ¹³C NMR (100 MHz) δ 0.5, 13.9, 13.9, 15.2, 16.2, 17.8, 20.8, 28.4, 29.7, 37.8, 58.0, 63.4, 82.2, 131.4, 131.7, 133.3, 141.9, 154.4, 171.5; [ζ]_D²⁵ = 48.30 (*c* 1.33, CHCl₃); HRMS calcd for C₁₈H₂₉NO₄Na ([M+Na]⁺). 346.1994. found 346.1992.

(4*S*)-3-[(2*E*,4*S*,5*S*)-5-Hydroxy-2,4-dimethyl-5-phenylpent-2-enoyl]-4-isopropylloxazolidin-2-one **11d**.

The title compound was prepared according to the General Procedure C, the reaction mixture was stirred at 430 °C, and isolated by column chromatography as a colorless oil (91%); IR (neat) 3503, 2966, 2931, 2875, 1771, 1686, 1301, 1211, 757 cm⁻¹; ¹H NMR (400 MHz) δ 0.79 (3H, d, *J*=6.8Hz), 0.95 (6H, d, *J*=6.8Hz), 2.02 (3H, d, *J*=1.5Hz), 2.37 (1H, sept d, *J*=6.8, 4.4Hz), 2.80-2.90 (1H, m), 3.74 (1H, br s) 4.22 (1H, dd, *J*=5.9, 9.0Hz), 4.32 (1H, t, *J*=9.0Hz), 4.37 (1H, d, *J*=9.0Hz), 4.57-4.65

(1H, m), 5.89 (1H, qd, *J*=1.5, 10.2Hz), 7.27-7.41 (5H, m); ¹³C NMR (100 MHz) δ 4.1, 15.2, 16.2, 17.9, 28.5, 42.4, 58.1, 63.5, 78.7, 127.2, 127.7, 128.3, 132.2, 141.4, 141.8, 154.6, 171.5; [ζ]_D²⁵ = 465.4 (*c* 0.87, CHCl₃); HRMS calcd for C₁₉H₂₅NO₄Na ([M+Na]⁺). 354.1681. found 354.1683.

Stereochemical Proofs

Aldol adduct **5a** was transformed to 1,3-diacetoxyoctane by ozonolysis, reduction and acetylation, and the sign of its optical rotation compared to the literature value ([ζ]_D²⁴ = 215.2 (*c* 1.7, CHCl₃)).¹ The absolute configuration of aldol aducts **5a-e** was determined by the modified Mosher's method.

Mosher Ester Analysis for 5a.

(S)#MTPA ester

^1H NMR (400 MHz) τ 0.87 (3H, t, $J=7.2\text{Hz}$), 0.87 (3H, d, $J=6.8\text{Hz}$), 0.92 (3H, d, $J=6.8\text{Hz}$), 1.20-1.43 (6H, m), 1.60-1.75 (2H, m), 1.88 (3H, d, $J=1.5\text{Hz}$), 2.35 (1H, sept d, $J=6.8$, 4.2Hz), 2.40-2.57 (2H, m), 3.54 (3H, s), 4.18 (1H, dd, $J=5.1$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.46-4.51 (1H, m), 5.17 (1H, tt, $J=5.6$, 6.8Hz), 5.91 (1H, qt, $J=1.5$, 4.9Hz), 7.37-7.43 (3H, m), 7.47-7.58 (2H, m).

(R)#MTPA ester

^1H NMR (400 MHz) τ 0.83 (3H, t, $J=6.8\text{Hz}$), 0.88 (3H, d, $J=7.1\text{Hz}$), 0.92 (3H, d, $J=7.1\text{Hz}$), 1.10-1.30 (6H, m), 1.57-1.68 (2H, m), 1.91 (3H, d, $J=1.5\text{Hz}$), 2.36 (1H, sept d, $J=6.8$, 4.2Hz), 2.47-2.64 (2H, m), 3.55 (3H, s), 4.18 (1H, dd, $J=5.1$, 9.0Hz), 4.32 (1H, t, $J=9.0\text{Hz}$), 4.46-4.52 (1H, m), 5.19 (1H, tt, $J=5.6$, 7.1Hz), 5.98 (1H, qt, $J=1.5$, 7.1Hz), 7.37-7.43 (3H, m), 7.50-7.53 (2H, m).

Mosher Ester Analysis for 5b.

(S)#MTPA ester

^1H NMR (400 MHz) τ 0.87 (3H, d, $J=6.8\text{Hz}$), 0.90 (3H, t, $J=6.8\text{Hz}$), 0.91 (3H, d, $J=6.8\text{Hz}$), 1.22-1.36 (18H, m), 1.55-1.74 (2H, m), 1.87 (3H, d, $J=1.5\text{Hz}$), 2.36 (1H, sept d, $J=6.8$, 2.8Hz), 2.40-2.57 (2H, m), 3.54 (3H, s), 4.17 (1H, dd, $J=5.1$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.45-4.51 (1H, m), 5.17 (1H, tt, $J=5.9$, 7.1Hz), 5.91 (1H, qt, $J=1.5$, 7.6Hz), 7.37-7.41 (3H, m), 7.50-7.54 (2H, m).

(R)#MTPA ester

^1H NMR (400 MHz) τ 0.88 (3H, d, $J=6.8\text{Hz}$), 0.90 (3H, t, $J=6.8\text{Hz}$), 0.92 (3H, d, $J=6.8\text{Hz}$), 1.14-1.37 (18H, m), 1.55-1.72 (2H, m), 1.91 (3H, d, $J=1.2\text{Hz}$), 2.36 (1H, sept d, $J=6.8$, 4.4Hz), 2.47-2.61 (2H, m), 3.54 (3H, s), 4.18 (1H, dd, $J=5.1$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.46-4.52 (1H, m), 5.19 (1H, tt, $J=6.1$, 6.4Hz), 5.98 (1H, qt, $J=1.2$, 7.3Hz), 7.36-7.41 (3H, m), 7.50-7.55 (2H, m).

Mosher Ester Analysis for 5c.

(S)#MTPA ester

^1H NMR (400 MHz) δ 0.87 (3H, d, $J=7.1\text{Hz}$), 0.92 (3H, d, $J=7.1\text{Hz}$), 0.93 (3H, d, $J=6.6\text{Hz}$), 0.95 (3H, d, $J=6.6\text{Hz}$), 1.88 (3H, d, $J=1.0\text{Hz}$), 2.00 (1H, sept d, $J=6.6$, 5.3Hz), 2.35 (1H, sept d, $J=7.1$, 4.4Hz), 2.43-2.59 (2H, m), 3.52 (3H, s), 4.18 (1H, dd, $J=5.2$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.45-4.51 (1H, m), 5.03 (1H, td, $J=5.3$, 5.8Hz), 5.90 (1H, qt, $J=1.0$, 7.1Hz), 7.37-7.42 (3H, m), 7.50-7.54 (2H, m).

(R)#MTPA ester

^1H NMR (400 MHz) δ 0.86 (3H, d, $J=6.8\text{Hz}$), 0.87 (6H, d, $J=6.8\text{Hz}$), 0.92 (3H, d, $J=6.8\text{Hz}$), 1.92 (3H, d, $J=1.2\text{Hz}$), 1.97 (1H, sept d, $J=6.8$, 5.4Hz), 2.36 (1H, sept d, $J=6.8$, 4.2Hz), 2.47-2.63 (2H, m), 3.54 (3H, s), 4.18 (1H, dd, $J=4.9$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.46-4.53 (1H, m), 5.06 (1H, td, $J=5.4$, 6.6Hz), 5.98 (1H, qt, $J=1.2$, 7.1Hz), 7.37-7.42 (3H, m), 7.51-7.55 (2H, m).

Mosher Ester Analysis for 5d.

(S)#MTPA ester

^1H NMR (400 MHz) δ 0.86 (3H, d, $J=6.8\text{Hz}$), 0.91 (3H, d, $J=6.8\text{Hz}$), 1.72 (3H, dd, $J=1.7$, 6.6Hz), 1.85 (3H, d, $J=1.4\text{Hz}$), 2.34 (1H, sept d, $J=7.1$, 4.4Hz), 2.44-2.54 (1H, m), 2.52-2.61 (1H, m), 3.53 (3H, s), 4.17 (1H, dd, $J=5.2$, 8.8Hz), 4.31 (1H, t, $J=8.8\text{Hz}$), 4.47-4.51 (1H, m), 5.55 (1H, qdd, $J=1.7$, 8.0 , 13.9Hz), 5.50-5.57 (1H, m), 5.89 (1H, qd, $J=6.6$, 14.4Hz), 5.90 (1H, qt, $J=1.4$, 7.3Hz), 7.35-7.43 (3H, m), 7.47-7.53 (2H, m).

(R)#MTPA ester

^1H NMR (400 MHz) δ 0.87 (3H, d, $J=6.8\text{Hz}$), 0.92 (3H, d, $J=6.8\text{Hz}$), 1.68 (3H, dd, $J=1.7$, 6.6Hz), 1.90 (3H, d, $J=1.5\text{Hz}$), 2.35 (1H, sept d, $J=6.8$, 4.2Hz), 2.54 (1H, ddd, $J=7.1$, 7.3 , 15.4Hz), 2.64 (1H, ddd, $J=6.6$, 7.3 , 15.4Hz), 3.53 (3H, s), 4.18 (1H, dd, $J=5.1$, 9.0Hz), 4.31 (1H, dd, $J=8.8$, 9.0Hz), 4.47-4.53 (1H, m), 5.42 (1H, qdd, $J=1.7$, 6.8 , 15.1Hz), 5.52 (1H, ddd, $J=6.6$, 6.8 , 7.1Hz), 5.82 (1H, qd, $J=6.6$, 15.1Hz), 5.97 (1H, qt, $J=1.5$, 7.3Hz), 7.36-7.43 (3H, m), 7.46-7.52 (2H, m).

Mosher Ester Analysis for 5e.

(S)-MTPA ester

¹H NMR (400 MHz) δ 0.88 (3H, d, J =6.6Hz), 0.92 (3H, d, J =6.6Hz), 0.95 (3H, t, J =7.6Hz), 1.62 (3H, s), 1.86 (3H, s), 1.96-2.08 (2H, m), 2.35 (1H, sept d, J =6.6, 4.2Hz), 2.50 (1H, ddd, J =6.8, 7.1, 15.3Hz), 2.61 (1H, ddd, J =6.8, 7.1, 15.3Hz), 3.52 (3H, s), 4.17 (1H, dd, J =3.9, 9.0Hz), 4.31 (1H, t, J =9.0Hz), 4.45-4.52 (1H, m), 5.48 (1H, t, J =7.1Hz), 5.59 (1H, t, J =7.1Hz), 5.84 (1H, t, J =6.8Hz), 7.29-7.39 (3H, m), 7.46-7.55 (2H, m).

(R)-MTPA ester

¹H NMR (400 MHz) δ 0.87 (3H, d, J =6.8Hz), 0.91 (3H, d, J =6.8Hz), 0.93 (3H, t, J =7.6Hz), 1.46 (3H, s), 1.90 (3H, d, J =1.5Hz), 1.94-2.07 (2H, m), 2.35 (1H, sept d, J =6.8, 4.2Hz), 2.53 (1H, td, J =7.1, 15.4Hz), 2.67 (1H, td, J =7.1, 15.4Hz), 3.57 (3H, s), 4.17 (1H, dd, J =4.2, 9.0Hz), 4.31 (1H, dd, J =8.8, 9.0Hz), 4.47-4.52 (1H, m), 5.45 (1H, t, J =7.1Hz), 5.51 (1H, t, J =7.1Hz), 5.91 (1H, qt, J =1.5, 7.1Hz), 7.33-7.39 (3H, m), 7.46-7.54 (2H, m).

Aldol adduct **5f** was transformed to 1,3-Diacetoxy-3-phenylpropane by three-steps sequence ((i) O₃ then Me₂S; (ii) NaBH₄; (iii) Ac₂O, Py, DMAP.), and the sign of its optical rotation compared to the literature value ($[\alpha]_D^{24} = 454.1$ (c 0.55, CHCl₃)).²

The absolute configuration of aldol adduct **8** was determined by the modified Mosher's method.

Mosher Ester Analysis for **8**.

(S)-MTPA ester

¹H NMR (400 MHz) δ 0.84 (3H, t, J =6.8Hz), 0.88 (3H, d, J =6.8Hz), 0.93 (3H, d, J =6.8Hz), 1.12-1.38 (6H, m), 1.56-1.73 (2H, m), 2.40 (1H, sept d, J =6.8, 2.4Hz), 2.65 (2H, dd, J =6.1, 6.4Hz), 3.55 (3H, s), 4.22 (1H, dd, J =2.4, 9.0Hz), 4.28 (1H, dd, J =8.3, 9.0Hz), 4.44-4.51 (1H, m), 5.23 (1H, quint, J =6.1Hz), 7.00-7.09 (1H, m), 7.34 (1H, t, J =15.4 Hz), 7.38-7.43 (3H, m), 7.49-7.54 (2H, m).

(R)-MTPA ester

¹H NMR (400 MHz) δ 0.85 (3H, t, J =7.1Hz), 0.87 (3H, d, J =7.1Hz), 0.93 (3H, d, J =7.1Hz), 1.17-1.40 (6H, m), 1.59-1.76 (2H, m), 2.39 (1H, sept d, J =7.1, 3.6Hz), 2.54-2.68 (2H, m), 3.53 (3H, s), 4.22 (1H, dd, J =2.9, 9.0Hz), 4.28 (1H, dd, J =8.3, 9.0Hz), 4.43-4.52 (1H, m), 5.20-5.26 (1H, m), 6.90-7.00 (1H, m), 7.26 (1H, t, J =16.3Hz), 7.36-7.41 (3H, m), 7.49-7.54 (2H, m).

Aldol adduct **11a** was transformed to methyl ester by three-steps sequence ((i) O₃ then Me₂S; (ii) NaClO₂, NaH₂PO₄; (iii) TMSCHN₂, MeOH), and the sign of its optical rotation compared to the literature value ($[\zeta]_D^{24} = 43.7$ (c 2.03, CHCl₃).³ The absolute configuration of aldol adduct **11a** was checked by the modified Mosher's method.

Mosher Ester Analysis for **11a**.

(*S*)-4-MTPA ester

¹H NMR (400 MHz) δ 0.85 (3H, t, $J=6.8$ Hz), 0.86 (3H, d, $J=6.8$ Hz), 0.90 (3H, d, $J=7.1$ Hz), 0.93 (3H, d, $J=6.8$ Hz), 1.24-1.34 (6H, m), 1.59-1.79 (2H, m), 1.94 (3H, d, $J=1.4$ Hz), 2.33 (1H, sept d, $J=6.8, 4.4$ Hz), 2.83-2.92 (1H, m), 3.56 (3H, s), 4.15 (1H, dd, $J=5.4, 9.0$ Hz), 4.30 (1H, dd, $J=8.8, 9.0$ Hz), 4.46-4.52 (1H, m), 5.06-5.12 (1H, m), 5.87 (1H, qd, $J=1.4, 10.0$ Hz), 7.37-7.42 (3H, m), 7.52-7.57 (2H, m).

(*R*)-4-MTPA ester

¹H NMR (400 MHz) δ 0.84 (3H, t, $J=6.8$ Hz), 0.88 (3H, d, $J=6.8$ Hz), 0.92 (3H, d, $J=7.1$ Hz), 1.03 (3H, d, $J=6.8$ Hz), 1.13-1.30 (6H, m), 1.50-1.70 (2H, m), 1.94 (3H, d, $J=1.5$ Hz), 2.37 (1H, sept d, $J=6.8, 4.4$ Hz), 2.83-2.96 (1H, m), 3.53 (3H, s), 4.17 (1H, dd, $J=5.1, 9.0$ Hz), 4.31 (1H, t, $J=9.0$ Hz), 4.46-4.53 (1H, m), 5.06-5.13 (1H, m), 5.84 (1H, qd, $J=1.5, 9.8$ Hz), 7.36-7.43 (3H, m), 7.52-7.57 (2H, m).

Aldol adduct **11b** was transformed to acetal by ozonolysis and literature procedure, and the ¹H NMR spectrum was identical to that reported.⁴ The absolute configuration of aldol adduct **11b** was determined by the modified Mosher's method.

(3) Watabu, H.; Ohkubo, M.; Matsubara, H.; Sakai, T.; Tsuboi, S.; Utaka, M. *Chem. Lett.* **1989**, 12, 2183.

(4) Harada, T.; Egusa, T.; Igarashi, Y.; Kinugasa, M.; Oku, A. *J. Org. Chem.* **2002**, 67, 7080.

Mosher Ester Analysis for 11b.

(S)-4MTPA ester

¹H NMR (400 MHz) δ 0.87 (3H, d, $J=7.1$ Hz), 0.88 (3H, d, $J=7.1$ Hz), 0.91 (3H, d, $J=7.1$ Hz), 0.94 (3H, d, $J=7.1$ Hz), 0.94 (3H, d, $J=6.8$ Hz), 1.91 (3H, d, $J=1.0$ Hz), 1.97-2.08 (1H, m), 2.35 (1H, sept d, $J=7.1$, 4.4Hz), 2.89-2.98 (1H, m), 3.54 (3H, s), 4.17 (1H, dd, $J=5.4$, 8.8Hz), 4.31 (1H, dd, $J=8.8$, 9.0Hz), 4.46-4.54 (1H, m), 4.95 (1H, dd, $J=3.2$, 8.3Hz), 5.96 (1H, qd, $J=1.0$, 10.0Hz), 7.35-7.42 (3H, m), 7.56-7.62 (2H, m).

(R)-4MTPA ester

¹H NMR (400 MHz) δ 0.83 (3H, d, $J=6.6$ Hz), 0.89 (3H, d, $J=7.1$ Hz), 0.90 (3H, d, $J=6.6$ Hz), 0.92 (3H, d, $J=7.1$ Hz), 1.00 (3H, d, $J=6.8$ Hz), 1.91 (3H, d, $J=1.4$ Hz), 1.90-2.04 (1H, m), 2.39 (1H, sept d, $J=7.1$, 4.4Hz), 2.90-3.00 (1H, m), 3.51 (3H, s), 4.18 (1H, dd, $J=5.1$, 9.0 Hz), 4.31 (1H, dd, $J=8.5$, 9.0Hz), 4.45-4.52 (1H, m), 4.97 (1H, dd, $J=4.2$, 7.6), 5.94 (1H, qd, $J=1.4$, 10.0Hz), 7.37-7.42 (3H, m), 7.54-7.59 (2H, m).

Aldol adduct **11c** was transformed to ketone by ozonolysis and literature procedure, and the ¹H NMR spectrum was identical to that reported.⁵ The absolute configuration of aldol adduct **11c** was determined by the modified Mosher's method.

Mosher Ester Analysis for 11c.

(S)-4MTPA ester

¹H NMR (400 MHz) δ 0.89 (3H, d, $J=6.8$ Hz), 0.92 (3H, d, $J=6.8$ Hz), 0.93 (3H, d, $J=7.1$ Hz), 0.97 (3H, t, $J=7.6$ Hz), 1.65 (3H, d, $J=1.2$ Hz), 1.77 (3H, d, $J=1.4$ Hz), 2.01-2.11 (2H, m), 2.39 (1H, sept d, $J=6.8$, 4.4Hz), 2.86-2.96 (1H, m), 3.52 (3H, s), 4.18 (1H, dd, $J=4.6$, 9.0Hz), 4.30 (1H, dd, $J=8.8$, 9.0Hz), 4.43-4.49 (1H, m), 5.31 (1H, d, $J=8.5$ Hz), 5.61 (1H, qt, $J=1.2$, 7.0Hz), 5.76 (1H, qd, $J=1.4$, 9.8Hz), 7.32-7.36 (3H, m), 7.46-7.50 (2H, m).

(5) Toshima, K.; Jyojima, T.; Yamaguchi, H.; Noguchi, Y.; Yoshida, T.; Murase, H.; Nakata, M.; Matsumura, *J. Org. Chem.* **1997**, 62, 3271.

(*R*)-**11d** MTPA ester

¹H NMR (400 MHz) δ 0.89 (3H, d, $J=7.1$ Hz), 0.91 (3H, d, $J=7.8$ Hz), 0.93 (3H, d, $J=7.1$ Hz), 0.96 (3H, t, $J=7.6$ Hz), 1.40 (3H, d, $J=1.2$ Hz), 1.88 (3H, d, $J=1.5$ Hz), 1.98-2.09 (2H, m), 2.39 (1H, sept d, $J=7.1, 4.2$ Hz), 2.84-2.92 (1H, m), 3.52 (3H, s), 4.19 (1H, dd, $J=4.6, 9.0$ Hz), 4.31 (1H, dd, $J=8.8, 9.0$ Hz), 4.44-4.50 (1H, m), 5.17 (1H, d, $J=9.2$ Hz), 5.58 (1H, qt, $J=1.2, 7.3$ Hz), 5.79 (1H, qd, $J=1.5, 10.0$ Hz), 7.34-7.38 (3H, m), 7.45-7.50 (2H, m).

Aldol adduct **11d** was transformed to diol by two-steps sequence ((i) O₃ then Me₂S; (ii) NaBH₄), and the sign of its optical rotation compared to the literature value ($[\zeta]_D^{25} = 440.8$ (c 1.04, CHCl₃).⁶

(6) Abiko, A.; Liu, J.; Masamune, S. *J. Am. Chem. Soc.* **1997**, *119*, 2586.

7

7

10

10

5a

5b

5c

5c

5d

5d

FILE C:\ALICE\DATA\SS_VMA\NMR\SS (04JACS)\ss667_TC.up02\ss6
 5e
 COMINT Wed Sep 1 14:14:06 2004
 DATIM 1H
 OBNUC 1H
 EXMOD non
 OBRQ 399.65 MHz
 OBSSET 130.00 KHz
 OBFIN 4300.00 Hz
 POINT 32768
 FREQU 8000.00 Hz
 SCANS 16
 ACQTM 4.0960 sec
 PD 2.9040 sec
 PW1 5.50 usec
 IRNUC 1H
 CTEMP 23.5 c
 SLVNT CDCL3
 EXREF 0.00 ppm
 BF 0.12 Hz
 RGAIN 24

5e

5f

11b

DILE	C:\ALICE\DATA\SS_VMA\NMR\SS (04JACS)*ss668_TC_up02*ss668
COMINT	11c
DATIM	Wed Sep 1 18:10:39 2004
OBNUC	1H
EXMOD	non
OFFRQ	399.65 MHz
OBSF1	130.00 KHz
OBSF1N	4300.00 Hz
POINT	32/768
FREQU	8000.00 Hz
SCANS	32
ACQTM	4.0960 sec
PD	2.9040 sec
PW1	5.50 usec
IRNUC	1H
CTEMP	23.6 c
SLVNT	CDCl3
EXREF	0.00 ppm
BF	0.12 Hz
RGAIN	22

11c

11c