SUPPORTING INFORMATION

Rhodium-catalyzed imination of sulfoxides and sulfides: Efficient preparation of *N*-unsubstituted sulfoximines and sulfilimines

Hiroaki Okamura and Carsten Bolm*

*Institut für Organische Chemie der RWTH Aachen, Professor-Pirlet-Straße 1, D-
52056 Aachen, Germany*

e-mail: Carsten.Bolm@oc.rwth-aachen.de

¹H and ¹³C NMR (400 and 100 MHz, respectively) spectra were recorded in CDCl₃ or DMSO-d₆ using TMS as internal standard. Chemical shifts are given in ppm and spin-spin coupling constants, *J*, are given in Hz. IR spectra were recorded as KBr pellets. The solvents (CH₂Cl₂ and MeOH) were used after single distillation without drying. All reactions were performed under atmospheric condition.

General procedure for the imination of sulfoxides (given in Table 1): To a suspension of the sulfoxide (1.0 mmol), sulfonylamide or trifluoroacetamide (2.0 mmol), MgO (160 mg, 4.0 mmol), and Rh₂(OAc)₄ (11 mg, 2.5 mol%) in CH₂Cl₂ (10 mL) was added PhI(OAc)₄ (483 mg, 1.5 mmol) at room temperature. The resulting mixture was stirred for the period of 6-12 h as indicated in Table 1. Purification by column chromatography (ca. 70 g of silica gel 60, 0.063-0.100 mm, column diameter 27 mm) afforded the sulfoximines.

***N*-(4-nitro-benzenesulfonyl)methylphenylsulfoximine (2a)**¹: Yield: 293 mg (86%); white solid, mp. 148-151 °C; *R*_f 0.30 (hexane : ethyl acetate = 1 : 1); ¹H NMR (DMSO-d₆): δ 8.37 (d, *J* = 8.5 Hz, 2H), 8.03 (bd, *J* = 7.9 Hz, 2H), 7.99 (bd, *J* = 8.5 Hz, 2H), 7.81 (t, *J* = 7.4 Hz, 1H), 7.70 (t, *J* = 7.9 Hz, 2H), 3.71 (s, 3H); ¹³C NMR (DMSO-d₆): δ 149.7 (C), 149.0 (C), 138.0 (C), 135.0 (CH), 130.1 (2 x CH), 128.1 (2 x CH), 127.7 (2 x CH), 124.8 (2 x CH), 45.3 (CH₃); IR (KBr): ν 1525, 1352, 1323, 1155, 1106, 1074, 746 cm⁻¹; MS (EI), *m/z* (relative intensity): 340 ([M]⁺, 4.5 %), 325 ([M-CH₃]⁺, 100 %); Calcd. for C₁₃H₁₂N₂O₅S₂: C, 45.87; H, 3.55; N, 8.23; found C, 45.85; H, 3.88; N, 8.26.

***N*-(4-methylsulfonyl)methylphenylsulfoximine (2b)**: Yield: 185 mg (79%); white solid, mp. 113-115 °C; *R*_f 0.52 (ethyl acetate); ¹H NMR (CDCl₃): δ 8.05 (bd *J* = 8.6 Hz, 2H), 7.74 (tt, *J* = 1.2, 7.6 Hz, 1H), 7.65 (bt, *J* = 8.0 Hz, 2H), 3.45 (s, 3H), 3.15 (s, 3H); ¹³C NMR (CDCl₃): δ 138.0 (C), 134.6 (CH), 129.8 (2 x CH), 127.4 (2 x CH), 46.5 (CH₃), 45.3 (CH₃); IR (KBr): ν 1294, 1227, 1136, 1100, 1076 cm⁻¹; MS (EI), *m/z* (relative intensity): 233 ([M]⁺, 2 %), 218 ([M-CH₃]⁺, 100 %); Calcd. for C₈H₁₁NO₃S₂: C, 41.18; H, 4.75; N, 6.00; found C, 41.26; H, 4.76; N, 5.96.

***N*-(trifluoroacetyl)methylphenylsulfoximine (2c)**: Yield: 211 mg (84%); white solid, mp. 76.5-77.5 °C; R_f 0.40 (hexane : ethyl acetate = 1 : 1); ^1H NMR (CDCl_3): δ 8.00 (bd, $J = 7.7$ Hz, 2H), 7.76 (tt, $J = 1.2, 7.4$ Hz, 1H), 7.66 (bt, $J = 7.4$ Hz, 2H), 3.46 (s, 3H); ^{13}C NMR (CDCl_3): δ 163.9 (CO, q, $J = 32$ Hz), 136.5 (C), 134.7 (CH), 129.9 (2 x CH), 126.9 (2 x CH), 115.8 (CF_3 , q, $J = 282$ Hz), 44.2 (CH_3); IR (KBr): ν 1663, 1242, 1189, 1131, 508 cm^{-1} ; MS (EI), m/z (relative intensity): 182 ($[\text{M}-\text{CF}_3]^+$, 100 %); Calcd. for $\text{C}_9\text{H}_8\text{NO}_2\text{F}_3\text{S}$: C, 43.03; H, 3.21; N, 5.58; found C, 43.31; H, 3.48; N, 5.52.

***N*-(4-nitro-benzenesulfonyl)diphenylsulfoximine (4a)**: Yield: 272 mg (68%); white solid, mp. 154-156 °C; R_f 0.71 (CH_2Cl_2 : ethyl acetate = 4 : 1); ^1H NMR ($\text{DMSO}-d_6$): δ 8.40-8.32 (m, 2H), 8.10-7.94 (m, 5H), 7.80-7.60 (m, 7H); ^{13}C NMR (CDCl_3): δ 149.6 (C), 149.1 (C), 139.3 (2 x C), 134.3 (2 x CH), 129.8 (4 x CH), 128.1 (2 x CH), 127.6 (4 x CH), 124.0 (2 x CH); IR (KBr): ν 1535, 1325, 1242, 1157, 1094, 1059 cm^{-1} ; MS (EI), m/z (relative intensity): 402 ($[\text{M}]^+$, 9 %), 125 ($[\text{Ph}-\text{SO}]^+$, 100 %); Calcd. for $\text{C}_{18}\text{H}_{14}\text{N}_2\text{O}_5\text{S}_2$: C, 53.72; H, 3.51; N, 6.96; found C, 53.78; H, 3.56; N, 6.77.

***N*-(4-nitro-benzenesulfonyl)dimethylsulfoximine (6a)²**: Yield: 140 mg (50%); white solid, mp. 226.5-228.5 °C, lit.² mp. 224-226 °C; R_f 0.33 (ethyl acetate); ^1H NMR ($\text{DMSO}-d_6$): δ 8.37 (d, $J = 8.8$ Hz, 2H), 8.11 (d, $J = 8.8$ Hz, 2H), 3.52 (s, 6H); ^{13}C NMR ($\text{DMSO}-d_6$): δ 149.7 (C), 149.6 (C), 128.1 (2 x CH), 124.7 (2 x CH), 43.4 (2 x CH_3); IR (KBr): ν 1532, 1351, 1300, 1221, 1161, 1081, 613 cm^{-1} ; MS (EI), m/z (relative intensity): 278 ($[\text{M}]^+$, 45 %).

***N*-(4-nitro-benzenesulfonyl)-*t*-butylmethylsulfoximine (8a)**: Yield: 99 mg (49%); White solid, mp. 156 °C (decomp.); R_f 0.49 (ethyl acetate); ^1H NMR (CDCl_3): δ 8.32 (dt, $J = 2.0, 9.1$ Hz, 2H), 8.15 (dt, $J = 2.0, 9.1$ Hz, 2H), 3.40 (s, 3H), 1.47 (s, 9H); ^{13}C NMR (CDCl_3): δ 149.5 (C), 149.3 (C), 127.9 (2 x CH), 124.0 (2 x CH), 62.5 (C), 35.4 (CH_3), 22.7 (3 x CH_3); IR (KBr): ν 1525, 1352, 1303, 1154, 1095, 1073, 743, 612 cm^{-1} ; MS (EI), m/z (relative intensity): 305 ($[\text{M}-\text{CH}_3]^+$, 1 %), 264 ($[\text{M}-\text{C}_4\text{H}_8]^+$, 17 %); HRMS of $[\text{M}-\text{C}_4\text{H}_8]^+$: Calcd. 263.9875; found 268.9875.

General procedure for the preparation of *N*-unsubstituted sulfoximines (given in Table 2): To a suspension of the sulfoxide (1.0 mmol), trifluoroacetamide (226 mg, 2.0 mmol), MgO (160 mg, 4.0 mmol), and $\text{Rh}_2(\text{OAc})_4$ (11 mg, 2.5 mol%) in CH_2Cl_2 (10 mL) was added $\text{PhI}(\text{OAc})_4$ (483 mg, 1.5 mmol) at room temperature, and the mixture was stirred for 6 h. The resulting suspension was filtered through a pad of celite, and the filtrate was concentrated under reduced pressure (rotary evaporator). To the solution of the resulting residue in MeOH (10 mL) was added K_2CO_3 (690 mg, 5 mmol) at room temperature, and the mixture was stirred for 30 min. The reaction mixture was concentrated under reduced pressure, and the remaining product purified by column chromatography (ca. 70 g of silica gel 60, 0.063-0.100 mm, column diameter 27 mm) to give the *N*-unsubstituted sulfoximines.

Methylphenylsulfoximine (9)³: Yield 110 mg (71%); colorless oil; R_f 0.40 (ethyl acetate : ethanol = 4 : 1); ^1H NMR (CDCl_3): δ 8.05-8.01 (m, 2H), 7.68-7.55 (m, 3H),

3,12 (s, 3H), 2.78 (br s, 1H, NH); ^{13}C NMR (CDCl_3): δ 143.3 (C), 132.8 (CH), 129.0 (2 x CH), 127.5 (2 x CH), 46.1 (CH_3).

Diphenylsulfoximine (10)⁴: Yield 143 mg (66%); white solid, mp. 100-102 °C, lit.⁴ mp. 103 °C; ^1H NMR (CDCl_3): δ 8.07-8.03 (m, 4H), 7.54-7.44 (m, 6H), 2.66 (br s, 1H for NH); ^{13}C NMR (CDCl_3): δ 143.3 (2 x C), 132.6 (2 x CH), 129.2 (4 x CH), 127.9 (4 x CH).

S-Oxo-S-imino-tetrahydrothiophene (12)⁵: Yield: 105 mg (88%); colorless oil; R_f 0.50 (ethyl acetate); ^1H NMR (CDCl_3): δ 3.17-3.11 (m, 2H), 2.70 (bs, 1H, NH), 2.29-2.25 (m, 2H); ^{13}C NMR (CDCl_3): δ 55.3 (2 x CH_2), 23.9 (2 x CH_2); IR (KBr): ν 3265, 1417, 1274, 1210, 1145, 1006 cm^{-1} ; MS (EI), m/z (relative intensity): 119 ($[\text{M}]^+$, 20 %).

(-)-(R)-Methyl-*p*-toluylsulfoximine (14)⁶: Yield: 129 mg (76%); colorless oil; $[\alpha]_{\text{D}}^{25} = -30$ (c 2.0, acetone), lit.⁶ $[\alpha]_{\text{D}}^{25} = -31.9$ (c 3.00, acetone); R_f 0.32 (ethyl acetate : ethanol = 10 : 1); ^1H NMR (CDCl_3): δ 7.89 (dd, $J = 1.7, 8.5$ Hz, 2H), 7.55 (dd, $J = 1.7, 8.5$ Hz, 2H), 3.10 (s, 3H), 2.73 (bs, 1H for NH), 2.46 (s, 3H); ^{13}C NMR (CDCl_3): δ 143.8 (C), 140.5 (C), 129.8 (2 x CH), 127.7 (2 x CH), 46.4 (CH_3), 21.6 (CH_3); Chiral HPLC (DAICEL CHIRALCEL OJ, 15 % *i*-PrOH in heptane, 0.5 mL/min) 99% ee, (*R*)-isomer (major): 32 min, (*S*)-isomer (minor): 45 min.

General procedure for the imination of sulfides (given in Table 3): To a suspension of the sulfide (1.0 mmol), sulfonylamide or trifluoroacetamide (2.0 mmol), MgO (160 mg, 4.0 mmol), and $\text{Rh}_2(\text{OAc})_4$ (2.5 mol%, 11 mg) in CH_2Cl_2 (10 mL) was added $\text{PhI}(\text{OAc})_4$ (483 mg, 1.5 mmol) at room temperature, and the mixture was stirred for 6 h. The resulting suspension was purified by column chromatography (ca. 70 g of silica gel 60, 0.063-0.100 mm, column diameter 27 mm) to give the corresponding sulfilimine.

***N*-(4-nitro-benzenesulfonyl)methylphenylsulfilimine (16a)**: Yield: 292 mg (90%); white solid, mp. 164.5-165.5 °C; R_f 0.42 (ethyl acetate); ^1H NMR (CDCl_3): δ 8.21 (dt, $J = 2.0, 9.0$ Hz, 2H), 8.00 (dt, $J = 2.0, 9.0$ Hz, 2H), 7.70 (bd, $J = 7.1$ Hz, 2H), 7.57 (tt, $J = 1.2, 7.1$ Hz, 1H), 7.51 (bt, $J = 7.3$ Hz, 2H), 2.94 (s, 3H); ^{13}C NMR (CDCl_3): δ 149.8 (C), 149.1 (C), 135.1 (C), 133.0 (CH), 130.2 (2 x CH), 127.4 (2 x CH), 125.9 (2 x CH), 124.0 (2 x CH), 39.3 (CH_3); IR (KBr): ν 1523, 1350, 1291, 1145, 1088, 998, 959, 615 cm^{-1} ; MS (EI), m/z (relative intensity): 324 ($[\text{M}]^+$, 10 %); Calcd. for $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_4\text{S}_2$: C, 48.14; H, 3.73; N, 8.64; found C, 48.19; H, 3.79; N, 8.56.

***N*-(trifluoroacetyl)methylphenylsulfilimine (16b)**: Yield: 205 mg (87%); white solid, mp. 84.5-85.5 °C; R_f 0.45 (ethyl acetate); ^1H NMR (CDCl_3): δ 7.83-7.77 (m, 2H), 7.67-7.54 (m, 2H), 2.95 (s, 3H); ^{13}C NMR (CDCl_3): δ 166.7 (CO, q, $J = 34$ Hz), 133.2 (CH), 133.0 (C), 130.4 (2 x CH), 127.0 (2 x CH), 117.0 (CF_3 , q, $J = 288$ Hz), 34.7 (CH_3); IR (KBr): ν 1628, 1190, 1165, 1136 cm^{-1} ; MS (EI), m/z (relative intensity): 235 ($[\text{M}]^+$, 38 %), 166 ($[\text{M}-\text{CF}_3]^+$, 78 %); Calcd. for $\text{C}_9\text{H}_8\text{F}_3\text{NOS}$: C, 45.95; H, 3.43; N, 5.95; found C, 46.12; H, 3.50; N, 6.00.

***N*-(4-nitro-benzensulfonyl)diphenylsulfilimine (18a)**: Yield: 378 mg (98%); white solid, mp. 160.5-161.5 °C; R_f 0.36 (hexane : ethyl acetate = 1 : 1); ^1H NMR (CDCl_3): δ 8.19 (dt, J = 2.1, 8.9 Hz, 2H), 8.02 (dt, J = 2.1, 8.9 Hz, 2H), 7.64 (m, 4H), 7.58-7.44 (m, 6H); ^{13}C NMR (CDCl_3): δ 150.1 (C), 149.1 (C), 135.8 (2 x C), 132.8 (2 x CH), 130.2 (4 x CH), 127.5 (2 x CH), 127.2 (4 x CH), 124.0 (2 x CH); IR (KBr): ν 1562, 1349, 1300, 1150, 1087, 962, 749, 687, 606 cm^{-1} ; MS (EI), m/z (relative intensity): 386 ($[\text{M}]^+$, 1 %), 200 ($[\text{M}-\text{Ns}]^+$, 100 %); Calcd. for $\text{C}_{18}\text{H}_{14}\text{N}_2\text{O}_4\text{S}_2$: C, 55.94; H, 3.65; N, 7.25; found C, 55.76; H, 3.55; N, 7.14.

***N*-(trifluoroacetyl)diphenylsulfilimine (18b)**⁷: Yield: 273 mg (98%); white solid, mp. 68-70 °C; R_f 0.62 (ethyl acetate); ^1H NMR (CDCl_3): δ 7.77 (m, 4H), 7.60-7.50 (m, 6H); ^{13}C NMR (CDCl_3): δ 163.5 (CO, q, J = 35 Hz), 133.7 (2 x C), 132.6 (2 x CH), 130.0 (4 x CH), 127.8 (4 x CH), 119.0 (CF_3 , q, J = 290 Hz); IR (KBr): ν 1634, 1181, 746, 683 cm^{-1} ; MS (EI), m/z (relative intensity): 297 ($[\text{M}]^+$, 71 %).

***N*-(4-nitro-benzenesulfonyl)-*t*-butylmetylsulfoximine (20a)**: Yield: 185 mg (61%); white solid, mp. 136 °C (decomp.); R_f 0.24 (ethyl acetate); ^1H NMR (CDCl_3): δ 8.31 (bd, J = 9.0 Hz, 2H), 8.09 (db, J = 9.0 Hz, 2H), 2.50 (s, 3H), 1.30 (s, 9H); ^{13}C NMR (CDCl_3): δ 150.1 (C), 149.1 (C), 127.5 (2 x CH), 124.0 (2 x CH), 55.7 (C), 28.0 (CH_3), 23.5 (3 x CH_3); IR (KBr): ν 1521, 1348, 1289, 1156, 1090, 986, 946, 608 cm^{-1} ; MS (EI), m/z (relative intensity): 247 ($[\text{M}-\text{C}_4\text{H}_9]^+$, 74 %); Calcd. for $\text{C}_{11}\text{H}_{16}\text{N}_2\text{O}_4\text{S}_2$: C, 43.41; H, 5.30; N, 9.20; found C, 43.84; H, 5.32; N, 9.16.

Preparation of diphenylsulfilimine (18c)⁸: To a solution of *N*-(trifluoroacetyl)-diphenylsulfilimine (**18b**) (297 mg, 1.0 mmol) in MeOH (10 mL) was added aq. KOH (2.5 mL, 2.0 N) at room temperature, and the mixture was stirred for 1 h. After concentration of the reaction mixture under reduced pressure, the residue was diluted with 1 N HCl (20 mL) and washed with CHCl_3 . The aqueous layer was made alkaline by adding NaOH, and subsequently extracted three times with CHCl_3 . The organic layer was dried (MgSO_4), filtered and concentrated under reduced pressure to give diphenylsulfilimine as a white solid. Yield: 177 mg (88%); mp. 65-66 °C, lit.⁸ mp. 70-71 °C; ^1H NMR (CDCl_3): δ 7.62-7.57 (m, 2H), 7.47-7.40 (m, 3H), 2.10 (bs, 1H); ^{13}C NMR (CDCl_3): δ 130.8 (C), 130.3 (CH), 129.1 (2 x CH), 125.8 (2 x CH).

References

1. Cren, S.; Kinahan, T. C.; Skinner, C. L.; Tye, H. *Tetrahedron Lett.* **2002**, 43, 2749.
2. (a) Heintzelman, R. W.; Swern, D. *Synthesis* **1976**, 731. (b) Huang, S.-L.; Swern, D., *J. Org. Chem.* **1979**, 44, 2510.
3. Brandt, J.; Gais, H.-J. *Tetrahedron: Asymmetry* **1997**, 8, 909.
4. Furukawa, N.; Akutagawa, K.; Yoshimura, T.; Oae, S. *Synthesis* **1982**, 77.
5. Mock, W. L.; Nugent, R. M. *J. Am. Chem. Soc.* **1975**, 97, 6526.
6. Johnson, C. R.; Kirchhoff, R. A.; Corkins, H. G. *J. Org. Chem.* **1974**, 39, 2458.
7. (a) Ketcha, D. M.; Abou-Gharbia, M.; Smith, F. X.; Swern, D. *Tetrahedron Lett.* **1983**, 24, 2811. (b) Abou-Gharbia, M.; Ketcha, D. M.; Zacharias, D. E.; Swern, D. *J. Org. Chem.* **1985**, 50, 2224.
8. (a) Furukawa, N.; Omata, T.; Yoshimura, T.; Aida, T.; Oae, S. *Tetrahedron Lett.* **1972**,

1619. (b) Yoshimura, T.; Omata, T.; Furukawa, N.; Oae, S. *J. Org. Chem.* **1976**, *41*, 1728.

¹H and ¹³C NMR

OKA 433 a3120231.b/h inova400/gas Dec 3 2003 (DMSO)

plot at 160 Hz/cm : 0.40 ppm/cm

12.6 12.0 11.4 10.8 ppm

10t at 805 Hz/cm : 8.00 ppm/cm

OKA 433 a3120231.b/c inova400/gas Dec 3 2003 (DMSO)

OKA 2B a4020968.b/h mercury300/gsw5 Feb 11 2004 (CDCI3)

plot at 120 Hz/cm : 0.40 ppm/cm

OKA 2B a4020968.b/c mercury300/gsw5 Feb 11 2004 (CDCI3)

plot at 604 Hz/cm : 8.00 ppm/cm

Oka 440 a3110637.b/h mercury300/gsw5 Nov 10 2003 (CDCl3)

plot at 120 Hz/cm : 0.40 ppm/cm

OKA 4401 a3110734.b/c inova400/gas Nov 10 2003 (CDCl3)

plot at 805 Hz/cm : 8.00 ppm/cm

OKA 4A a4020963.b/h mercury300/gsw5 Feb 10 2004 (DMSO)

plot at 120 Hz/cm : 0.40 ppm/cm

plot at 805 Hz/cm : 8.00 ppm/cm

OKA 442 a3111125.b/c inova400/gas Nov 12 2003 (CDCI3)

OKA 4431 a3111315.b/h inova400/gas Nov 14 2003 (DMSO)

plot at 160 Hz/cm : 0.40 ppm/cm

plot at 805 Hz/cm : 8.00 ppm/cm

OKA 4431 a3111315.b/c inova400/gas Nov 14 2003 (DMSO)

APT: C, CH2

CH, CH3

DEPT: CH

OKA 8A a4020966.b/h mercury300/gsw5 Feb 11 2004 (CDCl3)

plot at 120 Hz/cm : 0.40 ppm/cm

plot at 604 Hz/cm ; 8.00 ppm/cm

OKAMURA 455 a3120961.b/c mercury300/gsw5 Dec 11 2003 (CDCI3)

OCA 11 a2012222.b/h inova400/gas Jan 22 2002 (CDCl3)

plot at 160 Hz/cm : 0.40 ppm/cm

OKA-456 a4011230.b/h mercury300/gsw5 Jan 13 2004 (CDCl3)

plot at 120 Hz/cm : 0.40 ppm/cm

OKA 456 a4011230.b/c mercury300/gsw5 Jan 13 2004 (CDCI3)

plot at 604 Hz/cm : 8.00 ppm/cm

OKA 458 a3120514.b/h inova400/gas Dec 8 2003 (CDCl3)

plot at 160 Hz/cm : 0.40 ppm/cm

12.6 12.0 11.4 10.8 ppm

CHCl₃ →

plot at 805 Hz/cm : 8.00 ppm/cm

OKA 458 a3120514.b/c inova400/gas Dec 8 2003 (CDCI3)

DEPT: CH

Oka 450 a3121033.b/h inova400/gas Dec 12 2003 (CDCl3)

plot at 160 Hz/cm : 0.40 ppm/cm

12.6 12.0 11.4 10.8 ppm

OKA 450 a3121033.b/c inova400/gas Dec 12 2003 (CDCI3)

plot at 805 Hz/cm : 8.00 ppm/cm

OCA #36 a3110441.b/h inova400/gas Nov 5 2003 (CDCl3)

plot at 160 Hz/cm : 0.40 ppm/cm

OKA 436 a3110441.b/c inova400/gas Nov 5 2003 (CDCI3)

plot at 805 Hz/cm : 8.00 ppm/cm

APT: C, CH2

CH, CH3

DEPT: CH

plot at 604 Hz/cm : 8.00 ppm/cm

OKA 441 a3110699.b/c mercury300/gsw5 Nov 11 2003 (CDCI3)

OKA 448 a3111738.b/h mercury300/gsw5 Nov 18 2003 (CDCl3)

plot at 120 Hz/cm : 0.40 ppm/cm

12.6 12.0 11.4 10.8 ppm

OCA 448 a3111738.b/c mercury300/gsw5 Nov 18 2003 (CDCI3)

plot at 604 Hz/cm : 8.00 ppm/cm

OKA 18B a4020967.b/h mercury300/gsw5 Feb 11 2004 (CDCl3)

plot at 120 Hz/cm : 0.40 ppm/cm

OKA 461 a4011229.b/c mercury300/gsw5 Jan 13 2004 (CDCI3)

plot at 504 Hz/cm : 8.00 ppm/cm

OKA 459 a3121032.b/h inova400/gas Dec 12 2003 (CDCl3)

plot at 160 Hz/cm : 0.40 ppm/cm

OKA 459 a3121032.b/c inova400/gas Dec 12 2003 (CDCI3)

plot at 805 Hz/cm : 8.00 ppm/cm

APT: C, CH2

CH, CH3

DEPT: CH

plot at 120 Hz/cm : 0.40 ppm/cm

OKAMURA 452 a3120962.b/h mercury300/gsw5 Dec 11 2003 (CDCl3)