Rhodium-Catalyzed Multi-Component-Coupling Reactions Involving a Carborhodation–Cross-Coupling Sequence

Ryo Shintani,* Takafumi Yamagami, and Tamio Hayashi*

Department of Chemistry, Graduate School of Science, Kyoto University,
Sakyo, Kyoto 606-8502, Japan

Supporting Information

I. General

All air- and moisture-sensitive manipulations were carried out with standard Schlenk techniques under nitrogen or in a glove box under argon.

THF and toluene were purified by passing through a neutral alumina column under nitrogen. Benzene and 1,4-dioxane were distilled over benzophenone ketyl under nitrogen. Dichloromethane was distilled over CaH_2 under nitrogen. Triethylamine was distilled over KOH under vacuum.

1-Hexyne (Wako Chemicals), phenylacetylene (Wako Chemicals), 4-methyl-1-pentyne (TCI), 2-iodoaniline (Wako Chemicals), 2,4-diiodoaniline (Wako Chemicals), 4-iodoaniline (Wako Chemicals), iodomethane (Wako Chemicals), 3-bromo-2-methyl-1-propene (Aldrich), pyridine (Wako Chemicals), acetic anhydride (Wako Chemicals), phenyl isocyanate (Wako Chemicals), diphenylphosphoryl azide (TCI), *N,N'*-dicyclohexylcarbodiimide (Wako Chemicals), 2-iodobenzoic acid (Wako Chemicals), ZnCl_2 (Wako Chemicals), NaH (Kanto Chemicals; 60 wt% in mineral oil), *t*-BuLi (Kanto Chemicals; 1.46 M solution in pentane), MeLi (Kanto Chemicals; 1.04 M solution in Et_2O), 1,3-bis(diphenylphosphino)propane (Kanto Chemicals), 1,4-bis(diphenylphosphino)butane (TCI), and 1,1'-bis(diphenylphosphino)ferrocene (Fluka) were used as received.

5-Methyl-2-hexynoic acid, 4-methyl-2-pentynoic acid, and 2-heptynoic acid were prepared from the corresponding terminal alkyne following the literature procedure.¹ 5-Methoxy-1-pentyne,² $[\text{RhCl}(\text{cod})]_2$,³ $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$,⁴ and $\text{Pd}(\text{PPh}_3)_4$ ⁵ were synthesized following the literature procedures.

All other chemicals and solvents were purchased from Aldrich, Wako Chemicals, TCI, or Kanto Chemicals and used as received.

¹ Tsou, H.-R.; Mamuya, N.; Johnson, B. D.; Reich, M. F.; Gruber, B. C.; Ye, F.; Nilakantan, R.; Shen, R.; Discafani, C.; DeBlanc, R.; Davis, R.; Koehn, F. E.; Greenberger, L. M.; Wang, Y.-F.; Wissner, A. *J. Med. Chem.* **2001**, *44*, 2719.

² Jackson, W. R.; Perlmutter, P.; Smalridge, A. J. *Australian J. Chem.* **1988**, *41*, 251.

³ Giordano, G.; Crabtree, R. H. *Inorg. Synth.* **1979**, *19*, 218.

⁴ Cramer, R. *Inorg. Synth.* **1974**, *15*, 16.

⁵ Coulson, D. R. *Inorg. Synth.* **1972**, *13*, 121.

II. Synthesis of Substrates

The yields have not been optimized.

N-(2-Iodophenyl)-*N*-methyl-2-heptynamide (**1a**)

Diphenylphosphoryl azide (5.50 g, 20.0 mmol) was added to a solution of 2-iodobenzoic acid (4.96 g, 20.0 mmol) and triethylamine (2.02 g, 20.0 mmol) in benzene (20 mL) at 0 °C. After stirring for 2 h at 0 °C, the reaction mixture was washed with water and NaCl (saturated, aqueous), dried over MgSO₄, and filtered. The resulting solution was refluxed for 2 h and the volatiles were removed under vacuum. The residue was dissolved in benzene (20 mL) and was then added to a solution of 1-hexynylmagnesium bromide (20.0 mmol) in THF (8.0 mL) [prepared from ethylmagnesium bromide (20.0 mmol) and 1-hexyne (1.64 g, 20.0 mmol) at 0 °C]. The reaction mixture was stirred for 12 h at room temperature and quenched with NH₄Cl (saturated, aqueous). After extraction with Et₂O, the organic layer was washed successively with HCl (10% aqueous), NaHCO₃ (saturated aqueous), and NaCl (aqueous), and dried over MgSO₄, filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with CH₂Cl₂/hexane = 2/1 to afford *N*-(2-iodophenyl)-2-heptynamide as a pale yellow solid (1.79 g, 5.47 mmol).

A solution of this amide in THF (15 mL) was added to a suspension of NaH (240 mg, 6.00 mmol; 60 wt% in mineral oil) in THF (5 mL) at 0 °C, and the resulting mixture was stirred for 1 h at 0 °C. Iodomethane (851 mg, 6.00 mmol) was then added to it dropwise, and the mixture was stirred for 16 h at room temperature. The reaction was quenched with water and extracted with Et₂O. The organic layer was washed with NaCl (saturated, aqueous), dried over MgSO₄, filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with Et₂O/hexane = 1/1 to afford compound **1a** as a pale yellow oil (660 mg, 1.93 mmol; 10% overall yield). This compound exists as a 10:1 mixture of rotamers.

Major rotamer: ¹H NMR (CDCl₃): δ 7.91 (dd, ³J_{HH} = 7.9 Hz and ⁴J_{HH} = 1.3 Hz, 1H), 7.42 (ddd, ³J_{HH} = 7.8 and 7.4 Hz and ⁴J_{HH} = 1.3 Hz, 1H), 7.31 (dd, ³J_{HH} = 7.8 Hz and ⁴J_{HH} = 1.6 Hz, 1H), 7.09 (ddd, ³J_{HH} = 7.8 and 7.4 Hz and ⁴J_{HH} = 1.6 Hz, 1H), 3.22 (s, 3H), 2.11-2.01 (m, 2H), 1.20-1.14 (m, 2H), 1.07-0.98 (m, 2H), 0.73 (t, ³J_{HH} = 7.2 Hz, 3H). ¹³C NMR (CDCl₃): δ 154.2, 145.5, 139.6, 129.7, 129.4, 129.3, 99.5, 94.3, 74.7, 35.0, 29.2, 21.2, 18.1, 13.3. Anal. Calcd for C₁₄H₁₆INO: C, 49.28; H, 4.73. Found: C, 49.03; H, 4.51.

N-(2-Iodophenyl)-*N*-methyl-6-methoxy-2-hexynamide (**1b**)

This was synthesized from 2-iodobenzoic acid and 5-methoxy-1-pentyne, following the procedure for compound **1a**. Brown oil, 26% overall yield. This compound exists as a 9.5:1 mixture of rotamers.

Major rotamer: ^1H NMR (CDCl_3): δ 7.92 (dd, $^3J_{\text{HH}} = 7.9$ Hz and $^4J_{\text{HH}} = 1.2$ Hz, 1H), 7.43 (td, $^3J_{\text{HH}} = 7.6$ Hz and $^4J_{\text{HH}} = 1.3$ Hz, 1H), 7.32 (dd, $^3J_{\text{HH}} = 7.8$ Hz and $^4J_{\text{HH}} = 1.5$ Hz, 1H), 7.09 (td, $^3J_{\text{HH}} = 7.7$ Hz, $^4J_{\text{HH}} = 1.5$ Hz, 1H), 3.24 (s, 3H), 3.22 (s, 3H), 3.12-3.07 (m, 2H), 2.21-2.10 (m, 2H), 1.50-1.38 (m, 2H). ^{13}C NMR (CDCl_3): δ 154.2, 145.6, 140.0, 129.9, 129.5, 129.4, 99.6, 92.9, 74.9, 70.4, 58.5, 35.1, 27.6, 15.3. Anal. Calcd for $\text{C}_{14}\text{H}_{16}\text{INO}_2$: C, 47.08; H, 4.52. Found: C, 47.07; H, 4.45.

***N*-(2-Iodophenyl)-*N*,5-dimethyl-2-hexynamide (**1c**)**

A solution of DCC (1.90 g, 9.21 mmol) in CH_2Cl_2 (10 mL) was added to a solution of 2-iodoaniline (2.01 g, 9.18 mmol) and 5-methyl-2-hexynoic acid (1.16 g, 9.20 mmol) in CH_2Cl_2 (10 mL) at 0°C , and the mixture was stirred for 18 h at 0°C . The precipitate was filtered off through celite, and the solvent was removed under vacuum to afford *N*-(2-iodophenyl)-5-methyl-2-hexynamide as a yellow solid (1.00 g, 3.06 mmol).

A solution of this amide in THF (9 mL) was added to a suspension of NaH (134 mg, 3.35 mmol; 60 wt% in mineral oil) in THF (3 mL) at 0°C , and the resulting mixture was stirred for 1 h at 0°C . Iodomethane (851 mg, 3.36 mmol) was then added to it dropwise, and the mixture was stirred for 16 h at room temperature. The reaction was quenched with water and extracted with Et_2O . The organic layer was dried over MgSO_4 , filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with Et_2O /hexane = 3/2 to afford compound **1c** as a pale yellow solid (1.01 g, 2.96 mmol; 32% overall yield). This compound exists as an 11:1 mixture of rotamers.

Major rotamer: ^1H NMR (CDCl_3): δ 7.91 (d, $^3J_{\text{HH}} = 7.9$ Hz, 1H), 7.41 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 7.32 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 7.07 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 3.22 (s, 3H), 2.03-1.93 (m, 2H), 1.55-1.47 (m, 1H), 0.66 (d, $^3J_{\text{HH}} = 6.7$ Hz, 3H), 0.64 (d, $^3J_{\text{HH}} = 6.7$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 154.4, 145.7, 139.8, 129.9, 129.6, 129.5, 99.7, 92.6, 75.6, 35.1, 27.7, 27.2, 21.6, 21.5. Anal. Calcd for $\text{C}_{14}\text{H}_{16}\text{INO}$: C, 49.28; H, 4.73. Found: C, 49.40; H, 4.76.

***N*-(2-Iodophenyl)-*N*,4-dimethyl-2-pentynamide (**1d**)**

This was synthesized from 2-iodoaniline and 4-methyl-2-pentynoic acid, following the procedure for compound **1c**. Pale yellow oil, 32% overall yield. This compound exists as a 10:1 mixture of rotamers.

Major rotamer: ^1H NMR (CDCl_3): δ 7.91 (dd, $^3J_{\text{HH}} = 7.9$ Hz and $^4J_{\text{HH}} = 1.3$ Hz, 1H),

7.41 (td, $^3J_{\text{HH}} = 7.6$ Hz and $^4J_{\text{HH}} = 1.4$ Hz, 1H), 7.30 (dd, $^3J_{\text{HH}} = 7.8$ Hz and $^4J_{\text{HH}} = 1.6$ Hz, 1H), 7.09 (ddd, $^3J_{\text{HH}} = 7.7$ and 7.4 Hz and $^4J_{\text{HH}} = 1.6$ Hz, 1H), 3.22 (s, 3H), 2.37 (heptet, $^3J_{\text{HH}} = 6.8$ Hz, 1H), 0.88 (d, $^3J_{\text{HH}} = 6.8$ Hz, 3H), 0.85 (d, $^3J_{\text{HH}} = 6.8$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 154.2, 145.5, 139.4, 129.6, 129.3, 129.2, 99.5, 98.1, 73.8, 34.8, 21.3, 21.3, 20.0. Anal. Calcd for $\text{C}_{13}\text{H}_{14}\text{INO}$: C, 47.73; H, 4.31. Found: C, 47.74; H, 4.24.

***N*-(2-Iodophenyl)-*N*-methyl-3-phenylpropynamide (1e)**

This was synthesized from 2-iodobenzoic acid and phenylacetylene, following the procedure for compound **1a**. Orange solid, 10% overall yield. This compound exists as an 11:1 mixture of rotamers.

Major rotamer: ^1H NMR (CDCl_3): δ 7.97 (d, $^3J_{\text{HH}} = 7.9$ Hz, 1H), 7.46 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 7.39 (dd, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.3$ Hz, 1H), 7.32 (t, $^3J_{\text{HH}} = 7.4$ Hz, 1H), 7.22 (t, $^3J_{\text{HH}} = 7.7$ Hz, 2H), 7.14 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.3$ Hz, 1H), 7.08 (d, $^3J_{\text{HH}} = 7.4$ Hz, 2H), 3.30 (s, 3H). ^{13}C NMR (CDCl_3): δ 154.3, 145.5, 139.8, 132.5, 130.1, 130.0, 129.7, 129.5, 128.3, 120.2, 99.7, 90.6, 82.3, 35.2. Anal. Calcd for $\text{C}_{16}\text{H}_{12}\text{INO}$: C, 53.21; H, 3.35. Found: C, 53.24; H, 3.45.

***N*-(2-Iodophenyl)-*N*-methylacetamide (3)**

Pyridine (0.61 mL, 7.54 mmol) and acetic anhydride (0.71 mL, 7.51 mmol) were successively added to a solution of 2-iodoaniline (1.10 g, 5.02 mmol) in CH_2Cl_2 (10 mL) at 0 °C, and the mixture was stirred for 12 h at room temperature. The reaction was quenched with water and extracted with CH_2Cl_2 . The organic layer was washed with NaCl (saturated, aqueous), dried over MgSO_4 , filtered, and concentrated under vacuum to afford *N*-(2-iodophenyl)acetamide as a pale orange solid (1.02 g, 3.91 mmol).

A solution of this amide in THF (10 mL) was added to a suspension of NaH (175 mg, 4.38 mmol; 60 wt% in mineral oil) in THF (4 mL) at 0 °C, and the resulting mixture was stirred for 1 h at 0 °C. Iodomethane (625 mg, 4.40 mmol) was then added to it dropwise, and the mixture was stirred for 21 h at room temperature. The reaction was quenched with water and extracted with Et_2O . The organic layer was dried over MgSO_4 , filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with Et_2O /hexane = 1/1 to afford compound **3** as a white solid (680 mg, 2.74 mmol; 49% overall yield). This compound exists as a 33:1 mixture of rotamers.

Major rotamer: ^1H NMR (CDCl_3): δ 7.94 (dd, $^3J_{\text{HH}} = 7.9$ Hz and $^4J_{\text{HH}} = 1.4$ Hz, 1H), 7.43 (td, $^3J_{\text{HH}} = 7.6$ Hz and $^4J_{\text{HH}} = 1.4$ Hz, 1H), 7.28 (dd, $^3J_{\text{HH}} = 7.8$ Hz and $^4J_{\text{HH}} = 1.6$ Hz, 1H), 7.08 (td, $^3J_{\text{HH}} = 7.7$ Hz, $^4J_{\text{HH}} = 1.5$ Hz, 1H), 3.18 (s, 3H), 1.80 (s, 3H). ^{13}C NMR (CDCl_3): δ 170.1, 146.6, 140.2, 130.0, 129.8, 128.8, 99.4, 35.7, 22.5. Anal. Calcd for

C₉H₁₀INO: C, 39.30; H, 3.66. Found: C, 39.03; H, 3.64.

***N*-(4-Iodophenyl)-*N*-methyl-2-heptynamide (5)**

This was synthesized from 4-iodoaniline and 2-heptynoic acid, following the procedure for compound **1c**. White solid, 36% overall yield. This compound exists as a 5.9:1 mixture of rotamers.

Major rotamer: ¹H NMR (CDCl₃): δ 7.72 (d, ³*J*_{HH} = 8.5 Hz, 2H), 7.03 (d, ³*J*_{HH} = 8.4 Hz, 2H), 3.29 (s, 3H), 2.13 (t, ³*J*_{HH} = 6.9 Hz, 2H), 1.27 (quint, ³*J*_{HH} = 7.2 Hz, 2H), 1.10 (sextet, ³*J*_{HH} = 7.4 Hz, 2H), 0.80 (t, ³*J*_{HH} = 7.3 Hz, 3H). ¹³C NMR (CDCl₃): δ 154.0, 143.1, 138.2, 129.2, 94.5, 92.6, 74.8, 36.1, 29.3, 21.5, 18.4, 13.5. Anal. Calcd for C₁₄H₁₆INO: C, 49.28; H, 4.73. Found: C, 49.22; H, 4.65.

***N*-(2,4-Diiodophenyl)-*N*-methyl-2-heptynamide (8)**

This was synthesized from 2,4-diiodoaniline and 2-heptynoic acid, following the procedure for compound **1c**. White solid, 8% overall yield. This compound exists as a 10:1 mixture of rotamers.

Major rotamer: ¹H NMR (CDCl₃): δ 8.26 (d, ⁴*J*_{HH} = 1.7, 1H), 7.72 (dd, ³*J*_{HH} = 8.2 Hz and ⁴*J*_{HH} = 1.7 Hz, 1H), 7.03 (d, ³*J*_{HH} = 8.3 Hz, 1H), 3.19 (s, 3H), 2.17-2.06 (m, 2H), 1.26-1.17 (m, 2H), 1.11-1.02 (m, 2H), 0.80 (t, ³*J*_{HH} = 7.3 Hz, 3H). ¹³C NMR (CDCl₃): δ 154.0, 147.3, 145.4, 138.6, 130.9, 101.0, 94.2, 94.2, 74.7, 35.0, 29.3, 21.4, 18.3, 13.5. Anal. Calcd for C₁₄H₁₅I₂NO: C, 36.00; H, 3.24. Found: C, 35.93; H, 3.21.

***N*-(2-Methyl-2-propenyl)-*N*-phenyl-2-heptynamide (11a)**

A solution of phenyl isocyanate (1.19 g, 10.0 mmol) in benzene (10 mL) was added to a solution of 1-hexynylmagnesium bromide (10.0 mmol) in THF (8.0 mL) [prepared from ethylmagnesium bromide (10.0 mmol) and 1-hexyne (822 mg, 10.0 mmol) at 0 °C]. The reaction mixture was stirred for 21 h at room temperature, quenched with NH₄Cl (saturated, aqueous), and extracted with Et₂O. The organic layer was washed successively with HCl (10% aqueous), NaHCO₃ (saturated, aqueous), and NaCl (saturated, aqueous), and dried over MgSO₄, filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with

Et₂O/hexane = 2/1 to afford *N*-phenyl-2-heptynamide as a pale yellow oil (1.65 g, 8.20 mmol).

A solution of this amide (1.65 g, 8.20 mmol) in THF (10 mL) was added to a suspension of NaH (361 mg, 9.03 mmol; 60 wt% in mineral oil) in THF (10 mL) at 0 °C, and the resulting mixture was stirred for 1 h at 0 °C. 3-Bromo-2-methyl-1-propene (1.25 g, 9.22 mmol) was then added to it dropwise, and the mixture was stirred for 18 h at room temperature. The reaction was quenched with water and extracted with Et₂O. The organic layer was washed with NaCl (saturated, aqueous), dried over MgSO₄, filtered, and concentrated under vacuum. The residue was chromatographed on silica gel with Et₂O/hexane = 1/3 to afford compound **11a** as a yellow oil (1.50 g, 5.87 mmol; 59% overall yield). This compound exists as a 4.9:1 mixture of rotamers.

Major rotamer: ¹H NMR (CDCl₃): δ 7.39-7.22 (m, 5H), 4.83 (s, 1H), 4.74 (s, 1H), 4.32 (s, 2H), 2.08 (t, ³J_{HH} = 6.8 Hz, 2H), 1.75 (s, 3H), 1.21 (quint, ³J_{HH} = 7.2 Hz, 2H), 1.05 (sextet, ³J_{HH} = 7.4 Hz, 2H), 0.74 (t, ³J_{HH} = 7.3 Hz, 3H). ¹³C NMR (CDCl₃): δ 154.2, 141.8, 139.9, 128.6, 127.8, 127.5, 113.3, 94.2, 74.8, 54.1, 29.2, 21.3, 20.1, 18.2, 13.2. Anal. Calcd for C₁₇H₂₁NO: C, 79.96; H, 8.29. Found: C, 80.01; H, 8.38.

N-(2-Methyl-2-propenyl)-*N*-phenyl-5-methyl-2-hexynamide (**11b**)

This was synthesized from phenyl isocyanate and 4-methyl-1-pentyne, following the procedure for compound **11a**. Pink oil, 51% overall yield. This compound exists as a 5.9:1 mixture of rotamers.

Major rotamer: ¹H NMR (CDCl₃): δ 7.39-7.22 (m, 5H), 4.83 (s, 1H), 4.75 (s, 1H), 4.33 (s, 2H), 2.00 (d, ³J_{HH} = 6.3 Hz, 2H), 1.76 (s, 3H), 1.58-1.50 (m, 1H), 0.68 (d, ³J_{HH} = 6.7 Hz, 6H). ¹³C NMR (CDCl₃): δ 154.1, 141.7, 139.8, 128.7, 127.8, 127.5, 113.2, 93.1, 75.6, 54.0, 27.6, 27.0, 21.4, 20.0. Anal. Calcd for C₁₇H₂₁NO: C, 79.96; H, 8.29. Found: C, 79.91; H, 8.34.

General Procedure for the Preparation of Arylzinc Chlorides

t-BuLi (13.7 mL, 20.0 mmol; 1.46 M solution in pentane) was added dropwise to a solution of aryl bromide (10.0 mmol) in THF (10 mL) at -78 °C. The resulting mixture was stirred for 30 min at -78 °C, and then for 15 min at room temperature. The aryllithium solution thus obtained was added to a suspension of ZnCl₂ (1.45 g, 10.6 mmol) in THF (2.5 mL) at 0 °C. The reaction mixture was stirred for 15 min at room temperature, and was then concentrated under vacuum to remove pentane, affording a solution of arylzinc chloride in THF. The concentration was determined by acid-base titration with methyl orange (typically 0.6–1.2 M).

Only 3-chlorophenylzinc chloride was prepared from 3-chlorophenylmagnesium bromide (0.40 mL, 0.40 mmol; 1.00 M solution in THF) and ZnCl₂ (54.5 mg, 0.40 mmol) at 0 °C and was directly used in the rhodium-catalyzed reaction.

Methylzinc Chloride

MeLi (4.81 mL, 5.00 mmol; 1.04 M solution in Et₂O) was added dropwise to a suspension of ZnCl₂ (0.73 g, 5.36 mmol) in THF (2.0 mL) at 0 °C, and the resulting

mixture was stirred for 1 h at room temperature. The concentration was determined by acid-base titration with methyl orange (0.67 M).

III. Catalytic Reactions

General Procedure for Table 2.

A solution of $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$ (3.9 mg, 20 μmol Rh) and dppf (12.2 mg, 22.0 μmol) in dioxane (10 mL) was stirred for 10 min at room temperature. Substrate **1** (0.20 mmol) and ArZnCl (0.33–0.60 mL, 0.40–0.60 mmol; 0.67–1.22 M solution in THF) were added to it, and the mixture was stirred for 20 h at 40 °C. The reaction was quenched with water (80 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with CH_2Cl_2 to afford product **2**.

Entry 1. Yellow solid. 87% yield.

^1H NMR (CDCl_3): δ 7.49–7.42 (m, 3H), 7.26–7.23 (m, 2H), 7.11 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.0$ Hz, 1H), 6.73 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.61 (t, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.01 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 3.32 (t, $^3J_{\text{HH}} = 7.5$ Hz, 2H), 3.26 (s, 3H), 1.51–1.38 (m, 4H), 0.90 (t, $^3J_{\text{HH}} = 7.0$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.8, 159.7, 142.3, 141.7, 129.0, 128.2, 128.0, 126.9, 123.2, 122.8, 122.7, 121.3, 107.3, 34.8, 29.8, 25.7, 22.9, 13.9. Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{NO}$: C, 82.44; H, 7.26. Found: C, 82.27; H, 7.28.

Entry 3. Yellow solid. 84% yield.

^1H NMR (CDCl_3): δ 7.49–7.41 (m, 3H), 7.27–7.22 (m, 2H), 7.12 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 6.74 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 6.62 (t, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.04 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.48 (t, $^3J_{\text{HH}} = 6.8$ Hz, 2H), 3.40–3.33 (m, 2H), 3.29 (s, 3H), 3.26 (s, 3H), 1.82–1.74 (m, 2H). ^{13}C NMR (CDCl_3): δ 167.7, 158.7, 142.3, 141.3, 129.1, 128.3, 128.1, 126.9, 123.5, 122.9, 122.6, 121.3, 107.4, 72.5, 58.5, 31.6, 27.6, 25.7. Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{NO}_2$: C, 78.15; H, 6.89. Found: C, 78.09; H, 7.03.

Entry 4. Yellow solid. 94% yield.

^1H NMR (CDCl_3): δ 7.50–7.42 (m, 3H), 7.28–7.23 (m, 2H), 7.12 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.1$ Hz, 1H), 6.73 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 6.60 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} =$

1.1 Hz, 1H), 6.01 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.34 (d, $^3J_{\text{HH}} = 7.3$ Hz, 2H), 3.26 (s, 3H), 1.69-1.78 (m, 1H), 0.98 (d, $^3J_{\text{HH}} = 6.7$ Hz, 6H). ^{13}C NMR (CDCl_3): δ 168.0, 158.8, 142.2, 141.5, 129.0, 128.2, 128.1, 127.0, 124.3, 122.9, 122.8, 121.3, 107.3, 42.6, 27.1, 25.7, 22.3. Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{NO}$: C, 82.44; H, 7.26. Found: C, 82.28; H, 7.34.

Entry 5. Yellow solid. 57% yield.

^1H NMR (CDCl_3): δ 7.51-7.43 (m, 3H), 7.15-7.10 (m, 2H), 7.10 (td, $^3J_{\text{HH}} = 7.6$ Hz and $^4J_{\text{HH}} = 1.0$ Hz, 1H), 6.72 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 6.60 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 0.8$ Hz, 1H), 5.58 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 5.01 (septet, $^3J_{\text{HH}} = 6.9$ Hz, 1H), 3.26 (s, 3H), 1.07 (d, $^3J_{\text{HH}} = 6.8$ Hz, 6H). ^{13}C NMR (CDCl_3): δ 167.7, 164.3, 142.3, 138.0, 128.7, 127.9, 127.7, 127.3, 123.2, 123.0, 122.7, 121.4, 107.2, 28.4, 25.7, 20.7. Anal. Calcd for $\text{C}_{19}\text{H}_{19}\text{NO}$: C, 82.28; H, 6.90. Found: C, 82.03; H, 6.90.

Entry 6. Yellow solid. 85% yield.

^1H NMR (CDCl_3): δ 7.46-7.39 (m, 3H), 7.38-7.30 (m, 7H), 7.16 (t, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.76 (d, $^3J_{\text{HH}} = 7.8$ Hz, 1H), 6.68 (t, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.42 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.20 (s, 3H). ^{13}C NMR (CDCl_3): δ 166.8, 154.5, 143.3, 141.3, 139.9, 129.9, 129.3, 129.1, 129.0, 128.9, 128.7, 127.8, 124.2, 123.2, 123.1, 121.3, 107.6, 25.8. Anal. Calcd for $\text{C}_{22}\text{H}_{17}\text{NO}$: C, 84.86; H, 5.50. Found: C, 85.16; H, 5.78.

Entry 7. Yellow solid. 77% yield.

^1H NMR (CDCl_3): δ 7.39 (t, $^3J_{\text{HH}} = 7.9$ Hz, 1H), 7.12 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 0.9$ Hz, 1H), 6.98 (dd, $^3J_{\text{HH}} = 8.0$ Hz and $^4J_{\text{HH}} = 2.3$ Hz, 1H), 6.82 (d, $^3J_{\text{HH}} = 7.4$ Hz, 1H), 6.77 (dd, $^4J_{\text{HH}} = 2.4$ and 1.5 Hz, 1H), 6.74 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.64 (td, $^3J_{\text{HH}} = 7.6$ Hz and $^4J_{\text{HH}} = 0.8$ Hz, 1H), 6.11 (d, $^3J_{\text{HH}} = 7.5$ Hz, 1H), 3.82 (s, 3H), 3.41-3.33 (m, 1H), 3.26 (s, 3H), 3.27-3.18 (m, 1H), 1.52-1.38 (m, 4H), 0.91 (t, $^3J_{\text{HH}} = 7.1$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.8, 160.1, 159.4, 143.0, 142.3, 130.2, 128.0, 123.1, 123.0, 122.6, 121.4, 119.1, 113.7, 112.1, 107.3, 55.3, 34.7, 29.8, 25.7, 22.9, 13.9. Anal. Calcd for $\text{C}_{21}\text{H}_{23}\text{NO}_2$: C,

78.47; H, 7.21. Found: C, 78.49; H, 7.36.

Entry 8. Yellow solid. 80% yield.

^1H NMR (CDCl_3): δ 7.35 (t, $^3J_{\text{HH}} = 7.5$ Hz, 1H), 7.25 (d, $^3J_{\text{HH}} = 7.5$ Hz, 1H), 6.98 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.1$ Hz, 1H), 7.05 (s, 1H), 7.03 (d, $^3J_{\text{HH}} = 7.9$ Hz, 1H), 6.73 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.62 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.0$ Hz, 1H), 6.05 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.41-3.33 (m, 1H), 3.26 (s, 3H), 3.27-3.18 (m, 1H), 2.40 (s, 3H), 1.52-1.38 (m, 4H), 0.91 (t, $^3J_{\text{HH}} = 7.1$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.9, 160.1, 142.2, 141.6, 138.7, 128.9, 128.8, 127.9, 127.2, 123.9, 123.0, 122.9, 122.8, 121.3, 107.3, 34.8, 29.8, 25.6, 23.0, 21.5, 13.9. Anal. Calcd for $\text{C}_{21}\text{H}_{23}\text{NO}$: C, 82.58; H, 7.59. Found: C, 82.30; H, 7.58.

Entry 9. Yellow solid. 78% yield (contaminated with ~5% impurity).

^1H NMR (CDCl_3): δ 7.45-7.29 (m, 2H), 7.27-7.24 (m, 1H), 7.16-7.11 (m, 2H), 6.74 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.66 (td, $^3J_{\text{HH}} = 7.7$ Hz and $^4J_{\text{HH}} = 1.0$ Hz, 1H), 6.06 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.40-3.32 (m, 1H), 3.26 (s, 3H), 3.27-3.16 (m, 1H), 1.50-1.38 (m, 4H), 0.90 (t, $^3J_{\text{HH}} = 7.6$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.6, 157.3, 143.3, 142.4, 134.9, 130.4, 128.4, 128.3, 127.0, 125.2, 123.6, 122.8, 122.3, 121.4, 107.5, 34.5, 29.7, 25.7, 22.9, 13.9. Anal. Calcd for $\text{C}_{20}\text{H}_{20}\text{ClNO}$: C, 73.72; H, 6.19. Found: C, 73.99; H, 6.13.

Entry 10. Yellow solid. 77% yield.

^1H NMR (CDCl_3): δ 7.19 (d, $^3J_{\text{HH}} = 8.5$ Hz, 2H), 7.12 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 6.99 (d, $^3J_{\text{HH}} = 8.5$ Hz, 2H), 6.73 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 6.65 (t, $^3J_{\text{HH}} = 7.6$ Hz, 1H), 6.22 (d, $^3J_{\text{HH}} = 7.7$ Hz, 1H), 3.89 (s, 3H), 3.32-3.25 (m, 2H), 3.26 (s, 3H), 1.50-1.37 (m, 4H), 0.89 (t, $^3J_{\text{HH}} = 6.9$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.9, 160.0, 159.6, 142.2, 133.8, 128.5, 127.8, 123.1, 122.9, 122.7, 121.2, 114.3, 107.3, 55.3, 35.0, 30.0, 25.6, 22.9, 13.9. Anal. Calcd for

C₂₁H₂₃NO₂: C, 78.47; H, 7.21. Found: C, 78.51; H, 7.27.

Entry 11. Yellow solid. 75% yield.

¹H NMR (CDCl₃): δ 7.52 (d, ³J_{HH} = 7.5 Hz, 1H), 7.24 (td, ³J_{HH} = 8.0 Hz and ⁴J_{HH} = 1.1 Hz, 1H), 7.03 (td, ³J_{HH} = 7.6 Hz and ⁴J_{HH} = 1.0 Hz, 1H), 6.80 (d, ³J_{HH} = 7.7 Hz, 1H), 3.24 (s, 3H), 3.13-3.09 (m, 2H), 2.36 (s, 3H), 1.58-1.50 (m, 2H), 1.50-1.40 (m, 2H), 0.96 (t, ³J_{HH} = 7.3 Hz, 3H). ¹³C NMR (CDCl₃): δ 167.5, 159.5, 142.1, 127.4, 123.7, 123.4, 122.4, 121.5, 107.4, 35.5, 30.4, 25.5, 23.5, 23.0, 14.0. Anal. Calcd for C₁₅H₁₉NO₂: C, 78.56; H, 8.35. Found: C, 78.45; H, 8.55.

Procedure for Equation 2.

A solution of [RhCl(C₂H₄)₂]₂ (3.9 mg, 20 μmol Rh) and dppf (12.2 mg, 22.0 μmol) in dioxane (10 mL) was stirred for 10 min at room temperature. **1a** (68.2 mg, 0.20 mmol), **3** (55.0 mg, 0.20 mmol), and PhZnCl (0.41 mL, 0.40 mmol; 0.97 M solution in THF) were added to it, and the mixture was stirred for 20 h at 40 °C. The reaction was quenched with water (80 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with CH₂Cl₂ to afford product **2a** as a yellow solid (47.3 mg, 0.162 mmol; 81% yield) and **3** was recovered unchanged (82%).

Procedures for Equation 3.

Reaction with a palladium catalyst: PhZnCl (0.41 mL, 0.40 mmol; 0.97 M solution in THF) was added to a solution of Pd(PPh₃)₄ (11.6 mg, 10.0 μmol) and **5** (68.2 mg, 0.20 mmol) in THF (1.0 mL), and the mixture was stirred for 3 h at 60 °C. The reaction was quenched with water (80 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with Et₂O/hexane = 1/1 to afford product **6** (56.5 mg, 0.194 mmol; 97% yield).

Reaction with a rhodium catalyst: A solution of [RhCl(C₂H₄)₂]₂ (3.9 mg, 20 μmol Rh) and dppf (12.2 mg, 22.0 μmol) in dioxane (10 mL) was stirred for 10 min at room temperature. **5** (68.2 mg, 0.20 mmol) and PhZnCl (0.41 mL, 0.40 mmol; 0.97 M solution in THF) were added to it, and the mixture was stirred for 20 h at 40 °C. The reaction was quenched with water (80 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with CH₂Cl₂ to afford product **7** as a pale yellow oil (44 μmol; 22% yield, a mixture of *E/Z* isomers).

Procedure for Equation 4.

A solution of [RhCl(C₂H₄)₂]₂ (1.9 mg, 10 μmol Rh) and dppf (6.1 mg, 11 μmol) in dioxane (5.0 mL) was stirred for 10 min at room temperature. Substrate **8** (46.7 mg, 0.10 mmol) and PhZnCl (0.20 mL, 0.19 mmol; 0.97 M solution in THF) were added to it, and the mixture was stirred for 20 h at 40 °C. The reaction was quenched with water (40 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with CH₂Cl₂

to afford a mixture of **9** and **10** as a yellow solid (30.7 mg, 74.0 μ mol; 74% yield (**9**/**10** = 95/5)). Analytically pure **9** and **10** were obtained by further chromatographic purification on silica gel.

9: ^1H NMR (CDCl_3): δ 7.54-7.47 (m, 3H), 7.41 (dd, $^3J_{\text{HH}} = 8.1$ Hz and $^4J_{\text{HH}} = 1.6$ Hz, 1H), 7.22-7.19 (m, 2H), 6.51 (d, $^3J_{\text{HH}} = 8.1$ Hz, 1H), 6.19 (d, $^4J_{\text{HH}} = 1.7$ Hz, 1H), 3.35-3.28 (m, 2H), 3.23 (s, 3H), 1.50-1.38 (m, 4H), 0.90 (t, $^3J_{\text{HH}} = 7.1$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.1, 161.7, 141.7, 141.0, 136.3, 131.7, 129.1, 128.6, 126.6, 124.9, 122.3, 109.3, 84.0, 34.8, 29.7, 25.7, 22.9, 13.9. Anal. Calcd for $\text{C}_{20}\text{H}_{20}\text{INO}$: C, 57.57; H, 4.83. Found: C, 57.71; H, 4.90.

10: ^1H NMR (CDCl_3): δ 7.53-7.43 (m, 3H), 7.37 (dd, $^3J_{\text{HH}} = 8.1$ Hz and $^4J_{\text{HH}} = 1.8$ Hz, 1H), 7.31-7.26 (m, 4H), 7.23-7.19 (m, 1H), 7.17-7.14 (m, 2H), 6.81 (d, $^3J_{\text{HH}} = 7.0$ Hz, 1H), 6.24 (d, $^4J_{\text{HH}} = 1.7$ Hz, 1H), 3.37-3.32 (m, 2H), 3.29 (s, 3H), 1.55-1.40 (m, 4H), 0.92 (t, $^3J_{\text{HH}} = 7.2$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 167.9, 160.2, 141.7, 140.8, 134.1, 129.1, 128.5, 128.2, 126.9, 126.6, 126.5, 126.2, 123.3, 123.2, 121.8, 107.5, 34.6, 29.8, 25.8, 22.9, 13.9. Anal. Calcd for $\text{C}_{26}\text{H}_{25}\text{NO}$: C, 84.98; H, 6.86. Found: C, 84.90; H, 7.10.

General Procedure for Equation 5.

A solution of $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$ (3.9 mg, 20 μ mol Rh) and dppf (12.2 mg, 22.0 μ mol) in toluene (10 mL) was stirred for 10 min at room temperature. Substrate **11** (0.20 mmol), iodomethane (141 mg, 1.00 mmol), and PhZnCl (0.66 mL, 0.60 mmol; 0.91 M solution in THF) were added to it, and the mixture was stirred for 20 h at 40 $^\circ\text{C}$. The reaction was quenched with water (80 μL), passed through a pad of silica gel with EtOAc, and concentrated under vacuum. The residue was purified by silica gel preparative TLC with Et_2O /hexane = 1/3 to afford product **12** (contaminated with **13**).

R = *n*Bu. White solid. 70% yield (**12a**/**13a** = 96/4).

12a: ^1H NMR (CDCl_3): δ 7.72-7.67 (m, 2H), 7.41-7.28 (m, 5H), 7.17-7.12 (m, 3H), 3.50 (d, $^2J_{\text{HH}} = 9.4$ Hz, 1H), 3.38 (d, $^2J_{\text{HH}} = 9.4$ Hz, 1H), 3.23-3.20 (m, 1H), 2.90-2.82 (m, 1H), 1.47-1.32 (m, 5H), 1.28-1.18 (m, 1H), 0.86 (t, $^3J_{\text{HH}} = 7.2$ Hz, 3H), 0.85 (s, 3H), 0.82 (t, $^3J_{\text{HH}} = 7.4$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 168.4, 153.0, 142.0, 140.0, 134.0, 128.7, 127.8, 127.6, 127.0, 124.3, 120.0, 57.4, 41.0, 35.4, 34.1, 30.3, 26.7, 22.9, 14.0, 9.0. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{30}\text{NO}$ ($\text{M}+\text{H}^+$) 348.2322, found 348.2321.

R = *i*Bu. White solid. 84% yield (**12b**/**13b** = 97/3).

12b: ^1H NMR (CDCl_3): δ 7.70-7.66 (m, 2H), 7.40-7.27 (m, 5H), 7.17-7.12 (m, 3H), 3.48 (d, $^2J_{\text{HH}} = 9.4$ Hz, 1H), 3.37 (d, $^2J_{\text{HH}} = 9.4$ Hz, 1H), 3.34 (dd, $^2J_{\text{HH}} = 12.5$ Hz and $^3J_{\text{HH}} = 8.2$ Hz, 1H), 2.85 (dd, $^2J_{\text{HH}} = 12.5$ Hz and $^3J_{\text{HH}} = 6.6$ Hz, 1H), 1.63-1.52 (m, 1H), 1.49-1.40 (m, 1H), 1.29-1.18 (m, 1H), 0.98 (d, $^3J_{\text{HH}} = 6.7$ Hz, 3H), 0.94 (d, $^3J_{\text{HH}} = 6.6$ Hz, 3H), 0.83 (s, 3H), 0.83 (t, $^3J_{\text{HH}} = 7.5$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 168.4, 151.9, 141.7, 139.7, 135.3, 128.7, 128.1, 127.5, 127.0, 124.3, 120.1, 57.4, 42.6, 41.1, 34.4, 26.8, 26.4, 22.5, 21.9, 9.1. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{30}\text{NO}$ ($\text{M}+\text{H}^+$) 348.2322, found 348.2326.

IV. ¹H NMR Spectra