Supporting Information for

Room Temperature Palladium-Catalyzed Arylation of Indoles

Nicholas R. Deprez, Dipannita Kalyani, Andrew Krause, and Melanie S. Sanford*

University of Michigan Department of Chemistry, 930 N. University Ave, Ann Arbor, MI 48109

General Procedures: NMR spectra were obtained on a Varian Inova 400 (399.96 MHz for ^1H ; 100.57 for ^{13}C) spectrometer unless otherwise noted. ^1H NMR chemical shifts are reported in parts per million (ppm) relative to TMS, with the residual solvent peak used as an internal reference. Multiplicities are reported as follows: singlet (s), doublet (d), doublet of doublets (dd), doublet of doublets of doublets (ddd), triplet of doublets (td), quartet (q), multiplet (m), and broad resonances (br).

Materials and Methods: Substrates **1-5**, **8**, **10** and **11** were obtained from commercial sources and were used as received. Substrates **3** and **9** were prepared by N-methylation of the corresponding indoles.⁶ Substrate **6** was prepared by reaction of 5-hydroxyindole with acetyl chloride (1.1 equiv) and Et_3N (1.1 equiv) in CH_2Cl_2 . Substrate **7** was prepared according to a literature procedure.¹ Aryliodonium salts with $\text{Ar} = \text{Ph}$, *m*- CF_3 , *p*-F, *p*-Cl, *p*-Br, *p*-Me, *o*-Me, and 1-naphthyl were prepared in two steps – (i) conversion of the aryl iodide to the corresponding diacetoxiodoarene² and (ii) reaction of $\text{ArI}(\text{OAc})_2$ with $\text{ArB}(\text{OH})_2$ in the presence of $\text{BF}_3 \cdot \text{OEt}_2$.³ The aryliodonium salt $[(p\text{-MeOC}_6\text{H}_4)_2\text{I}]\text{BF}_4$ was prepared by the reaction of anisole and trifluoromethanesulfonic acid with $p\text{-MeOC}_6\text{H}_4\text{I}(\text{OAc})_2$ ² according to a literature procedure.⁴ $\text{IMesPd}(\text{OAc})_2$ was prepared by reaction of $(\text{IMesPd}(\text{Cl})_2)_2$ ⁵ with AgOAc . Other palladium catalysts were obtained from commercial sources (Strem or Pressure Chemicals) and used as received. Solvents were obtained from Fisher Chemical and used without further purification. Flash chromatography was performed on EM Science silica gel 60 (0.040-0.063 mm particle size, 230-400 mesh) and thin layer chromatography was performed on Merck TLC plates pre-coated with silica gel 60 F_{254} . Control reactions (in the absence of Pd catalyst) were run for each substrate and generally showed no reaction under our standard conditions.

Experimental Procedures

1-Methylindole (131.2 mg, 1.00 mmol, 1.0 equiv) and IMesPd(OAc)₂ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (735.4 mg, 2.0 mmol, 2.0 equiv) was added, and the reaction was stirred for 18 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.33 in 96% hexanes/4% EtOAc). The product (**1a**) was obtained as a white solid (178 mg, 86% yield, average of three runs). Mp = 99-102 °C. ¹H NMR (500 MHz, CDCl₃): δ 7.65 (d, *J* = 7.9 Hz, 1H), 7.53-7.51 (multiple peaks, 2H), 7.48 (t, *J* = 8.0 Hz, 2H), 7.43-7.39 (m, 1H), 7.38 (d, *J* = 8.2 Hz, 1H), 7.26 (td, *J* = 7.6, 1.4 Hz, 1H), 7.15 (td, *J* = 7.9, 1.0 Hz, 1H), 6.57 (s, 1H), 3.76 (s, 3H). ¹³C NMR (125 MHz, CDCl₃): δ 141.7, 138.5, 132.9, 129.5, 128.6, 128.1, 127.9, 121.8, 120.6, 120.0, 109.8, 101.8, 31.3. Anal. Calcd for C₁₅H₁₃N: C, 86.92, H, 6.32, N, 6.76; Found; C, 86.87, H, 6.54, N, 6.71. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₅H₁₃N, 207.1048; found, 207.1044.

Indole (117.2 mg, 1.00 mmol, 1.0 equiv) and IMesPd(OAc)₂ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (735.4 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.1 in 96% hexanes/4% EtOAc). The product (**2a**) was obtained as a white solid (155.9 mg, 81% yield). Mp = 188-189 °C. ¹H NMR (400 MHz, CDCl₃): δ 8.31 (br s, 1H), 7.65-7.69 (multiple peaks, 3H), 7.39-7.49 (multiple peaks, 3H), 7.35 (t, *J* = 7.4 Hz, 1H), 7.23 (ddd, *J* = 7.8, 7.0, 1.2 Hz, 1H), 7.16 (ddd, *J* = 7.7, 7.0, 1.0 Hz, 1H), 6.86 (s, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 137.8, 136.7, 132.2, 129.2, 129.0, 127.7, 125.1, 122.3, 120.6, 120.2, 110.9, 99.9. Anal. Calcd for C₁₄H₁₁N: C, 87.01, H, 5.74, N, 7.25; Found; C, 87.03, H, 5.71, N, 7.16. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₄H₁₁N, 193.0891; found, 193.0892.

1,3-Dimethylindole (150.0 mg, 1.03 mmol, 1.0 equiv) and IMesPd(OAc)₂ (27.3 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (759.7 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.2 in 98% hexanes/2% EtOAc). The product (**3a**) was obtained as a light yellow viscous oil (203 mg, 89% yield). ¹H NMR (500 MHz, CDCl₃): δ 7.61 (d, *J* = 7.84 Hz, 1H), 7.50 (t, *J* = 7.93 Hz, 2H), 7.44-7.40 (multiple peaks, 3H), 7.34 (d, *J* = 8.2 Hz, 1H), 7.28-7.25 (m, 1H), 7.16 (td, *J* = 7.0, 1.0 Hz, 1H), 3.62 (s, 3H), 2.29 (s, 3H). ¹³C NMR (125 MHz, CDCl₃): δ 137.6, 137.2, 132.1, 130.6, 128.4, 128.3, 127.7, 121.7, 119.1, 118.8, 109.2, 108.5, 30.9, 9.3. Anal. Calcd for C₁₆H₁₅N: C, 86.84, H, 6.83, N, 6.33; Found; C, 86.71, H, 6.92, N, 6.33. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₅N, 221.1204; found, 221.1202.

2-Methylindole (150 mg, 1.03 mmol, 1.0 equiv) and IMesPd(OAc)₂ (27.3 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (758 mg, 2.07 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was taken up in CH₂Cl₂ and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel in 80% hexanes/20% benzene (R_f = 0.25) followed by a second column in 97.5% hexanes/2.5% EtOAc (R_f = 0.18). The two columns were required to remove a high molecular weight side product (R_f = 0.18 in 80% hexanes/20% benzene) as well as unreacted starting material (R_f = 0.22 in 97.5% hexanes/2.5% EtOAc). The product (**4a**) was obtained as a white solid (66 mg, 29% yield). Mp = 108.9-111.2 °C. ¹H NMR (400 MHz, CDCl₃) δ 7.71 (d, *J* = 7.9 Hz, 1H), 7.55-7.49 (multiple peaks, 4H), 7.37-7.33 (multiple peaks, 2H), 7.25 (t, *J* = 8.1 Hz, 1H), 7.15 (t, *J* = 7.9 Hz, 1H), 3.77 (s, 3H), 2.53 (s, 3H). ¹³C NMR (125 MHz, CDCl₃): δ 136.5, 135.7, 133.3, 129.7, 128.4, 126.9, 125.6, 121.1, 119.6, 118.7, 113.9, 108.7, 29.6, 11.0. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₅N, 221.1204; found, 221.1201.

5-Methoxyindole (147.2 mg, 1.00 mmol, 1.0 equiv) and IMesPd(OAc)₂ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄

(735.4 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.28 in 90% hexanes/10% EtOAc). The product (**5a**) was obtained as white solid (130 mg, 58% yield). Mp = 166-169 °C. ¹H NMR (400 MHz, CDCl₃): δ 8.24 (br s, 1H), 7.65 (d, *J* = 7.2 Hz, 2H), 7.41-7.47 (multiple peaks, 2H), 7.25-7.36 (multiple peaks, 2H), 7.11 (d, *J* = 2.4 Hz, 1H), 6.88 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.77 (d, *J* = 1.9 Hz, 1H), 3.88 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 154.4, 138.5, 132.4, 131.9, 129.7, 129.0, 127.6, 125.0, 112.6, 111.6, 102.2, 99.8, 55.8. Anal. Calcd for C₁₅H₁₃NO: C, 80.69, H, 5.87, N, 6.27; Found; C, 80.61, H, 5.93, N, 6.19. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₅H₁₃NO, 223.0997; found, 223.0996.

5-Acetoxyindole (100.0 mg, 0.57 mmol, 1.0 equiv) and IMesPd(OAc)₂ (15.1 mg, 0.028 mmol, 0.05 equiv) were dissolved in AcOH (5.7 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (419.7 mg, 1.14 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.22 in 78% hexanes/22% EtOAc). The product (**6a**) was obtained as a light orange crystalline solid (101 mg, 71% yield). Mp = 170.7-172.2 °C. ¹H NMR (500 MHz, CDCl₃): δ 8.39 (br s, 1H), 7.65 (d, *J* = 7.3 Hz, 2H), 7.44 (t, *J* = 7.5 Hz, 2H), 7.35-7.30 (multiple peaks, 3H), 6.89 (dd, *J* = 8.6, 2.2 Hz, 1H), 6.79 (s, 1H), 2.33 (s, 3H). ¹³C NMR (125 MHz, CDCl₃): δ 170.6, 144.6, 139.3, 134.7, 132.1, 129.5, 129.1, 127.9, 125.2, 116.3, 112.6, 111.4, 100.1, 21.2. IR(KBr): 3392, 1742, 1371 cm⁻¹. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₃NO₂, 251.0946; found, 251.0938.

1-Methyl-5-morpholine-indole (175.0 mg, 0.81 mmol, 1.0 equiv) and IMesPd(OAc)₂ (42.8 mg, 0.81 mmol, 0.1 equiv) were dissolved in AcOH (8.1 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (893.3 mg, 2.43 mmol, 3.0 equiv) was added, and the reaction was stirred for 15 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.25 in 70% hexanes/30% EtOAc). The product (**7a**) was obtained as a white solid (204 mg, 86% yield). Mp = 143-145 °C. ¹H NMR (400 MHz, C₆D₆): δ 7.27-7.32 (multiple peaks, 2H), 6.93-7.16 (multiple peaks, 6H), 6.57

(s, 1H), 3.63-3.67 (multiple peaks, 4H), 3.12 (s, 3H), 2.84-2.89 (multiple peaks, 4H). ^{13}C NMR (100 MHz, C_6D_6): δ 146.9, 141.8, 134.9, 133.6, 129.5, 129.2, 128.7, 115.3, 110.3, 107.9, 102.1, 67.4, 52.3, 30.7. Two carbon resonances are coincidentally overlapping. Anal. Calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}$: C, 78.05, H, 6.89, N, 9.58; Found; C, 78.18, H, 7.03, N, 9.61. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}$, 292.1576; found, 292.1575. Small amounts of product (~7% by GC) were observed in the control reaction (in the absence of Pd catalyst), which is believed to result from traces of palladium in the starting material (which was prepared via Pd catalyzed cross coupling).

5-Bromoindole (117.6 mg, 0.60 mmol, 1.0 equiv) and $\text{IMesPd}(\text{OAc})_2$ (15.8 mg, 0.03 mmol, 0.05 equiv) were dissolved in AcOH (6 mL), and the resulting solution was stirred at 25 °C for ~5 min. $[\text{Ph}_2\text{I}]\text{BF}_4$ (441 mg, 1.20 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH_2Cl_2 (50 mL) and extracted with aqueous NaHCO_3 (2 x 30 mL). The organic layer was dried with MgSO_4 , concentrated, and the product was purified by chromatography on silica gel (R_f = 0.13 in 90% hexanes/10% ethyl acetate). The product (**8a**) was obtained as a white solid (121 mg, 74% yield). Mp = 193-196 °C. ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 11.75 (s, 1H), 7.86 (d, J = 7.2 Hz, 2H), 7.71 (d, J = 1.9 Hz, 1H), 7.44-7.50 (multiple peaks, 2H), 7.31-7.38 (multiple peaks, 2H), 7.2 (dd, J = 8.6, 2.0 Hz, 1H), 6.89 (d, J = 1.4 Hz, 1H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 139.1, 135.7, 131.5, 130.4, 128.9, 127.8, 125.1, 123.9, 122.0, 113.1, 111.8, 98.2. Anal. Calcd for $\text{C}_{14}\text{H}_{10}\text{NBr}$: C, 61.79, H, 3.70, N, 5.15; Found; C, 61.86, H, 3.82, N, 5.10. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{14}\text{H}_{10}\text{NBr}$, 270.9997; found, 270.9993.

1-Methyl-5-nitroindole (150.0 mg, 0.85 mmol, 1.0 equiv) and $\text{IMesPd}(\text{OAc})_2$ (22.5 mg, 0.043 mmol, 0.05 equiv) were dissolved in AcOH (8.5 mL), and the resulting solution was stirred at 25 °C for ~5 min. $[\text{Ph}_2\text{I}]\text{BF}_4$ (1.88 g, 5.11 mmol, 6.0 equiv) was added, and the reaction was stirred for 24 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH_2Cl_2 (950 mL) and extracted with aqueous NaHCO_3 (2 x 30 mL). The organic layer was dried with MgSO_4 , concentrated, and the product was purified by chromatography on silica gel (R_f = 0.2 in 92% hexanes/8% ethyl acetate). The product (**9a**) was obtained as a yellow powder (150 mg, 70% yield). Mp = 180-183 °C. ^1H NMR (500 MHz, CDCl_3): δ 8.59 (d, J = 2.2 Hz, 1H), 8.16 (dd, J = 9.0, 2.2 Hz, 1H), 7.52-7.45 (multiple peaks, 5H), 7.38 (d, J = 9.0 Hz, 1H), 6.73 (s, 1H), 3.81 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 144.8, 141.8, 140.9, 131.4, 129.3, 128.8, 128.7, 127.1, 117.6, 117.3, 109.4, 103.8, 31.6. Anal. Calcd for

C₁₅H₁₂N₂O₂: C, 71.42, H, 4.79, N, 11.10; Found; C, 71.63, H, 4.88, N, 10.85. HRMS electrospray (m/z): [M⁺] calcd for C₁₅H₁₂N₂O₂, 252.0899; found, 252.0894.

Pyrrole (939 mg, 14.0 mmol, 10 equiv) and IMesPd(OAc)₂ (37.0 mg, 0.07 mmol, 0.05 equiv) were dissolved in AcOH (13 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (515 mg, 1.40 mmol, 1.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.20 in 95% hexanes/5% EtOAc). The product (**10a**) was obtained as a white solid (137 mg, 69% yield). Mp = 125-128 °C. ¹H NMR (400 MHz, CDCl₃): δ 8.45 (br s, 1H), 7.48 (d, *J* = 7.3 Hz, 2H), 7.34-7.40 (multiple peaks, 2H), 7.21 (t, *J* = 7.3 Hz, 1H), 6.88 (m, 1H), 6.54 (m, 1H), 6.31 (td, *J* = 3.5, 2.6 Hz, 1H). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 132.7, 132.0, 128.8, 126.1, 123.8, 118.8, 110.0, 105.9. HRMS electrospray (m/z): [M⁺] calcd for C₁₀H₉N, 143.0735; found, 143.0735.

1-Methylpyrrole (1.136 g, 14.0 mmol, 10 equiv) and IMesPd(OAc)₂ (37.0 mg, 0.07 mmol, 0.05 equiv) were dissolved in AcOH (14 mL), and the resulting solution was stirred at 25 °C for ~5 min. [Ph₂I]BF₄ (515 mg, 1.40 mmol, 1.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.27 in 82% hexanes/18% benzene). The product (**11a**) was obtained as a white solid (148 mg, 67% yield, 0.94 mmol) which was 96% pure by GC analysis. This product was inseparable from 4% of the corresponding diphenylated product 1-methyl-2,5-diphenylindole (6.6 mg, 0.028 mmol). Mp = 45-48 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.37-7.50 (multiple peaks, 4H), 7.27-7.34 (m, 1H), 6.72 (t, *J* = 2.0 Hz, 1H), 6.19-6.26 (multiple peaks, 2H), 3.67 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 134.5, 133.3, 128.5, 128.2, 126.6, 123.6, 108.6, 107.7, 34.9. HRMS electrospray (m/z): [M⁺] calcd for C₁₁H₁₁N, 157.0891; found, 157.0888.

1-Methylindole (78.7 mg, 0.60 mmol, 1.0 equiv) and IMesPd(OAc)₂ (15.9 mg, 0.03 mmol, 0.05 equiv) were dissolved in AcOH (6 mL), and the resulting solution was stirred at 25 °C for ~5 min. [(*m*-CF₃C₆H₅)₂I]BF₄ (604.8 mg, 1.2 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.26 in 90% hexanes/10% benzene). The product (**1b**) was obtained as a pale yellow oil (106 mg, 64% yield). ¹H NMR (400 MHz, CDCl₃): δ 7.82 (s, 1H), 7.67-7.74 (multiple peaks, 3H), 7.59-7.65 (multiple peaks, 1H), 7.39-7.43 (m, 1H), 7.32 (ddd, *J* = 8.2, 7.0, 1.2 Hz, 1H), 7.21 (ddd, *J* = 7.9, 7.0, 1.1 Hz, 1H), 6.66 (s, 1H), 3.78 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 139.7, 138.6, 133.6, 132.4 (q, ⁴*J*_{C-F} = 1.2 Hz), 131.0 (q, ²*J*_{C-F} = 32 Hz), 129.0, 127.8, 125.9 (q, ³*J*_{C-F} = 3.9 Hz), 124.4 (q, ³*J*_{C-F} = 3.8 Hz), 124.0 (q, ¹*J*_{C-F} = 273 Hz), 122.2, 120.7, 120.1, 109.7, 102.6, 31.1. Anal. Calcd for C₁₆H₁₂F₃N: C, 69.81, H, 4.39, N, 5.09; Found; C, 69.89, H, 4.46, N, 5.05. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₂F₃N, 275.0922; found, 275.0920.

1-Methylindole (131.2 mg, 1.00 mmol, 1.0 equiv) and IMesPd(OAc)₂ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [(*p*-FC₆H₅)₂I]BF₄ (808.0 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (R_f = 0.43 in 90% hexanes/10% benzene). The product (**1c**) was obtained as a light orange solid (180 mg, 80% yield). Mp = 119-122 °C. ¹H NMR (400 MHz, CD₃OD): δ 7.51-7.57 (multiple peaks, 3H), 7.36 (d, *J* = 8.3 Hz, 1H), 7.15-7.25 (multiple peaks, 3H), 7.05 (ddd, *J* = 7.8, 7.0, 1.0 Hz, 1H), 6.48 (s, 1H), 3.71 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 162.4 (d, ¹*J*_{C-F} = 247 Hz), 140.3, 138.2, 130.9 (d, ³*J*_{C-F} = 7.4 Hz), 128.8 (d, ⁴*J*_{C-F} = 4.4 Hz), 127.8, 121.7, 120.4, 119.9, 115.4 (d, ²*J*_{C-F} = 22 Hz), 109.6, 101.6, 30.9. Anal. Calcd for C₁₅H₁₂FN: C, 79.98, H, 5.37, N, 6.22; Found; C, 80.33, H, 5.43, N, 6.26. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₅H₁₂FN, 225.0954; found, 225.0953.

1-Methylindole (131.2 mg, 1.00 mmol, 1.0 equiv) and IMesPd(OAc)₂ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. [(*p*-ClC₆H₅)₂I]BF₄ (873.6 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was

dried with MgSO_4 , concentrated, and the product was purified by chromatography on silica gel ($R_f = 0.25$ in 98% hexanes/2% ethyl acetate). The product (**1d**) was obtained as white solid (219 mg, 90% yield). Mp = 115-118 °C. ^1H NMR (400 MHz, CDCl_3): δ 7.65 (d, $J = 7.8$ Hz, 1H), 7.38-7.45 (m, 4H), 7.35 (d, $J = 7.9$ Hz, 1H), 7.27 (ddd, $J = 8.2, 7.0, 1.2$ Hz, 1H), 7.17 (ddd, $J = 7.9, 6.9, 1.1$ Hz, 1H), 6.56 (s, 1H), 3.69 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 140.1, 138.4, 133.8, 131.2, 130.4, 128.7, 127.8, 121.9, 120.5, 120.0, 109.6, 101.9, 31.1. Anal. Calcd for $\text{C}_{15}\text{H}_{12}\text{ClN}$: C, 74.53, H, 5.00, N, 5.79; Found; C, 74.23, H, 5.00, N, 5.61. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{15}\text{H}_{12}\text{ClN}$, 241.0658; found, 241.0655.

Indole (88.1 mg, 0.75 mmol, 1.00 equiv) and $\text{IMesPd}(\text{OAc})_2$ (19.8 mg, 0.037 mmol, 0.05 equiv) were dissolved in AcOH (7.5 mL), and the resulting solution was stirred at 25 °C for ~5 min. $[(p\text{-BrC}_6\text{H}_5)_2\text{I}]\text{BF}_4$ (788.6 mg, 1.50 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 80 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH_2Cl_2 (50 mL) and extracted with aqueous NaHCO_3 (2 x 30 mL). The organic layer was dried with MgSO_4 , concentrated, and the product was purified by chromatography on silica gel ($R_f = 0.13$ in 95% hexanes/5% ethyl acetate). The product (**2b**) was obtained as a white solid (135 mg, 66% yield). Mp = 208-212 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 11.59 (s, 1H), 7.82 (d, $J = 8.6$ Hz, 2H), 7.65 (d, $J = 8.6$ Hz, 2H), 7.54 (d, $J = 7.9$ Hz, 1H), 7.40 (d, $J = 8.0$ Hz, 1H), 7.11 (ddd, $J = 8.1, 7.0, 1.1$ Hz, 1H), 7.00 (ddd, $J = 7.8, 7.0, 1.0$ Hz, 1H), 6.94 (d, $J = 0.9$ Hz, 1H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 137.1, 136.3, 131.7, 131.4, 128.4, 126.8, 121.8, 120.2, 120.1, 119.4, 111.3, 99.2. Anal. Calcd for $\text{C}_{14}\text{H}_{10}\text{BrN}$: C, 61.79, H, 3.70, N, 5.15; Found; C, 61.59, H, 3.77, N, 5.07. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{14}\text{H}_{10}\text{BrN}$, 270.9997; found, 271.0002.

1-Methylindole (131.2 mg, 1.00 mmol, 1.0 equiv) and $\text{IMesPd}(\text{OAc})_2$ (26.5 mg, 0.05 mmol, 0.05 equiv) were dissolved in AcOH (10 mL), and the resulting solution was stirred at 25 °C for ~5 min. $[(p\text{-CH}_3\text{C}_6\text{H}_5)_2\text{I}]\text{BF}_4$ (791.9 mg, 2.00 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 25 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH_2Cl_2 (50 mL) and extracted with aqueous NaHCO_3 (2 x 30 mL). The organic layer was dried with MgSO_4 , concentrated, and the product was purified by chromatography on silica gel ($R_f = 0.21$ in 90% hexanes/10% benzene). The product (**1e**) obtained was a white solid (155 mg, 70% yield). Mp = 96-98 °C. ^1H NMR (400 MHz, CDCl_3): δ 7.65 (d, $J = 7.8$ Hz, 1H), 7.43 (d, $J = 8.2$ Hz, 2H), 7.38 (d, $J = 8.2$ Hz, 1H), 7.31-7.24 (multiple peaks, 3H), 7.16 (td, $J = 8.0, 1.0$ Hz, 1H), 6.55 (s, 1H), 3.76 (s, 3H), 2.45 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 141.5, 138.2, 137.6, 129.8, 129.2, 129.1, 127.9, 121.4, 120.3, 119.7, 109.5, 101.2, 31.0, 21.2. Anal. Calcd for $\text{C}_{16}\text{H}_{15}\text{N}$: C, 86.84, H, 6.83, N, 6.33; Found; C, 86.69, H, 6.89, N, 6.40. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{16}\text{H}_{15}\text{N}$, 221.1204; found, 221.1197.

1-Methylindole (111 mg, 0.84 mmol, 1.0 equiv) and IMesPd(OAc)₂ (22.3 mg, 0.042 mmol, 0.05 equiv) were dissolved in AcOH (8.5 mL), and the resulting solution was stirred at 25 °C for ~5 min. [(*p*-OMeC₆H₅)₂I]BF₄ (721 mg, 1.68 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (*R*_f = 0.23 in 70% hexanes/30% benzene). The product (**1f**) was obtained as a light brown solid (161 mg, 80% yield). Mp = 117-120 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.76 (d, *J* = 7.8 Hz, 1H), 7.54 (d, *J* = 8.8 Hz, 2H), 7.45 (d, *J* = 8.2 Hz, 1H), 7.36 (ddd, *J* = 7.9, 7.0, 1.1 Hz, 1H), 7.27 (ddd, *J* = 7.7, 6.9, 1.0 Hz, 1H), 7.10 (d, *J* = 8.8 Hz, 2H), 6.63 (s, 1H), 3.95 (s, 3H), 3.80 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 159.3, 141.3, 138.1, 130.5, 127.9, 125.1, 121.3, 120.2, 119.7, 113.9, 109.5, 100.9, 55.2, 30.9. Anal. Calcd for C₁₆H₁₅NO: C, 80.98, H, 6.37, N, 5.90; Found; C, 80.95, H, 6.37, N, 5.92. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₅NO, 237.1154; found, 237.1151.

1-Methylindole (116 mg, 0.88 mmol, 1.0 equiv) and IMesPd(OAc)₂ (23.4 mg, 0.044 mmol, 0.05 equiv) were dissolved in AcOH (8.5 mL), and the resulting solution was stirred at 25 °C for ~5 min. [(*p*-OMe₃C₆H₅)₂I]BF₄ (700 mg, 1.77 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH₂Cl₂ (50 mL) and extracted with aqueous NaHCO₃ (2 x 30 mL). The organic layer was dried with MgSO₄, concentrated, and the product was purified by chromatography on silica gel (*R*_f = 0.17 in 95% hexanes/5% benzene). The product (**1h**) was obtained as a light orange solid (121 mg, 62% yield). Mp = 90-92 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.66 (d, *J* = 7.8 Hz, 1H), 7.23-7.40 (multiple peaks, 6H), 7.16 (ddd, *J* = 7.8, 6.9, 1.0 Hz, 1H), 6.45 (s, 1H), 3.53 (s, 3H), 2.22 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 140.5, 138.0, 137.3, 132.5, 131.1, 130.0, 128.6, 128.0, 125.5, 121.2, 120.4, 119.6, 109.4, 101.5, 30.3, 20.0. Anal. Calcd for C₁₆H₁₅N: C, 86.84, H, 6.83, N, 6.33; Found; C, 86.57, H, 6.87, N, 6.26. HRMS electrospray (*m/z*): [M⁺] calcd for C₁₆H₁₅N, 221.1204; found, 221.1198.

Indole (94 mg, 0.80 mmol, 1.0 equiv) and $\text{IMesPd}(\text{OAc})_2$ (21.2 mg, 0.04 mmol, 0.05 equiv) were dissolved in AcOH (8.0 mL) and the resulting solution was stirred at 25 °C for ~5 min. $[(1\text{-Naphthyl})_2\text{I}]\text{BF}_4$ (749 mg, 1.60 mmol, 2.0 equiv) was added, and the reaction was stirred for 15 h at 60 °C. The reaction mixture was filtered through a plug of Celite and then evaporated to dryness. The resulting oil was dissolved in CH_2Cl_2 (50 mL) and extracted with aqueous NaHCO_3 (2 x 30 mL). The organic layer was dried with MgSO_4 , concentrated, and the product (**2c**) was purified by chromatography on silica gel (R_f = 0.20 in 95% hexanes / 5% ethyl acetate). The product was obtained as a white solid (131 mg, 67% yield). Mp = 99-102 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 11.59 (s, 1H), 8.31-8.37 (m, 1H), 7.96-8.05 (multiple peaks, 2H), 7.72 (dd, J = 7.1, 1.2 Hz, 1H), 7.56-7.65 (multiple peaks, 4H), 7.47 (d, J = 8.1 Hz, 1H), 7.16 (ddd, J = 8.0, 7.0, 1.2 Hz, 1H), 7.07 (ddd, J = 7.8, 7.0, 1.0 Hz, 1H), 6.75 (d, J = 2.1 Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 136.7, 136.4, 133.5, 130.9, 130.7, 128.3, 128.0, 127.2, 126.6, 126.0, 125.4, 121.3, 120.0, 119.2, 111.3, 102.4. Note: two of the ^{13}C peaks for this compound appear to be coincidentally overlapping. HRMS electrospray (m/z): $[\text{M}^+]$ calcd for $\text{C}_{18}\text{H}_{13}\text{N}$, 243.1048; found, 243.1042.

General Procedure for *in situ* Oxidant Generation for **1a and **1e**.** $\text{Pd}(\text{OAc})_2$ (8.5 mg, 0.04 mmol, 0.05 equiv), $\text{ArI}(\text{OAc})_2$ (1.52 mmol, 2 equiv) and $\text{ArB}(\text{OH})_2$ (1.52 mmol, 2 equiv) were combined in AcOH (7.6 mL). The resulting mixture was stirred for 15 min at room temperature. 1-Methylindole (100 mg, 0.76 mmol, 1 equiv) was then added and the reaction was stirred for 12 h at room temperature. The reactions were worked up and purified as described above for products **1a** and **1e** to afford the corresponding 2-arylindoles in 80% and 81% isolated yields, respectively.

General Procedure for *in situ* Oxidant Generation for **1d.** $\text{Pd}(\text{OAc})_2$ (8.5 mg, 0.04 mmol, 0.05 equiv), $p\text{-ClC}_6\text{H}_4\text{I}(\text{OAc})_2$ (1.09 g, 3.05 mmol, 4 equiv) and $p\text{-ClC}_6\text{H}_4\text{B}(\text{OH})_2$ (477 mg, 3.05 mmol, 4 equiv) were combined in AcOH (7.6 mL). The resulting mixture was stirred for 15 min at room temperature. 1-Methylindole (100 mg, 0.76 mmol, 1 equiv) was then added and the reaction was stirred for 12 h at room temperature. The reaction was worked up and purified as described above for product **1d**, to afford the corresponding product in 67% isolated yield.

General Procedure for *in situ* Oxidant Generation with $\text{MesI}(\text{OAc})_2$. $\text{Pd}(\text{OAc})_2$ (1.7 mg, 0.008 mmol, 0.05 eq), $\text{MesI}(\text{OAc})_2$ (114 mg, 0.31 mmol, 2.0 equiv) and $\text{ArB}(\text{OH})_2$ (0.31 mmol, 2.0 equiv) were combined in AcOH (1.56 mL), and the resulting mixture was stirred for 15 min at room temperature. 1-Methylindole (20.5 mg, 0.16 mmol, 1 equiv) was then added and the reaction was stirred for 12 h at room temperature. The reactions were analyzed by GC, and yields were determined by integration relative to an internal standard. Products **1a**, **1d** and **1e**, were obtained in 44%, 39% and 62% GC yields respectively.

Table S1. Catalyst Effects in the Pd-Catalyzed Arylation of 1-Methylindole with [Ph₂I]BF₄

ENTRY	CATALYST	YIELD (GC)
1	IMesPd(OAc) ₂	80%
2	Na ₂ PdCl ₄	66%
3	PdCl ₂ (BnCN) ₂	64%
4	PdCl ₂	59%
5	Pd(TFA) ₂	59%
6	PdBr ₂	58%
7	Pd(OAc) ₂	49%
8	PdI ₂	20%

References:

1. Wolfe, J. P.; Tomori, H.; Sadighi, J. P.; Yin, J.; Buchwald, S. L. *J. Org. Chem.* **2000**, *65*, 1158.
2. McKillop, A.; Kemp, D. *Tetrahedron* **1988**, *45*, 3299.
3. Chen, D. W.; Ochiai, M. J. *J. Org. Chem.* **1999**, *64*, 6804.
4. Shah, A.; Pike, V. W.; Widdowson, D. A. *J. Chem. Soc., Perkin Trans.* **1997**, 2463.
5. Jensen, D. R.; Sigman, M. S. *Org. Lett.* **2003**, *5*, 63.
6. Lane, B. S.; Sames, D. *Org. Lett.* **2004**, *6*, 2897.