Supporting information for:

Ru^{II}-SDP-Complexes Catalyzed Asymmetric Hydrogenation of ketones. The Effect of Alkali Metal Cation in The Reaction

**Jian-Hua Xie, Sheng Liu, Xiang-Hong Huo, Xu Cheng, Hai-Feng Duan, Bao-Ming Fan,
Li-Xing Wang, and Qi-Lin Zhou***

*State Key Laboratory and Institute of Elemento-organic Chemistry, Nankai University,
Tianjin 300071, China*

qlzhou@nankai.edu.cn

Table of contents

	Page
General	S2
Synthesis of 2-Arylcyclohexanone	S2-4
General Procedure for Asymmetric Hydrogenation of Aromatic Ketones	S4-5
General Procedure for Asymmetric Hydrogenation of 2-Arylcycloalkanones	S5-7
General Procedure for Kinetic Study on The Effect of Alkali Metal Cation in Hydrogenation of Acetophenone	S7-11
Reference	S11

General

All reactions and manipulations were performed in an argon-filled glovebox and using standard Schlenk techniques. ^1H and ^{13}C NMR spectra were recorded on 300 M spectrometer, and chemical shifts was reported in ppm downfield from internal $\text{Si}(\text{CH}_3)_4$. Anhydrous benzene and diethyl ether were distilled over sodium benzophenone ketyl under nitrogen. 2-Propanol and DMF were freshly distilled from calcium hydride before use. The ketone substrates were stirred over K_2CO_3 to remove acidic impurities and distilled under nitrogen prior to use. 2-arylcyclohexanones (**4a-j**),¹ 2-phenylcyclopentanones (**6**),¹ 2-phenylcycloheptanone (**7**)² and 1-phenyl-2-tetralone (**8**),² the catalyst **1a-d**³ and **9a-c**⁴ were prepared by the reported methods.

Synthesis of 2-Arylcyclohexanones¹

General procedure. Synthesis of 2-phenylcyclohexanone (4a). To a new prepared 80 mL phenylmagnesium bromide solution in ether (3.2 M, 0.26 mol) was added 2-chlorocyclohexanone (24 mL, 0.21 mol) in 88 mL of anhydrous ether via drop funnel under nitrogen at a rate only caused gentle refluxing. After the addition, the ether was completely distilled out, 70 mL of anhydrous benzene was added, and the mixture was refluxed for 8 hours. The reaction mixture was cooled to room temperature, hydrolyzed with water and extracted with 150 mL EtOAc. The extract was concentrated and distilled (106-108 °C/1 mmHg) to give 25 g of α -phenylcyclohexanone. Further purification by recrystallization with EtOAc provide the pure α -phenylcyclohexanone (**4a**) (22 g, 61 %) as colorless solid, mp 54-56 °C. ^1H NMR (300 MHz, CDCl_3) δ 1.78-1.90 (m, 2H), 1.98-2.07 (m, 2H), 2.13-2.20 (m, 1H), 2.22-2.30 (m, 1H), 2.41-2.56 (m, 2H), 3.62 (dd, J = 12.0, 5.4 Hz, 1H), 7.14 (d, J = 6.6 Hz, 2H), 7.28-7.36 (m, 3H).

2-(4-Methyl)phenylcyclohexanone (4b):⁵ oil; 45 % yield; ^1H NMR (300 MHz, CDCl_3) δ 1.78-1.84 (m, 2H), 1.95-2.05 (m, 2H), 2.22-2.28 (m, 1H), 2.12-2.17 (m, 1H), 2.31 (s, 3H), 2.39-2.56 (m, 2H), 3.57(dd, J = 12.0, 5.4 Hz, 1H), 7.03 (d, J = 7.8 Hz, 2H), 7.15 (d, J = 7.8 Hz, 2H).

2-(4-Methoxy)phenylcyclohexanone (4c):⁶ solid, m.p. 70-72 °C; 39 % yield; ^1H NMR (300 MHz, CDCl_3) δ 1.77-1.89 (m, 2H), 1.94-2.06 (m, 2H), 2.09-2.19 (m, 1H), 2.22-2.30 (m, 1H), 2.39-2.55 (m, 2H), 3.57 (dd, J = 12.0, 6.0 Hz, 1H), 3.80 (s, 3H), 6.88 (m, J = 9.0 Hz, 2H), 7.07 (m, J = 9.0 Hz, 2H).

2-(4-Chloro)phenylcyclohexanone (4d):⁶ solid, m.p. 76-78 °C; 51 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.79-1.91 (m, 2H), 1.96-2.03 (m, 2H), 2.15-2.29 (m, 2H), 2.40-2.56 (m, 2H), 3.59 (dd, *J* = 12.0, 5.4 Hz, 1H), 7.07 (m, *J* = 9.0 Hz, 2H), 7.30 (m, *J* = 9.0 Hz, 2H).

2-(4-Trifluoromethyl)phenylcyclohexanone (4e):⁷ solid, m.p. 99-101 °C; 31 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.81-1.92 (m, 2H), 1.95-2.07 (m, 2H), 2.18-2.33 (m, 2H), 2.45-2.59 (m, 2H), 3.67 (dd, *J* = 12.3, 6.0 Hz, 1H), 7.25 (m, *J* = 6.0 Hz, 2H), 7.59 (m, *J* = 6.0 Hz, 2H).

2-(3-Methyl)phenylcyclohexanone (4f):⁶ oil, 82-84 °C/0.015 mmHg; 54 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.78-1.87 (m, 2H), 1.98-2.07 (m, 2H), 2.09-2.18 (m, 1H), 2.23-2.30 (m, 1H), 2.34 (s, 3H), 2.40-2.57 (m, 2H), 3.57 (dd, *J* = 12.6, 5.1 Hz, 1H), 6.94 (d, *J* = 6.3 Hz, 2H), 7.07 (d, *J* = 7.8 Hz, 1H), 7.1 (d, *J* = 7.8 Hz, 1H).

2-(3-Methoxy)phenylcyclohexanone (4g):⁶ oil, 104-106 °C/0.015 mmHg; 35 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.78-1.89 (m, 2H), 1.95-2.06 (m, 2H), 2.09-2.18 (m, 1H), 2.23-2.30 (m, 1H), 2.39-2.56 (m, 2H), 3.58 (dd, *J* = 11.4 Hz, 5.4 Hz, 1H), 3.79 (s, 1H), 6.69 (t, *J* = 3.0 Hz, 1H), 6.73 (d, *J* = 7.5 Hz, 1H), 6.79 (dd, *J* = 7.5 Hz, 3.0 Hz, 1H), 7.26 (d, *J* = 7.5 Hz, 1H).

2-(3-Chloro)phenylcyclohexanone (4h):⁸ oil, 100-102 °C/0.015 mmHg; 45 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.73-1.90 (m, 2H), 1.92-2.06 (m, 2H), 2.13-2.21 (m, 1H), 2.24-2.31 (m, 1H), 2.40-2.58 (m, 2H), 3.59 (dd, *J* = 12.0, 5.4 Hz, 1H), 7.02 (d, *J* = 6.3 Hz, 1H), 7.13 (s, 1H), 7.27 (d, *J* = 6.3 Hz, 2H).

2-(3,5-Dimethyl)phenylcyclohexanone (4i): oil, 91-93 °C/0.015 mmHg; 40 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.73-1.90 (m, 2H), 1.95-2.20 (m, 3H), 2.20-2.28 (m, 1H), 2.30 (s, 6H), 2.42-2.60 (m, 3H), 3.53 (dd, *J* = 12.0, 5.4 Hz, 1H), 6.75 (s, 2H), 6.89 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 21.4, 25.3, 27.8, 34.9, 42.2, 126.3, 128.7, 137.7, 138.6, 210.7; MS (EI) *m/z* 202 (*M*⁺). Anal. Calcd for C₁₄H₁₈O: C, 83.12; H, 8.97. Found: C, 83.17; H, 9.03.

2-(2-Methyl)phenylcyclohexanone (4j):⁶ solid, m.p. 53-55 °C; 41% yield; ¹H NMR (300 MHz, CDCl₃) δ 1.73-1.90 (m, 3H), 1.95-2.17 (m, 2H), 2.20 (s, 6H), 2.20-2.29 (m, 3H), 3.78 (dd, *J* = 13.5, 5.1 Hz, 1H), 7.20 (m, 4H).

2-Phenylcyclopentanone (6):⁹ liquid, b.p. 118-120 °C/2 mmHg; 21 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.89-2.40 (m, 5H), 2.42-2.54 (m, 2H), 3.36-3.29 (m, 1H), 7.18-7.36 (m, 5H).

2-Phenylcycloheptanone (7):¹⁰ liquid, 123-125 °C/2 mmHg; 27 % yield; ¹H NMR (300 MHz, CDCl₃) δ 1.43-1.71 (m, 3H), 1.92-2.19 (m, 5H), 2.49-2.57 (m, 1H), 2.65-2.74 (m, 1H), 3.71 (dd, J = 11.4, 3.6 Hz, 1H), 7.18-7.34 (m, 5H).

1-Phenyl-2-tetralone (8):² solid, m.p. 36-38 °C; 68% yield; ¹H NMR (300 MHz, CDCl₃) δ 2.51-2.72 (m, 2H), 2.82-3.16 (m, 2H), 4.75 (s, 1H), 7.00 (d, J = 6.9 Hz, 1H), 7.08-7.12 (m, 2H), 7.18-7.36 (m, 6H).

Asymmetric Hydrogenation of Aromatic Ketones

General Procedure. Hydrogenation of acetophenone at S/C = 5000. The catalyst (0.002 mmol) was placed in a 20 mL hydrogenation vessel. The vessel was purged with hydrogen by pressurizing to 10 atm and releasing the pressure. The procedure was repeated three times. Anhydrous *i*-PrOH (3.0 mL) was introduced with a syringe and the vessel was purged with hydrogen and pressurized to 20 atm for 5 minutes. After releasing the pressure, acetophenone (1.2 g, 10 mmol) and a solution of *t*-BuOK in *i*-PrOH (0.2 mmol/mL, 0.7 mL, 0.14 mmol) was added. The vessel was purged with hydrogen and pressurized to 50 atm. After stirring at room temperature for several hours, the reaction was stopped. The reaction mixture was filtered through a short silica gel column, and the filtrate was diluted with acetone and analyzed by GC to determine the conversion and enantioselectivity. The ee of 1-phenylethanol was determined by GC using Suplco β -DEXTM 120 column, N₂ 2 mL/min, 100 °C then 1 °C/min to 160 °C, t_R (*R*) = 15.51 min, t_R (*S*) = 16.01 min.

Asymmetric Hydrogenation of 2-Arylcycloalkanones

General Procedure. Hydrogenation of 2-arylcycloalkanone at S/C = 2000, S/B = 10, at room temperature under 50 atm of hydrogen. The catalyst RuCl₂[(*S*)-Xyl-SDP][(*R,R*)-DPEN] (**1d**) (2.2 mg, 0.002 mmol) was placed in a 20 mL hydrogenation vessel. The vessel was purged with hydrogen by pressurizing to 10 atm and releasing the pressure. The procedure was repeated three times. Anhydrous *i*-PrOH (1.0 mL) was introduced with a syringe and the vessel was purged with hydrogen and pressurized to 20 atm for 5 minutes. After releasing the pressure, 2-arylcycloalkanone (4 mmol in 2.5 mL *i*-PrOH) and a solution of *t*-BuOK in *i*-PrOH (1 mmol/mL, 0.4 mL, 0.4 mmol) were added. The vessel was purged with hydrogen and pressurized to 50 atm. After stirring at room temperature for several hours, the reaction was stopped. The reaction mixture was filtered through a short silica

gel column, and the filtrate was diluted with acetone and analyzed by GC to determine the conversion, the *cis/trans*-selectivity and enantioselectivity.

(*S,S*)-*cis*-2-Phenylcyclohexanol (5a):¹⁰ 99% ee, $[\alpha]_{\text{D}}^{20} +97$ (*c* 1.06, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.26-1.48 (m, 3H), 1.52-1.74 (m, 3H), 1.82-2.10 (m, 3H), 2.72-2.83 (m, 1H), 4.03 (s, 1H), 7.24-7.37 (m, 5H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.8 °C/min to 180 °C) t_{R} (*R,R*) = 51.83 min, t_{R} (*S,S*) = 52.52 min.

***cis*-2-(4-Methyl)phenylcyclohexanol (5b):**¹¹ 99.5% ee, $[\alpha]_{\text{D}}^{20} +78$ (*c* 0.95, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.25-1.70 (m, 6H), 1.82-2.10 (m, 3H), 2.33 (s, 3H), 2.68-2.76 (m, 1H), 4.00 (s, 1H), 7.16 (s, 4H). GC (Suplco β -DEXTM 225 column: 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 180 °C) t_{R} (minor) = 93.78 min, t_{R} (major) = 94.55 min.

(*S S*)-*cis*-2-(4-Methoxy)phenylcyclohexanol (5c):¹¹ 97% ee, $[\alpha]_{\text{D}}^{20} +89$ (*c* 1.04, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.24-1.71 (m, 6H), 1.81-2.08 (m, 3H), 2.66-2.74 (m, 1H), 3.80 (s, 3H), 3.99 (s, 1H), 6.88 (d, *J* = 9.3 Hz, 2H), 7.19 (d, *J* = 9.3 Hz, 2H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.8 °C/min to 180 °C) t_{R} (*R,R*) = 126.05 min, t_{R} (*S,S*) = 127.07 min.

***cis*-2-(4-Chloro)phenylcyclohexanol (5d):**¹² 99% ee, $[\alpha]_{\text{D}}^{20} +77$ (*c* 1.00, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.18-1.70 (m, 6H), 1.82-2.08 (m, 3H), 2.68-2.74 (m, 1H), 4.01 (s, 1H), 7.32-7.19 (m, 4H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 180 °C) t_{R} (minor) = 128.06 min, t_{R} (major) = 129.06 min.

(*S,S*)-*cis*-2-(4-Trifluoromethyl)phenylcyclohexanol (5e):¹¹ 98% ee, $[\alpha]_{\text{D}}^{20} +96$ (*c* 1.10, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.20-1.70 (m, 6H), 1.82-2.10 (m, 3H), 2.69-2.75 (m, 1H), 4.00 (s, 1H), 7.19-7.22 (m, 2H), 7.29-7.33 (m, 2H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 180 °C) t_{R} (*R,R*) = 128.31 min, t_{R} (*S,S*) = 129.26 min.

***cis*-2-(3-Methyl)phenylcyclohexanol (5f):**¹³ 98% ee, $[\alpha]_{\text{D}}^{20} +100$ (*c* 0.99, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.24-1.74 (m, 6H), 1.82-2.11 (m, 3H), 2.35 (s, 3H), 2.68-2.75 (m, 1H), 4.01 (s, 1H), 7.00-7.08 (m, 3H), 7.21-7.22(d, *J* = 7.8 Hz, 1H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 80 °C then 0.5 °C/min to 180 °C) t_{R} (minor) = 132.13 min, t_{R} (major) = 132.84 min.

***cis*-2-(3-Methoxy)phenylcyclohexanol (5g):** 99.9% ee, $[\alpha]_{\text{D}}^{20} +92$ (*c* 1.03, CH₃OH); ¹H NMR

(300 MHz, CDCl₃) δ 1.23-1.72 (m, 6H), 1.73-2.21 (m, 3H), 2.68-2.78 (m, 1H), 3.81 (s, 1H), 4.02 (s, 1H), 6.77-6.87 (m, 3H), 7.23-7.29 (d, $J = 7.5$ Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 38.8, 43.6, 45.4, 52.1, 67.3, 74.3, 89.7, 130.8, 123.8, 139.2, 148.7, 165.0, 178.9; MS (EI) m/z 206 (M⁺). Anal. Calcd for C₁₃H₁₈O₂: C, 75.69; H, 8.80. Found: C, 75.74; H, 8.80. GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 80 °C then 0.5 °C/min to 180 °C) t_R (minor) = 160.90 min, t_R (major) = 161.56 min.

cis-2-(3-Chloro)phenylcyclohexanol (5h): ¹³ 98% ee, $[\alpha]_D^{20} +86$ (c 0.98, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.24-1.76 (m, 6H), 1.82-2.10 (m, 3H), 2.66-2.74 (m, 1H), 4.02 (s, 1H), 7.14-7.29 (m, 4H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 180 °C) t_R (minor) = 110.07 min, t_R (major) = 111.07 min.

cis-2-(3,5-Dimethyl)phenylcyclohexanol (5i): 94% ee, $[\alpha]_D^{20} +90$ (c 1.12, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.31-1.72 (m, 6H), 1.82-2.10 (m, 3H), 2.31 (s, 6H), 2.64-2.72 (m, 1H), 3.99 (s, 1H), 6.88 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 19.5, 24.5, 26.1, 33.2, 47.8, 70.4, 126.0, 126.6, 128.1, 129.7, 134.3, 146.3; MS (EI) m/z 204 (M⁺). Anal. Calcd for C₁₄H₂₀O: C, 82.30; H, 9.87. Found: C, 82.24; H, 9.64. HPLC (Chiralcel OJ column 25 cm, *i*-PrOH/Hex = 1:99, 1.0 mL/min) t_R (minor) = 5.11 min, t_R (major) = 5.72 min.

cis-2-(2-Methyl)phenylcyclohexanol (5j): ¹² 89% ee, $[\alpha]_D^{20} +102$ (c 1.14, CH₃OH). ¹H NMR (300 MHz, CDCl₃) δ 1.24-1.75 (m, 6H), 1.74-2.05 (m, 2H), 2.09-2.23 (m, 1H), 2.33 (s, 3H), 3.94-3.12 (m, 1H), 3.95 (s, 1H), 7.11-7.30 (m, 4H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 180 °C) t_R (minor) = 115.63 min, t_R (major) = 116.48 min.

(S,S)-cis-2-Phenylcyclopentanol: ¹¹ 60% ee, $[\alpha]_D^{20} +55$ (c 1.07, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.59-2.13 (m, 7H), 2.93-3.03 (m, 1H), 4.19-4.24 (m, 1H), 7.12-7.31 (m, 5H). GC (Suplco β -DEXTM 225 column 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.8 °C/min to 180 °C) t_R (minor) = 42.56 min, t_R (major) = 43.04 min.

(S,S)-cis-2-Phenylcycloheptanol: ¹¹ 87% ee, $[\alpha]_D^{20} +83$ (c 1.00, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ 1.43-1.64 (m, 4H), 1.65-1.98 (m, 6H), 2.81-2.90 (m, 1H), 3.98-4.03 (m, 1H), 7.18-7.35 (m, 5H). GC (Suplco β -DEXTM 225 column: 25 m $\times$ 0.25 mm $\times$ 0.25 μ m; N₂ 1 mL/min, 100 °C then 0.5 °C/min to 200 °C) t_R (minor) = 125.20 min, t_R (major) = 125.95 min.

(1*R*,2*S*)-cis-1-Phenyl-3,4-dihydro-1*H*-naphthalen-2-ol:² 72% ee, $[\alpha]_{\text{D}}^{20}$ –65 (*c* 1.01, CHCl₃); ¹HNMR (300 MHz, CDCl₃) δ 1.82-2.03 (m, 1H), 2.81-2.95 (dt, *J* = 6.9 Hz, 17.4 Hz, 1H), 3.05-3.18 (dt, *J* = 6.9 Hz, 17.4 Hz, 1H), 4.2 (m, 1H), 4.34 (d, *J* = 4.8 Hz, 1H), 6.92 (d, *J* = 7.5 Hz, 1H), 7.05-7.40 (m, 8H). HPLC (Chiralcel OD column 25 cm; *i*-PrOH/Hex = 5:95, 0.8 mL/min) *t*_R (*R,S*) = 10.65 min, *t*_R (*S,R*) = 13.01 min.

Kinetic Study on The Effect of Alkali Metal Cation in Hydrogenation of Acetophenone

General Procedure. A 120 mL stainless steel hydrogenation vessel was charged with (0.008 mmol) of **1** or **9**, hydrogen was introduced into the vessel to replace the argon for three times, and 15.4 mL *i*-PrOH was introduced under a hydrogen atmosphere. Then the vessel was closed and stirred for 5 min to ensure the catalyst **1** or **9** was completely dissolved. The vessel was opened and acetophenone 4.8 g (40 mmol), *t*-BuOM in *i*-PrOH 4.8 mL (0.1 mmol/mL, 4.8 mmol) were introduced into the vessel. The vessel was reclosed and pressurized to 50 atm. The reaction was vigorously stirred at 20 °C, and the samples were withdrawn by means of the sampling tube at regular time intervals. The conversions of acetophenone were determined by GC (HP-1 column 25 m × 0.25 mm × 0.25 μ m; N₂ 1 mL/min, 80 °C then 1 °C/min to 160 °C) *t*_R(subs) = 8.69 min, *t*_R(prod) = 8.96 min.

Table 1. Asymmetric hydrogenation of acetophenone using RuCl₂[(*S*)-SDP][(*R,R*)-DPEN] (1a**) as catalyst**

Entry	Time (min)	Conv. (%) (<i>t</i> -BuOK)	Conv. (%) (<i>t</i> -BuONa)	Conv. (%) (<i>t</i> -BuOLi)
1	5	1.3	1.2	
2	10	7.6	3.1	2.8
3	20	26.5	11.6	3.5
4	30	59.1	26.0	5.2
5	40	80.6	49.0	9.3
6	50	96.4	69.1	17.0
7	60	99.3	86.7	29.3
8	70	99.7	93.6	39.7
9	80	-	97.1	49.7

10	90	-	98.2	62.7
11	100	-	99.0	72.7
12	110	-	-	79.2
13	120	-	-	84.7

Table 2. Asymmetric hydrogenation of acetophenone using RuCl₂[(*S*)-An-SDP][(*R,R*)-DPEN] (1b) as catalyst

Entry	Time (min)	Conv. (%) (<i>t</i> -BuOK)	Conv. (%) (<i>t</i> -BuONa)	Conv. (%) (<i>t</i> -BuOLi)
1	10	5.4	6.8	1.7
2	20	24.3	31.0	5.5
3	30	55.0	58.9	17.3
4	40	88.2	87.5	31.5
5	50	98.2	96.2	48.4
6	60	99.6	98.8	63.3
7	70		99.6	74.9
8	80			83.1
9	90	-		89.2
10	100	-		93.1
11	110	-		95.8
12	120	-	-	97.5

Table 3. Asymmetric hydrogenation of acetophenone using RuCl₂[(*S*)-Tol-SDP][(*R,R*)-DPEN] (1c) as catalyst

Entry	Time (min)	Conv. (%) (<i>t</i> -BuOK)	Conv. (%) (<i>t</i> -BuONa)	Conv. (%) (<i>t</i> -BuOLi)
1	10	2.8	16.8	5.7
2	20	25.1	41.8	16.4
3	30	47.8	64.5	31.1
4	40	68.8	79.5	47.1
5	50	84.5	91.3	61.9
6	60	91.8	95.3	69.4
7	70	96.6	98.0	76.8

8	80	97.1	99.1	83.6
9	90	-	-	87.0
10	100	-	-	89.6
11	110	-	-	91.8
12	120	-	-	93.4

Table 4. Asymmetric hydrogenation of acetophenone using $\text{RuCl}_2[(S)\text{-Xyl-SDP}][(R,R)\text{-DPEN}]$ (1d) as catalyst

Entry	Time (min)	Conv. (%) (<i>t</i> -BuOK)	Conv. (%) (<i>t</i> -BuONa)	Conv. (%) (<i>t</i> -BuOLi)
1	5	3.2	4.3	1.0
2	10	10.1	21.1	2.4
3	20	28.1	54.7	10.2
4	30	46.2	81.4	22.6
5	40	59.6	89.0	34.5
6	50	72.2	94.9	45.7
7	60	79.2	97.9	56.8
8	70	86.4	99.1	64.5
9	80	89.9	99.7	70.6
10	90	94.4	-	75.6
11	100	96.4	-	80.8
12	110	98.5	-	85.6
13	120	-	-	88.5

Table 5. Asymmetric hydrogenation of acetophenone Using $\text{RuCl}_2[(R)\text{-Xyl-BINAP}][(R,R)\text{-DPEN}]$ (9c) as catalyst

Entry	Time (min)	Conv. (%) (<i>t</i> -BuOK)	Conv. (%) (<i>t</i> -BuONa)	Conv. (%) (<i>t</i> -BuOLi)
1	20	2.5	2.9	3.4
2	40	14.3	22.2	12.3
3	60	34.7	49.6	30.2
4	80	62.2	79.1	48.6

5	100	86.1	96.4	66.7
6	120	96.9	99.4	80.9
7	140	99.6	-	89.9
8	160	-	-	94.8
9	180	-	-	97.0

Table 6. The ee vs the conversion in the hydrogenation of acetophenone using $\text{RuCl}_2[(S)\text{-Xyl-SDP}][(R,R)\text{-DPEN}]$ (1d) as catalyst and *t*-BuOK as a base

Entry	Time (min)	Conv.(%)	Ee (%)
1	5	1.4	61.2
2	10	11.4	92.4
3	20	24.6	97.1
4	30	40.5	98.2
5	40	50.4	98.4
6	50	58.9	98.8
7	60	64.8	98.8
8	80	71.7	98.8
9	100	77.1	98.9
10	120	81.8	98.8
11	140	84.8	98.8
12	160	88.1	98.8
13	180	90.4	98.8
14	200	91.7	98.7

Reference:

- 1 Newman, M. S.; Farbman, M. D. *J. Am. Chem. Soc.* **1944**, *66*, 1550.
- 2 Alcock, N. J.; Mann, I.; Peach, P.; Wills, M. *Tetrahedron: Asymmetry* **2002**, *13*, 2485.
- 3 Xie, J.-H.; Wang, L.-X.; Fu, Y.; Zhu, S.-F.; Fan, B.-M.; Duan, H.-F.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2003**, *125*, 4404;
- 4 Doucet, H.; Ohkuma, T.; Murata, K.; Yokozawa, T.; Kozawa, M.; Katayama, E.; England, A. F.; Ikariya, T.; Noyori, R. *Angew. Chem. Int. Ed.* **1998**, *37*, 1703.

-
- 5 Huitric, A. C.; Kumler, W. D. *J. Am. Chem. Soc.* **1956**, 78, 614.
- 6 Mueller, G. P.; May, R. *J. Am. Chem. Soc.* **1949**, 71, 3313.
- 7 Blazejewski, J. C.; Cantacuzene, D.; Wakselman, C. *Tetrahedron* **1973**, 29, 4233.
- 8 Cecchi, P.; Pizzabiocca, A.; Renzi, G.; Chini, M.; Crotti, P.; Macchia, F.; Speranza, M. *Tetrahedron* **1989**, 45, 4227.
- 9 Hori, M.; Kataoka, T.; Shimizu, H.; Imai, E.; Iwamura, T.; Maeda, K. *Chem. Pharm. Bull.* **1986**, 34, 3599.
- 10 Ohkuma, T.; Li, J.; Noyori, R. *Synlett*, **2004**, 1383.
- 11 Huitric, A. C.; Carr, J. B. *J. Org. Chem.* **1961**, 26, 2648.
- 12 Huitric, A. C.; Clarke, W. G.; Leigh, K. J.; Staiff, D. C. *J. Org. Chem.* **1962**, 27, 715.
- 13 Timmerman, H.; Rekker, R. F.; Nauta, W. T. *Recl. Trav. Chim. Pays-Bas* **1965**, 84, 1348.