SUPPORTING INFORMATION

Scytalidamides A and B, New Cytotoxic Cyclic Heptapeptides from a Marine Fungus of the Genus *Scytalidium*.

Lik Tong Tan, Xing C. Cheng, Paul R. Jensen and William Fenical*

*Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,
University of California, San Diego, La Jolla, CA 92093-0204*

wfenical@ucsd.edu

Supporting Information:

Figure S1	¹ H NMR for scytalidamide A (1).
Figure S2	¹³ C NMR for scytalidamide A (1).
Figure S3	COSY plot for scytalidamide A (1).
Figure S4	HMQC plot for scytalidamide A (1).
Figure S5	HMBC plot for scytalidamide A (1).
Figure S6	ROESY plot for scytalidamide A (1).
Figure S7	¹ H NMR for scytalidamide B (2).
Figure S8	¹³ C NMR for scytalidamide B (2).
Figure S9	HMQC plot for scytalidamide B (2).
Figure S10	HMBC plot for scytalidamide B (2).
Figure S11	ROESY plot for scytalidamide B (2).
Figure S12	¹ H NMR for (2 <i>S</i> ,3 <i>S</i>)-3-MePro.
Figure S13	CD spectrum of (2 <i>S</i> ,3 <i>S</i>)-3-MePro.

Figure S1. ¹H NMR for scytalidamide A (1).

150.15
15191
12387

Figure S3. COSY plot for scytalidamide A (1).

F2
(ppm)

S4

Figure S5. HMBC plot for scytalidamide A (1).

scytalidamide A (colg)

700 MHz

0.4 M in CDCl₃

20034

Figure S6. ROESY plot for scytalidamide A (1).

¹H NMR of B (pyridine)
300 MHz

Figure S7. ¹H NMR for scytalidamide B (2).

S8

Figure S8. ^{13}C NMR for scytalidamide B (2).

S10

HMBC of B.

Figure S10. HMBC plot for scytalidamide B (2).

Figure S11. ROESY plot for scytalidamide B (2).

Figure S12. ^1H NMR for (2*S*,3*S*)-3-MePro.

S13

Figure S13. CD spectrum of (2*S*,3*S*)-3-MePro.

File Name : noname02.jws
 Date : 10/15/98 11:15AM [9/29/98 2:14PM]
 Sample : 3-methylproline
 Cell Length : 0.2 cm
 Concentration 0.0058 M
 Solvent : H₂O 0.5% TFA
 Temperature : 20 C
 Operator : Bruce & Jill
 Organization : UCD Chemistry
 Comment :