Predictive seafloor mapping by using high-resolution multibeam backscatter: a case study from the Palinuro Seamount (Eastern Tyrrhenian Sea)

Sara Innangi*, Salvatore Passaro*, Renato Tonielli*, Girolamo Milano***, Guido Ventura**, Stella Tamburrino*

* Istituto per l'Ambiente Marino Costiero, Consiglio Nazionale delle Ricerche, Calata Porta di Massa, 80133 Napoli, Italy
** Osservatorio Vesuviano - INGV - Via Diocleziano, 328 - 80124 - Napoli
*** Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata 605, 00143 Roma, Italy

Coordinate System: UTM Zone 33 North
Projection: Transverse Mercator
false easting: 500000
false northing: 0
central meridian: 15
scale factor: 0.9996
latitude of origin: 0
Linear Unit: Meter
Map limits: xmin= 478830.823
ymin= 4363324.541
xmax= 493968.261
ymax= 4379151.35
Geographic Coordinate System
Angular Unit: Degree (0.017453292519943299)
Prime Meridian: Greenwich
Datum: WGS84

Legend

- Acoustic Facies**
- BR - BedRock
 - BS - Biodeutritic Sand
 - CC - Coralligenous Concretion
 - CS - Coarse Sand
 - M - Mud
 - S - Sand