Supporting Information

Selective Dimerization of Arylalkynes to (*E*)-1,4-Diaryl Enynes Catalyzed by the [Ru(*p*-cymene)Cl₂]₂/Acetic Acid System, under Phosphine Free Conditions.

Mauro Bassetti,^{*} Chiara Pasquini, Antonino Raneri, Deborah Rosato

CNR, Istituto di Metodologie Chimiche, Dipartimento di Progettazione Molecolare, and Dipartimento di Chimica, Università La Sapienza, 00185 Roma, Italy

Table of Contents

General Experimental Section	page S2
Recycling of the Catalytic System	S3
Dimerization of phenylacetylene	S3
Dimerization of 1-ethynyl-4-nitrobenzene	S4
Dimerization of 1-ethynyl-4-chlorobenzene	S5
Dimerization of 1-ethynyl-3-fluorobenzene	S5
Dimerization of 1-ethynyl-4-methoxybenzene	S6
Dimerization of 1-ethynyl-2-methoxybenzene	S6
Dimerization of 4-ethynyl- α,α,α -trifluorotoluene	S7
Dimerization of 2-Ethynyl- α,α,α -trifluorotoluene	S8
Dimerization of ethynylferrocene	S8
Charts of ¹ H, ¹³ C NMR spectra and GC-MS analyses of the dimerization products	S9-S22

General Experimental Section. NMR spectra were recorded on either 300 or 200 MHz instruments. Chemical shifts are reported in δ values relative to tetramethylsilane, with reference to internal solvent (CHCl_3 , $\delta = 7.25$ ppm), and coupling constants (J) are given in Hz. Multiplicities are abbreviated as follows: singlet (s), doublet (d), quartet (q), multiplet (m) and virtual triplet (vt). The reported ^1H NMR data refer to the major olefin isomer unless stated otherwise. GC analyses were carried out with a 5% phenylsilicon, $30\text{m} \times 0.25$ mm, capillary column. FT-IR spectra were obtained in 0.1 mm CaF_2 solution cells or from KBr windows.

The dimerization reactions were carried out under argon using Schlenk-line techniques. Analytical thin-layer chromatography (TLC) was performed using silica gel 60 F254 pre-coated glass or aluminium plates. Flash chromatography was performed using silica gel 60 (0.040-0.063 mm). Petroleum ether (40° - 70°) for chromatography was distilled at atmospheric pressure before use. Glacial acetic acid was used without further purification. 1-Ethynyl-2-methoxybenzene and 1-ethynyl-4-chlorobenzene were available in the laboratory from previous work and were purified by column chromatography before use (Ricci, A.; Angelucci, F.; Bassetti, M.; Lo Sterzo, C. *J. Am. Chem. Soc.* **2002**, *124*, 1060). The other ethynylbenzene derivatives and ethynylferrocene were purchased from commercial suppliers and used as received.

The yields of 1,4-diphenylbut-1-ene-3-yne and of acetophenone in the gas-chromatographic analyses were determined using 1,2-diphenylethane as internal standard, with values of conversion factor 0.90 and 0.45, respectively. The addition of NBu_4PF_6 (30%), employed as a promoter in some ruthenium catalyzed reactions (Trost, B. M.; Rhee, Y. *J. Am. Chem. Soc.* **1999**, *121*, 11680-11683), gave 60% of calculated yield of 1,4-diphenylbut-1-ene-3-yne, under the conditions of Table 1, entry 11.

Recycling of the Catalytic System. The solvent mixture composed of acetic acid:water (4:1) was chosen in the attempt to make the catalytic system recyclable. In fact, two different phases separates out upon the addition of hexane to the reaction mixture (24 h). Upon stirring, the hexane layer extracts

the organic products and can be easily removed via cannula from the Schlenk tube, under argon, while the catalytic systems remains dissolved in the aqueous phase. Additional aliquots of phenylacetylene were added to perform subsequent reaction runs. However, the efficiency of the system was maintained for the first two runs, with conversion of **1** into the dimeric products around 40%, while it dropped significantly in the third run.

(E)-1,4-Diphenylbut-1-ene-3-yne (2a). [Ru(*p*-cymene)Cl₂]₂ (27 mg, 0.0455 mmol) and a magnetic bar were introduced into a 50 mL Schlenk tube, and 3 argon/vacuum cycles were performed under stirring. A solution of phenylacetylene **1** (100 μ L, 93 mg, 0.91 mmol) in acetic acid (5 mL) was degassed three times under vacuum/argon, then transferred via cannula into the tube. The amber solution (**I**, 9.1×10^{-3} M; **1**, 0.18 M) was kept under stirring at room temperature for 48 h. After removing the solvent under vacuum, a GC analysis of a filtered (silica) solution of the reaction mixture showed the following composition (GC percentages, in order of increasing retention time): phenylacetylene (6%), *p*-cymene (10%), acetophenone (3%), (*Z*)-1,4-diphenylbut-1-ene-3-yne (1%), (*E*)-1,4-diphenylbut-1-ene-3-yne (66%) (*E*:*Z* > 50). The thick brown oil was purified by chromatography over silica using petroleum ether as eluant. After elution of traces of starting material, a main fraction of product was collected (42 mg, yield 45%). The spectroscopic data are in agreement with the structure of (*E*)-1,4-diphenylbut-1-ene-3-yne.^{10,11a} Further elution with diethyl ether yielded a second fraction (32 mg), which analyzed by GC-MS showed various peaks tentatively assigned to the following products (in order of increasing retention times): Ph(CO)CH₂C $\equiv$ CPh (*m/z* = 220), PhCH₂(CO)C $\equiv$ CPh (*m/z* = 220), Ph(CO)CH₂CH=CHPh (*m/z* = 222), two isomers of PhC(OCOMe)=CH-CH=CHPh (*m/z* = 263), and other species with peaks at *m/z* = 206 and 204, resulting from fragmentation of oligomeric compounds. ¹H NMR: δ = 7.35 (m, 10H), 7.05 (d, *J* = 16.4 Hz, 1H), 6.39 (d, *J* = 16.2 Hz, 1H) ppm. ¹³C NMR: δ = 141.2, 136.3, 131.5, 128.7, 128.6, 128.3, 128.2, 126.2, 123.4, 108.1, 91.7, 88.8 ppm. GC-MS (*m/z*): 204, 189, 176, 126, 101, 89, 51 (99.64 % *E*).

Reaction of Phenylacetylene in AcOH/H₂O (4:1). After 3 argon/vacuum cycles performed on complex **I** (600 mg, 0.98 mmol) in a Schlenk tube, PhC≡CH (2.0 g, 19.6 mmol) and a mixture of acetic acid:water (4:1, 100 mL) were added to the tube. After additional argon/vacuum cycles, the solution was kept at 80 °C overnight. The mixture was extracted from diethyl ether/water, the solvent was removed from the organic phase, and the residue was purified by column chromatography over silica using distilled petroleum ether as eluant to give (*E* + *Z*)-1,4-diphenylbut-1-ene-3-yne (720 mg, 36% yield).

(*E*)-1,4-Di-*p*-nitrophenylbut-1-ene-3-yne (4a). 1-Ethynyl-4-nitrobenzene **3** (0.255 g, 1.73 mmol) and complex **I** (54 mg, 0.088 mmol) were dissolved in acetic acid (10 mL), yielding a dark orange solution. Consistent precipitation of a brown powder was observed during the course of the reaction. After 48 h, some hexane was added to the reaction mixture to assist further precipitation, and the precipitate was allowed to deposit in the flask upon standing in the fridge. The mother liquor was separated via cannula from the precipitate, which was then washed with small portions of methanol (× 3), and once with pentane. The residue was kept under vacuum, then purified by column chromatography, using dichloromethane as eluant. The product of interest was collected as a yellow-orange band. Removal of the solvent under vacuum yielded a brown powder (98 mg, 98% *E*). The mother liquor was poured into water/chloroform, the water phase extracted twice with chloroform, the collected organic phases washed until neutrality, and dried over sodium sulphate. After removal of the solvent, the residue was purified by column chromatography (dichloromethane/petroleum ether, 4:1), yielding additional material (50 mg, 92% *E*) (overall yield 58%). The spectroscopic data are in agreement with the structure of (*E*)-1,4-di-*p*-nitrophenylbut-1-ene-3-yne^{10b} and copies of ¹H and ¹³C NMR spectra are in this SI (see SFigure3). ¹H NMR: δ = 8.21 (m, 4H), 7.58 (m, 4H), 7.13 (d, *J* = 16.2 Hz), 6.53 (d, *J* = 16.5 Hz), 6.14 (d, *J* = 12.0) ppm (*E*:*Z* = 70:1). ¹³C NMR: δ = 147.8, 147.2, 141.9,

140.5, 132.3, 129.6, 127.0, 124.2, 123.7, 111.8, 92.9, 92.0 ppm. FT-IR (dichloromethane): 2198 (C≡C), 1595, 1521 (NO₂), 1344 (NO₂). GC-MS (m/z): 294, 231, 202, 189.

(*E*)-1,4-Di-*p*-chlorophenylbut-1-ene-3-yne (6a). Complex **I** (78 mg, 0.128 mmol) and 1-ethynyl-4-chlorobenzene (**5**, 350 mg, 2.56 mmol) were introduced into a Schlenk tube and dissolved in acetic acid (14 mL). A suspension started to form after few hours, then becoming an abundant precipitate. After 48 hours, the flask was cooled, the supernatant was removed via cannula and the solid washed twice with acetic acid. The crude product was purified by column chromatography using petroleum ether as eluant, to obtain 155 mg of compound **6a** (yield 44%, 99.5% of *E*). The spectroscopic data are in agreement with the structure of (*E*)-1,4-di-*p*-chlorophenylbut-1-ene-3-yne.^{6b} Copies of ¹H, ¹³C NMR spectra and of the GC-MS spectrum are in this SI (see SFigure4 and SFigure5). ¹H NMR: δ = 7.32 (m, 8H), 6.96 (d, *J* = 16.4 Hz, 1H), 6.31 (d, *J* = 16.1 Hz, 1H) ppm. ¹³C NMR: δ = 140.2, 134.6, 134.4, 134.3, 132.7, 129.0, 128.7, 127.5, 121.7, 108.4, 91.1, 89.5 ppm. FT-IR (dichloromethane): 2197 (C≡C), 1642, 1590, 1488, 1405, 1397. GC-MS (m/z): 272, 236, 202, 147, 100, 73 (0.30 %, *Z*); 272, 236, 202, 100 (99.47 %, *E*).

(*E*)-1,4-Di-*m*-fluorophenylbut-1-ene-3-yne (8a). A solution of complex **I** (78 mg) and 1-ethynyl-3-fluorobenzene **7** (305 mg, 2.54 mmol) in glacial acetic acid (15 mL) was kept under stirring for 24 h. The dark orange solution was poured into water/chloroform, the water phase extracted twice with chloroform, the collected organic phases washed with water until neutrality, and dried over sodium sulphate. After removal of the solvent, the remaining brown oil was purified by column chromatography, using hexane/chloroform 3/1, and the expected product collected as a yellowish oil (153 mg, 50% yield, 94% *E*). ¹H NMR: δ = 7.33-6.95 (m, 9H), 6.35 (d, *J* = 16.1 Hz, 1H) ppm. ¹³C NMR: δ = 163.0 (d, *J* = 245 Hz), 162.4 (d, *J* = 246 Hz), 140.5, 138.4 (d, *J* = 7 Hz), 130.1 (d, *J* = 24 Hz), 130.0 (d, *J* = 24 Hz), 127.4 (d, *J* = 4 Hz), 125.0, 122.3 (d, *J* = 4 Hz), 118.2 (d, *J* = 23 Hz), 115.6 (d, *J* = 21 Hz), 115.4 (d, *J* = 21 Hz), 112.7 (d, *J* = 23 Hz), 109.1, 91.2 (d, *J* = 3 Hz), 89.3 ppm. FT-IR

(dichloromethane): 2198 (C≡C), 1939, 1861, 1702, 1582, 1486, 1447. GC-MS (m/z): 240 (6.3%, *Z*), 240, 220, 144, 119 (93.7%, *E*). Anal. Calcd. for C₁₆H₁₀F₂: C, 79.99; H, 4.20. Found: C, 79.78; H, 4.25.

(*E*)-1,4-Di-*p*-methoxyphenylbut-1-ene-3-yne (10a). 1-Ethynyl-4-methoxybenzene **9** (196 μL, 200 mg, 1.51 mmol) and complex **I** (46 mg, 0.076 mmol) were dissolved in acetic acid (10 mL), yielding a dark orange solution. After 48 h, a yellow powder was allowed to deposit in the flask upon standing in the fridge, then separated via cannula from the reaction mixture and washed with acetic acid and then with pentane (× 3). The residue was dried under vacuum and purified on a short chromatographic column using petroleum ether/dichloromethane 7/3 as eluant and yielding 60 mg of **10a**, as a white yellowish powder (97% of isomer *E*). The reaction solution was poured into water and extracted with chloroform. Purification by column chromatography was carried out using as eluant a mixture of petroleum ether/dichloromethane 9/1, in order to remove traces of unreacted starting material. The composition in dichloromethane was increased progressively and the desired product eluted from a solvent mixture 6/4 (26 mg, 87% of isomer *E*). Overall yield 43%. R_f = 0.71 (hexane/chloroform 7/3). The spectroscopic data are in agreement with the structure of (*E*)-1,4-di-*p*-methoxyphenylbut-1-ene-3-yne^{6b} and copies of ¹H and ¹³C NMR spectra are in this SI (see SFigure8). ¹H NMR: δ = 7.29 (m, 4H), 6.85 (d, *J* = 16.2 Hz, 1H), 6.76 (m, 4H), 6.13 (d, *J* = 16.2 Hz, 1H), 3.715, 3.717 (2 s, 6H) ppm. ¹³C NMR: δ = 159.9, 159.4, 140.0, 132.8, 130.1, 129.3, 127.5, 115.7, 114.0, 105.9, 91.0, 87.9, 55.3, 55.2 ppm. FT-IR (dichloromethane): 2193 (C≡C), 1604, 1568, 1507, 1249.

1,4-Di-*o*-methoxyphenylbut-1-ene-3-yne (12). 1-Ethynyl-2-methoxybenzene **11** (224 mg, 1.7 mmol) and complex **I** (52 mg, 0.0847 mmol) were dissolved into 9.5 mL of acetic acid in a Schlenk tube and kept under stirring for 60 h at room temperature. The solvent was removed under vacuum, the residue was purified by column chromatography using a mixture of petroleum ether/dichloromethane (7/3). The dimeric product was obtained as a yellow oil (119 mg, 54% yield). The spectroscopic data

are in agreement with the structure of 1,4-di-*o*-methoxyphenylbut-1-ene-3-yne^{10a} and copies of ¹H and ¹³C NMR spectra and of GC-MS spectrum are in this SI (see SFigure9 and SFigure10).

¹H NMR: δ = 7.44-7.40 (m), 7.35 (d, J = 16.4 Hz), 7.11 (d, J = 12.3 Hz), 6.95-6.85 (m), 6.51 (d, J = 16.4 Hz), 5.96 (d, J = 12.0 Hz), 3.90, 3.89, 3.85, 3.83 ppm. ¹³C NMR: δ = 159.7, 156.9, 136.3, 133.4, 133.2, 132.1, 129.6, 129.5, 128.8, 126.8, 125.5, 120.4, 112.7, 110.9, 110.53, 110.50, 110.3, 109.0, 107.0, 92.6 and 93.8, 87.5 and 92.1, 55.7 and 55.8, 55.5, 55.4 ppm. FT-IR (dichloromethane): 2192, 1595, 1492, 1435. GC-MS (m/z): retention time 26.61 m (*Z*, 16.9%), m/z 264, 218, 205, 189, 176, 119, 91, 76; retention time 30.59 m (*E*, 82.7%) 264, 218, 205, 189, 176, 119, 91, 76.

(*E*)-1,4-Di-*p*-trifluoromethylphenylbut-1-ene-3-yne (14a). 4-Ethynyl- α,α,α -trifluorotoluene **13** (317 mg, 1.86 mmol) and complex **I** (58 mg, 0.093 mmol) were dissolved in acetic acid (10 mL), and the solution left under stirring for 24 h. The resulting solution was poured in water and extracted with chloroform. After washing the organic solution with water and drying over sodium sulphate, the solvent was removed under vacuum. The crude product was filtered on a short chromatographic column using hexane/chloroform (7/3) as eluant. The main fraction was further purified by column chromatography and eluted using petroleum ether/chloroform (9/1), to give, after removal of the solvent under vacuum, a yellow powder (101 mg, R_f = 0.76, yield 32%). The powder was crystallized from petroleum. The spectroscopic data are in agreement with the structure of 1,4-di-*p*-trifluoromethylphenylbut-1-ene-3-yne^{10b,6b} and copies of ¹H and ¹³C NMR spectra are in this SI (see SFigure11). ¹H NMR: δ = 7.63-7.50 (m, 8H), 7.09 (d, J = 16.3 Hz, 1H), 6.46 (d, J = 16.3 Hz, 1H) ppm. ¹³C NMR: δ = 140.7, 139.3, 131.7, 131.8, 130.8, 130.2, 126.6, 125.8 (q, J_{CF} = 3.8 Hz), 125.4 (q, J_{CF} = 3.8 Hz), 110.2, 91.5, 90.5, 77.6, 76.4 ppm; the two lower frequency peaks of the quartets due to the CF₃ groups are observed at 121.3, 115.9 (J_{CF} = 272 Hz), and 121.2, 115.8 (J_{CF} = 273 Hz) ppm. FT-IR (dichloromethane): 2199, 1923, 1617, 1575, 1410, 1321.

1,4-Di-*o*-trifluoromethylphenylbut-1-ene-3-yne (16). 2-Ethynyl- α,α,α -trifluorotoluene **15** (322 mg, 1.89 mmol) and complex **I** (58 mg, 0.093 mmol) in acetic acid (10 mL) were kept under stirring at room temperature for 40 h. The solution was then poured in water and extracted with chloroform ($\times 3$). The organic solution was washed with water ($\times 3$) and dried over sodium sulphate. The solvent was removed from the filtrate under vacuum and the crude product was purified by column chromatography. The main fraction was collected from petroleum ether/chloroform (9/1) as a light orange oil after removal of the solvent (77 mg). ^1H NMR and GC-MS are in agreement with a mixture of dimeric products (*E* + *Z*)-1,4-di-*o*-trifluoromethylphenylbut-1-ene-3-yne (**16a,b**) (77%) and of the three isomers of tris-*o*-trifluoromethylphenylbenzene (23%). Copies of ^1H and of GC-MS spectra are in this SI (see SFigure12 and SFigure13). ^1H NMR: δ = 7.71-7.24, 7.03-6.89 (m), 6.41 (d, J = 16.1 Hz), 6.11 (d, J = 12.1 Hz), 5.85 (C_6H_3 of trimers) ppm. GC-MS (m/z): 340 (5%, *Z*); 340, 271, 251, 202, 169, 150, 125, 69 (95%, *E*); 510, 401, 296, 190, 151 ($(2\text{-CF}_3\text{C}_6\text{H}_4)_3\text{C}_6\text{H}_3$).

(*E*)-1,4-Diferrocenylbut-1-ene-3-yne (18a). Ethynylferrocene **17** (0.40 g, 1.90 mmol) and complex **I** (59 mg, 0.095 mmol) were dissolved in acetic acid (5 mL), yielding a red brown solution. After 24 h, about $\frac{1}{4}$ of the solvent is removed under vacuum from the reaction mixture, allowing for further precipitation of a fine powder. After separating the solution from the powder, the solid residue was washed with small portions of acetic acid ($\times 3$), and of pentane ($\times 2$), then purified by column chromatography, starting with a mixture petroleum ether/ CH_2Cl_2 10/1 as eluant, and eluting the compound of interest with a 4/1 mixture. Removal of the solvent under vacuum yielded a crystalline orange solid (107 mg, 27% yield) of **18a**. The spectroscopic data match those reported for (*E*)-1,4-diferrocenyl-1-ene-3-yne in *E:Z* (1:1) mixture.²³ The compound can be further crystallized from a solution in CDCl_3 layered with heptane. ^1H NMR: δ = 6.72 (d, J = 15.9 Hz, 1H), 5.84 (d, J = 15.9 Hz, 1H), 4.42 (vt, J = 1.8 Hz, 2H) 4.37 (vt, J = 1.8 Hz, 2H) 4.27 (vt, J = 1.8 Hz, 2H), 4.22 (s, 5H), 4.20 (vt, J = 1.8 Hz, 2H), 4.14 (s, 5H) ppm. ^{13}C NMR: δ = 139.0, 105.9, 89.0, 85.4, 71.1, 70.0, 68.8, 67.0, 66.1

ppm. FT-IR (dichloromethane): 3100, 3031, 2195 ($\text{C}\equiv\text{C}$), 1612, 1411, 1106, 1043, 1027, 1003, 937 cm^{-1} ; (mineral oil): 948, 817, 722 cm^{-1} . Anal. Calcd. for $\text{C}_{24}\text{H}_{20}\text{Fe}_2$: C, 68.62; H, 4.80. Found: C, 68.19; H, 5.16. The mother liquor was poured into water/chloroform, the water phase extracted twice with chloroform, the collected organic phases washed until neutrality and dried over sodium sulphate. The residue was purified by column chromatography eluting first with petroleum ether/chloroform to recover traces of starting material and of **18a**. The chloroform content of the eluting solvent mixture was progressively increased, and a dark red band corresponding to acetylferrocene (163 mg, 38% yield) was then collected from pure chloroform.

Sample Name: analisi dimero purificato a T=120°C
Misc Info :
Vial Number: 1

SFigure1. GC spectrum of (E)-1,4-diphenylbut-1-ene-3-yne (2a).

a) ^1H NMR

b) ^{13}C NMR

Figure 2. ^1H and ^{13}C NMR spectra of (E)-1,4-diphenylbut-1-ene-3-yne (2a) in CDCl_3 .

a) ^1H NMR

b) ^{13}C NMR

Figure 3. ^1H and ^{13}C NMR spectra of (E)-1,4-di-*p*-nitrophenylbut-1-ene-3-yne (**4a**) in CDCl_3 .

a) ^1H NMR

b) ^{13}C NMR

Figure 4. ^1H and ^{13}C NMR spectra of (E)-1,4-di-*p*-chlorophenylbut-1-ene-3-yne (**6a**) in CDCl_3 .

Sample Name: dimero p-ClPhCCH purificato
 Misc Info :
 Vial Number: 1

Area Percent Report

Data File : C:\HPCHEM\1\DATA\NIESP43.D Vial: 1
 Acq On : 3 May 2005 10:51 Operator: ninor
 Sample : dimero p-ClPhCCH purificato Inst : GC/MS Ins
 Misc : Multiplr: 1.00
 Sample Amount: 0.00

MS Integration Params: autoint1.e

Method : C:\HPCHEM\1\METHODS\OSVALDO.M (Chemstation Integrator)
 Title :

Signal : TIC

peak #	R.T. min	first scan	max scan	last scan	PK TY	peak height	peak area	peak % max.	% of total
1	20.474	1572	1590	1598	BV	1455	45349	0.22%	0.220%
2	27.141	2190	2198	2214	VV 3	1033	63109	0.31%	0.306%
3	30.508	2468	2505	2519	PV	245142	20547898	100.00%	99.475%

Sum of corrected areas: 20656356

SFigure5. GC-MS spectrum of (E)-1,4-di-p-chlorophenylbut-1-ene-3-yne (6a).

SFigure6. GC-MS spectrum of (*E*)-1,4-di-*m*-fluorophenylbut-1-ene-3-yne (8a).

a) ^1H NMR

b) ^{13}C NMR

Figure 7. ^1H and ^{13}C NMR spectra of (*E*)-1,4-di-*m*-fluorophenylbut-1-ene-3-yne (**8a**) in CDCl_3 .

a) ^1H NMR

b) ^{13}C NMR

SFigure8. ^1H and ^{13}C NMR spectra of (*E*)-1,4-di-*p*-methoxyphenylbut-1-ene-3-yne (**10a**) in CDCl_3 .

a) ^1H NMR

b) ^{13}C NMR

Figure 9. ^1H and ^{13}C NMR spectra of 1,4-di-o-methoxyphenylbut-1-ene-3-yne (12) in CDCl_3 .

Sample Name: dimero o-OMePhCCH purificato
 Misc Info :
 Vial Number: 1

Area Percent Report

Data File : C:\HPCHEM\1\DATA\REG3.D
 Acq On : 3 May 2005 13:27
 Sample : dimero o-OMePhCCH purificato
 Misc :

Vial: 1
 Operator: ninor
 Inst : GC/MS Ins
 Multiplr: 1.00
 Sample Amount: 0.00

MS Integration Params: autoint1.e

Method : C:\HPCHEM\1\METHODS\OSVALDO.M (Chemstation Integrator)
 Title :

Signal : TIC

peak #	R.T. min	first scan	max scan	last scan	PK TY	peak height	peak area	peak % max.	% of total
1	11.604	769	781	802	PV 3	3814	167744	0.27%	0.224%
2	14.883	1070	1080	1100	BV 4	2132	129878	0.21%	0.174%
3	26.616	2119	2150	2172	VV	189751	12662057	20.47%	16.923%
4	30.586	2452	2512	2535	PV 3	400212	61862159	100.00%	82.679%

Sum of corrected areas: 74821837

SFigure10. GC-MS spectrum of 1,4-di-o-methoxyphenylbut-1-ene-3-yne (12).

a) ^1H NMR

b) ^{13}C NMR

Figure 11. ^1H and ^{13}C NMR spectra of (*E*)-1,4-di-*p*-trifluoromethylphenylbut-1-ene-3-yne (**14a**) in CDCl_3 .

a) ^1H NMR

SFigure12. ^1H spectrum of the mixture of 1,4-di-*o*-trifluoromethylphenylbut-1-ene-3-yne (16) and of (*o*-CF₃C₆H₄)₃C₆H₃, in CDCl₃.

SFigure13. GC-MS spectra of 1,4-di-*o*-trifluoromethylphenylbut-1-ene-3-yne (16) and of the 3 isomers of (*o*-CF₃C₆H₄)₃C₆H₃ (retention times 30.60, 33.46, 33.91 min).

a) ^1H NMR

SFigure14. ^1H and ^{13}C NMR spectra of (E)-1,4-diferrocenylbut-1-ene-3-yne (18a) in CDCl_3 .