Selective Formation of Six Membered Cyclic Sulfones and Sulfonates by C-H Insertion

Jinu P. John and Alexei Novikov*

Supporting Information

Experimental procedures, spectroscopic and analytical data for compounds 2-4, 6-9, 11-13.

General Information. All reactions were carried out under an inert atmosphere of dry nitrogen in oven or flame-dried glassware. Proton magnetic resonance spectra were recorded at 500 MHz on an Avance 500 Bruker spectrometer. Carbon magnetic resonance spectra were recorded at 125 MHz on an Avance 500 Bruker spectrometer. All chemical shifts were reported in δ units relative to tetramethylsilane. Optical rotations were measured on a Perkin-Elmer 341 Polarimeter. Infrared spectra were recorded on ATI Mattson Genesis Series FTIR spectrometer. High resolution mass spectral data were obtained on the Agilent 61969A TOF high resolution mass spectrometer using electrospray ionization, direct infusion, 10ml/min in 50% MeOH 5mM ammonium formate. Melting points were determined on a MEL-TEMP melting point apparatus. Analytical thin-layer chromatography (TLC) was performed using pre-coated TLC plates with silica Gel 60 F254 (E. Merck no. 5715-7). Flash column chromatography was performed using 40-63 μ m silica gel (Merck, Geduran, no. 11567-1) as the stationary phase. Acetone was purchased from Fisher (HPLC grade) and used without additional purification. Tetrahydrofuran (THF) was dried by distilling from sodium-benzophenone in a continuous still under an atmosphere of nitrogen. Dichloromethane, di-*iso*-propylethylamine, triethylamine, and acetonitrile were distilled from calcium hydride under an atmosphere of nitrogen.

Ethyl alkylthioacetates (2a-c). General Procedure. The corresponding bromide (**1a-c**) (1 equivalent, 0.5-5 mmol) was dissolved in acetone (20 ml/mmol), K_2CO_3 (1.5 equivalent) was added, followed by of ethyl-2-mercaptoacetate (1.1 equivalent) and catalytic amount (0.05 equivalent) of NaI. The reaction mixture was then refluxed for 12 h. After cooling to room temperature the volatiles were evaporated under reduced pressure. The residue was taken up in water (30 ml) and the crude product was extracted with EtOAc (2x30 ml/mmol). The combined extract was then washed with water and brine. The organic layer was dried over anhydrous Na_2SO_4 . The volatiles were evaporated under reduced pressure to yield the crude mixture that was purified on silica column (Ethyl acetate-Hexanes, 0 to 10%) to yield the pure product.

Thus prepared:

2a

Yield: 90%. **1H NMR** ($CDCl_3$, 500 MHz): 4.20 (q, $J=7$ Hz, 2H), 3.21 (s, 2H), 2.64 (t, $J=7$ Hz, 2H), 1.61 (quintet, $J=7$ Hz, 2H), 1.31-1.41 (m, 4H), 1.29 (t, $J=7$ Hz, 3H), 0.90 (t, $J=7$ Hz, 3H). **^{13}C NMR** ($CDCl_3$, 125 MHz): δ 170.8, 61.5, 33.9, 32.9, 31.1, 28.9, 22.4, 14.4, 14.1. **HRMS** (ESI) calcd for $C_9H_{19}O_2S$ [$M+H$] 191.1106, found 191.1105.

2b

Yield: 85%. **1H NMR** ($CDCl_3$, 500 MHz): δ 4.20 (q, $J=7$ Hz, 2H), 3.21 (s, 2H), 2.62 (t, $J=7$ Hz, 2H), 1.51-1.66 (m, 3H), 1.24-1.35 (m, 5H), 1.30 (t, $J=7$ Hz), 0.89 (d, $J=7$ Hz, 6H). **^{13}C NMR** ($CDCl_3$, 125 MHz): δ 170.9, 61.5, 38.2, 34.0, 33.2, 27.9, 27.1, 22.7, 14.4. **HRMS** (ESI) calcd for $C_{10}H_{24}NO_2S$ [$M+NH_4$] 222.1528, found 222.1522.

2c

Yield: 88%. **¹H NMR** (CDCl₃, 500 MHz): δ 8.17 (d, J=9 Hz, 1H), 7.88 (d, J=8 Hz, 1H), 7.81 (d, J=8 Hz, 1H), 7.57 (t, J=8 Hz, 1H), 7.52 (t, J=8 Hz, 1H), 7.39-7.47 (m, 2H), 4.33 (s, 2H), 4.22 (q, J=7 Hz, 2H), 3.13 (s, 2H), 1.31 (t, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 170.7, 134.3, 132.6, 131.5, 129.0, 128.6, 128.0, 126.4, 126.1, 125.2, 124.2, 61.6, 34.2, 33.0, 14.4. **HRMS** (ESI) calcd for C₁₅H₂₀NO₂S [M+NH₄]⁺ 278.1215, found 278.1209.

Sulfones 3a-c. General Procedure. The corresponding sulfide (**2a-c**) (1 equivalent, 0.5-5 mmol) was dissolved in CH₂Cl₂ (5 ml/mmol) and *m*-CPBA (2.5 equivalent) was added to it at 0°C (ice bath). The reaction mixture was allowed warm up to rt and stirred for 8 h at rt. Volatiles were evaporated under reduced pressure. The residue was taken up in EtOAc (50 ml) and washed with 10% Na₂SO₃, saturated NaHCO₃, and brine. The organic layer was then dried over anhydrous Na₂SO₄. The volatiles were evaporated under reduced pressure to yield the crude mixture that was purified on silica column (Ethyl Acetate-Hexanes, 0 to 50%) to yield the pure product.

Thus prepared:

3a

Yield: 95%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.24 (q, J=7 Hz, 2H), 3.88 (s, 2H), 3.17-3.21 (m, 2H), 1.78-1.86 (m, 2H), 1.36-1.43 (m, 2H), 1.29-1.36 (m, 2H), 1.27 (t, J=7 Hz, 3H), 0.87 (t, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 163.3, 62.9, 57.6, 54.0, 30.6, 22.3, 21.8, 14.2, 13.9. **HRMS** (ESI) calcd for C₉H₂₂NO₄S [M+NH₄]⁺ 240.1270, found 240.1264.

3b

Yield: 90%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.29 (q, J=7 Hz, 2H), 3.95 (s, 2H), 3.22-3.27 (m, 2H), 1.84-1.94 (m, 2H), 1.61 (nonet, J=7 Hz, 1H), 1.30-1.38 (m, 5H), 0.92 (d, J=7 Hz, 6H). **¹³C NMR** (CDCl₃, 125 MHz): δ 163.3, 62.9, 57.6, 53.9, 37.6, 27.8, 22.5, 20.0, 14.2. **HRMS** (ESI) calcd for C₁₀H₂₄NO₄S [M+NH₄]⁺ 254.1426, found 254.1420.

3c

Yield: 85%. **¹H NMR** (CDCl₃, 500 MHz): δ 8.21 (d, J=9 Hz, 1H), 7.92 (d, J=8 Hz, 1H), 7.89 (d, J=8 Hz, 1H), 7.71 (d, J=7 Hz, 1H), 7.61 (t, J=8 Hz, 1H), 7.47-7.56 (m, 2H), 5.06 (s, 2H), 4.35 (q, J=7 Hz, 2H), 3.93 (s, 2H), 1.37 (t, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 163.7, 134.2, 132.5, 130.9, 130.5, 129.0, 127.2, 126.5, 125.5, 124.2, 124.0, 62.9, 56.7, 55.6,

14.1. **HRMS** (ESI) calcd for $C_{15}H_{20}NO_4S$ $[M+NH_4]$ 310.1113, found 310.1107.

Sulfonates 5a-d. General Procedure. The corresponding alcohol (1 equivalent) was dissolved in CH_2Cl_2 (5 ml/mmol), and chlorosulfonyl acetic acid ethyl ester (1.1 mmol) was added. The reaction mixture was cooled to $-10\text{ }^{\circ}C$, followed by addition of Et_3N (1.5 equivalent). The reaction mixture was kept at $-10\text{ }^{\circ}C$ for 12 h. After warming up to room temperature, the reaction mixture was diluted with EtOAc (40 ml), washed with 1M HCl, saturated $NaHCO_3$ and brine, and dried over anhydrous Na_2SO_4 . The volatiles were evaporated under reduced pressure to yield the crude mixture that was purified on silica column (Ethyl Acetate-Hexanes, 0 to 50%) to yield the pure product.

Thus prepared:

6a

Yield: 80%. **1H NMR** ($CDCl_3$, 500 MHz): δ 4.39 (t, $J=7$ Hz, 2H), 4.29 (q, $J=7$ Hz, 2H), 4.08 (s, 2H), 1.77 (nonet, $J=7$ Hz, 1H), 1.66 (q, $J=7$ Hz, 2H), 1.33 (t, $J=7$ Hz, 3H), 0.95 (d, $J=7$ Hz, 6H). **^{13}C NMR** ($CDCl_3$, 125 MHz): δ 162.2, 71.2, 62.8, 54.9, 37.9, 24.6, 22.4, 14.1. **HRMS** (ESI) calcd for $C_9H_{22}NO_5S$ $[M+NH_4]$ 256.1219, found 256.1213.

6b

Yield: 85%. **1H NMR** ($CDCl_3$, 500 MHz): δ 4.96-5.06 (m, 1H), 4.28 (q, $J=7$ Hz, 2H), 4.06 (s, 2H), 1.70-1.78 (m, 2H), 1.32-1.50 (m, 7H), 1.46 (d, $J=6$ Hz), 1.33 (t, $J=7$ Hz), 0.96 (d, $J=7$ Hz, 3H), 0.93 (d, $J=6$ Hz, 3H). **^{13}C NMR** ($CDCl_3$, 125 MHz): δ 162.3, 81.4, 62.8, 56.2, 46.0, 24.7, 22.8, 22.5, 21.7, 14.2. **HRMS** (ESI) calcd for $C_{10}H_{24}NO_5S$ $[M+NH_4]$ 270.1375, found 270.1369.

6c

Yield: 30%. **1H NMR** ($CDCl_3$, 500 MHz): δ 8.21 (d, $J=8$ Hz, 1H), 7.91 (d, $J=8$ Hz, 1H), 7.84 (d, $J=8$ Hz, 1H), 7.55-7.66 (m, 3H), 7.49 (t, $J=8$ Hz, 1H), 4.30-4.37 (m, 4H), 4.34 (q, $J=7$ Hz), 4.35 (s), 1.34 (t, $J=7$ Hz, 3H). **^{13}C NMR** ($CDCl_3$, 125 MHz): δ 161.7, 145.4, 135.1, 128.1, 127.9, 127.5, 127.3, 127.2, 125.5, 121.8, 118.5, 63.3, 55.1, 14.5. **HRMS** (ESI) calcd for $C_{14}H_{18}NO_5S$ $[M+NH_4]$ 312.0906, found 312.0900.

6d

Yield: 85%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.66 (td, J=11, 4 Hz, 1H), 4.27 (q, J=7 Hz, 2H), 4.06 (s, 2H), 2.25-2.31 (m, 1H), 2.10 (septet of d, J=7, 3 Hz, 1H), 1.65-1.76 (m, 2H), 1.37-1.54 (m, 2H), 1.25-1.35 (m, 4H), 1.33 (t, J=7 Hz), 1.06 (qd, J=13, 3H, 1H), 0.80-1.00 (m, 10H), 0.94 (d, J=6Hz), 0.92 (d, J=7Hz), 0.83 (d, 7Hz). **¹³C NMR** (CDCl₃, 125 MHz): δ 162.2, 85.3, 62.8, 56.4, 47.7, 42.2, 33.9, 31.9, 25.8, 23.2, 22.0, 21.0, 15.7, 14.2. **HRMS** (ESI) calcd for C₁₄H₃₀NO₅S [M+NH₄]⁺ 324.1845, found 324.1839. m.p. (°C): 43-44.

Preparation of Diazo Compounds (4a-c, 7a,b,d). General Procedure. The corresponding sulfone or sulfonate (1 equivalent, 0.2-2) was dissolved in MeCN (5 ml/mmol) along with *p*-nitrosulfonylazide (1.1 equivalent), and cooled to -10°C. *i*-PrNEt₂ (1 equivalent) was added, and the reaction mixture was kept at -10°C for 12 h, then warmed up to rt at stirred for additional 5 h. After evaporation of the volatiles, the residue was taken up in chloroform (10 ml). After several minutes *p*-nitrosulfonamide precipitated and was filtered off, and washed with additional portions of chloroform. The volatiles were evaporated under reduced pressure to yield the crude product that was purified on silica column (Ethyl Acetate-Hexanes, 0 to 40%) to yield the pure product.

Note: the peak for the carbon next to diazo group has not been detected in the ¹³C NMR spectra for any of the prepared diazo compounds even when the acquisition time was raised to 15 sec. IR data are provided, showing characteristic peak at circa 2130.

Thus prepared:

4a

Yield: 70%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.33 (q, J=7 Hz, 2H), 3.35-3.39 (m, 2H), 1.79-1.86 (m, 2H), 1.35-1.45 (m, 4H), 1.33 (t, J=7 Hz, 3H), 0.91 (t, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 160.3, 62.7, 56.7, 30.3, 22.5, 22.3, 14.4, 13.8. **IR** (CHCl₃, cm⁻¹): 2131, 1712, 1336, 1147.

4b

Yield: 70%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.34 (q, J=7 Hz, 2H), 3.34-3.39 (m, 2H), 1.79-1.88 (m, 2H), 1.58 (nonet, J=7 Hz, 1H), 1.27-1.38 (m, 5H), 1.34 (t, J=7 Hz), 0.91 (d, J=7 Hz, 6H). **¹³C NMR** (CDCl₃, 125 MHz): δ 160.3, 62.7, 57.1, 37.2, 27.8, 22.5, 20.8, 14.5. **IR** (CHCl₃, cm⁻¹): 2131, 1712, 1338, 1147.

4c

Yield: 75%. **¹H NMR** (CDCl₃, 500 MHz): δ 8.14 (d, J=8 Hz, 1H), 7.94 (d, J=8 Hz, 1H), 7.91 (d, J=8 Hz, 1H), 7.48-7.63 (m, 4H), 5.12 (s, 2H), 4.36 (q, J=7 Hz, 2H), 1.35 (t, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 160.4, 134.1, 132.4, 130.9, 130.6, 129.1, 127.3, 126.5, 125.4, 123.9, 123.7, 62.8, 59.4, 14.5. **IR** (CHCl₃, cm⁻¹): 2133, 1710, 1344, 1265, 1160.

7a

Yield: 65%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.39 (t, J=7 Hz, 2H), 4.34 (q, J=7 Hz, 2H), 1.77 (nonet, J=7 Hz, 1H), 1.65 (q, J=7 Hz, 2H), 1.34 (t, J=7 Hz, 3H), 0.94 (d, J=7 Hz, 6H). **¹³C NMR** (CDCl₃, 125 MHz): δ 159.8, 72.4, 62.9, 37.7, 24.7, 22.4, 14.5. **IR** (CHCl₃, cm⁻¹): 2135, 1720, 1376, 1214.

7b

Yield: 70%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.98 (sex, J=6 Hz, 1H), 4.34 (q, J=7 Hz, 2H), 1.69-1.78 (m, 2H), 1.30-1.50 (m, 7H), 1.45 (d, J=6 Hz), 1.34 (t, J=7 Hz), 0.96 (d, J=6 Hz, 3H), 0.94 (d, J=6 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 159.8, 83.0, 62.8, 46.0, 24.6, 22.9, 22.4, 21.5, 14.5. **IR** (CHCl₃, cm⁻¹): 2135, 1722, 1382, 1214.

7d

Yield: 75%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.66 (td, J=11, 4 Hz, 1H), 4.34 (qd, J=11, 1 Hz, 2H), 2.28-2.35 (m, 1H), 2.10 (septet of d, J=7, 2 Hz, 1H), 1.65-1.78 (m, 2H), 1.41-1.55 (m, 2H), 1.24-1.40 (m, 4H), 1.34 (t, J=7 Hz), 1.06 (qd, J=13, 3H, 1H), 0.80-1.00 (m, 10H), 0.95 (d, J=7Hz), 0.93 (d, J=7Hz), 0.81 (d, 7Hz). **¹³C NMR** (CDCl₃, 125 MHz): δ 159.8, 87.1, 62.8, 47.8, 41.9, 33.8, 32.00, 25.6, 23.2, 22.1, 21.1, 15.7, 14.5. **IR** (CHCl₃, cm⁻¹): 2135, 1724, 1382, 1214. M.p. (°C): 87-88.

Cyclization. General Procedure. To the suspension of Rh₂(OAc)₄ (2 mol %) in CH₂Cl₂ (4 ml/mmol) a solution of the corresponding diazo compound (1 equivalent, 0.1-1 mmol) in CH₂Cl₂ (2 ml/mmol) was added at rt over a period of 5 h using a syringe pump. Upon completion of the addition, the reaction mixture was stirred at rt for additional 10 h. The volatiles were removed under reduced pressure and the crude reaction mixture was purified on silica column (Ethyl Acetate-Hexanes, 0 to 40%). In many cases, additional purification on a silica gel column (Ether-Hexanes, 0 to 50%) was required to yield the pure product.

Cyclization products:

8

Yield: 55%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.29-4.37 (m, 2H), 3.55 (d, J=11 Hz, 1H), 3.17 (dt, J=14, 4 Hz, 1H), 2.91 (td, J=13, 4 Hz, 1H), 2.53-2.62 (m, 1H), 2.09-2.24 (m, 2H), 1.97 (dq, J=14, 4 Hz, 1H), 1.22-1.40 (m, 4H), 1.34 (t, J=7Hz), 1.08 (d, J=7 Hz, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 164.0(C), 72.7(CH), 62.7(CH₂), 52.0(CH₂), 34.5(CH), 32.0(CH₂), 22.8(CH₂), 19.9(CH₃), 14.3(CH₃). **HRMS** (ESI) calcd for C₉H₂₀NO₄S [M+NH₄] 238.1113, found 238.1118. m.p. (°C): 94-95.

9

Yield: 56%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.23 (qd, J=7, 1 Hz, 2H), 3.56 (d, J=3 Hz, 1H), 3.50 (td, J=14, 4 Hz, 1H), 2.91-2.98 (m, 1H), 2.17-2.30 (m, 1H), 2.00-2.11 (m, 2H), 1.26-1.41 (m, 7H), 1.36 (s), 1.32 (td, J=7, 1Hz), 1.07 (s, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 166.8(C), 74.3(CH), 62.3(CH₂), 48.7(CH₂), 36.9(C), 33.0(CH₂), 29.7(CH₃), 26.8(CH₃), 20.0(CH₂), 14.2(CH₃). **HRMS** (ESI) calcd for C₁₀H₂₂NO₄S [M+NH₄] 252.1270, found 252.1264.

11

Yield: 53%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.58-4.68 (m, 2H), 4.26-4.36 (m, 2H), 3.81 (s, 1H), 2.25-2.35 (m, 1H), 1.53 (ddd, J=14, 5, 3 Hz, 1H), 1.38 (s, 3H), 1.33 (dt, J=8, 1 Hz, 3H), 1.23 (s, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 164.6(C), 70.9(CH), 70.4 (CH₂), 62.6(CH₂), 36.5(C), 34.7 (CH₂), 27.7(CH₃), 26.7(CH₃), 14.2(CH₃). **HRMS** (ESI) calcd for C₉H₂₀NO₅S [M+NH₄] 254.1062, found 254.1056.

12

Yield: 55%. **¹H NMR** (CDCl₃, 500 MHz): δ 4.93 (dq, J=12, 6, 2 Hz, 1H), 4.23-4.35 (m, 2H), 3.74 (s, 1H), 2.31 (dd, J=14, 12Hz, 1H), 1.49 (d, J=6 Hz, 3H), 1.44 (s, 3H), 1.31-1.40 (m, 4H), 1.33 (t, J=7 Hz), 1.12 (s, 3H). **¹³C NMR** (CDCl₃, 125 MHz): δ 165.2(C), 79.4(CH), 69.2(CH), 62.5(CH₂), 40.3(CH₂), 36.0(C), 29.5(CH₃), 27.2(CH₃), 21.3(CH₃), 14.3(CH₃). **HRMS** (ESI) calcd for C₁₀H₂₂NO₅S [M+NH₄] 268.1219, found 268.1213.

13a (major)

Yield: 80% (**13a**+**13b**). ¹H NMR (CDCl₃, 500 MHz): δ 4.56 (td, J=11, 4 Hz, 1H), 4.27 (qd, J=7, 2 Hz, 2H), 3.77 (s, 1H), 2.23 (td, J=11, 3 Hz, 1H), 2.11-2.17 (m, 1H), 1.72-1.83 (m, 2H), 1.49-1.58 (m, 1H), 1.42 (dd, J=12, 11 Hz, 1H), 1.31 (s, 3H), 1.32 (t, J=7 Hz, 3H), 0.95-1.16 (m, 8H), 1.11 (s), 0.99 (d, J=6Hz). ¹³C NMR (CDCl₃, 125 MHz): δ 165.3(C), 84.3(CH), 71.6 (CH), 62.5(CH₂), 43.2(CH), 40.5(CH₂), 38.7(C), 34.1(CH₂), 31.8(CH), 27.1(CH₃), 23.9(CH₂), 22.7(CH₃), 22.0(CH₃), 14.2(CH₃). HRMS (ESI) calcd for C₁₄H₂₈NO₅S [M+NH₄] 322.1688, found 322.1682.

13b (minor)

¹H NMR (CDCl₃, 500 MHz): δ 4.57 (td, J=11, 4 Hz, 1H), 4.32 (q, J=7 Hz, 2H), 3.86 (s, 1H), 2.10-2.15 (m, 1H), 1.87 (dq, J=14, 3 Hz, 1H), 1.74-1.81 (m, 1H), 1.49-1.60 (m, 1H), 1.36-1.44 (m, 1H), 1.23-1.36 (m, 7H), 1.34 (t, J=7Hz), 1.29 (s), 1.09-1.17 (m, 4H), 1.14 (s), 0.91-1.03 (m, 4H), 0.98 (d, J=6Hz). ¹³C NMR (CDCl₃, 125 MHz): δ 163.9(C), 83.6(CH), 72.3 (CH), 62.6(CH₂), 50.5(CH), 40.6(CH₂), 40.4(C), 34.0(CH₂), 31.7(CH), 27.6(CH₃), 24.2(CH₂), 21.9(CH₃), 16.3(CH₃), 14.3(CH₃). HRMS (ESI) calcd for C₁₄H₂₈NO₅S [M+NH₄] 322.1688, found 322.1682.