-SUPPORTING INFORMATION FOR-

**Selective Synthesis of (2*Z*,4*E*)-Dienyl Esters by
Ene-diene Cross Metathesis**

Gustavo Moura-Letts and Dennis P. Curran

Department of Chemistry, University of Pittsburgh, Pittsburgh, PA, 15260

curran@pitt.edu

Table of Contents

General	2
Procedures for Diene Reagents and Cross Metathesis.....	2
Spectroscopic Data.....	3
Copies of NMR Spectra	12

General: DCM was distilled from CaH. Commercially available reagents were used and received from supplier. Reactions were monitored using thin layer chromatography (TLC) using Kieselgel 60 F₂₅₄ silica gel plates. Flash chromatography was performed over silica gel 60, 230-400 mesh, with the designated solvents. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DPX-300 or a Bruker DPX-500 spectrometer using residual solvent peaks as internal standard: CDCl₃ 7.23 ppm (¹H NMR) and 77.00 ppm (¹³C NMR). Chemical shifts are reported in ppm (δ) downfield from tetramethylsilane and proton-proton coupling constants (*J*) in Hz. Infrared spectra (IR) were recorded on a ATI Mattson genesis series FTIR spectrometer and are reported in reciprocal centimetres (cm⁻¹). Optical rotations were measured with Perkin-Elmer 241 digital polarimeter with a sodium lamp at ambient temperature and are reported as follows: [α]_D (c = g/100mL). Low and high resolution mass spectra were obtained on a VG 70-G or Micromass Autospec double focusing instrument using EI, ESI, or CI.

Methyl (2Z,4E)-hexa-2,4-dienoate 7: This known compound¹ was prepared by a Still-Gennari reaction.² To a solution of methyl 2-((bis(perfluoroethoxy))phosphoryl)acetate (200 mg, 0.63 mmol) in THF (8 mL) was added 18-crown-6 (412 mg, 1.56 mmol), and the solution was cooled to -78 °C. KHMDS (1.26 mL, 0.5 M) was added dropwise, and the solution was allowed to warm to -45 °C and then stirred for 1.5 h. Crotonaldehyde (36.5 mg, 0.52 mmol) was added at -78 °C and then the solution stirred for 3 h. The reaction was quenched with sat. aqueous NH₄Cl and extracted with diethyl ether. The organic phase was dried over MgSO₄, concentrated and purified by flash chromatography to give 62.3 mg (96%) of **7** as a colorless oil.

Methyl (2Z,4E)- 4-methylhexa-2,4-dienoate 11: This known compound (CAS# 6804-81-5) was prepared by the same procedure as for **7**.

Methyl (2Z,4E)-3-methylhexa-2,4-dienoate 12: This compound was prepared by the literature procedure.³

Typical procedure for small-scale ene-diene metathesis: To a solution of (3*R*,4*S*)-4,6-*bis*(*tert*-butyldimethylsilyloxy)-3-methylhex-1-ene **6** (100 mg, 0.279 mmol) in DCM (1.5 mL) was added (2*Z*,4*E*)-methyl hexa-2,4-dienoate **7** (35.2 mg, 0.279 mmol), and the solution was heated to 40 °C. Catalyst **2** (12.1 mg, 0.014 mmol) was dissolved in DCM (0.4 mL) and then added. The mixture was heated at 40 °C for 10 h. The solvent was removed by rotary evaporation, and the residue was purified by flash chromatography to give 71.6 mg (58%) **8** as a colorless oil.

Typical procedure for the large scale ene-diene metathesis: To a solution of (3*S*,4*S*)-4,6-*bis*(*tert*-butyldimethylsilyloxy)-3-methylhex-1-ene **15** (15 g, 42 mmol) in DCM (260 mL) was added (2*Z*,4*E*)-methyl hexa-2,4-dienoate **7** (5.3 mg, 42 mmol), and the solution was heated to 40 °C. Catalyst **5** (1.37 g, 1.26 mmol) was dissolved in DCM (29mL) and then added. The mixture was heated at 40 °C for 10 h. The solvent was removed by rotary evaporation, the residue dissolved in a minimal amount of DCM (20 mL) and then purified by fluorous solid phase extraction over a 50 g cartridge by eluting with 100 mL of 9:1 MeOH, followed by 100 mL of pure MeOH to give the organic fraction. Flash chromatography of the product from these

fractions with 20:1 hexanes/EtOAc furnished 10.9 g (59%) **16** as a colorless oil. The f-GH catalyst was recovered by flushing the cartridge with 200 mL of THF. The THF fraction was concentrated and the dark green residue was recrystallized from hexanes/DCM mixture to furnish 1005 mg (74%) of pure catalyst **5**.

Fluorous Grubbs Hoveyda catalyst 5: This was prepared by the same procedure as described for the lower homolog (ethylene spacer) catalyst.⁴ Green solid; mp: 140-141 °C; IR (thin film) 1625, 1490, 1315, 1242, 1215, 1115, 752, 720, 561 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 17.01 (s, 1H), 7.27 (dd, *J* = 8.4, 2 Hz, 1H), 7.04 (s, 4H), 6.70-6.67 (m, 2H), 4.83 (q, *J* = 6.1 Hz, 1H), 4.15 (s, 4H), 2.66 (t, *J* = 7.4 Hz, 1H), 2.49-2.34 (m, 18H), 2.15-1.95 (m, 2H), 1.87-1.79 (m, 1H), 1.22 (d, *J* = 6.1 Hz, 1H); ¹³C-NMR (75 MHz, CDCl₃) δ 211.4, 150.9, 145.3, 138.8, 134.1, 129.4, 129.2, 122.2, 112.9, 75.0, 33.6, 21.0. ¹⁹F-NMR (282 MHz, CDCl₃) δ -81.1 (3F), -114.4 (2F), -122.3 (6F), -123.9 (2F), -123.7 (2F), -126.5 (3F) ppm.

(3*R*,4*S*)-4,6-bis(*tert*-Butyldimethylsilyloxy)-3-methylhex-1-ene 6: This compound was prepared by literature procedure.⁵

Methyl-(2*Z*,4*E*,6*R*,7*S*)-7,9-bis(*tert*-butyldimethylsilyloxy)-6-methylnona-2,4-dienoate 8:

$[\alpha]_D -6.1$ (c 0.4, CHCl_3); IR (thin film) 2954, 2857, 1721, 1424, 1165, 1101, 830, 751 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.35 (dd, $J = 15.4, 11.3$ Hz, 1H), 6.58 (t, $J = 11.3$ Hz, 1H), 6.04 (dd, $J = 15.4, 8$ Hz, 3H), 5.59 (d, $J = 11.3$ Hz, 1H), 3.82-3.77 (m, 1H), 3.73 (s, 3H), 3.64 (t, $J = 6.6$ Hz, 1H), 2.51-2.46 (m, 1H), 1.66-1.54 (m, 2H), 1.07 (d, $J = 6.9$ Hz, 3H), 0.89 (s, 18H), 0.06 (s, 6H), 0.04 (s, 6H); ^{13}C -NMR (75 MHz, CDCl_3) δ 167.1, 147.6, 145.8, 127.1, 115.6, 72.6, 60.0, 51.2, 42.9, 37.4, 26.1, 18.5, 8.3, 15.7, -4.1, -5.0 ppm; MS (ESI) m/z 303(100), 147(80), 385(64), 145(32), 427(22), 442(5); HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{46}\text{O}_4\text{Si}_2$ 442.2897, found 442.2897.

COMPOUNDS FROM TABLE 1

Methyl-(2Z,4E,6R,7S)-7-(tert-Butyldimethylsilyloxy)-9-hydroxy-6-methylnona-2,4-dienoate:

$[\alpha]_D -14.9$ (c 0.25, CHCl_3); IR (thin film) 3421, 2953, 2857, 1719, 1412, 1190, 1167, 837 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.37 (dd, $J = 15.4, 11.3$ Hz, 1H), 6.56 (t, $J = 11.3$ Hz, 1H), 6.01 (dd, $J = 15.4, 8$ Hz, 3H), 5.61 (d, $J = 11.3$ Hz, 1H), 3.87-3.83 (m, 1H), 3.75-3.71 (m, 5H), 2.59-2.52 (m, 1H), 1.85 (bs, 1H), 1.70-1.66 (m, 2H), 1.08 (d, $J = 6.8$ Hz, 3H), 0.90 (s, 9H), 0.09 (s, 6H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.8, 147.2, 145.5, 126.8, 115.5, 73.2, 59.4, 51.0, 42.7, 36.0, 25.8, 18.0, 15.0, -4.4 ppm; MS (ESI) m/z 189(100), 147(60), 119(32), 271(15), 328(4); HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$ 328.2097, found 328.2097.

Methyl-(2Z,4E,6R,7S)- 9-(tert-butyldimethylsilyloxy)-7-(methoxymethoxy)-6-methylnona-2,4-dienoate:

$[\alpha]_D -7.8$ (c 0.5, CHCl_3); IR (thin film) 3053, 2955, 2930, 2857, 2254, 1713, 1637, 1463, 1439, 1265, 1198, 1096, 909, 736 cm^{-1} ; ^1H -NMR (600 MHz, CDCl_3) δ 7.36 (dd, $J = 15.4, 11.3$ Hz, 1H), 6.54 (t, $J = 11.3$ Hz, 1H), 6.02 (dd, $J = 15.4, 7.7$ Hz, 1H), 5.58 (d, $J = 11.3$ Hz, 1H), 4.64 (dd, $J = 21.3, 6.8$ Hz, 2H), 3.71 (s, 3H), 3.68-3.64 (m, 4H), 3.36 (s, 3H), 2.63-2.60 (m, 1H), 1.67-1.58 (m, 2H), 1.10 (d, $J = 6.8$ Hz, 3H), 0.86 (s, 9H), 0.02 (s, 6H); ^{13}C -NMR (125.8 MHz, CDCl_3) δ 166.8, 146.9, 145.4, 127.0, 115.6, 96.5, 78.6, 78.4, 59.6, 55.7, 51.1, 40.9, 34.9, 25.9, 18.2, 15.3, 5.4 ppm; MS (ESI) m/z 108(100), 119(95), 153(86), 184(80), 253(75), 357(24); HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{33}\text{O}_5\text{Si}$ 357.2097, found 357.2098.

Methyl-(2Z,4E,6R,7S)-7-(4-Methoxybenzyloxy)-9-(tert-butyldimethylsilyloxy)-6-methylnona-2,4-dienoate:

$[\alpha]_D -13.3$ (c 0.41, CHCl_3), IR (thin film) 3055, 2985, 2927, 2854, 1465, 1390, 1421, 1265 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.44 (dd, $J = 15.1, 11.4$ Hz, 1H), 7.07 (d, $J = 6.7$ Hz, 2H), 6.80 (d, $J = 6.7$ Hz, 2H), 6.53 (t, $J = 11.4$ Hz, 1H), 6.11 (dt, $J = 15.2, 7.2$ Hz, 1H), 5.59 (d, $J = 11.4$ Hz, 2H), 4.72 (d, $J = 6.7$ Hz, 1H), 3.77-3.75 (m, 1H), 3.72 (s, 3H), 3.65-3.58 (m, 2H), 3.41 (s, 3H), 2.53-2.49 (m, 1H), 1.60-1.50 (m, 2H), 1.03 (d, $J = 6.9$ Hz, 3H), 0.83 (s, 9H), 0.02 (s, 6H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.7, 158.0, 144.9, 143.7, 130.9, 129.5, 127.1, 115.8, 113.8, 73.2, 59.5, 55.0, 51.0, 42.7, 36.1, 25.9, 18.0, 15.0, -4.5 ppm. MS (ESI) m/z 108(100), 342(80), 153(86), 147(50), 448(16); HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{40}\text{O}_5\text{Si}$ 448.2613, found 448.2628.

Methyl-(2Z,4E)-6-(4-Methoxyphenyl)hexa-2,4-dienoate:

IR (thin film) 3052, 2986, 2959, 2872, 2253, 1720, 1610, 1511, 1439, 1265, 1175, 909, 734 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.44 (dd, $J = 15.1, 11.4$ Hz, 1H), 7.07 (d, $J = 6.7$ Hz, 2H), 6.80 (d, $J = 6.7$ Hz, 2H), 6.53 (t, $J = 11.4$ Hz, 1H), 6.11 (dt, $J = 15.2, 7.2$ Hz, 1H), 5.59 (d, $J = 11.4$ Hz, 2H), 3.74 (s, 3H), 3.69 (s, 3H), 3.43 (d, $J = 6.7$ Hz, 2H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.7, 158.0, 144.9, 143.7, 130.9, 129.4, 127.2, 115.8, 113.8, 55.0, 50.9, 38.4 ppm; MS (ESI) m/z 111(100), 173(30), 232(20); HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{O}_3$ 232.1099, found 232.1094.

Methyl-(2Z,4E)-6-Phenylhexa-2,4-dienoate:

IR (thin film) 3053, 2985, 2959, 2928, 2253, 1717, 1456, 1265, 909, 735 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.40 (dd, $J = 15.2, 11.4$ Hz, 1H), 7.23-7.10 (m, 5H), 6.49 (t, $J = 11.4$ Hz, 1H), 6.09 (dt, $J = 15.2, 7.2$ Hz, 2H), 5.54 (d, $J = 11.4$, 1H), 3.64 (s, 3H), 3.45 (d, $J = 7.2$ Hz, 2H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.8, 144.9, 143.2, 139.0, 128.5, 127.5, 126.3, 116.0, 51.0, 39.3. MS (ESI) m/z 111(100), 128(75), 171(50), 202(16); HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{14}\text{O}_2$ 202.0993, found 202.0984.

Methyl-(2Z,4E)-6-Chlorohexa-2,4-dienoate:

IR (thin film) 3053, 2986, 2953, 2249, 1712, 1638, 1602, 1439, 1265, 1177, 908 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.58 (ddt, $J = 15.2, 11.3, 1.2$ Hz, 1H), 6.57 (t, $J = 11.3$ Hz, 2H), 6.09 (ddt, $J = 15.2, 7.2, 0.7$ Hz, 1H), 4.10 (d, $J = 7.2$ Hz, 2H), 3.73 (s, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.3, 142.8, 137.2, 129.5, 119.0, 51.3, 44.1. MS (ESI) m/z 65(100), 111(100), 125(95), 59(86), 129(57), 160(24); HRMS (ESI) calcd for $\text{C}_7\text{H}_9\text{O}_2\text{Cl}$ 160.0291, found 232.0286.

Methyl-2Z,4E)-6-Acetoxyhexa-2,4-dienoate:

IR (thin film) 3053, 2986, 2955, 2872, 2254, 1739, 1648, 1607, 1438, 1377, 1265, 1232, 909, 735 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.54 (dd, $J = 15.5, 11.3$ Hz, 1H), 6.54 (t, $J = 11.3$ Hz, 1H), 6.04 (dt, $J = 15.5, 5.9$ Hz, 1H), 5.68 (d, $J = 11.3$ Hz, 1H), 4.67 (d, $J = 5.8$ Hz, 2H), 3.69 (s, 3H), 2.05 (s, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 170.5, 166.3, 143.2, 136.1, 128.7, 127.2, 118.2, 63.8, 51.1, 20.7 ppm; MS (ESI) m/z 111(100), 184(55), 124(48); HRMS (ESI) calcd for $\text{C}_9\text{H}_{12}\text{O}_4$ 184.0736, found 184.0741.

(E)-Methyl 1-Benzoyl-5-ethylidene-4-((1E,3Z)-5-methoxy-5-oxopenta-1,3-dienyl)-2-methyl-1,2,5,6-tetrahydropyridine-2-carboxylate:

IR (thin film) 3606, 2971, 2306, 1741, 1712, 1422, 1265, 1160, 1128, 948, 738 cm^{-1} ; ^1H -NMR (600 MHz, CDCl_3) δ 7.67 (dd, $J = 15.5, 11.4$ Hz, 1H), 7.43-7.41 (m, 5H), 6.60 (t, $J = 11.4$ Hz, 1H), 6.46 (d, $J = 15.5$ Hz, 1H), 5.83 (q, $J = 7.0$ Hz, 1H), 5.74 (s, 1H), 5.69 (d, $J = 11.4$ Hz, 1H), 3.70 (s, 3H), 3.74 (s, 3H), 3.66 (d, $J = 14.6$ Hz, 1H), 3.14 (d, $J = 14.6$ Hz, 1H), 1.72 (s, 3H), 1.61 (d, $J = 7.0$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 171.8, 171.0, 166.6, 144.4, 138.0, 135.8, 135.7, 130.3, 129.4, 128.5, 127.9, 127.6, 127.4, 123.6, 117.9, 63.0, 52.9, 51.3, 43.6, 20.7, 13.5 ppm. MS (ESI) m/z 105(100), 77(86), 336(82), 395(20); HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{25}\text{NO}_5$ 395.1733, found 395.1717.

COMPOUNDS FROM TABLE 2

Methyl-(2Z,4E)-6-(4-Methoxyphenyl)-4-methylhexa-2,4-dienoate:

IR (thin film) 3052, 2986, 2959, 2872, 2254, 1720, 1610, 1511, 1439, 1265, 1175, 909, 734 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.27 (d, $J = 8.6$ Hz, 1H), 7.01 (d, $J = 8.6$ Hz, 1H), 6.60 (d, $J = 12.5$ Hz, 1H), 6.02 (t, $J = 7.4$ Hz, 1H), 5.82 (d, $J = 12.5$ Hz, 1H), 4.67 (d, $J = 5.8$ Hz, 2H), 3.95 (s, 3H), 3.87 (s, 3H), 3.60 (d, $J = 7.4$ Hz, 2H), 2.12 (s, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 166.9, 157.8, 145.4, 136.4, 132.8, 131.8, 129.1, 116.6, 113.7, 54.9, 51.0, 33.5, 14.9 ppm; MS (ESI) m/z 125(100), 246(50), 121(35); HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{O}_3$ 246.1256, found 246.1259.

Methyl-(2Z,4E)-4-Methyl-6-phenylhexa-2,4-dienoate:

IR (thin film) 3053, 2986, 2959, 2929, 2872, 2253, 1710, 1626, 1453, 1265, 903, 735 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.38-7.21 (m, 5H), 6.47 (d, $J = 12.5$ Hz, 1H), 5.90 (t, $J = 7.4$ Hz, 2H), 5.70 (d, $J = 12.5$ Hz, 1H), 3.72 (s, 3H), 3.52 (d, $J = 7.4$ Hz, 2H); ^{13}C -NMR (75 MHz, CDCl_3) δ 167.1, 145.5, 140.0, 135.9, 133.3, 128.5, 126.1, 116.9, 51.3, 34.6, 15.1 ppm; MS (ESI) m/z 125(100), 91(35), 216(25); HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{O}_2$ 216.1150, found 216.1145.

Ethyl-(2Z,4E)-6-(4-Methoxyphenyl)-3-methylhexa-2,4-dienoate:

IR (thin film) 3051, 2991, 2879, 2872, 2249, 1720, 1610, 1512, 1439, 1265, 1175, 912, 734 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.06 (d, $J = 8.7$ Hz, 2H), 6.82 (d, $J = 8.7$ Hz, 2H), 6.22 (dt, $J = 15.6, 6.5$ Hz, 1H), 6.08 (d, $J = 15.4$ Hz, 1H), 5.69 (s, 1H), 4.12 (q, $J = 7.1$ Hz, 2H), 3.78 (s, 3H), 3.40 (d, $J = 6.5$ Hz, 2H), 2.2 (s, 3H), 1.25 (t, $J = 7.1$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 167.2, 158.2, 152.2, 135.8, 134.4, 131.3, 129.6, 118.4, 114.0, 59.6, 55.2, 38.4, 29.7 ppm; MS (ESI) m/z 125(100), 246(50), 121(35); HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{20}\text{O}_3$ 260.1256, found 260.1257.

Ethyl-(2Z,4E)-3-Methyl-6-phenylhexa-2,4-dienoate:

IR (thin film) 3055, 2986, 2959, 2929, 2872, 2253, 1710, 1626, 1453, 1265, 903, 735 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.38-7.21 (m, 5H), 6.24 (dt, $J = 15.6, 6.5$ Hz, 1H), 6.12 (d, $J = 15.4$ Hz, 1H), 5.69 (s, 1H), 4.27 (q, $J = 7.1$ Hz, 2H), 3.54 (d, $J = 6.5$ Hz, 2H), 2.02 (s, 3H), 1.25 (t, $J = 7.1$ Hz, 3H); ^{13}C -NMR (75 MHz, CDCl_3) δ 167.1, 152.2, 135.8, 135.8, 134.3, 129.6, 121.7, 114.4, 55.3, 38.4, 29.7, 14.3 ppm; MS (ESI) m/z 125(100), 91(35), 216(25); HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{O}_2$ 230.1150, found 230.1143.

(3S,4S)-4,6-bis(tert-Butyldimethylsilyloxy)-3-methylhex-1-ene 15: This compound was prepared by literature procedure.⁶

Methyl-(2Z,4E,6R,7S)-7,9-bis(tert-Butyldimethylsilyloxy)-6-methylnona-2,4-dienoate 16:

$[\alpha]_D +10.1$ (c 0.87, CHCl_3); IR (thin film) 2954, 2857, 1721, 1424, 1165, 1101, 830, 751 cm^{-1} ; ^1H -NMR (600 MHz, CDCl_3) δ 7.31 (dd, $J = 15.4, 11.3$ Hz, 1H), 6.54 (t, $J = 11.3$ Hz, 1H), 6.13 (dd, $J = 15.4, 8$ Hz, 3H), 5.55 (d, $J = 11.3$ Hz, 1H), 3.76-3.72 (m, 1H), 3.69 (s, 3H), 3.64-3.58 (m, 2H), 2.49-2.44 (m, 1H), 1.62-1.50 (m, 2H), 1.02 (d, $J = 6.9$ Hz, 3H), 0.84 (s, 18H), 0.02 (s, 4H), 0.01 (s, 8H); ^{13}C -NMR (125.8 MHz, CDCl_3) δ 166.9, 148.3, 145.3, 126.3, 115.3, 72.7, 59.8, 50.9, 42.5, 36.8, 25.9, 18.2, 14.5, -4.4, -5.3 ppm; MS (ESI) m/z 303(100), 147(80), 385(64), 145(32), 427(22), 442(5); HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{46}\text{O}_4\text{Si}_2$ 442.2897, found 442.2897.

REFERENCES:

- 1) Bougioukou, D. J.; Simonou, I. *Tetrahedron. Lett.* **2002**, *43*, 339-342.
- 2) Still, W. C.; Gennari, C., *Tetrahedron Lett.* **1983**, *24*, 4405-4408.
- 3) Posner, G. H.; Crouch, D.; Kinter, C. H.; Carry, J. C. *J. Org. Chem.* **1991**, *56*, 6981-6987.
- 4) Matsugi, M.; Curran, D. P. *J. Org. Chem.* **2005**, *70*, 1636-1642.
- 5) White, J. D.; Hong, J.; Robarge, L. A. *J. Org. Chem.* **1999**, *64*, 6206-6216.
- 6) Wang, P.; Kim, Y. J.; Navarro-Villalobos, M.; Rohde, B. D.; Gin, D. Y. *J. Am. Chem. Soc.* **2005**, *127*, 3256-3257

gm1453211-recrystallized

137.04
 136.17
 135.52
 132.63
 131.54
 128.68
 126.51
 126.48
 126.42
 123.67
 123.64
 122.31
 122.28
 116.39
 114.49
 114.43
 114.37
 111.08
 109.42
 109.28
 107.41
 106.76
 104.11
 103.72
 103.52
 99.79
 99.56
 98.78
 98.75
 96.97
 96.38
 94.37
 92.94
 91.29
 88.44
 88.14
 85.59
 83.55
 81.16
 81.13
 81.09
 79.08
 76.47
 76.11
 73.45
 71.32
 66.74
 64.77
 64.37
 62.02
 61.99
 ppm

41 S

gm1338p1

1.25078
1.22700
1.20320

2.00834

4.07637
3.74005
3.70130
3.68845
3.45303
3.42947

7.43910
7.08826
7.05935
6.82747
6.79853
6.57063
6.53275
6.13976
5.60647
5.56869

10

ppm

S
17

2.7881
2.7245
2.3420

1.2803
1.9247
2.0000
1.2552
1.2429
1.0972

Integral

ppm

Current Data Parameters	
NAME	gm139011
EXPNO	1
PROCNO	1

F2 - Acquisition Parameters			
Date_	20051207		
Time	19.46		
INSTRUM	spect		
PROBHD	5 mm TBI 1H/		
PULPROG	zg		
TD	65536		
SOLVENT	CD2Cl2		
NS	8		
DS	0		
SWH	8992.806 Hz		
FIDRES	0.137219 Hz		
Acq	3.6438515 sec		
RG	25.4		
QW	55.600 usec		
DE	6.00 usec		
TE	290.0 K		
D1	1.00000000 sec		

```
===== CHANNEL f1 =====
NUC1      1H
P1        9.00 us
PL1       0.00 dB
SF01     600.8336050 MHz
```

F2 - Processing parameters	
SI	65536
SF	600.8300409 MHz
WDW	EM
SSB	0
LB	0 10 Hz
GB	0
GC	1.00

1D NMR plot parameters	
CX	20.00 cm
F1P	13.416 ppr
F1	8060.47 Hz
F2P	-1.552 ppr
F2	-932.34 Hz
PPMCM	0.74837 ppr
HZCM

Table 3 entry 2

Current Data Parameters
 NAME gm139011
 EXPNO 2
 PROCNO 1

gm139011

F2 - Acquisition Parameters

Date_ 500000
 Time 15.04
 INSTRUM spect
 PROBHD 5 mm TXI 13C
 PULPROG c13wznoe
 TD 32768
 SOLVENT CDCl3
 NS 17766
 DS 0
 SWH 32679.738 Hz
 FIDRES 0.997306 Hz
 AQ 0.5014004 sec
 RG 32768
 DW 15.300 usec
 DE 6.00 usec
 TE 290.0 K
 D3 0.00100000 sec
 PL12 6.00 dB
 D1 8.00000000 sec
 CPDPRG2 waltz16
 PCPD2 100.00 usec
 SF02 500.1330008 MHz
 NUC2 1H
 PL2 120.00 dB
 P1 21.60 usec
 DE 6.00 usec
 SF01 125.7715724 MHz
 NUC1 13C
 PL1 0.00 dB

F2 - Processing parameters

SI 8192
 SF 125.7577961 MHz
 WDW EM
 SSB 0
 LB 4.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters

CX 20.00 cm
 F1P 239.478 ppm
 F1 30116.16 Hz
 F2P -20.385 ppm
 F2 -2563.57 Hz
 PPMCM 12.99313 ppm/cm
 HZCM 1

166.87
 157.25
 155.20
 146.89
 145.36
 143.23
 139.08
 127.01
 126.77
 115.62
 96.55
 78.60
 78.40
 77.25
 77.00
 76.74
 59.65
 55.72
 51.05
 40.88
 40.62
 34.94
 34.52
 25.90
 18.22
 15.22
 -5.36

Table 3 entry 2

S
20

-4.531
 -4.476
 15.051
 18.007
 25.856
 36.074
 42.695
 51.001
 55.011
 59.457
 73.232
 76.574
 76.998
 77.423

113.735
 113.766
 113.802
 113.870
 115.745
 127.212
 129.330
 129.361
 129.397
 129.464
 130.945
 143.670
 144.899

158.025
 166.662

Table 3 entry 3

Table 3 entry 4

Table 3 entry 4

Table 3 entry 6

3.68331
3.64265
3.46905
3.44534

7.44572
7.22771
7.20287
7.12990
7.12488
6.52698
6.48855
6.11469
6.06397
5.56685
5.52904

gm1328p1

S
25

ppm

2.4709
1.7548

0.7586

0.8607

0.9155

3.9602

1.0000

Integral

ppm

Table 3 entry 6

gm1343p1

7.54405
7.53998
7.53018
6.57820
6.54041
6.50257
6.06684
6.01529
5.70142
5.66360
4.67534
4.67136
4.65588
3.69311
2.11751
2.06227
2.05100
2.03824

Table 3 entry 7

S
27

Table 3 entry 7

gm1329111

7.63226
7.62842
7.59462
7.59081
7.58627
7.58170
7.57786
7.54406
7.54022
6.50514
6.56747
6.52973
6.14270
6.11885
6.11658
6.09487
6.09235
6.06828
6.06600
6.04450
5.75599
5.71825
5.29210
4.17786
4.17442
4.15414
4.15061
4.11835
4.09450
3.74407
3.72686
3.47979

Table 3 entry 8

ppm

S
29

1.0000
1.1779
1.0896
1.0486
2.2566
3.1602

Integral

ppm

Table 3 entry 8

gm1384211

1.73682
1.72247
1.62990
1.60622
1.21915

4.45674
4.42913
4.40802
3.74350
3.70391
3.68826
3.62281
3.17247
3.10207

7.67822
7.42972
7.41891
7.41160
7.22978
6.60446
6.48052
6.42892
5.82823
5.73836
5.71203
5.67441

Table 3 entry 9

S
13

2.2447
2.3600

2.1748
2.0750
1.2371
0.6820

1.0282
1.0570
1.0102
0.7938
0.9590

0.9820
4.8856

Integral

ppm

171.755
171.014
166.650
165.036
144.348
137.996
135.795
135.694
130.302
129.375
128.519
127.864
127.588
127.441
123.581
120.581
120.467
117.894
77.825
77.763
77.703
77.644
77.583
77.426
77.205
77.002
76.579
76.301
76.150
63.050
62.894
62.822
62.389
62.317
53.035
52.855
52.698
51.264
43.583
34.677
32.759
29.683
25.098
20.706
13.531

Table 3 entry 9

32

ppm

gm133941

3.98930
3.95183
3.87131
3.61944
3.59466
2.12499

7.28969
7.26106
7.02698
6.99822
5.85073

Table 4 entry 1

ppm

14.885
14.842

33.532

54.941
51.031

77.424
76.999
76.573

113.737
116.543
116.599
116.655
129.122
131.781
132.814
136.441
145.379

157.844

166.903

Table 4 entry 1

gm135422

0.14319	
0.15397	
0.15648	
0.93791	
0.96217	
1.29156	
1.29882	
1.31340	
1.96778	
1.99228	
2.07393	
2.72057	
2.74783	
3.39928	
3.51504	
3.53977	
3.72281	
3.73497	
4.25999	
4.26647	

5.30067	
5.57460	
5.71633	
5.87639	
5.90125	
5.44576	
6.48763	
7.20827	
7.23478	
7.30405	
7.32750	
7.56920

Table 4 entry 2

S
35

ppm

Table 4 entry 2

Table 4 entry 3

gm134011221

ppm

Integral

ppm

gm1340p111

Current Data Parameters
NAME gm1340p111
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20060504
Time 9.30
INSTRUM spect
PROBHD 5 mm TBI 1H/
PULPROG c13winoe
TD 65536
SOLVENT CDCl3
NS 28479
DS 0
SWH 37878.789 Hz
FIDRES 0.577984 Hz
AQ 0.8651252 sec
RG 32768
DW 13.200 usec
DE 6.00 usec
TE 290.0 K
D1 8.00000000 sec
D3 0.00100000 sec

===== CHANNEL f1 =====

NUC1 13C
P1 13.50 usec
PL1 0.00 dB
SF01 151.0953827 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 0.00 dB
PL12 12.00 dB
SF02 600.8336050 MHz

F2 - Processing parameters

SI 65536
SF 151.0788304 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 20.00 cm
F1P 234.922 ppm
F1 35491.70 Hz
F2P -15.800 ppm
F2 -2387.09 Hz
PPMCM 12.53610 ppm/cm
HZCM 1897.00000 Hz/cm

Table 4 entry 3

gm135422

7.56920	7.32750
7.30405	7.23478
7.20827	6.24783
6.22590	6.19617
6.17430	6.15312
6.11556	6.06360
5.68517	4.26647
4.25999	3.73497
3.72281	3.53977
3.51504	3.39928
2.74783	2.72057
2.07393	1.99228
1.96778	1.27046
1.24673	1.22283
0.96217	0.93791
0.15848	0.15397
0.14319

Table 4 entry 4

69 S

Current Data Parameters
 NAME gml340p111
 EXPHO 2
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20060504
 Time 9.30
 INSTRUM spect
 PROBHD 5 mm TBI 1H/
 PULPROG zgpg30proe
 TD 65536
 SOLVENT CDCl3
 NS 28479
 DS 0
 SWH 37878.789 Hz
 FIDRES 0.577984 Hz
 AQ 0.8651252 sec
 RG 32768
 DW 13.200 usec
 DE 6.00 usec
 TE 290.0 K
 D1 8.0000000 sec
 D3 0.00100000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 13.50 usec
 PL1 0.00 dB
 SF01 151.0953827 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 12.00 dB
 SF02 600.8336050 MHz

F2 - Processing parameters
 SI 65536
 SF 151.0788304 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters

CX 20.00 cm
 F1P 234.922 ppm
 F1 35491.70 Hz
 F2P -15.800 ppm
 F2 -2387.09 Hz
 PFCH 12.52
 HZCH 1893.9

Table 4 entry 4

S
40