Sequential Cycloaddition Approach to the Tricyclic Core of Vibsanin E. Total Synthesis of 5-*epi*-10-*epi*-Vibsanin E

Huw M. L. Davies*, Øystein Loe and Douglas G. Stafford

Department of Chemistry, University at Buffalo, The state University of New York, Buffalo, NY 14260-300

Supporting Information

Section A: General Information	S-2
Section B: Experimental Procedures	S-3
Section C: Relative Stereochemistry of 5- <i>epi</i> -10- <i>epi</i> -Vibsanin E	S-16
Section D: ¹ H NMR Spectra	S-18

Section A: General Information.

Commercial reagents from Aldrich Chemical Company and Acros Organics were purchased at the highest commercial purity and used without further purification unless otherwise noted. All moisture sensitive reactions were performed using glassware flame dried under vacuum and purged with argon prior to use. Tetrahydrofuran, hexanes and dichloromethane were used directly from the solvent purification system, where solvent was passed through two columns of activated alumina, purchased from MBRAUN. All other solvents were of reagent grade. Solvents used in rhodium carbenoid transformations were degassed with argon for 10-15 min. prior to use. Organic reaction mixtures were concentrated on a Buchi rotary evaporator. Analytical thin layer chromatography was performed on 250 μ m Whatman silica gel (Aluminum backing, UV 254 nm) plates using UV light or 50% sulfuric acid in methanol as visualizing agents. Column chromatography was carried out using E. Merck silica 60 (230 - 400 mesh) or ICN 60 (32 – 64 mesh) following the method of Still.¹

¹H NMR spectra were recorded on a Varian Nuclear Magnetic Resonance spectrometer at 400 or 500 MHz and ¹³C NMR spectra were recorded at 75 or 125 MHz, with the sample solvent being CDCl₃ or C₆D₆. The CDCl₃ samples were calibrated at the solvent peak at 7.27 ppm for the ¹H NMR spectra, for the ¹³C NMR spectra the middle solvent peak was set at 77 ppm. The C₆D₆ samples were calibrated at 0 ppm with TMS as internal standard or at the solvent peak at 7.157 ppm for the ¹H NMR spectra. The ¹³C NMR spectra were set likewise at 0 ppm with TMS or at 128.026 ppm for the middle peak. The following abbreviations are used to explain the multiplicities: s, singlet; d, doublet; t, triplet; q,

¹ Still, W. C.; Kahn, M.; Mitra, A. J. *J. Org. Chem.* **1978**, *43*, 2923

quartet; dd, doublet of doublets; dt, doublet of triplets; ddd, doublet of doublet of doublets. Infrared spectra were recorded on a Nicolet Impact 420 FT-IR spectrometer. High resolution mass spectra were obtained from the Mass Spectroscopy Facility at the University at Buffalo, The State University of New York.

Section B: Experimental Procedures:

Synthesis of 4-Methyl-4-(4-methyl-pent-3-enyl)-cyclohepta-1,5-dienecarboxylic acid methyl ester (**5**)

The catalyst $\text{Rh}_2(\text{OOct})_2$ (741 mg, 0.952 mmol, 0.5 mol%) was dissolved in the triene **3** (157 g, 1.05 mol) (due to much better solubility than in hexane). This mixture was diluted with hexane (1.90 L) to furnish a homogenous dark green solution that was degassed by bubbling argon through (20 min.). The diazo compound **4** (24.0 g, 0.190 mol) was dissolved in cold ($-20\text{ }^{\circ}\text{C}$) hexane (0.30 L) and added dropwise to the alkene/ rhodium solution from an acetone/ dry-ice cooled addition funnel. After the addition was complete (1 h) the reaction was left stirring overnight at room temperature. The organics were concentrated under reduced pressure and the catalyst was removed by filtration through a short silica plug (10% diethyl ether in hexane). The residue was dissolved in toluene (150 mL) and heated at reflux (5 h) in order to force the Cope rearrangement to completion.

The toluene was removed under reduced pressure and the product was purified by silica gel chromatography (5% diethyl ether in hexane). This furnished the cycloheptadiene **5** in 62% yield (29.2 g, 0.118 mol).

Alternatively,

To a stirred solution of the triene **3** (906 mg, 6.0 mmol), and $\text{Rh}_2(R\text{-DOSP})_4$ (22 mg, 0.012 mmol) in hexane (30 ml) at 23 °C under argon, vinyl diazoacetate **4** (148 mg, 1.17 mmol) in hexane (25 ml) at 0 °C was added via cannula over 20 min. After 40 h. the solvent was removed under reduced pressure. Purification by silica gel chromatography (pentane/diethyl ether 19:1) gave **5** (202 mg, 0.811 mmol) in 69 % yield and 64 % ee (determined by GC: α -TA; 100 °C, 21.4 psi, 2.1 mL/min; T_R = 185.4 min (minor), 189.7 min (major). $[\alpha]_D^{25} = -11^\circ$ (c 2.16, CHCl_3) ^1H NMR (CDCl_3 , 400MHz, δ) 7.08 (dd, 1H, J = 7.4, 7.4 Hz), 5.50 (dt, 1H, J = 11.6, 5.2 Hz), 5.29 (d, 1H, J = 11.6 Hz), 5.06 (t, 1H, J = 6.6 Hz) 3.71 (s, 3H), 3.11 (d, 2H, J = 5.2 Hz), 2.50 (dd, 1H, J = 13.2, 7.4 Hz), 2.24 (dd, 1H, J = 13.2, 7.4 Hz), 1.95 (ddd, 2H, J = 11.2, 6.6, 6.4Hz), 1.65 (s, 3H), 1.58 (s, 3H), 1.30 (dd, 2H, J = 11.2, 6.4 Hz), 0.98 (s, 3H). ^{13}C NMR (CDCl_3 , 75MHz, DEPT, δ) 167.5 (4°), 141.7 (3°), 139.6 (3°), 135.2 (4°), 131.3 (3°), 124.6 (3°), 123.2 (3°), 51.7 (1°), 43.2 (2°), 37.7 (4°), 37.2 (2°), 27.9 (1°), 25.7 (2°), 25.6 (1°), 23.1 (2°), 17.5 (1°). IR (neat) 3009, 2956, 2924, 2860, 1715, 1651 cm^{-1} . MS m/z (relative intensity) 248 (3), 233 (6), 205 (12), 173 (16), 145 (25), 133 (27), 119 (46), 105 (100), 91 (78). Anal. Calcd. for $\text{C}_{16}\text{H}_{24}\text{O}_2$: C, 77.38; H, 9.74. Found : C, 77.17; H, 9.66

Synthesis of 4-Methyl-4-(4-methyl-pent-3-enyl)-cyclohepta-1,5-dienecarbaldehyde

(6)

The ester **5** (19.0 g, 76.6 mmol) was dissolved in THF (500 mL) and DIBAL-H (200 mL, 1.0 M, 0.20 mol) was added dropwise from an addition funnel (30 min.) at 0 °C. The reaction was stirred over night and allowed to warm to room temperature. The crude mixture was quenched with a water/ ice solution (500 mL) and diluted with HCl (100 mL, 10%). The aqueous phase was extracted with diethyl ether (3 X 150 mL). The organics were back washed with water (200 mL) and brine (200 mL) before dried over magnesium sulfate and the solvent was evaporated. This gave the crude alcohol (17 g) that was used in the Swern oxidation without further purification. ¹H NMR (CDCl₃, 500 MHz, δ) 5.77 (t, 1H, *J* = 7.0 Hz), 5.51 (dt, 1H, *J* = 11.5, 6.0 Hz), 5.30 (d, 1H, *J* = 11.5 Hz), 5.10 (t, 1H, *J* = 6.5 Hz), 4.03 (s, 2H), 2.85 (dd, 1H, *J* = 5.0, 18.5 Hz), 2.75 (dd, 1H, *J* = 5.0, 18.5 Hz), 2.38 (dd, 1H, *J* = 6.5, 16.5 Hz), 2.11 (dd, 1H, *J* = 6.5, 16.5 Hz), 2.00-1.95 (m, 2H), 1.69 (s, 3H), 1.61 (s, 3H), 1.49 (s, broad, 1H), 1.32 (m, 2H), 0.99 (s, 3H).

Oxalyl chloride (13 mL, 18.9 g, 0.149 mol) was dissolved in dichloromethane (500 mL) and cooled to -78 °C. Then DMSO (17 mL, 18.7 g, 0.24 mol) was added and the solution was stirred for 15 min at -78 °C. The alcohol (17 g, 77 mmol) dissolved in dichloromethane (200 mL) was added dropwise at -78 °C and then stirred for 30 min. Triethylamine (120 mL) was added over 10 min and then the reaction was stirred for 5

min at $-78\text{ }^{\circ}\text{C}$. The mixture was allowed to warm to room temperature and the reaction mixture was extracted with water (100 mL) and the organics were dried over sodium sulfate. After removal of the solvent and purification by silica gel chromatography (5% ethyl ether in hexane) the aldehyde **6** (15.1 g, 69.3 mmol) was obtained in 90% yield. ^1H NMR (CDCl_3 , 500 MHz, δ) 9.39 (s, 1H), 6.84 (t, 1H, $J = 6.7$ Hz), 5.52 (dt, 1H, $J = 11.6$, 17.1 Hz), 5.38 (d, 1H, $J = 11.6$ Hz), 5.09 (t, 1H, $J = 7.0$ Hz), 3.08 (dd, 1H, $J = 5.2$, 19.5 Hz), 3.03 (dd, 1H, $J = 5.2$, 19.5 Hz), 2.68 (dd, 1H, $J = 6.7$, 13.7 Hz), 2.45 (dd, 1H, $J = 6.7$, 13.7 Hz), 2.02 - 1.97 (m, 2H), 1.69 (s, 3H), 1.61 (s, 3H), 1.40 - 1.33 (m, 2H), 1.05 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz, δ) 193.0, 153.3, 145.4, 139.4, 131.5, 124.4, 122.8, 43.2, 38.2 (2C), 28.0, 25.6, 23.1, 23.0, 17.5. IR (neat, cm^{-1}) 3350 (overtone, Fermi), 3004, 2962, 2919, 2860, 2722, 1683, 1651, 1454, 1380, 1162, 725. MS (EI) m/z (relative intensity) 218 (45), 175 (54), 107 (100), 105 (69), 91 (75), 79 (55), HRMS calcd. for $\text{C}_{15}\text{H}_{22}\text{O}$ 218.1665, found 218.1676.

Synthesis of 1,5,5-Trimethyl-6-oxa-tricyclo[6.3.2.0^{4,13}]trideca-7,10-diene (**7**)

The aldehyde **6** (15.1 g, 69.3 mmol) was dissolved in dichloromethane (500 mL) and cooled to $-78\text{ }^{\circ}\text{C}$ and $\text{BF}_3 \cdot \text{OEt}_2$ (30 mL, 0.24 mol) was added. The reaction was stirred at $-78\text{ }^{\circ}\text{C}$ for 20 min. and quenched by addition of water (500 mL). The reaction mixture was extracted with dichloromethane (3 X 250 mL) and the organics were dried over

magnesium sulfate. The crude product was purified on a silica gel column (4% diethyl ether in hexane) to afford the tricyclic system **7** in 84% yield (13.0 g, 59.6 mmol). ^1H NMR (CDCl_3 , 500 MHz, δ) 6.03 (d, 1H, $J = 2.1$ Hz), 5.74 (ddd, 1H, $J = 1.8, 11.1, 11.1$ Hz), 4.99 (d, 1H, $J = 11.1$ Hz), 2.97 (d, 1H, $J = 15.6$ Hz), 2.79 (s, broad, 1H), 2.32 (m, 2H), 1.55 - 1.47 (m, 2H), 1.39 (dd, 1H, $J = 4.0, 12.8$ Hz), 1.31 - 1.21 (m, 2H), 1.25 (s, 3H), 1.24 (s, 3H), 1.14 (dt, 1H, $J = 4.0, 12.8$ Hz), 0.96 (s, 3H) ^{13}C NMR (CDCl_3 , 125 MHz, δ (multiplicity)) 137.6 (d), 136.7 (d), 129.1 (d), 112.0 (s), 76.1 (s), 41.0 (d), 38.1 (2C) (t,t), 37.1 (s), 33.1 (q), 31.2 (d), 28.3 (t), 27.7 (q), 26.0 (q), 19.8 (t). IR (neat, cm^{-1}) 3053, 2928, 2866, 1660, 1453, 1370, 1230, 1147, 940, 805. MS (EI) m/z (relative intensity) 218 (100), 136 (67), 107 (59), 105 (65), 91 (72), 69 (53). HRMS calcd. for $\text{C}_{15}\text{H}_{22}\text{O}$ 218.1665, found 218.1663.

Synthesis of 1,5,5-Trimethyl-6-oxa-tricyclo[6.3.2.0^{4,13}]tridec-10-ene (**8**)

To a stirred solution of the diene **7** (13.0 g, 59.6 mmol) in acetic acid (500 mL) at room temperature was added NaCNBH_3 (17 g, 270 mmols) and the solution was stirred for 20 min. The same quantity of NaCNBH_3 (17 g, 270 mmol) was added again and stirred for 20 minutes. At this point the reaction was quenched with water (1 L) and extracted with hexane (2 x 250 mL). The organics were dried over magnesium sulfate and evaporated to yield the crude product. The mixture was purified on a silica column with 5% diethyl ether in hexane to yield the desired compound **8** in 80% yield (11.0 g, 50.0 mmol). ^1H

NMR (CDCl₃, 400 MHz, δ) 5.54 (ddd, 1H, J = 2.4, 7.3, 12.2 Hz), 5.42 (ddd, 1H, J = 1.5, 2.7, 12.2 Hz), 3.83 (dd, 1H, J = 2.6, 10.6 Hz), 3.35 (d, 1H, J = 10.6 Hz), 2.85 (ddd, 1H, J = 3.3, 7.0, 14.4 Hz), 2.39 (d, 1H, broad, J = 4.4 Hz), 2.23 (ddd, 1H, J = 7.0, 11.7, 14.4), 1.77 - 1.68 (m, 3H), 1.57 - 1.47 (m, 3H), 1.31 (s, 3H), 1.30 - 1.21 (m, 2H), 1.20 (s, 3H), 1.02 (s, 3H). ¹³C NMR (CDCl₃, 125 MHz, δ) 140.1, 127.7, 74.1, 67.6, 43.6, 43.3, 41.3, 35.9, 35.4, 34.2, 33.4, 33.1, 27.7, 24.6, 23.2. IR (neat, cm⁻¹) 2946, 2892, 1662, 1454, 1364, 1087. HRMS calcd. for C₁₅H₂₄O 220.1822, found 220.1824.

Synthesis of 1,5,5-Trimethyl-6-oxa-tricyclo[6.3.2.0^{4,13}]tridec-10-en-9-one (9)

To a stirred solution of the alkene 8 (11.0 g, 50.0 mmol) in 1,4-dioxane (450 mL) was added SeO₂ (6.66 g, 60 mmol). The reaction was stirred at reflux (2 h) and then allowed to cool to room temperature. The reaction was added water (1 L) and extracted with ether (2 X 200 mL). The organics were dried over MgSO₄ and the solvent was evaporated. The crude product was dissolved in dichloromethane (500 mL) and 4 Å molecular sieves were added with pyridinium chlorochromate (25 g, 0.12 mol). The reaction was stirred at room temperature (2 h) before the mixture was concentrated down and diluted with diethyl ether (1L), The mixture was extracted with water (2 X 200 mL) and brine (2 X 200 mL). The organics were dried over magnesium sulfate. The crude product was purified on a silica column with 10% acetone in hexane to yield the ketone 9 in 72% (8.5 g, 36 mmol) yield. ¹H NMR (CDCl₃, 500 MHz, δ) 6.07 (s, 2H), 4.57 (d, 1H, J = 11.7 Hz), 3.60, (dd,

1H, $J = 3.1, 11.7$ Hz), 2.62 (d, 1H, $J = 4.4$ Hz), 2.16 (t, 1H, $J = 4.4$ Hz), 2.08 (d, 1H, $J = 13.7$ Hz), 1.73 (dd, 1H, $J = 5.8, 13.7$ Hz), 1.56 (m, 2H), 1.30 (s, 3H), 1.27 (m, 2H), 1.17 (s, 3H), 1.15 (m, 1H), 1.12 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz, δ) 203.0, 151.9, 131.9, 73.6, 61.4, 50.4, 43.8, 42.9, 38.0, 36.9, 32.2, 31.9, 28.0, 24.1, 19.7. IR (neat, cm^{-1}) 2895, 1647, 1455, 1088. HRMS calcd. for $\text{C}_{15}\text{H}_{22}\text{O}_2$ 234.1614, found 234.1623.

Synthesis of 1,5,5,11-Tetramethyl-6-oxa-tricyclo[6.3.2.0^{4,13}]tridecan-9-one (**10**)

To a stirred solution of activated CuI (activated under vacuum at 130 °C for 30 min.) (0.24 g, 1.3 mmol) in diethyl ether (5 mL) at 0 °C was added methyl lithium (1.9 mL, 1.4 M, 2.6 mmol). This was stirred for 20 min. at 0 °C and a gray homogenous solution resulted. To this solution the ketone **9** (91.8 mg, 0.392 mmol) in diethyl ether (1.5 mL) was added dropwise over 2 min. The reaction was stirred at 0 °C and allowed to warm slowly to room temperature and stirred overnight before quenched and extracted with a 1:10 solution of $\text{NH}_4\text{OH}/\text{NH}_4\text{Cl}$ (2 X 25 mL), followed by brine (25 mL). The organics were dried over magnesium sulfate and the crude product was purified on a silica column with 10% acetone in hexane to yield the desired ketone **10** (74.1 mg, 0.296 mmol) in 76% yield. ^1H NMR (CDCl_3 , 500 MHz, δ) 4.28 (d, 1H, $J = 11.9$ Hz), 3.59 (dd, 1H, $J = 3.4, 11.9$ Hz), 2.62 (m, 2H), 2.48 (dd, 1H, $J = 3.4, 12.8$ Hz), 2.20 (m, 2H), 1.72 (m, 1H), 1.60 (m, 1H), 1.50 (m, 1H), 1.45 (dd, 1H, $J = 5.5, 14.6$ Hz), 1.29 (m, 1H), 1.28 (s, 3H), 1.23 (m, 1H), 1.18 (m, 1H), 1.13 (s, 3H), 0.99 (d, 3H, $J = 7.3$ Hz), 0.95 (s, 3H). ^{13}C NMR

(CDCl₃, 125 MHz, δ) 213.2, 73.3, 60.1, 50.6, 50.1, 43.1, 41.8, 40.5, 37.5, 33.0, 30.6, 30.2, 27.6, 23.5, 20.2, 19.9. IR (neat, cm⁻¹) 2959, 2895, 1698, 1450, 1111. HRMS calcd. for C₁₆H₂₆O₂ 250.1927 found 250.1928

Synthesis of 1,5,12,12-Tetramethyl-11-oxa-tetracyclo[7,6,2,0^{2,7},0^{13,17}]heptadec-4-en-8-one (13) and 1,4,12,12-Tetramethyl-11-oxa-tetracyclo[7,6,2,0^{2,7},0^{13,17}]heptadec-4-en-8-one (14)

The conjugated ketone **9** (2.21 g, 9.44 mmol) was dissolved in isoprene (200 mL) and exposed to UV light for 15 h at room temperature. The excess isoprene was removed under reduced pressure and the crude product was purified on a silica gel column with 1 : 30 : 69 ratio of isopropanol: diethyl ether: hexane as eluent. The product was isolated in 91% (2.60 g, 8.61 mmol) yield as a mixture of regioisomers favoring **13** in a 2:1 ratio.

Spectroscopic data for major product, alkene **13**: ¹H NMR (500 MHz, CDCl₃) δ 5.36 (s, broad, 1H), 4.10 (dd, 1H, J = 2.4, 12.0 Hz), 3.73 (dd, 1H, J = 4.6, 12.0 Hz), 2.93 (ddd, 1H, J = 5.2, 11.6, 17.7 Hz), 2.57 (m, broad, 1H), 2.50 (m, 1H), 2.22 (m, broad, 1H), 2.15 - 2.12 (m, broad, 2H), 1.98 (m, 1H), 1.93 (ddd, 1H, J = 2.7, 5.8, 14.6 Hz), 1.82 (ddd, 1H, J = 2.4 (two), 14.3 Hz), 1.71 (dd, 1H, J = 5.5, 14.3 Hz), 1.66 (m, broad, 1H), 1.64 (s, 3H), 1.56 (m, broad, 1H), 1.46 (ddd, 1H, J = 4.6, 11.6, 16.2 Hz), 1.38 - 1.31 (m, 2H), 1.30 (s,

3H), 1.13 (s, 3H), 1.05 (m, 1H), 0.91 (s, 3H). ^{13}C NMR (125 MHz, CDCl_3) δ 214.8, 132.8, 120.3, 73.6, 61.3, 51.6, 50.6, 46.9, 46.0, 42.0, 37.2, 34.0, 33.1, 32.9, 29.7, 28.7, 27.1, 24.3, 22.6, 22.2. IR (neat, cm^{-1}) 2906, 1680, 1455, 1367, 1090. HRMS (EI) calcd. for $\text{C}_{20}\text{H}_{30}\text{O}_2$ 302.2240, found 302.2242.

Spectroscopic data for minor product, alkene **14**: ^1H NMR (500 MHz, CDCl_3) δ 5.37 (s, broad, 1H), 4.10 (dd, 1H, $J = 2.1, 11.9$ Hz), 3.71 (dd, 1H, $J = 4.3, 11.9$ Hz), 2.82 (ddd, 1H, $J = 5.2, 11.0, 17.1$ Hz), 2.56 (s, broad, 1H), 2.48 (m, 1H), 2.22 (m, 2H), 1.95 (m, 3H), 1.83 (d, broad, 1H, $J = 14.4$ Hz), 1.72 (dd, 1H, $J = 5.5, 14.4$ Hz), 1.66 (s, broad, 3H), 1.55 (m, 2H), 1.35 (m, 2H), 1.29 (s, 3H), 1.13 (s, 3H), 1.06 (m, 1H), 0.93 (s, 3H). ^{13}C NMR (125 MHz, CDCl_3) δ 214.8, 133.2, 119.8, 73.5, 61.3, 51.4, 50.6, 47.0, 46.2, 42.2, 34.0, 33.6, 33.0, 32.9, 32.8, 29.6, 27.1, 24.3, 23.2, 22.2. IR (neat, cm^{-1}) 2906, 1679, 1452, 1367, 1090. HRMS (EI) calcd. for $\text{C}_{20}\text{H}_{30}\text{O}_2$ 302.2240, found 302.2244.

Synthesis of 4,5-dihydroxy-1,5,12,12-Tetramethyl-11-oxa-tetracyclo[7,6,2,0^{2,7},0^{13,17}]heptadec-8-one (15) and 4,5-dihydroxy-1,4,12,12-Tetramethyl-11-oxa-tetracyclo[7,6,2,0^{2,7},0^{13,17}]heptadec-8-one (16)

A mixture of the two alkene regioisomers **13** and **14** (125 mg, 0.414 mmol) was dissolved in acetone (4 mL) and water (0.5 mL) followed by addition of NMO (134 mg, 1.18 mmol) and OsO_4 (0.67 mL, 2.5 wt % in *t*-butanol, 0.05 mmol). The reaction was stirred at

room temperature for 3 h and was then diluted with diethyl ether (20 mL). The organics were extracted with aq. HCl solution (20 mL, 10%) followed by sat. NaHCO₃ solution (2 x 50 mL). The organics were dried over MgSO₄ and the drying agent removed by filtration before the mixture was concentrated under reduced pressure. The crude product was purified on a silica column with 4% methanol in a 1:1 mixture of diethyl ether and dichloromethane as eluent and gave the major diol **15** in 46% yield (64 mg, 0.19 mmol) and the minor diol **16** in 25% yield (35 mg, 0.10 mmol) after two columns, as well as 2 % of mixed fractions (3 mg, 0.089 mmol).

Spectroscopic data for **15**: ¹H NMR (500 MHz, CDCl₃) δ 3.96 (dd, 1H, *J* = 7.3, 12.7 Hz), 3.68 (dd, 1H, *J* = 2.0, 12.7 Hz), 3.43 - 3.36 (m, 2H), 2.69 (ddd, 1H, *J* = 2.0, 7.3, 7.8 Hz), 2.55 (m, 1H), 2.30 (s, broad, 2H), 2.0 (d, 1H, *J* = 13.7 Hz), 1.82 - 1.55 (m, 8H), 1.46 (ddd, 1H, *J* = 5.9, 12.2, 18.3 Hz), 1.30 (s, 3H), 1.28 (s, 3H), 1.21 (s, 3H), 1.12 (m, 1H), 1.00 (ddd, 1H, *J* = 4.4, 14.2, 18.3 Hz), 0.90 (s, 3H). ¹³C NMR (75 MHz, CDCl₃) δ 215.0, 74.9, 73.8, 70.9, 59.9, 51.9, 48.8, 47.0, 42.8, 40.8, 39.5, 34.6, 34.5, 32.6, 32.5, 26.9, 26.7, 24.8, 24.6, 23.2. IR (neat, cm⁻¹) 3380, 2959, 2849, 1687, 1159, 1097, 1064, 1012 HRMS (ESI) calcd. for C₂₀H₃₂O₄Na 359.2193 found 359.2189.

Spectroscopic data for **16**: ¹H NMR (500 MHz, CDCl₃) δ 3.99 (dd, 1H, *J* = 7.3, 13.1 Hz), 3.72 (dd, 1H, *J* = 2.1, 13.1 Hz), 3.48 (ddd, 1H, *J* = 4.9, 6.1, 11.6 Hz), 3.11 (ddd, 1H, *J* = 3.4, 11.9, 15.0 Hz), 2.66 (1H, ddd, *J* = 2.1, 7.3, 9.2 Hz), 2.58 (m, 1H), 1.98 (m, 1H), 1.92 - 1.87 (m, 2H), 1.86 (s, 1H, (OH)), 1.73 (d, 1H, *J* = 6.1 Hz, (OH)), 1.70 - 1.62 (m, 5H), 1.51 (ddd, 1H, broad), 1.44 (ddd, 1H, broad), 1.31 (1H, hidden by CH₃), 1.31 (s, 6H),

1.20 (s, 3H), 1.01 (m, 1H), 0.90 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3) δ 213.7, 74.0, 73.6, 70.8, 60.3, 51.3, 50.7, 44.1, 43.3, 39.9, 39.5, 34.3, 34.0, 33.9, 32.6, 27.3, 27.0, 25.2, 24.2, 23.1. IR (neat, cm^{-1}) 3457, 3397, 2918, 1687, 1452, 1367, 1303, 1088, 1059. HRMS (ESI) calcd. for $\text{C}_{20}\text{H}_{32}\text{O}_4\text{Na}$ 359.2193 found 359.2196.

Synthesis of [1,5,5-Trimethyl-9-oxo-10-(2-oxo-propyl)-6-oxa-tricyclo[6.3.2.0^{4,13}]tridec-11-yl]-acetaldehyde (17)

To a stirred solution of the diol **15** (46 mg, 0.13 mmol) in methanol (5.8 mL) at room temperature was added a mixture of sodium periodate (88 mg, 0.41 mmol) in water (3.5 mL). The reaction was stirred at room temperature for 12 h then quenched by adding ether (25 mL) and extracted with a saturated brine solution (2 x 25 mL). The organic layer was dried over magnesium sulfate and concentrated under reduced pressure. The product was purified on a silica column with 20% acetone in hexane as eluent and gave the aldehyde in 81% (37 mg, 0.11 mmol) yield. ^1H NMR (500 MHz, CDCl_3) δ 9.80 (d, 1H, $J = 2.2$ Hz), 4.16 (dd, 1H, $J = 12.8, 5.5$ Hz), 3.85 (dd, 1H, $J = 12.8, 6.6$ Hz), 3.37 (ddd, 1H, $J = 16.8, 10.3, 3.7$ Hz) 2.97 (dd, 1H, $J = 17.6, 10.3$ Hz), 2.85 (ddd, 1H, $J = 9.9, 9.9, 5.9$ Hz), 2.58 (dd, 1H, $J = 18.3, 2.2$ Hz), 2.54 – 2.40 (m, 3H), 2.07 (s, 3H), 1.92 (m, 1H), 1.86 – 1.81 (m, 2H), 1.70 – 1.63 (m, 2H), 1.41 (m, 1H), 1.32 – 1.26 (m, 1H), 1.28 (s, 3H), 1.20 (s, 3H), 1.06 (m, 1H), 0.89 (s, 3H). ^{13}C NMR(125 MHz, C_6D_6) δ 212.4, 205.9, 200.3, 73.0, 60.2, 52.1, 50.1, 47.5, 46.3, 42.9, 42.2, 41.2, 35.2, 34.6, 33.7, 29.5, 26.4,

26.2, 26.0, 24.2. IR (neat, cm^{-1}) 2969, 2919, 2720, 1722, 1713, 1691, 1367, 1215, 1096, 1076 HRMS (ESI) calcd. for $\text{C}_{20}\text{H}_{30}\text{O}_4\text{Na}$ 357.2036 found 357.2030.

Synthesis of 5-*epi*-10-*epi*-vibsanin E (19**) and 3-Methyl-but-2-enoic acid (*Z*)-2-[1,5,5-trimethyl-9-oxo-10-(2-oxo-propyl)-6-oxa-tricyclo[6.3.2.0^{4,13}]tridec-11-yl]-vinyl ester (**18**)**

To a stirred solution of the aldehyde **17** (13.2 mg, 0.0395 mmol) in pyridine (1 mL) was added 3,3-dimethylacrylic anhydride (10.6 mg, 0.0581 mmol) and 4-pyrrolidinopyridine (8.5 mg, 0.0577 mmol). The mixture was heated to reflux for 3 h and then allowed to cool to room temperature and ether (20 mL) was added to the crude mixture. The organic layer was extracted with water (2 X 20 mL) followed by a saturated sodium bicarbonate solution (2 x 20 mL). The organics were dried over magnesium sulfate, filtered and the solvents were evaporated to yield 22 mg of crude product. This was purified on silica column with 1 % methanol and 3 % acetone in dichloromethane to yield a 2:1 mixture of *trans* and *cis* 5-*epi*-10-*epi*-vibsanin E (4.8 mg, 0.012 mmol,) favoring the *trans* isomer. This mixture was separated on a preparative HPLC system to deliver the *trans* isomer **19** in 19 % (3.1 mg, 0.075 mmol) yield and the *cis* isomer **18** in 10 % (1.6 mg, 0.038 mmol) yield. HPLC system: 1 % isopropanol in hexane, 15 mL/min at 276 psi. Preparative column: Dynamax 60A, 30 cm long, 2.5 cm outer diameter.

Spectroscopic data for 5-*epi*-10-*epi*-vibsanin E (**86**): ^1H NMR (500 MHz, C_6D_6) δ 7.26 (d, 1H, $J = 12.4$ Hz), 5.71 (qq, 1H, $J = 1.5$ Hz), 5.15 (dd, 1H, $J = 12.4, 10.9$ Hz), 4.51 (dd, 1H, $J = 12.7, 4.9$ Hz), 3.95 (dd, 1H, $J = 12.7, 7.3$ Hz), 3.54 (ddd, 1H, $J = 11.8, 9.8, 3.7$ Hz), 2.91 (ddd, 1H, $J = 7.3, 6.7, 4.4$ Hz), 2.84 (dd, 1H, $J = 18.0, 3.7$ Hz), 2.37 (dd, 1H, $J = 18.0, 9.8$ Hz), 2.12 (dddd, 1H, $J = 10.1, 4.7$ Hz), 2.07 (d, 3H, $J = 1.5$ Hz), 1.71 (m, 2H) 1.62 (s, 3H) 1.57 (dd, 1H, $J = 11.8, 10.9$ Hz), 1.51 (d, br, 1H, $J = 15.0$ Hz), 1.38 (d, 3H, $J = 1.5$ Hz), 1.34 (m, 1H), 1.28 (dd, 1H, $J = 14.6, 6.4$ Hz), 1.14 (s, 3H), 1.06 (s, 3H), 0.97 (ddd, 1H, $J = 9.4, 4.7$ Hz), 0.72 (s, 3H), 0.69 (dd, 1H, $J = 14.6, 4.9$ Hz). ^{13}C NMR (125 MHz, C_6D_6) δ 212.4, 206.2, 163.3, 160.1, 137.7, 115.0, 114.8, 73.1, 60.4, 53.4, 50.5, 47.7, 47.5, 42.4, 40.6, 34.9, 34.3, 33.9, 29.5, 27.1, 26.4, 26.1, 25.8, 24.4, 20.3 IR (neat, cm^{-1}) 2962, 2925, 1728, 1712, 1687, 1642, 1450, 1356, 1221, 1136, 1091, 943. HRMS (ESI) calcd. for $\text{C}_{25}\text{H}_{36}\text{O}_5\text{Na}$ 439.2455 found 439.2458

Spectroscopic data for 3-Methyl-but-2-enoic acid (Z)-2-[1,5,5-trimethyl-9-oxo-10-(2-oxo-propyl)-6-oxa-tricyclo[6.3.2.0^{4,13}]tridec-11-yl]-vinyl ester **85**: ^1H NMR (500 MHz, C_6D_6) δ 7.44 (d, 1H, $J = 6.6$ Hz), 5.66 (qq, 1H, $J = 0.9, 0.6$ Hz), 4.58 (dd, 1H, $J = 12.5, 5.1$ Hz), 4.33 (dd, 1H, $J = 11.0, 6.6$ Hz), 3.95 (dd, 1H, $J = 12.5, 7.0$ Hz), 3.43 (ddd, 1H, $J = 12.5, 9.8, 3.7$ Hz), 3.01 (dd, 1H, $J = 18.3, 9.8$ Hz), 2.96 (ddd, 1H, $J = 12.2, 7.0, 5.1$ Hz), 2.58 (dd, 1H, $J = 12.5, 11.0$ Hz), 2.45 (dd, 1H, $J = 18.3, 3.7$ Hz), 2.14 (m, 1H), 2.05 (d, 3H, $J = 0.6$ Hz), 1.79-1.71 (m, 1H), 1.68 (broad dd, 1H, $J = 14.6$ Hz), 1.62 (s, 3H), 1.40 (m, 1H), 1.36 (d, 3H, $J = 0.9$ Hz), 1.33 (dd, 1H, $J = 14.6, 6.4$ Hz), 1.17 (s, 3H), 1.08 (s, 3H), 1.06 (dd, 1H, $J = 6.4$ Hz), 1.00 (m, 1H), 0.89 (s, 3H), 0.75 (m, 1H). ^{13}C NMR (125 MHz, C_6D_6) δ 212.8, 206.3, 163.1, 160.4, 135.8, 115.0, 114.0, 73.1, 60.5, 53.5, 47.5,

47.14, 47.05, 42.9, 40.8, 35.5, 34.1, 33.8, 29.4, 27.0, 26.42, 26.37, 26.1, 24.5, 20.4. IR (neat, cm^{-1}) 2970, 2905, 1728, 1687, 1642, 1221, 1131. HRMS (ESI) calcd. for $\text{C}_{25}\text{H}_{36}\text{O}_5\text{Na}$ 439.2455 found 439.2457

Section C: Relative Stereochemistry for 5-*epi*-10-*epi*-Vibsanin E (19)

Compounds **13** – **17** did not give characteristic NOE correlations in order to establish the relative stereochemistry of the two alkenes **13** and **14**. Furthermore, none of these compounds were suited for X-ray crystallography either. Because of this the products were taken through to the next steps in the synthesis. In order to establish the correct relative stereochemistry for the synthesized compound **18** and **19** nOe studies were undertaken and this data was compatible with a *trans* addition for the photochemical addition of isoprene across the enone **7**, but epimeric at the 5 and 10 position compared to the natural product. This is clearly seen by inspecting the nOe correlations for **19**. The proton at position 5 is quite characteristic for vibsanin E type of compounds as it is a ddd typically in the 3.0 – 3.6 ppm range with C_6D_6 as the NMR solvent. For the synthesized compound this proton interacts with position 12, 18 and the dd signal from the proton on the enol ether. It also has to be “down” because of the interaction with the proton at position 18, since the geminal proton at this position interacts with the hydrogen at position 2, which is known to be “up” due to the convex shape of these diterpenes. A very powerful indication for the suggested structure is the correlation between position 5 and 12. The relevant proton at position 12 is found at 1.70 ppm because it is shielded by the carbonyl group just above it in space. This correlation can only be observed if the

proton at position 5 is “down”. Since this hydrogen also shows an nOe from the proton on enol ether, the proton at position 10 has to be “up”, Figure 14.

Figure 1. nOe Correlations for 5-*epi*-10-*epi*-Vibsanin E **19**

Section D: ^1H NMR Spectra