Sequential Hydrocarbon Functionalization: Allylic C—H Oxidation, Vinylic C—H Arylation

Jared H. Delcamp and M. Christina White*

General Information: All commercially obtained reagents for the allylic oxidation, Heck arylation reaction were used as received: 1,4-benzoquinone (Sigma-Aldrich); 2-methoxyphenylboronic acid, 4-methoxyphenylboronic acid, 4-chlorophenylboronic acid, 4-fluorophenylboronic acid, 4-formylphenylboronic acid, 3-chlorophenylboronic acid, *o*-tolylboronic acid, 2-methoxyphenyl boronic acid, 3-cyanophenylboronic acid, and 4-methoxycarbonylphenylboronic acid (Frontier Scientific); phenylboronic acid (Sigma-Aldrich); Pd(OAc)₂ (Strem Chemicals). Pd(OAc)₂ was stored in a glove box under an argon atmosphere and weighed out in the air prior to use. Bottles of Pd(OAc)₂ that were found to give suboptimal yields for the Heck arylation were recrystallized according to the procedure provided below. Solvents tetrahydrofuran (THF), diethyl ether (Et₂O), and methylene chloride (CH₂Cl₂) were purified prior to use by passage through a bed of activated alumina (Glass Contour, Laguna Beach, California). Anhydrous N,N-dimethylformamide (DMF) (Sure Seal) was obtained from Sigma-Aldrich and used as received. All allylic oxidation, Heck arylation reactions were run under air with no precautions taken to exclude moisture. An exception to this is the 1-butene reaction run under an atmosphere of O₂. Thin-layer chromatography (TLC) was conducted with E. Merck silica gel 60 F254 precoated plates (0.25 mm) and visualized with UV and potassium permanganate staining. Flash column chromatography was performed as described by Still et al.¹ using EM reagent silica gel 60 (230-240 mesh). ¹H NMR spectra were recorded on a Varian Unity-400 (400 MHz) or Varian Unity-500 (500 MHz) spectrometer and are reported in ppm using solvent as an internal standard (CDCl₃ at 7.26 ppm). Data reported as: s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, m = multiplet, b = broad, ap = apparent; coupling constant(s) in Hz; integration. Proton-decoupled ¹³C NMR spectra were recorded on a Varian Unity-400 (100 MHz) or Varian Unity-500 (125 MHz) spectrometer and are reported in ppm using solvent as an internal standard (CDCl₃ at 77.23 ppm). Regioselectivity of the Heck addition was determined by GC analysis with authentic terminal olefin standard. IR spectra were recorded as thin films on NaCl plates on a Perkin-Elmer Spectrum BX and are reported in frequency of absorption (cm⁻¹). High-resolution mass spectra were obtained at the University of Illinois Mass Spectrometry Laboratory.

Synthesis of Catalyst 1

1,2-bis(phenylsulfinyl)ethane: A 50 mL round bottomed flask (RBF) was charged with a stir bar, 2 g (8.12 mmol, 1 equiv.) of 1,2-bis(phenylthio)ethane, and 12.2 mL of acetic acid. A solution of H₂O₂ (50 wt%, 31.08 mmol, 2.114 mL, 2 equiv.) in acetic acid (6.7 mL) was added dropwise at room temperature. After approximately 15 minutes the solution became homogeneous and turned a pale yellow. An additional 8 mL of acetic acid was then added and the solution allowed to stir for 24 hours at room temperature. The acetic acid was removed with mild heating (45°C) under high vacuum. The pale yellow solid was emulsified in cold ethanol and cold filtered to yield a mixture of the meso and racemic 1,2-bis(phenylsulfinyl)ethane in 92% yield (2.088g). *Meso*-1,2-bis(phenylsulfinyl)ethane ¹H NMR (500 MHz, CDCl₃) δ 7.56-7.52 (m, 10H), 3.05 (s, 4H). ¹³C NMR (500 MHz, CDCl₃) δ 142.29, 131.55, 129.63, 124.10, 47.06. IR (neat) 3048.84, 2970.01, 2922.41, 1442.10, 1036.34, 745.45, 695.70 cm⁻¹; *racemic*-1,2-bis(phenylsulfinyl)ethane. ¹H NMR (500 MHz, CDCl₃) δ 7.51-7.48 (m, 10H), 3.40 (m, 2H), 2.74 (m, 2H). ¹³C NMR (125 MHz, CDCl₃) δ 142.55, 131.53, 129.64, 124.08, 47.94. IR (neat, cm⁻¹) 3053.16, 2911.39, 1443.77, 1084.88, 1042.50, 748.52. HRMS (ESI) *m/z* calculated for C₁₄H₁₄O₂S₂Na [M+Na]⁺: 301.0333, found 301.0320.

Catalyst 1: A flame dried 250 mL flask was charged with 2.53g (9.1 mmol) of 1,2-bis(phenylsulfinyl)ethane, 101 mL of CH₂Cl₂, and 2.04g (9.1 mmol) of Pd(OAc)₂. The mixture was stirred at 40°C for 24h. The reaction becomes a dark red homogenous reaction during the reaction time. The solution was concentrated *in vacuo* and dried with a stream of N₂ for 6 h to give a dark red solid used without further purification. **Note: The catalyst must be stored at below 4°C.** The

catalyst slowly decomposes at ambient temperature; however, may be stored for prolonged periods (months) at reduced temperatures. ^1H NMR and IR data of this catalyst look like 1,2-bis(phenylsulfinyl)ethane ligand and $\text{Pd}(\text{OAc})_2$. Trace amounts of phenyl vinyl sulfoxide can be observed by ^1H NMR (usually ~10% but can vary by +/- 5%). We have noted suboptimal yields (e.g. 70% vs 90%) of Heck arylation with old bottles of $\text{Pd}(\text{OAc})_2$. Recrystallization of $\text{Pd}(\text{OAc})_2$ prior to complexation using the following procedure restores full activity of the catalyst.

$\text{Pd}(\text{OAc})_2$ Recrystallization: “Old” $\text{Pd}(\text{OAc})_2$ was dissolved in minimal refluxing benzene. A black precipitate was removed by hot Acrodisc® filtration. The resulting solution was cooled to room temperature without further manipulation. Amber crystals began to form after ~2 hours. After 24 hours the solution was filtered to give the recrystallized $\text{Pd}(\text{OAc})_2$. A difference in NMR purity was noted between “old” and recrystallized $\text{Pd}(\text{OAc})_2$ samples. Reported hydrogen values are normalized ratios of the smallest peak in the acetate region. “Old” $\text{Pd}(\text{OAc})_2$ (see attached) ^1H NMR (500 MHz, CDCl_3) δ 2.17 (s, 1H), 2.10 (s, 3.6H), 2.07 (s, 6.1H), 2.06 (s, 6.1H), 2.03 (m, 15.3H), 2.00 (m, 95.7H), 1.97 (s, 5.7H), 1.95 (s, 6.3), 1.89 (s, 9.4H). Recrystallized $\text{Pd}(\text{OAc})_2$ (see attached) ^1H NMR (500 MHz, CDCl_3) δ 2.10 (s, 1H), 2.03 (s, 2.8H), 2.00 (s, 40.1H), 1.97 (s, 1.2H), 1.90 (s, 2.3H).

General Procedure for Sequential Allylic Oxidation/Vinyl Arylation: To a 40 mL borosilicate vial was added catalyst **1** (10 mol%, 0.1 mmol) and benzoquinone (2 equiv, 2 mmol). Olefin (1 equiv, 1 mmol) in dioxane (0.33M, 3mL), carboxylic acid (2-4 equiv, 2-4 mmol), and a stir bar were added sequentially. The olefin was weighed out in a 1 dram vial and transferred via dioxane (3 x 1 mL). The vial was capped and stirred at 45°C for 24-48 hours. The allylic oxidation was run until complete conversion of the α -olefin starting material was observed by GC or TLC. The vial was cooled to room temperature, and boronic acid (1.5 equiv, 1.5 mmol) was added to the reaction mixture using weighing paper. The vial was capped and stirred at room temperature for the indicated time. The reaction mixture was diluted with saturated aqueous NH_4Cl and extracted with CH_2Cl_2 or hexane (2 times). The combined organics were washed with 5% aqueous K_2CO_3 (1 time), H_2O (1 time), and dried over MgSO_4 . The mixture was filtered and concentrated *in vacuo*. Purification by flash chromatography (EtOAc/hexanes mixtures) provided the pure product.

d), 127.7, 126.8, 72.2, 60.0, 37.5, 27.0, 21.5, 19.4. IR (neat, cm^{-1}) 3070.72, 3050.07, 3027.54, 2957.36, 2931.20, 2857.85, 1738.12. HRMS (ESI) m/z calculated for $\text{C}_{29}\text{H}_{34}\text{O}_3\text{SiNa}$ $[\text{M}+\text{Na}]^+$: 481.2175, found 481.2171.

(*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-methoxyphenyl)pent-1-en-3-yl acetate (3): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 1 equiv.), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), *tert*-butyl(pent-4-enyloxy)diphenylsilane (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 4-Methoxyphenylboronic acid (1.5 mmol, 228 mg, 1.5 equiv.) was

added and stirred at room temperature for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 10% ethyl acetate/hexanes as eluent to yield (*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-methoxyphenyl)pent-1-en-3-yl acetate as a clear oil. Run 1 (0.248 g, 0.508 mmol, 51%); run 2 (0.252 g, 0.515 mmol, 52%). **Average yield = 52%.** ^1H NMR (500 MHz, CDCl_3) δ 7.66 (m, 4H), 7.45-7.33 (m, 6H), 7.30 (ap d, $J = 9$ Hz, 2H), 6.85 (ap d, $J = 8.5$ Hz, 2H), 6.58 (d, $J = 16$ Hz, 1H), 5.99 (dd, $J = 16, 8$ Hz, 1H), 5.64 (ap q, $J = 7$ Hz, 1H), 3.82 (s, 3H), 3.73 (t, $J = 6.5$ Hz, 2H), 2.02 (s, 3H), 1.96-1.88 (m, 2H), 1.06 (s, 9H). ^{13}C NMR (125 MHz, CDCl_3) δ 170.5, 159.6, 135.8 (ap d), 133.9, 133.8, 133.6, 132.6, 129.8 (ap d), 129.3, 128.0, 127.8, 125.4, 114.1, 72.5, 60.1, 55.5, 37.6, 27.0, 21.6, 19.4. IR (neat, cm^{-1}) 3071.10, 3046.59, 2956.80, 2931.89, 2857.75, 1737.61, 1608.13. HRMS (ESI) m/z calculated for $\text{C}_{30}\text{H}_{36}\text{O}_4\text{SiNa}$ $[\text{M}+\text{Na}]^+$: 511.2281, found 511.2276.

(*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-chlorophenyl)pent-1-en-3-yl acetate (4): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), *tert*-butyl(pent-4-enyloxy)diphenylsilane (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 4-Chlorophenylboronic acid (1.5 mmol, 235 mg, 1.5 equiv.) was added

and stirred at 45°C for 24 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 3% ethyl acetate/hexanes as eluent to yield (*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-chlorophenyl)pent-1-en-3-yl acetate as a clear oil. Run 1 (0.305 g, 0.618 mmol, 62%); run 2 (0.313 g, 0.635 mmol, 64%). **Average yield = 63%.** ^1H NMR (400 MHz, CDCl_3) δ 7.65 (m, 4H), 7.45-7.33 (m, 6H), 7.27 (bs, 4H), 6.56 (d, $J = 15.5$ Hz, 1H), 6.10 (dd, $J = 16, 7.5$ Hz, 1H), 5.64 (ap q, $J = 7.0$ Hz, 1H), 3.72 (t, $J = 5.5$ Hz, 2H), 2.03 (s, 3H), 2.20-1.88 (m, 2H), 1.05 (m, 9H). ^{13}C NMR (100 MHz, CDCl_3) δ 170.4, 135.8, 135.7, 135.0, 133.8, 133.7, 131.5, 129.9, 129.8, 128.9, 128.4, 128.0, 127.9, 72.0, 59.9, 37.4, 27.0, 21.5, 19.4. IR (neat, cm^{-1}) 3074.93, 3049.09, 2931.36, 2858.11, 1738.37. HRMS (ESI) m/z calculated for $\text{C}_{29}\text{H}_{33}\text{O}_3\text{ClSiNa}$ $[\text{M}+\text{Na}]^+$: 515.1785, found 515.1775.

(*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-fluorophenyl)pent-1-en-3-yl acetate (5): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), *tert*-butyl(pent-4-enyloxy)diphenylsilane (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 4-Fluorophenylboronic acid (1.5 mmol, 210 mg, 1.5 equiv.) was added

and stirred at 45°C for 7 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (125 mL SiO_2) with 5% ethyl acetate/hexanes as eluent to yield (*E*)-5-(*tert*-butyldiphenylsilyloxy)-1-(4-fluorophenyl)pent-1-en-3-yl acetate as a clear oil. Run 1 (0.3482 g, 0.732 mmol, 73%); run 2 (0.3508 g, 0.737 mmol, 74%). **Average yield = 74%.** ^1H NMR (500 MHz, CDCl_3) δ 7.05 (m, 4H), 6.79-6.66 (m, 8H), 6.35 (ap t, $J = 8.5$ Hz, 2H), 5.97 (d, $J = 16$ Hz, 1H), 5.43 (dd, $J = 16, 7.5$ Hz, 1H), 5.05 (ap q, $J = 7$ Hz, 1H), 3.12 (t, $J = 6$ Hz, 2H), 1.43-1.30 (m, 2H), 1.39 (s, 3H), 0.45 (s, 9H). ^{13}C NMR (125 MHz, CDCl_3) δ 170.3, 163.6, 161.6, 135.7 (ap d), 133.7 (ap d), 132.6 (ap d), 131.7, 129.8, 128.3 (ap d), 127.8, 127.4,

115.7, 115.5, 72.1, 59.9, 37.4, 27.0, 21.4, 19.3. IR (neat, cm^{-1}) 3071.40, 3043.92, 2950.90, 2931.20, 2858.02, 1737.30, 1601.84. HRMS (ESI) m/z calculated for $\text{C}_{29}\text{H}_{33}\text{O}_3\text{FSiNa}$ $[\text{M}+\text{Na}]^+$: 499.2081, found 499.2086.

(E)-5-(tert-butyl diphenylsilyloxy)-1-(4-formylphenyl)pent-1-en-3-yl acetate (6): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), tert-butyl(pent-4-enyloxy)diphenylsilane (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 4-formylphenylboronic acid (1.5 mmol, 225 mg, 1.5 equiv.) was added and heated to 45°C for 5 hours. The mixture was diluted with 40 mL of sat. aq.

NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 15% ethyl acetate/hexanes as eluent to yield (E)-5-(tert-butyl diphenylsilyloxy)-1-(4-formylphenyl)pent-1-en-3-yl acetate as a clear oil. Run 1 (0.293 g, 0.602 mmol, 60%); run 2 (0.286 g, 0.588g, 59%). **Average yield = 60%.** ^1H NMR (500 MHz, CDCl_3) δ 9.99 (s, 1H), 7.83 (d, J = 8.5 Hz, 2H), 7.66 (d, J = 7 Hz, 4H), 7.49 (d, J = 8.5 Hz, 2H), 7.43-7.34 (m, 6H), 6.65 (d, J = 16.5 Hz, 1H), 6.31 (dd, J = 16.0, 7.0 Hz, 1H), 5.68 (ap q, J = 7.0 Hz, 1H), 3.75 (t, J = 5.5 Hz, 2H), 2.05 (s, 3H), 2.06-1.91 (m, 2H), 1.06 (s, 9H). ^{13}C NMR (125 MHz, CDCl_3) δ 191.9, 170.4, 142.6, 135.8, 135.7, 133.7, 133.6, 131.6, 131.3, 130.3, 129.9 (ap d), 127.9, 127.3, 71.8, 59.9, 37.4, 27.0, 21.4, 19.4. IR (neat, cm^{-1}) 3071.13, 3048.95, 2957.46, 2931.26, 2857.81, 2737.53, 1738.21, 1698.58. HRMS (ESI) m/z calculated for $\text{C}_{30}\text{H}_{34}\text{O}_4\text{SiNa}$ $[\text{M}+\text{Na}]^+$: 509.2124, found 509.2125.

(E)-1-(3-Chlorophenyl)-7-methylocta-1,6-dien-3-yl acetate (7): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 7-methyl-1,6-octadiene (1 mmol, 124 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. 3-Chlorophenylboronic acid (1.5 mmol, 235 mg, 1.5 equiv.), benzoquinone (1 mmol, 183 mg, 1 equiv.), acetic

acid (1 mmol, 60 mg, 1 equiv.) were added and heated to 45°C for 6 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 4% ethyl acetate/hexanes as eluent to yield (E)-1-(3-chlorophenyl)-7-methyl-1,6-octadien-3-yl acetate as a clear oil. Run 1 (0.178 g, 0.606 mmol, 61%); run 2 (0.193 g, 0.654 mmol, 65%). **Average yield = 63%.** ^1H NMR (400 MHz, CDCl_3) δ 7.36 (s, 1H), 7.22 (m, 3H), 6.35 (d, J = 15.6 Hz, 1H), 6.13 (dd, J = 16.4, 7.6 Hz, 1H), 5.38 (ap q, J = 6.4 Hz, 1H), 5.10 (t, J = 7.2 Hz, 1H), 2.08 (s, 3H), 2.05 (m, 2H), 1.77 (m, 2H), 1.69 (s, 3H), 1.58 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 170.6, 138.5, 134.7, 132.7, 131.2, 130.0, 129.5, 128.0, 126.6, 125.0, 123.3, 74.2, 34.6, 25.9, 23.9, 21.5, 17.9. IR (neat, cm^{-1}) 2967.04, 2928.00, 2858.35, 1738.54, 1594.59, 1566.65. HRMS (ESI) m/z calculated for $\text{C}_{17}\text{H}_{21}\text{O}_2\text{ClNa}$ $[\text{M}+\text{Na}]^+$: 315.1128, found 315.1133.

(E)-5-(1,3-dioxoisindolin-2-yl)-1-o-tolylpent-1-en-3-yl acetate (8): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 2-(pent-4-enyl)isindoline-1,3-dione (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. *o*-Tolylphenylboronic acid (2 mmol, 272 mg, 1.5 equiv.), benzoquinone (1 mmol, 108 mg, 1 equiv.), and acetic

acid (1 mmol, 60 mg, 1 equiv.) were added and heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 20% ethyl acetate/hexanes as eluent to yield (E)-5-(1,3-dioxoisindolin-2-yl)-1-o-tolylpent-1-en-3-yl acetate as a white powder. Run 1 (0.274 g, 0.749 mmol, 75%); run 2 (0.258 g, 0.706 mmol, 71%). **Average yield = 73%.** ^1H NMR (400 MHz, CDCl_3) δ 7.83 (m, 2H), 7.69 (m, 2H), 7.36 (m, 1H), 7.12 (m, 3H), 6.87 (d, J = 16 Hz, 1H), 6.00 (dd, J = 16, 7.2 Hz, 1H), 5.44 (ap q, J = 6.4 Hz, 1H), 3.81 (m, 2H), 2.34 (s, 3H), 2.15 (m, 2H), 2.07 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 170.4, 168.4, 136.0, 135.3, 134.2, 132.2, 131.3, 130.5, 128.1, 127.9, 126.2,

125.8, 123.4, 72.6, 34.5, 33.1, 21.4, 20.0. IR (neat, cm^{-1}) 3061.90, 3020.60, 2946.33, 1772.19, 1738.23, 1714.39. HRMS (ESI) m/z calculated for $\text{C}_{22}\text{H}_{21}\text{O}_4\text{NNa}$ $[\text{M}+\text{Na}]^+$: 386.1368, found 386.1375.

(E)-5-(1,3-dioxoisindolin-2-yl)-1-(2-methoxyphenyl)pent-1-en-3-yl acetate (9):

To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 2-(pent-4-enyl)isoindoline-1,3-dione (1 mmol, 325 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. 2-Methoxyphenylboronic acid (1.5 mmol, 228 mg, 1.5 equiv.), benzoquinone (1 mmol, 108 mg, 1 equiv.), and acetic acid (1 mmol, 60 mg, 1 equiv.) were added and stirred at room temperature for 5 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (125 mL SiO_2) with 20% ethyl acetate/hexanes as eluent to yield (E)-5-(1,3-dioxoisindolin-2-yl)-1-(2-methoxyphenyl)pent-1-en-3-yl acetate as a clear oil. Run 1 (0.201 g, 0.530 mmol, 53%); run 2 (0.211 g, 0.556 mmol, 56%). **Average yield = 55%.** ^1H NMR (500 MHz, CDCl_3) δ 7.81 (dd, J = 5.0, 3.0 Hz, 2H), 7.68 (dd, J = 5.0, 3.0 Hz, 2H), 7.36 (d, J = 6.5 Hz, 1H), 7.19 (t, J = 8 Hz, 1H), 6.93 (d, J = 16.5 Hz, 1H), 6.87 (t, J = 7.5 Hz, 1H), 6.82 (d, J = 8.5 Hz, 1H), 6.19 (dd, J = 16, 7.5 Hz, 1H), 5.44 (ap q, J = 5.5 Hz, 1H), 3.87-3.74 (m, 5H), 2.16-2.07 (m, 5H). ^{13}C NMR (125 MHz, CDCl_3) δ 170.4, 168.4, 157.1, 134.1, 132.2, 129.2, 128.1, 127.3, 127.2, 125.1, 123.3, 120.6, 110.9, 72.8, 55.5, 34.5, 33.0, 21.4. IR (neat, cm^{-1}) 2941.06, 2838.34, 1772.05, 1738.07, 1714.00, 1598.14. HRMS (ESI) m/z calculated for $\text{C}_{22}\text{H}_{21}\text{NO}_5\text{Na}$ $[\text{M}+\text{Na}]^+$: 402.1317, found 402.1327.

(E)-5-(4-methoxybenzyloxy)-1-phenylpent-1-en-3-yl 3,5-dinitrobenzoate (10):

To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 5-(4-methoxybenzyloxy)-1-pentene (1 mmol, 206 mg, 1 equiv.) in dioxane (3mL), 3,5-dinitrobenzoic acid (4 mmol, 848 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. Phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), benzoquinone (1 mmol, 108 mg, 1 equiv.), and acetic acid (1 mmol, 60 mg, 1 equiv.) were then added to the reaction, followed by heating to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with methylene chloride (2 x 75 mL). The organic layers were combined and rinsed with 5% K_2CO_3 (75 mL), H_2O (75 mL), and dried with MgSO_4 . The filtered organic solution was concentrated via rotary evaporator. The crude yellow oil was further purified via silica chromatography (400 mL SiO_2) with 14% ethyl acetate/14% methylene chloride/hexanes as eluent to give (E)-1-Phenyl-5-(4-methoxybenzyloxy)-1-penten-3-(3,5-dinitrobenzoate) as a yellow solid. Run 1 (0.269 g, 0.55 mmol, 55%); run 2 (0.253 g, 0.51 mmol, 51%). **Average yield = 53%.** ^1H NMR (500 MHz, CDCl_3) δ 9.16 (m, 1H), 9.02 (d, J = 1.5 Hz, 2H), 7.38 (d, J = 8.5 Hz, 2H), 7.32 (t, J = 8 Hz, 2H), 7.27 (m, 1H), 7.14 (d, J = 8.5 Hz, 2H), 6.74 (d, J = 16 Hz, 1H), 6.66 (d, J = 8.5 Hz, 2H), 6.23 (dd, J = 16, 7.5 Hz, 1H), 5.93 (m, 1H), 4.40 (d, J = 11.5 Hz, 1H), 4.34 (d, J = 11 Hz, 1H), 3.70 (s, 3H), 3.61 (m, 2H), 2.27 (m, 1H), 2.15 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ 162.0, 159.1, 148.5, 135.9, 134.5, 134.4, 130.1, 129.7, 129.5, 128.8, 128.6, 126.9, 126.1, 122.2, 113.6, 75.8, 72.9, 65.7, 55.3, 34.8. IR (neat, cm^{-1}) 3101.83, 2934.64, 2863.07, 1730.84, 1544.89. HRMS (ESI) m/z calculated for $\text{C}_{26}\text{H}_{24}\text{O}_8\text{N}_2\text{Na}$ $[\text{M}+\text{Na}]^+$: 515.1430, found 515.1436.

(E)-1-(3-cyanophenyl)-6-(methoxy(methyl)amino)-6-oxohex-1-en-3-yl 4-nitrobenzoate (11):

To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), *N*-methoxy-*N*-methylhex-5-enamide (1 mmol, 157 mg, 1 equiv.) in dioxane (3mL), *p*-nitrobenzoic acid (2 mmol, 334 mg, 2 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. 3-cyanophenylboronic acid (1.5 mmol, 220 mg, 1.5 equiv.), benzoquinone (1 mmol, 108 mg, 1 equiv.), and acetic acid (1 mmol, 60 mg, 1 equiv.) were added and heated to 45°C for 7 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with CH_2Cl_2 (2 x 75 mL). The organic layers were combined, rinsed with 5% K_2CO_3 (75mL), H_2O (75 mL) and dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 50% ethyl

acetate/hexanes as eluent to yield (*E*)-1-(3-cyanophenyl)-6-(methoxy(methyl)amino)-6-oxohex-1-en-3-yl 4-nitrobenzoate as a clear oil. Run 1 (0.320 g, 0.757 mmol, 76%); run 2 (0.328 g, 0.776 mmol, 78%). **Average yield = 77%.** ^1H NMR (500 MHz, CDCl_3) δ 8.30 (d, J = 8.5 Hz, 2H), 8.23 (d, J = 9 Hz, 2H), 7.66 (s, 1H), 7.60 (d, J = 8 Hz, 1H), 7.34 (d, J = 7.5 Hz, 1H), 7.43 (ap t, J = 7.5 Hz, 1H), 6.72 (d, J = 16 Hz, 1H), 6.31 (dd, J = 16, 7 Hz, 1H), 5.75 (ap q, J = 7 Hz, 1H), 3.63 (s, 3H), 3.14 (s, 3H), 2.60 (t, J = 7, 2H), 2.27 (m, 2H). ^{13}C NMR (125 MHz, CDCl_3) δ 173.1, 164.1, 150.8, 137.4, 135.7, 131.6, 131.5, 131.0 (ap d), 130.3, 129.7, 129.5, 123.8, 118.7, 113.1, 75.8, 61.4, 32.4, 29.3, 27.7. IR (neat, cm^{-1}) 3110.89, 3078.75, 2938.96, 2230.86, 1724.36, 1660.20. HRMS (ESI) m/z calculated for $\text{C}_{22}\text{H}_{22}\text{O}_6\text{N}_3$ $[\text{M}+\text{H}]^+$: 424.1509, found 424.1505.

(*E*)-methyl 4-(3-(2-bromoacetoxy)-3-cyclohexylprop-1-enyl)benzoate (12): To a 40 mL borosilicate vial was added catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), allylcyclohexane (1 mmol, 124 mg, 1 equiv.) in dioxane (3mL), bromoacetic acid (4 mmol, 556 mg, 4 equiv.), and a stir bar. The mixture was heated to 45°C for 48 hours. 4-methoxycarbonylphenylboronic acid (1.5 mmol, 270 mg, 1.5 equiv.), benzoquinone (1 mmol, 108 mg, 1 equiv.), and acetic acid (1 mmol, 60 mg, 1 equiv.) were added and heated to 45°C for 24 hours.

The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL). The organic layers were dried over MgSO_4 . After concentration, the crude product was purified via silica chromatography (140 mL SiO_2) with 10% ethyl acetate/hexanes as eluent to yield (*E*)-methyl 4-(3-(2-bromoacetoxy)-3-cyclohexylprop-1-enyl)benzoate as a clear oil. Run 1 (0.247 g, 0.627 mmol, 63%); run 2 (0.234 g, 0.595 mmol, 60%). **Average yield = 62%.** ^1H NMR (500 MHz, CDCl_3) δ 7.99 (d, J = 8 Hz, 2H), 7.44 (d, J = 8 Hz, 2H), 6.66 (d, J = 15.5 Hz, 1H), 6.23 (dd, J = 15.5, 7.5 Hz, 1H), 5.26 (ap t, J = 7.5 Hz, 1H), 3.91 (s, 3H), 3.87 (d, J = 3 Hz, 2H), 1.86-1.67 (m, 6H), 1.29-1.02 (m, 5H). ^{13}C NMR (125 MHz, CDCl_3) δ 167.0, 166.8, 140.8, 133.1, 130.1, 129.6, 128.4, 126.7, 80.9, 52.3, 42.0, 28.9, 28.7, 26.4 (ap d), 26.0 (ap d). IR (neat, cm^{-1}) 2929.53, 2853.45, 1720.77. HRMS (ESI) m/z calculated for $\text{C}_{19}\text{H}_{23}\text{O}_4\text{BrNa}$ $[\text{M}+\text{Na}]^+$: 417.0677, found 417.0683.

(*E*)-1-(benzo[*d*][1,3]dioxol-5-yl)undec-1-en-3-yl 2-(*tert*-butoxycarbonyl)acetate (13): To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 1-undecene (1 mmol, 154 mg, 1 equiv.) in dioxane (3mL), *N*-Boc-glycine (2 mmol, 350 mg, 2 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 3,4-Methylenedioxyphenylboronic acid (1.5 mmol, 249 mg, 1.5

equiv.) was added and stirred at room temperature for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL). The organic layers were combined, rinsed with 60 mL of 5% K_2CO_3 (aq.) solution, and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (125 mL SiO_2) with 15% ethyl acetate/5% triethylamine/hexanes as eluent to yield (*E*)-1-(benzo[*d*][1,3]dioxol-5-yl)undec-1-en-3-yl 2-(*tert*-butoxycarbonyl)acetate as a pale oil. NOTE: Product was found to be slightly unstable to silica gel unless buffered. Run 1 (0.3106 g, 0.695 mmol, 70%); run 2 0.3001 g, 0.671 mmol, 67%). **Average yield = 69%.** ^1H NMR (500 MHz, CDCl_3) δ 6.92 (s, 1H), 6.81 (d, J = 8 Hz, 1H), 6.76 (d, J = 8 Hz, 1H), 6.53 (d, J = 16 Hz, 1H), 5.96 (s, 2H), 5.93 (dd, J = 16, 8Hz, 1H), 5.42 (ap q, J = 7 Hz, 1H), 5.01 (m, 1H), 3.96 (dd, J = 18.5, 6 Hz, 1H), 3.89 (dd, J = 18, 5.5 Hz, 1H), 1.75-1.65 (m, 2H), 1.45 (s, 9H), 1.35-1.22 (m, 12H), 0.88 (t, J = 7.5 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 170.0, 155.9, 148.2, 147.8, 133.1, 130.8, 125.5, 121.8, 108.5, 105.9, 101.3, 80.1, 76.6, 42.8, 34.8, 32.0, 29.6, 29.5, 29.4, 28.5, 25.4, 22.9, 14.3. IR (neat, cm^{-1}) 3392.35, 2928.61, 2857.15, 1744.28, 1715.64, 1505.27. HRMS (ESI) m/z calculated for $\text{C}_{25}\text{H}_{37}\text{NO}_6\text{Na}$ $[\text{M}+\text{Na}]^+$: 470.2519, found 470.2509.

(E)-1-(4-Bromophenyl)undec-1-en-3-yl 2-(tert-butoxycarbonyl) acetate

(14): To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 1-undecene (1 mmol, 154 mg, 1 equiv.) in dioxane (3mL), *N*-Boc-glycine (2 mmol, 350 mg, 2 equiv.), and a stir bar. The mixture was heated to 45°C for 24 hours. 4-Bromophenylboronic acid (1.5 mmol, 301 mg, 1.5 equiv.) was added and heated to 45°C for 5 hours. The mixture was diluted with 40 mL of

sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (140 mL SiO_2) with 10% ethyl acetate/hexanes as eluent to yield (*E*)-1-(4-Bromophenyl)-1-undecen-3-*N*-boc-glycine as a pale oil. Run 1 (0.359 g, 0.746 mmol, 75%); run 2 0.360 g, 0.749 mmol, 75%). **Average yield = 75%.** ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, J = 8.4 Hz, 2H), 7.23 (d, J = 8.4 Hz, 2H), 6.54 (d, J = 15.6 Hz, 1H), 6.09 (dd, J = 15.6, 7.2 Hz, 1H), 5.42 (ap q, J = 6.8 Hz, 1H), 5.02 (bs, 1H), 3.92 (m, 2H), 1.70 (m, 2H), 1.44 (s, 9H), 1.34-1.85 (m, 12H), 0.87 (t, J = 6.8 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 169.9, 155.8, 135.2, 131.9, 131.8, 128.3, 128.1, 122.0, 80.1, 76.1, 42.7, 34.6, 32.0, 29.6, 29.5, 29.4, 28.5, 25.3, 22.8, 14.3. IR (neat, cm^{-1}) 3369.70, 2928.07, 2856.25, 1747.79, 1716.11, 1508.35. HRMS (ESI) m/z calculated for $\text{C}_{24}\text{H}_{37}\text{NO}_4\text{Br}$ $[\text{M}+\text{H}]^+$: 482.1906, found 482.1911.

(E)-1-(3-Methoxyphenyl)-1-buten-3-ol reaction scheme.

(E)-1-(3-Methoxyphenyl)-1-buten-3-ol (15): 1-Butene (0.112 g, 2 mmol, 0.178 mL, 1 equiv.) was condensed in a 25 mL graduated tube at -78°C. The liquid was transferred via cannula under a positive pressure of oxygen to a 40 mL sealed tube (cooled to -78°C). Dioxane (3mL, purged with O_2 for 30 minutes prior to use) was added to the graduated tube at -78°C (to trap any residual butene in the cannula), then warmed to the point when it could

be transferred with cannule (with a positive pressure of O_2) to the sealed tube. To the sealed tube (maintained at -78°C) was added the following solids: Benzoquinone (0.432 g, 4 mmol, 2 equiv.), catalyst **1** (101 mg, 0.2 mmol, 10 mol%), 4-nitrobenzoic acid (1.00 g, 6 mmol, 3 equiv.). After addition of each solid, 1 mL of oxygenated dioxane was added and used to rinse the solid off of the sides of the sealed tube (3 mL total). The tube was sealed under an atmosphere of O_2 , warmed to room temperature with stirring, and then heated to 45°C for 20h. The mixture was cooled to room temperature, N_2 was bubbled through it for 5 minutes (to remove unreacted 1-butene). 3-Methoxyphenyl boronic acid (0.456 g, 3 mmol, 1.5 equiv.) was then added and heated in the sealed tube at 45°C. Reaction progress was monitored by TLC until disappearance of the allylic ester was observed (~4h). Upon completion the mixture was cooled to room temperature and diluted with 100 mL of a saturated aqueous potassium carbonate/methanol mixture (1:1) to effect hydrolysis of the *p*-nitrobenzyl ester. The reaction progress was again monitored by TLC. Upon completion (~4h) the mixture was diluted with 100 mL H_2O and extracted 2 x 100 mL CH_2Cl_2 . The organic layers were combined and dried over MgSO_4 . Concentration via rotary evaporator yielded a crude mixture which was purified via silica gel chromatography (130 mL SiO_2) in 20% ethyl acetate/hexanes as eluent to afford (*E*)-1-(3-Methoxyphenyl)-1-buten-3-ol as a clear oil. Run 1 (0.276 g, 1.55 mmol, 77%); run 2 (0.269 g, 1.34 mmol, 76%). **Average yield = 77%.** ^1H NMR (400 MHz, CDCl_3) δ 7.23 (t, J = 8 Hz, 1H), 6.97 (d, J = 8 Hz, 1H), 6.91 (t, J = 1.6, 1H), 6.80 (ddd, J = 8, 2.4, .4, 1H), 6.53 (d, J = 15.6, 1H), 6.26 (dd, J = 15.6, 6, 4.48 (ap dp, J = 6.4, .4, 1H), 3.81 (s, 3H), 1.78 (bs, 1H), 1.37 (d, J = 6.4, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 159.9, 138.3, 134.0, 129.8, 129.4, 119.3, 113.5, 111.9, 69.1, 55.4, 23.6. IR (neat, cm^{-1}) 3368.94, 2970.20, 2930.23, 2835.91, 1598.23, 1580.15. HRMS (ESI) m/z calculated for $\text{C}_{11}\text{H}_{14}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$: 201.0891, found 201.0889.

Table 2.

$ \begin{array}{ccc} \text{C}_8\text{H}_{17} \text{---} \text{CH(R)---CH=CH}_2 & \xrightarrow[\text{BQ (0-2 equiv.), air, dioxane, 4h, 45}^\circ\text{C}]{\text{Pd(II)L}_n \text{ (10 mol\%)} \\ \text{PhB(OH)}_2 \text{ (1.5 equiv.)} \\ \text{AcOH (0-4 equiv.)}} & \text{C}_8\text{H}_{17} \text{---} \text{CH(R)---CH=CHPh} \\ \mathbf{16} & & \mathbf{17} \end{array} $						
entry	R	Pd(II)L _n	AcOH	BQ	isolated ^a yield	internal:terminal ^{b,c}
1a	OAc	1	4 equiv.	2 equiv.	98%	41:1 ^d
1b	OBn				92%	>20:1 ^e
1c	OMe				95%	>20:1 ^e
1d	Me ^g				75%	8:1 ^f
1e	OTBDPS				94%	>20:1 ^e
2		Pd(OAc) ₂	4 equiv.	2 equiv.	37%	---
3		1	4 equiv.	---	8%	---
4		1	---	2 equiv.	71%	---
5		1	1 equiv.	2 equiv.	94%	---

^aAverage yields for 2 runs done at 1.0 mmol. ^bE:Z and olefin isomer ratios determined on crude material. ^cE:Z = >20:1, ¹H NMR. ^dGC using authentic terminal olefin. ^e¹H NMR. ^fInternal:trisubstituted:terminal = 8:1:1 (¹H NMR). ^g3-methyl-1-hexene.

Table 2.

1.0 mmol scale:

(E)-1-phenylundec-1-en-3-yl acetate (17): To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%) or palladium acetate (0.1 mmol, 22.4 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), undec-1-en-3-yl acetate (1 mmol, 212 mg, 1 equiv.) in dioxane (3mL), acetic acid 4 equiv. (4 mmol, 24.0 mg, 0.231 mL) or 1 equiv. (1 mmol, 60.0 mg, 58 μ L), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.) and a stir bar. After heating to 45°C for 4 hours, the mixture was diluted with 40 mL of sat. aq. NH₄Cl and extracted with hexanes (2 x 75 mL). The organic layers were combined, rinsed with 60 mL of 5% K₂CO₃ (aq.) solution, and dried over MgSO₄. After concentration, the crude product was purified via silica gel chromatography (125 mL SiO₂) with 5% ethyl acetate/hexanes as eluent to yield (E)-1-phenylundec-1-en-3-yl acetate (**17**) as a colorless oil. All yields are the average of 2-3 runs. ¹H NMR (400 MHz, CDCl₃) δ 7.38 (m, 2H), 7.31 (m, 2H), 7.25 (m, 1H), 6.59 (d, *J* = 16 Hz, 1H), 6.12 (dd, *J* = 15.6, 7.2 Hz, 1H), 5.39 (ap q, *J* = 7.2 Hz, 1H), 2.08 (s, 3H), 1.76-1.61 (m, 2H), 1.35-1.22 (m, 12H), 0.87 (t, *J* = 6.8 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 170.7, 136.6, 132.7, 128.8, 128.1 (ap d), 126.8, 75.1, 34.8, 32.1, 29.7, 29.6, 29.5, 25.4, 22.9, 21.6, 14.4. IR (neat, cm⁻¹) 2926.47, 2856.91, 1737.89, 1237.51. HRMS (EI) *m/z* calculated for C₁₉H₂₈O₂ [M]⁺; 288.2089, found 288.2088.

NOTE: 2,6-Dimethylbenzoquinone (2 mmol, 272.0 mg, 2 equiv.) may be used instead of benzoquinone to yield (E)-1-phenylundec-1-en-3-yl acetate as a colorless oil (0.268 g, 0.932 mmol, 93%) with >20:1 E:Z selectivities by crude ¹H NMR and 66:1 internal:terminal olefin selectivities by crude GC analysis.

2-phenylundec-1-en-3-yl acetate: Terminal olefin standard. ¹H NMR (400 MHz, CDCl₃) δ 7.43 (m, 2H), 7.35-7.29 (m, 3H), 5.67 (t, *J* = 6 Hz, 1H), 5.28 (ap d, *J* = 6 Hz, 2H), 2.12 (s, 3H), 1.61 (ap q, *J* = 6.4 Hz, 2H), 1.30-1.20 (m, 12H), 0.86 (t, *J* = 7.2 Hz, 3H). ¹³C (125 MHz, CDCl₃) δ 170.6, 148.8, 139.8, 128.5, 127.9, 127.2, 113.8, 75.8, 34.1, 32.0, 29.6, 29.4 (ap d), 25.6, 22.8, 21.5, 14.3. IR (neat, cm⁻¹) 2953.63, 2927.47, 2856.50, 1741.83, 1237.16. HRMS (EI) *m/z* calculated for C₁₉H₂₈O₂ [M]⁺; 288.2089, found 288.2091.

(E)-1-((1-phenylundec-1-en-3-yloxy)methyl)benzene: To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 1-((undec-1-en-3-yloxy)methyl)benzene (1 mmol, 260 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar. The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH₄Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K₂CO₃ (1 x 75 mL). The organic layers were combined and dried over MgSO₄. After concentration, the crude product was purified via silica gel chromatography (125 mL SiO₂) with 3% ethyl acetate/hexanes as eluent to yield (E)-1-((1-phenylundec-1-en-3-

ylxy)methyl)benzene as a clear oil. Run 1 (0.308 g, 0.923 mmol, 92%); run 2 (0.305 g, 0.914 mmol, 91%). **Average yield = 92%.** The Heck arylation may also be performed at room temperature for 4h: run 1 (0.309 g, 0.924 mmol, 92%); run 2 (0.311 g, 0.931, 93%). **Average yield = 93%.** By crude ^1H NMR all selectivities were >20:1 E:Z and >20:1 internal olefin : terminal olefin. ^1H NMR (400 MHz, CDCl_3) δ 7.42 (d, J = 7.2 Hz, 2H), 7.36-7.26 (m, 8H), 6.53 (d, J = 15.6 Hz, 1H), 6.13 (dd, J = 16, 8 Hz, 1H), 4.64 (d, J = 12 Hz, 1H), 4.41 (d, J = 12 Hz, 2H), 3.90 (q, J = 8 Hz, 1H), 1.80-1.71 (m, 1H), 1.63-1.54 (m, 1H), 1.46-1.26 (m, 12H), 0.88 (t, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 139.1, 136.9, 132.5, 131.04, 128.9, 128.6, 128.0, 127.9, 127.7, 126.7, 80.4, 70.3, 36.2, 32.1, 29.9, 29.8, 29.6, 25.8, 22.9, 14.4. IR (neat, cm^{-1}) 3025.86, 2952.73, 2926.76, 2852.95. HRMS (CI) m/z calculated for $\text{C}_{24}\text{H}_{33}\text{O}$ $[\text{M}+\text{H}]^+$: 337.2532, found 337.2538.

(E)-1-(3-methoxyundec-1-enyl)benzene: To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 3-methoxyundec-1-ene (1 mmol, 184 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar.

The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K_2CO_3 (1 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (125 mL SiO_2) with 3% ethyl acetate/hexanes as eluent to yield (E)-1-(3-methoxyundec-1-enyl)benzene as a clear oil. Run 1 (0.246 g, 0.946 mmol, 95%); run 2 (0.247 g, 0.951 mmol, 95%). **Average yield = 95%.** Heck arylation may also be performed at room temperature for 4h: run 1 (0.243 g, 0.936 mmol, 94%); run 2 (0.235 g, 0.904 mmol, 90%). **Average yield = 92%.** By crude ^1H NMR all selectivities were >20:1 E:Z and >20:1 internal olefin : terminal olefin. ^1H NMR (400 MHz, CDCl_3) δ 7.42 (d, J = 7.2 Hz, 2H), 7.34 (t, J = 7.6 Hz, 2H), 7.26 (t, J = 7.2 Hz, 1H), 6.54 (d, J = 16 Hz, 1H), 6.06 (d, J = 15.6, 8 Hz, 1H), 3.70 (q, J = 7.6 Hz, 1H), 3.33 (s, 3H), 1.74-1.66 (m, 1H), 1.60-1.52 (m, 1H), 1.44-1.20 (m, 12H), 0.89 (t, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 136.9, 132.4, 130.7, 128.8, 127.8, 126.6, 82.9, 56.4, 36.0, 32.1, 29.9, 29.8, 29.5, 25.6, 22.9, 14.3. IR (neat, cm^{-1}) 2957.02, 2928.01, 2923.33, 2854.66. HRMS (CI) m/z calculated for $\text{C}_{18}\text{H}_{29}\text{O}$ $[\text{M}+\text{H}]^+$: 261.2218, found 261.2216.

To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 3-methylhex-1-ene (1 mmol, 98.2 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar. The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K_2CO_3 (1 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (125 mL SiO_2) with 100% hexanes as eluent to yield an inseparable mixture of isomers as a clear oil. Note: Due to volatility concerns, the hexanes was removed via rotary evaporator at 0°C. By crude ^1H NMR the ratio of internal olefin : trisubstituted olefin : terminal olefin at 45°C is 8:1:1. Run 1 (0.129 g, 0.742 mmol, 74%); run 2 (0.132 g, 0.759 mmol, 76%). **Average yield = 75%.** Heck arylation may also be performed at room temperature for 4h: run 1 (0.134 g, 0.777 mmol, 78%). **Yield = 78%.** By crude ^1H NMR the room temperature ratio of internal olefin : trisubstituted olefin : terminal olefin, 10:7:1. By crude ^1H NMR all selectivities were >20:1 E:Z. ^1H NMR (400 MHz, CDCl_3) δ Internal olefin: 6.34 (d, J = 16 Hz, 1H), 6.10 (dd, J = 16, 8 Hz, 1H), 2.31 (m, 1H). Trisubstituted olefin: 5.35 (t, J = 7.2 Hz, 1H), 3.37 (d, J = 7.2 Hz, 2H). Terminal olefin: 5.18 (s, 1H), 5.03 (s, 1H), 2.12 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ Internal olefin: 138.2, 137.3, 128.7, 128.1, 127.0, 126.2, 39.6, 37.3, 20.9, 20.7, 14.4. HRMS (CI) m/z calculated for $\text{C}_{13}\text{H}_{19}$ $[\text{M}+\text{H}]^+$: 175.1487, found 175.1488.

(E)-tert-butylphenyl(1-phenylundec-1-en-3-yloxy)silane: To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), tert-butylphenyl(undec-1-en-3-yloxy)silane (1 mmol, 409 mg, 1

equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar. The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K_2CO_3 (1 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (125 mL SiO_2) with 3% ethyl acetate/hexanes as eluent to yield (*E*)-*tert*-butyldiphenyl(1-phenylundec-1-en-3-yloxy)silane as a pale yellow oil. Run 1 (0.459 g, 0.948 mmol, 95%); run 2 (0.451 g, 0.930 mmol, 93%). **Average yield = 94%.** Heck arylation may also be performed at room temperature for 4h: run 1 (0.478 g, 0.989 mmol, 99%), run 2 (0.479 g, 0.991 mmol, 99%). **Average yield = 99%.** By crude ^1H NMR all selectivities were >20:1 E:Z and >20:1 internal olefin : terminal olefin. ^1H NMR (400 MHz, CDCl_3) δ 7.70 (dd, J = 13.6, 7.2 Hz, 4H), 7.43-7.21 (m, 11H), 6.21 (d, J = 16 Hz, 1H), 6.13 (dd, J = 15.6, 6.4 Hz, 1H), 4.29 (ap q, J = 5.6 Hz, 1H), 1.65-1.47 (m, 2H), 1.30-1.18 (m, 12H), 1.09 (s, 9H), 0.88 (t, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 137.4, 136.2, 136.1, 135.7, 134.7, 134.5, 133.0, 129.8, 129.7, 129.6, 128.6, 127.7, 127.6, 127.4, 126.6, 74.8, 38.2, 32.1, 29.7 (app d), 29.5, 27.3, 24.9, 22.9, 19.6, 14.4. IR (neat, cm^{-1}) 2957.98, 2958.12, 2927.73, 2856.35. HRMS (CI) m/z calculated for $\text{C}_{34}\text{H}_{45}\text{OSi}$ $[\text{M}+\text{H}]^+$: 485.3240, found 485.3238.

Entry 1a:

Run 1 (287.1 mg, 0.997 mmol, 100%, 99:1 internal:terminal); run 2 (283.4 mg, 0.984 mmol, 98%, 97:3 internal:terminal); run 3 (272.6 mg, 0.947 mmol, 95%, 98:2 internal:terminal). **Average yield = 99%, 98:2.**

Entry 2:

Run 1 (104.3 mg, 0.362 mmol, 36%); run 2 (107.3 mg, 0.372 mmol, 37%). **Average yield = 37%.**

Entry 3:

Run 1 (20.2 mg, 0.070 mmol, 7%); run 2 (25.0 mg, 0.0868 mmol, 9%). **Average yield = 8%.**

Entry 4:

Run 1 (193.5 mg, 0.672 mmol, 67%); run 2 (202.3 mg, 0.702 mmol, 70%); run 3 (219.6 mg, 0.762 mmol, 76%). **Average yield = 71%.** Note: Variation was observed for this entry based on catalyst batch. Batches containing move phenylvinyl sulfoxide (and presumably therefore acetic acid) show heightened reactivity toward the arylation without acetic acid added.

Entry 5:

Run 1 (265.2 mg, 0.921 mmol, 92%); run 2 (272.6 mg, 0.947 mmol, 95%). **Average yield = 94%.**

Footnote 13:

To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 1-undecene (1 mmol, 98.2 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar. The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH_4Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K_2CO_3 (1 x 75 mL). The organic layers were combined and dried over MgSO_4 . After concentration, the crude product was purified via silica gel chromatography (125 mL SiO_2) with 100% hexanes as eluent to yield (*E*)-1-(3-methylhex-1-enyl)benzene as a clear oil (0.137 g, 0.603 mmol, 60%). **Yield = 60%.** By crude ^1H NMR the ratio of products was determined to be 4:1:1. Heck arylation may also be preformed at room temperature for 4h (0.140 g, 0.612 mmol, 61%). **Yield = 61%.** By crude ^1H NMR the ratio of internal olefin : allylic benzene : terminal olefin was found to be 4:3:1. By crude ^1H NMR E:Z selectivities were found to be >20:1 in all cases. ^1H NMR (400 MHz, CDCl_3) δ Internal olefin: 6.43 (d, J = 15.6 Hz, 1H), 6.28 (dt, J = 16, 6.8 Hz, 1H), 2.25 (q, J = 7.6 Hz, 2H). Allylic benzene: 5.66-5.52 (m, 2H), 3.38 (d, J = 6 Hz, 2H), 2.07 (q, J = 6.4 Hz, 2H). Terminal olefin: 5.31 (s, 1H), 5.10 (s, 1H), 2.55 (t, J = 6.8 Hz, 2H).

(E)-1-(3,3-dimethylbut-1-enyl)benzene: To a 40 mL borosilicate vial was added the following: catalyst **1** (0.1 mmol, 50.3 mg, 10 mol%), benzoquinone (2 mmol, 216.2 mg, 2 equiv.), 3,3-dimethyl-1-butene (1 mmol, 84.2 mg, 1 equiv.) in dioxane (3mL), acetic acid (4 mmol, 240 mg, 4 equiv.), phenylboronic acid (1.5 mmol, 183 mg, 1.5 equiv.), and a stir bar. The mixture was heated to 45°C for 4 hours. The mixture was diluted with 40 mL of sat. aq. NH₄Cl and extracted with hexanes (2 x 75 mL) and rinsed with 5% K₂CO₃ (1 x 75 mL). The organic layers were combined and dried over MgSO₄. After concentration, the crude product was purified via silica gel chromatography (125 mL SiO₂) with 100% hexanes as eluent to yield (E)-1-(3,3-dimethylbut-1-enyl)benzene as a clear oil. Run 1 (0.158 g, 0.999 mmol, quantitative); run 2 (0.158 g, 1.000 mmol, quantitative). **Average yield = quantitative.** By crude ¹H NMR all selectivities were >20:1 E:Z and >20:1 internal olefin : terminal olefin. ¹H NMR (400 MHz, CDCl₃) δ 7.39 (d, *J* = 7.6 Hz, 2H), 7.32 (t, *J* = 7.6 Hz, 2H), 7.22 (t, *J* = 7.2 Hz, 1H), 6.34 (d, *J* = 16 Hz, 1H), 6.28 (d, *J* = 16 Hz, 1H), 1.16 (s, 1H). ¹³C NMR (100 MHz, CDCl₃) δ 142.0, 138.2, 128.7, 126.9, 126.2, 124.7, 33.6, 29.8. IR (neat, cm⁻¹) 3025.38, 2960.48, 2903.23, 2866.92. HRMS (CI) *m/z* calculated for C₁₂H₁₇ [M+H]⁺: 161.1330, found 161.1330.

References for Supporting Information:

1. Still, W.C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

Reference 9 in text:

9. Edmondson, S.D.; Mastracchio, A.; Duffy, J.L.; Eiermann, G.J.; He, H.; Ita, I.; Leiting, B.; Leone, J.F.; Lyons, K.A.; Makarewicz, A.M.; Patel, R.A.; Petrov, A.; Wu, J.K.; Thornberry, N.A.; Weber, A.E. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 3048.