Supporting Information

Sequential Ruthenium-Catalyzed Hydroamination and Rhenium-Catalyzed C-H Bond Activation Leading to Indene Derivatives

Yoichiro Kuninobu,* Yuta Nishina, and Kazuhiko Takai*

*Division of Chemistry and Biochemistry, Graduate School of Natural Science and Technology,
Okayama University, Tsushima, Okayama 700-8530, Japan*

General. All reactions were carried out in dry toluene under an argon atmosphere. Toluene was purchased from Wako Pure Chemical Industries and was dried and degassed before use. $\text{Ru}_3(\text{CO})_{12}$ and NH_4PF_6 were purchased from Aldrich Co., and were used without any purification. $[\text{ReBr}(\text{CO})_3(\text{thf})]_2$ was prepared by heating a THF solution of $\text{ReBr}(\text{CO})_5$ at reflux temperature for 16 h.^[1] The resulting solution was concentrated in vacuo and then recrystallized from THF/hexane to give $[\text{ReBr}(\text{CO})_3(\text{thf})]_2$ as a white solid in 75% yield.^[2] Acetylenes, α,β -unsaturated carbonyl compounds, and anilines were purchased from Wako Pure Chemical Industries, Tokyo Kasei Kogyo Co., Nacalai Tesque, or Aldrich Co., and were used after distillation.

^1H (400 MHz) and ^{13}C (100 MHz) NMR spectra were recorded using a JEOL JNM-LA400 spectrometer. Proton chemical shifts are reported relative to tetramethylsilane (CDCl_3) at δ 0.00 ppm or the residual solvent peak (CDCl_3 at δ 7.26 ppm). Carbon chemical shifts are reported relative to CDCl_3 at δ 77.00 ppm. IR spectra were recorded on a Nicolet Protégé 460 spectrometer.

Indene derivatives **3a-g** are already known. The structures of the reaction products were determined by comparing of the spectrum data of the products with the spectrum data of **3a**,^[3] **3b**,^[4] and **3c-g**.^[5]

General Procedure. A mixture of an aromatic acetylene (0.500 mmol), aniline (0.500 mmol), $\text{Ru}_3(\text{CO})_{12}$ (0.500 μmol), NH_4PF_6 (1.50 μmol) and toluene (0.2 mL) was stirred at 100 °C for 3 h. After cooling, an α,β -unsaturated carbonyl compound (0.750 mmol) and $[\text{ReBr}(\text{CO})_3(\text{thf})]_2$ (0.0150 mmol) were added, and then the reaction mixture was stirred at 180 °C for 24 h. The crude product was purified by column chromatography on silica gel to give the desired indene derivatives.

3-Methyl-1H-indene-2-carboxylic acid ethyl ester (3h). Compounds: ethynylbenzene (51.1 mg), ethyl vinyl ketone (63.0 mg). Yield: 45.6 mg (49%)

yield). ^1H NMR (400 MHz, CDCl_3) δ 1.38 (t, $J = 7.2$ Hz, 3H), 2.56 (m, 3H), 3.70 (m, 2H), 4.32 (q, $J = 7.2$, 2H), 7.57 (d, $J = 8.1$, 1H), 7.63 (d, $J = 8.1$, 1H), 7.72 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 7.87 (1C), 12.90 (1C), 35.21 (1C), 38.72 (1C), 121.46 (1C), 123.92 (1C), 126.70 (1C), 127.86 (1C), 137.04 (1C), 142.93 (1C), 145.40 (1C), 149.64 (1C), 199.69 (1C); IR (nujol, ν / cm^{-1}) 2923 (s), 2853 (s), 1977 (m), 1734 (w), 1700 (w), 1653 (s), 1636 (w), 1559 (m), 1457 (s), 1418 (m), 1375 (m), 1227 (m), 1207 (m), 1170 (w), 1153 (w), 1118 (w), 758 (w), 720 (w), 695 (w); Anal Calcd for $\text{C}_{13}\text{H}_{14}\text{O}$: C, 83.83; H, 7.58; O, 8.59. Found: C, 83.83; H, 7.58.

Ethyl 3,7-dimethyl-1H-indene-2-carboxylate (4e). Compounds:

1-ethynyl-3-methylbenzene (58.1 mg), ethyl acrylate (75.1 mg). ^1H NMR (400 MHz, CDCl_3) δ 1.37 (t, $J = 7.2$ Hz, 3H), 2.36 (s, 1H), 2.52 (m, 3H), 3.51 (m, 2H), 7.14 (m, 1H), 7.27 (m, 1H), 7.32 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 12.38 (1C), 14.07 (1C), 22.59 (1C), 37.63 (1C), 59.84 (1C), 118.68 (1C), 126.86 (1C), 128.64 (1C), 129.41 (1C), 133.17 (1C), 142.17 (1C), 144.90 (1C), 151.68 (1C), 165.98 (1C).

Ethyl 3-methyl-1H-cyclopenta[a]naphthalene-2-carboxylate (4f).

Compounds: 2-ethynylnaphthalene (76.1 mg), ethyl acrylate (75.1 mg). ^1H NMR (400 MHz, CDCl_3) δ 1.39 (t, $J = 7.2$ Hz, 3H), 2.61 (m, 3H), 3.91 (m, 2H), 7.43 (m, 1H), 7.49 (m, 1H), 7.56 (d, $J = 8.7$ Hz, 1H), 7.83 (m, 2H), 7.96 (d, $J = 8.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 12.74 (1C), 15.23 (1C), 37.68 (1C), 59.88 (1C), 119.19 (1C), 124.03 (1C), 126.50 (1C), 127.54 (1C), 128.74 (1C), 129.14 (1C), 129.63 (1C), 132.94 (1C), 141.15 (1C), 142.41 (1C), 152.22 (1C), 165.93 (1C).

- [1] Du, S.; Kautz, J. A.; McGrath, T. D.; Stone, F. G. A. *Organometallics* **2003**, 22, 2842.
- [2] Vitali, D.; Calderazzo, F. *Gazz. Chim. Ital.* **1972**, 102, 587.
- [3] Jalander, L. *Synth. Commun.* **1988**, 18, 343.
- [4] Ghoshal, P. N.; Pathak, B. *J. Ind. Chem. Soc.* **1976**, 53, 1126.
- [5] Kuninobu, Y.; Nishina, Y.; Shouho, M.; Takai, K. *Angew. Chem., Int. Ed.* **2006**, 45, 2766.

3a

3b

3c

3c

3f

3f

+

+

3h

3h