SUPPORTING INFORMATION

Silver Catalyzed Asymmetric Allylation of Simple Ketones

Manabu Wadamoto and Hisashi Yamamoto*

General Procedures.

All non aqueous reactions were carried out in oven- or flame-dried glassware under an atmosphere of dry argon unless otherwise noted. Except as otherwise indicated, all reactions were magnetically stirred and monitored by analytical thin-layer chromatography using Whatman pre-coated silica gel flexible plates (0.25 mm) with F₂₅₄ indicator. Visualization was accomplished by UV light (256 nm), potassium permanganate. Flash column chromatography was performed using silica gel 60 (mesh 230-400) supplied by E. Merck. Yields refer to chromatographically and spectrographically pure compounds, unless otherwise noted. Infrared spectra were recorded as thin films on sodium chloride plates using a Nicolet 20 SXB FTIR. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker Avance 400 (400 MHz ¹H, 100 MHz ¹³C), and a Bruker Avance 500 (500 MHz ¹H, 125 MHz ¹³C). Chemical shift values (δ) are reported in ppm relative to Me₄Si (δ 0.0 ppm). ³¹P NMR chemical shifts are reported as downfield from H₃PO₄ (δ 0.0 ppm) by pre-referencing of the spectrometer. The proton spectra are reported as follows δ (multiplicity, number of protons, coupling constant *J*). Multiplicities are indicated by s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), h (heptet), m (multiplet) and b (broad). Optical rotations were measured on a JASCO DIP-140 digital polarimeter. GC analysis was done with a Shimadzu model 17A instrument with a flame-ionization detector and a capillary column of chiral column Chiraldex B-DP (20 m x 0.25 mm) using nitrogen as a carrier gas. Analytical high-performance liquid chromatography (HPLC) was done with a Varian ProSter instrument using a chiral column (4.6 mm x 25 cm, Daicel CHIRALCEL OD-H, AD-H or OJ-H). Preparative high performance liquid chromatography was done with a Varian ProSter instrument using a Varian Dynamax preparative HPLC column (21.4 mm x 25 cm). Commercial grade reagents and solvents were used without further purification except as indicated below. Tetrahydrofuran (in Pure-Pac TM stainless steel containers, from Aldrich) were purified

by both A2 alumina and Q5 reactant using a MBRAUN MBSP series solvent purification system. (*R*)-BINAP (guaranteed reagent) was used as purchased from Aldrich. (*R*)-SEGPPOS was donated by the Takasago International Corporation. (*R*)-MeO-BIPHEP (guaranteed reagent) was used as purchased from Fluka. (*R*)-DIFLUORPHOS•CHCl₃ (guaranteed reagent) was used as purchased from STREM. Allyltrimethoxysilane (Aldrich) and all ketones were purified by distillation before use. 2-Chlorocyclohexenone,¹ 2-iodocyclohexenone,² and 2-iodocyclopentenone² were prepared according to literature procedures. (*E*)-crotyltrimethoxysilane (*E/Z* < 1/99) was prepared (*E*)-crotyltrichlorosilane with MeOH in the presence of triethylamine in CH₂Cl₂ and distilled before use. The (*E*)-crotyltrichlorosilane was synthesized from trichlorosilane, crotylchloride, triethylamine, and CuCl according to the usual procedure.³ (*Z*)-Crotyltrimethoxysilane (*E/Z* < 1/99), cyclohex-2-enyltrimethoxysilane, (*Z*)-cyclooct-2-enyltrimethoxysilane and 3-trimethoxysilylcyclooctane was prepared by reaction of (*Z*)-crotyltrichlorosilane 3-trichlorosilylcyclohexene and 3-trichlorosilylcyclooctene with MeOH in the presence of triethylamine and purified by distillation before use. These trichlorosilanes were synthesized from trichlorosilane, corresponding diene, and Pd(PPh₃)₄ in a sealed tube according to the literature procedure.⁴

Starting Material

(±)-Cyclohex-2-enyltrimethoxysilane

¹H NMR (400 MHz, CD₃Cl) δ 1.52 (m, 1H), 1.74 (m, 4 H), 1.99 (m, 2 H), 3.58 (s, 1 H), 5.68 (m, 2 H); ¹³C NMR (100 MHz, CD₃Cl) δ 21.9, 22.1, 22.7, 24.8, 50.9, 125.6, 126.6; Exact Mass Calcd for C₉H₂₀O₃Si (M+H)⁺: 203.1 Found: 203.1

(±)- (*Z*)-cyclooct-2-enyltrimethoxysilane

^1H NMR (400 MHz, CD_3Cl) δ 1.30 (m, 2H), 1.56 (m, 3 H), 1.66 (m, 3 H), 2.08 (m, 2 H), 2.55 (m, 1 H), 5.52 (dt, $J = 1.4$ Hz, $J = 10.2$ Hz, 1 H), 5.75 (m, 1 H); ^{13}C NMR (100 MHz, CD_3Cl) δ 20.6, 26.1, 26.5, 29.2, 29.4, 50.9, 128.2, 131.4; Exact Mass Calcd for $\text{C}_{11}\text{H}_{22}\text{O}_3\text{Si}$ ($\text{M}+\text{H}$) $^+$: 231.1 Found: 231.1

General Procedure for Silver Catalyzed Asymmetric Allylation.

Into a Schlenk tube were added AgF (3 mg, 0.025 mmol), (*R*)-DIFLUORPHOS (17 mg, 0.025 mmol) and MeOH (1.0 ml). The mixture was stirred for 10 min and dried under vacuum (5 mmHg) for 1 h. After cooling of the resulting white solid to -78°C , MeOH (20 μl , 0.5 mmol) and THF (1.5 ml) were added and stirred for 10 min. To this clear solution, allyltrimethoxysilane (1.0 mmol) and ketone (0.5 mmol) were added successively and stirred at -78°C for 12 h. The reaction mixture was passed through a short silica gel column quickly to remove the catalyst with elution by ether. After evaporation of the solvent, the crude mixture was purified by flash column chromatography.

(2*R*)-(+)-2-Phenyl-4-pentene-2-ol (3aa)⁵

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a white crystal (96% yield, 82% ee): $[\alpha]_{\text{D}}^{28} = +66.0$ ($c = 0.27$, CHCl_3 , 82 %ee); ^1H NMR (400 MHz, CD_3Cl) δ 1.55 (s, 3 H), 2.05 (s, 1 H), 2.50 (dd, $J = 8.32$ Hz, 13.6 Hz, 1 H), 2.69 (dd, $J = 8.40$ Hz, 13.7 Hz, 1 H), 5.14 (m, 2 H), 5.62 (m, 1 H), 7.24 (m, 1 H), 7.35 (m, 2 H), 7.44 (m, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel OJ-H column (98:2 hexane:2-propanol), 0.8 mL/min; major enantiomer $t_{\text{r}} = 13.6$ min, minor enantiomer $t_{\text{r}} = 17.3$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁵

(+)-2-(3-(Trifluoromethyl)phenyl)pent-4-en-2-ol (3ba)⁵

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (92% yield, 90% ee): $[\alpha]_D^{31} = +34.3$ ($c = 0.26$, CHCl_3 , 90 %ee); ^1H NMR (400 MHz, CD_3Cl) δ 1.56 (s, 3 H), 2.13 (bs, 1 H), 2.52 (dd, $J = 8.0$ Hz, 16.0 Hz, 1 H), 2.68 (dd, $J = 8.0$ Hz, 16.0 Hz, 1 H), 5.15 (m, 2 H), 5.60 (m, 1 H), 7.47 (m, 2 H), 7.61 (m, 1 H), 7.72 (m, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel AD-H column (99:1 hexane:2-propanol), 0.5 mL/min; major enantiomer $t_r = 20.4$ min, minor enantiomer $t_r = 21.5$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁵

(R)-(+)-2-(4-Methoxyphenyl)pent-4-en-2-ol (3ca)⁵

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (63% yield, 79% ee): $[\alpha]_D^{31} = +43.1$ ($c = 0.57$, CHCl_3 , 79 %ee); ^1H NMR (500 MHz, CD_3Cl) δ 1.50 (s, 3 H), 2.14 (bs, 1 H), 2.46 (dd, $J = 8.2$ Hz, 13.7 Hz 1 H), 2.46 (dd, $J = 6.5$ Hz, 13.7 Hz 1 H), 3.77 (s, 3 H), 5.09 (m, 2 H), 5.59 (m, 1 H), 6.85 (m, 2 H), 7.33 (m, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (99:1 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 15.7$ min, minor enantiomer $t_r = 17.2$ min.⁵

(+)-2-o-Tolylpent-4-en-2-ol (3da)⁶

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (62% yield, 92% ee): $[\alpha]_D^{31} = +43.1$ (c

= 0.24, CHCl₃, 92 %ee); ¹H NMR (400 MHz, CD₃Cl) δ 1.62 (s, 3 H), 2.04 (bs, 1 H), 2.37 (m, 4 H), 2.88 (ddt, *J* = 5.3 Hz, 12.7 Hz, 1.2 Hz, 1 H), 5.15 (m, 2 H), 5.66 (m, 1 H), 7.16 (m, 3 H), 7.44 (m, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (99.7:0.3 hexane:2-propanol), 1.0mL/min; major enantiomer *t_r* = 7.3 min, minor enantiomer *t_r* = 7.9 min. Other spectral data (TLC, IR and ¹³C NMR) indicated good agreement with reported data.⁶

(+)-2-(Naphthalen-1-yl)pent-4-en-2-ol (3ea)⁷

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (98% yield, 95% ee): [α]_D³¹ = +20.7 (*c* = 0.18, CHCl₃, 95 %ee); ¹H NMR (400 MHz, CD₃Cl) δ 1.83 (s, 3 H), 2.29 (bs, 1 H), 2.83 (dd, *J* = 7.9 Hz, 14.7 Hz, 1 H), 3.13 (dd, *J* = 6.7 Hz, 13.9 Hz, 1 H), 5.13 (m, 2 H), 5.67 (m, 1 H), 7.41 (q, *J* = 8.4 Hz, 1 H), 7.50 (m, 2 H), 7.59 (dd, *J* = 1.2 Hz, 7.3 Hz, 1 H), 7.77 (d, *J* = 8.2 Hz, 1 H), 7.87 (d, *J* = 8.6 Hz, 1 H), 8.74 (dd, *J* = 1.0 Hz, *J* = 8.3 Hz, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (98:2 hexane:2-propanol), 0.75 mL/min; minor enantiomer *t_r* = 19.9 min, major enantiomer *t_r* = 23.6 min. Other spectral data (TLC, IR and ¹³C NMR) indicated good agreement with reported data.⁷

(S)-(+)-3-Methyl-1-phenylhex-5-en-3-ol (3fa)⁷

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (36% yield, 63% ee): [α]_D²⁹ = +6.8 (*c* = 0.12, CHCl₃, 63 %ee); ¹H NMR (400 MHz, CD₃Cl) δ 1.26 (s, 3 H), 1.43 (bs, 1 H), 1.78 (m, 2 H), 2.30 (d, *J* = 7.2 Hz, 2 H), 2.70 (m, 2 H), 5.16 (m, 2 H), 5.89 (m, 1 H), 7.18 (m, 3 H), 7.28 (m, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H

column (95:5 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 7.6$ min, minor enantiomer $t_r = 8.2$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁷

(+)-1-Allyl-2,3-dihydro-1*H*-inden-1-ol (3ga)⁶

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (89% yield, 91% ee): $[\alpha]_D^{31} = +6.2$ ($c = 0.25$, CHCl_3 , 91 %ee); ^1H NMR (400 MHz, CD_3Cl) δ 1.94 (bs, 1 H), 2.19 (m, 1 H), 2.35 (m, 1 H), 2.45 (m, 1 H), 2.63 (m, 1 H), 2.81 (m, 1 H), 3.05 (m, 1 H), 5.17 (m, 2 H), 5.90 (m, 1 H), 7.27 (m, 3 H), 7.35 (m, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (99:1 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 14.6$ min, minor enantiomer $t_r = 19.9$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁶

(2R)-(+)-1-Allyl-1,2,3,4-tetrahydronaphthalene-1-ol (3ha)⁵

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (63% yield, 93% ee): $[\alpha]_D^{31} = +38.3$ ($c = 0.24$, CHCl_3 , 93 %ee); ^1H NMR (400 MHz, CD_3Cl) δ 1.88 (m, 4 H), 2.05 (m, 1 H), 2.61 (m, 1 H), 2.78 (m, 2 H), 5.14 (m, 2 H), 5.82 (m, 1 H), 7.09 (m, 1 H), 7.18 (m, 2 H), 7.56 (m, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (98:2 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 12.6$ min, minor enantiomer $t_r = 16.4$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁵

(+)-5-Allyl-6,7,8,9-tetrahydro-5*H*-benzo[7]annulen-5-ol (3ia)

The crude mixture was purified by flash column chromatography with elution by hexane: ether (10:1) provided as a colorless liquid (78% yield, 92% ee): $[\alpha]_D^{31} = +35.5$ ($c = 0.25$, CHCl_3 , 92 %ee); FTIR (neat) ν_{max} 3459, 3070, 2928, 2855, 1636, 1446, 1022, 995, 916, 759, 747 cm^{-1} ; ^1H NMR (400 MHz, CD_3Cl) δ 1.51 (m, 1 H), 1.88 (m, 4 H), 2.02 (m, 1 H), 2.16 (bs, 1 H), 2.64 (dd, $J = 7.9$ Hz, 14.0 Hz, 1 H), 2.74 (dd, $J = 6.7$ Hz, $J = 14.3$ Hz, 1 H), 2.86 (m, 1 H), 2.99 (m, 1 H) 5.17 (m, 2 H), 5.70 (m, 1 H), 7.09 (d, $J = 7.4$ Hz, 1 H), 7.14 (dt, $J = 1.4$ Hz, $J = 7.3$ Hz, 1 H), 7.24 (dt, $J = 1.5$ Hz, $J = 7.4$ Hz, 1 H) 7.66 (dd, $J = 1.3$ Hz, $J = 7.9$ Hz, 1 H); ^{13}C NMR (100 MHz, CD_3Cl) δ 19.7, 29.7, 47.0, 71.9, 118.7, 126.2, 126.3, 127.2, 128.9, 134.0, 136.8 MS (CI) Exact Mass Calcd for $\text{C}_{14}\text{H}_{18}\text{O}$ ($\text{M}+\text{H}^+$): 185.1. Found: 185.1. Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (98:2 hexane:2-propanol), 0.8 mL/min; minor enantiomer $t_r = 13.6$ min, major enantiomer $t_r = 17.3$ min.

(-)-4-Allylchroman-4-ol (3ja)⁶

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (74% yield, 92% ee); TLC R_f 0.7 (EtOAc/Hexanes, 1:3): $[\alpha]_D^{28} = -15.8$ ($c = 0.71$, CHCl_3 , 92 %ee); ^1H NMR (500 MHz, CD_3Cl) δ 1.88 (dt, $J = 4.5$ Hz, 13.5 Hz, 1 H), 2.14 (m, 2 H), 2.61 (dd, $J = 6.0$ Hz, 14.0 Hz, 1 H), 2.72 (dd, $J = 7.5$ Hz, 14.0 Hz, 1 H), 4.20 (m, 2 H), 5.24 (m, 2 H), 5.75 (m, 2 H), 6.81 (dd, $J = 1.0$ Hz, 8.5 Hz, 1 H), 6.92 (dt, $J = 1.0$ Hz, 7.5 Hz, 1 H), 7.16 (dd, $J = 1.0$ Hz, 7.5 Hz, 1 H), 7.43 (dd, $J = 1.0$ Hz, 8.5 Hz, 1 H). Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (98:2 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 12.5$ min, minor enantiomer $t_r = 17.0$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁶

(-)-1-Allyl-2-chlorocyclohex-2-enol (3ka)

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (74% yield, 96% ee): $[\alpha]_D^{31} = -15.3$ ($c = 0.26$, CHCl_3 , 96 %ee); FTIR (CD_3Cl) ν_{max} 3446, 2940, 2868, 1733, 1640, 1437, 9992, 916, 791 cm^{-1} ; ^1H NMR (400 MHz, CD_3Cl) δ 1.66 (m, 2 H), 1.85 (m, 2 H), 2.05 (m, 2 H), 2.16 (bs, 1 H), 2.49 (m, 2 H), 5.12 (m, 2 H), 5.78 (m, 1 H), 5.96 (dd, $J = 3.5$ Hz, 5.0 Hz, 1 H); ^{13}C NMR (100 MHz, CD_3Cl) δ 18.6, 26.7, 35.3, 44.3, 72.4, 118.9, 128.8, 133.1, 136.7 MS (CI) Exact Mass Calcd for $\text{C}_9\text{H}_{13}\text{ClO}$ (M-OH^+): 155.1. Found: 155.1. Enantiometric excess was determined by GC with a Chiraldex β -DP column (20m x0.25mm, 90°C, 100kPa); major enantiomer $t_r = 32.0$ min, minor enantiomer $t_r = 35.1$ min.

(-)-1-Allyl-2-iodocyclohex-2-enol (3la)

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (97% yield, 96% ee): $[\alpha]_D^{29} = -15.1$ ($c = 0.50$, CHCl_3 , 96 %ee); FTIR (CD_3Cl) ν_{max} 2936, 1700, 1653, 1558, 1540, 1540, 1506, 1456, 1081, 965 cm^{-1} ; ^1H NMR (500 MHz, CD_3Cl) δ 1.70 (m, 2 H), 1.87 (m, 1 H), 1.95 (bs, 1 H), 2.00 (m, 3 H), 2.43 (m, 2 H), 5.15 (m, 2 H), 5.78 (m, 1 H), 6.55 (dd, $J = 1.0$ Hz, 4.5 Hz, 1 H); ^{13}C NMR (126 MHz, CD_3Cl) δ 18.8, 29.6, 34.1, 47.1, 72.9, 111.7, 118.9, 132.9, 141.8 MS (CI) Exact Mass Calcd for $\text{C}_9\text{H}_{13}\text{IO}$ (M-OH^+): 247.0. Found: 247.0. Enantiometric excess was determined by GC with a Chiraldex β -DP column (20m x0.25mm, 120°C, 80kPa); major enantiomer $t_r = 12.6$ min, minor enantiomer $t_r = 13.7$ min.

(-)-1-Allyl-2-iodocyclopent-2-enol (3ma)²

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (92% yield, 94% ee): $[\alpha]_D^{29} = -9.7$ ($c = 0.33$, CHCl_3 , 94 %ee); FTIR (neat) ν_{max} 3392, 3074, 2924, 1652, 1558, 1436, 1315, 1065, 917 cm^{-1} ; ^1H NMR (500 MHz, CD_3Cl) δ 1.94 (m, 2 H), 2.32 (m, 6 H), 5.16 (m, 2 H), 5.74 (m, 1 H), 6.25 (m, 1 H). Enantiometric excess was determined by GC with a Chiraldex β -DP column (20mx0.25mm, 120°C, 80kPa); major enantiomer $t_r = 28.8$ min, minor enantiomer $t_r = 29.7$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.²

(+)-(E)-1-Allyl-2-benzylidenecyclohexanol (3na)⁶

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (96% yield, 84% ee): $[\alpha]_D^{30} = +69.0$ ($c = 0.56$, CHCl_3 , 84 %ee); δ 1.30 (m, 1 H), 1.52 (s, 2 H), 1.57 (m, 1 H), 1.72 (m, 2 H), 1.82 (bs, 1 H), 1.92 (m, 2 H), 2.31 (dd, $J = 8.5$ Hz, 14.0 Hz, 1 H), 2.69 (dd, $J = 6.2$ Hz, 14.0 Hz, 1 H), 2.83 (m, 1 H), 5.14 (m, 2 H), 5.79 (m, 1 H), 6.58 (m, 1 H), 7.17 (m, 3 H), 7.28 (m, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel AD-H column (98:2 hexane:2-propanol), 1.0 mL/min; minor enantiomer $t_r = 11.7$ min, major enantiomer $t_r = 15.7$ min. Other spectral data (TLC, IR and ^{13}C NMR) indicated good agreement with reported data.⁶

(+)-2-Cyclohexenylpent-4-en-2-ol (3oa)⁶

The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid (56% yield, 95% ee): $[\alpha]_D^{30} = +32.5$ ($c = 0.59$, CHCl_3 , 95 %ee); δ 1.25 (s, 3 H), 1.55 (m, 2 H), 1.61 (m, 2 H), 1.98 (m, 2 H), 2.04 (m, 2 H), 2.22 (m, 1 H), 2.37 (bs, 1 H), 2.40 (m, 1 H), 5.08 (m, 2 H), 5.70 (m, 2 H). Enantiometric excess was determined by GC with a Chiraldex β -DP column (20mx 0.25mm, 110°C, 100kPa); major enantiomer $t_r = 11.3$ min, minor enantiomer $t_r = 11.7$ min.⁶

(-)-(2*R*,3*S*)-3-Methyl-2-phenylpent-4-en-2-ol (3ab)⁷

The α/γ and *anti/syn* ratios were determined to be <1/99 and 10/90, respectively, by ^1H NMR analysis. The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid. *Syn* isomer was isolated by preparative HPLC (97:3 hexane: ethylacetate). (92% yield, 93% ee): $[\alpha]_D^{31} = -7.32$ ($c = 0.26$, CHCl_3 , 93 %ee); ^1H NMR (500 MHz, CD_3Cl) δ 0.78 (d, $J = 5.0$ Hz, 3 H), 1.44 (s, 3 H), 1.79 (bs, 1 H), 2.46 (t, $J = 10.0$ Hz, 1 H), 5.02 (m, 2 H), 5.71 (m, 1 H), 7.15 (m, 1 H), 7.25 (m, 2 H), 7.32 (m, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel AD-H column (98:2 hexane:2-propanol), 0.8 mL/min; major enantiomer $t_r = 13.0$ min, minor enantiomer $t_r = 13.7$ min.⁷

(-)-(R*)-1-((S*)-(-)-Cyclohex-2-enyl)-1-phenylethanol (3ac)⁹

The *anti/syn* ratios were determined to be 4/96, respectively, by ^1H NMR analysis. The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid. *Syn* isomer was isolated by preparative HPLC (97:3 hexane: ethylacetate). (72% yield, 98% ee): $[\alpha]_D^{28} = -50.9$ ($c = 0.34$, CHCl_3 , 98 %ee); ^1H NMR (400 MHz, CD_3Cl) δ 1.17 (m, 1 H), 1.28 (m, 2 H), 1.41 (m, 2 H), 1.52 (s, 3 H), 1.65 (m, 2H) 1.76 (s, 1 H), 2.09 (m, 1H), 2.26 (m, 1 H), 2.92 (m, 1H), 5.61(dt, J

= 1.1 Hz, 10.0 Hz, 1 H), 5.79 (m, 1 H), 7.24 (t, $J = 7.3$ Hz, 1 H), 7.35 (t, $J = 7.9$ Hz, 2 H), 7.45 (d, $J = 7.9$ Hz, 2 H). Enantiometric excess was determined by HPLC with a Chiralcel AD-H column (99:1 hexane:2-propanol), 0.5 mL/min; minor enantiomer $t_r = 18.0$ min, major enantiomer $t_r = 22.2$ min.⁹

(-)-(R*)-1-((S*,Z)-(-)-Cyclooct-2-enyl)-1-phenylethanol (3ad)

The *anti/syn* ratios were determined to be 1/>99, respectively, by ¹H NMR analysis. The crude mixture was purified by flash column chromatography with elution by (10:1 hexane: ether) provided as a colorless liquid. *Syn* isomer was isolated by preparative HPLC (97:3 hexane: ethylacetate). (72% yield, 98% ee): $[\alpha]_D^{28} = -61.9$ (c = 0.43, CHCl₃, 98 %ee); FTIR (CD₃Cl) $\nu_{\max}$ 3473, 2924, 2853, 1445, 1373, 1136, 759, 700 cm⁻¹; ¹H NMR (500 MHz, CD₃Cl) δ 1.17 (m, 1 H), 1.28 (m, 3 H), 1.42 (m, 2 H), 1.51 (s, 3 H) 1.64 (m, 2H) 1.76 (bs, 1 H), 2.08 (m, 1H), 2.24 (m, 1 H), 2.92 (m, 1H), 5.60 (dt, $J = 1.0$ Hz, 9.5 Hz, 1 H), 5.77 (m, 1 H), 7.23 (t, $J = 7.3$ Hz, 1 H), 7.35 (t, $J = 7.9$ Hz, 2 H), 7.44 (d, $J = 7.9$ Hz, 2 H); ¹³C NMR (126 MHz, CD₃Cl) δ 22.6, 25.5, 26.8, 27.1, 29.4, 29.6, 31.6, 46.5, 76.3, 125.0, 126.3, 128.0, 129.8, 130.8, 147.8 MS (CI) Exact Mass Calcd for C₁₁H₁₃N₂O (M-H)⁺: 213.1. Found: 213.1. Enantiometric excess was determined by HPLC with a Chiralcel OD-H column (99:1 hexane:2-propanol), 1.0 mL/min; major enantiomer $t_r = 9.9$ min, minor enantiomer $t_r = 11.8$ min.

1. Montalban, A. G.; Wittenberg, L.-O.; McKillop, A.; *Tetrahedron Lett.* **1999**, 40, 5893
2. Negishi, E.; Tan, Z.; Liou, S.; Liao, B. *Tetrahedron* **2000**, 56, 10197.
3. Hosomi, A.; Kohara, S.; Ogata, K.; Yanagi, T.; Tominaga, Y. *J. Org. Chem.* **1990**, 55, 2415.
4. Tsuji, J.; Hara, M.; Ohno, K. *Tetrahedron* **1974**, 30, 2143.
5. Casolari, S.; D'Addario, D.; Tagliavine, E. *Org. Lett.* **1999**, 1, 1061.
6. Kim, J. G.; Waltz, K. M.; Garcia, I. F.; Kwiatkowski, G. D.; Walsh, P. J. *J. Am. Chem. Soc.* **2004**, 126, 12580.
7. Wada, R.; Oisaki, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2004**, 126, 8910.

8. Kamble, R. M.; Singh, V. K. *Tetrahedron Lett.* **2001**, 42, 7525.
9. Yasuda, M.; Hirata, K.; Nishino, M.; Yamamoto, A.; Baba, A. *J. Am. Chem. Soc.* **2002**, 126, 13442.

³¹P-NMR Experiment

AgF (0.025 mmol) and ligand (0.025 mmol) in CD₃OD (750 μl) was stirred at r.t. for 10min. The resulting clear solution was transferred to a NMR sample tube, then cooled to -78 °C. All ³¹P-NMR experiments were done at -78 °C. The coordination number can be assigned by coupling constant of ¹²⁷Ag and ³¹P. The molar ratio of three BINAP-Silver complexes was calculated by integration value in ³¹P NMR.