Supporting Online Material

Solution structure of a double mutant of the carboxy-terminal dimerization domain of HIV-1 capsid protein

Hing C. Wong^{*}, Ronald Shin^{*}, and N. Rama Krishna^{*,†}

^{*}Comprehensive Cancer Center and [†]the Department of Biochemistry & Molecular Genetics, The University of Alabama at Birmingham, Birmingham, AL 35294-2041, USA

Captions for Supplementary Figures:

- Figure 1S: Comparison of HSQC spectra of the double mutant CTD at 0.75 mM (blue) and 3 mM (red).
- Figure 2S: HNCA Panels showing sequential connectivities in the stretch of residues 148-156.
- Figure 3S: HNCACB panels showing connectivities in the stretch of residues 215-222.
- Figure 4S: HNCACB panels showing connectivities for residues 166-172
- Figure 5S: Typical HCCH-TOCSY panel showing side chain assignments for a few residues
- Figure 6S: ¹³C-HSQC-NOESY panels showing inter-helical NOE contacts between Leu190 (helix-2) and Val165 and Phe168 (helix-1).
- Figure 7S: Stereo (cross-eye) view of the W184A/M185A-CTD showing the polar (blue) and non-polar (red) residue side chains.

Figure 1S:

Figure 2S:

HNCA

Figure 3S:

HNCACB

Figure 4S:

HNCACB

Figure 5S:

HCCH-
TOCSY

Figure 6S:

^{13}C -HSQC-NOESY

Figure 7S:

Distribution of polar (blue) and non-polar (red) residue side chains in the double mutant CTD.