Solvent Dependent Diastereoselectivities in Reductions of β -Hydroxyketones by SmI_2

Pramod R. Chopade,^a Todd A. Davis,^a Edamana Prasad,^b Robert A. Flowers, II^{*,b}

^a*Department of Chemistry & Biochemistry, Texas Tech University, Box 41061, Lubbock, TX 79409-1061*

^b*Department of Chemistry, Lehigh University, Bethlehem, PA 18015*

rof2@lehigh.edu

Supporting Information

<u>Contents</u>	<u>Page No.</u>
1. Preparation of SmI_2 in THF	S2
2. Preparation of SmI_2 in CH_3CN	S2
3. Preparation of SmI_2 in DME	S2
4. General Procedure for Aldol Reactions	S2
5. General Procedure for Proline Catalyzed Aldol Reactions	S3
6. General Procedure for Reduction of β -Hydroxyketones using SmI_2 in THF, CH_3CN , and DME	S4
7. General Procedure for Determination of Major Diastereomer	S5
8. Diastereoselection of SmI_2 in THF vs. SmI_2 in CH_3CN	S6
9. Diastereoselection of SmI_2 in THF vs. SmI_2 in DME	S7
10. Diastereoselectivity in the Reduction of 4-Hydroxy-5,5-dimethyl-hexan-2-one utilizing SmI_2 in THF, CH_3CN , and DME	S8
11. ^1H NMR of <i>Syn</i> - and <i>Anti</i> - 5,6-Dimethyl-hexan-2,4-diol	S9
12. Diastereoselectivity in the Reduction of (1R)-Hydroxy-1-phenyl-butan-3-one utilizing SmI_2 in THF, CH_3CN , and DME	S10
13. NMR Data for the Reduction of (1R)-Hydroxy-1-phenyl-butan-3-one	S11
14. Diastereoselectivity in the Reduction of 1-Phenyl-3-hydroxy-butan-1-one utilizing SmI_2 in THF, CH_3CN , and DME	S12

15. ^{13}C NMR Trace of <i>Syn</i> Acetal of 5,5-Dimethyl hexan-2,4-diol	S13
16. ^{13}C NMR Trace of <i>Anti</i> Acetal of 1-Phenyl-butan-1,3-diol	S14
17. ^{13}C NMR Trace of <i>Syn</i> Acetal of 1-Phenyl-butan-1,3-diol	S15

Preparation of SmI_2 in THF:

A typical preparation of SmI_2 in THF utilized 6.4g (25 mmol) of molecular iodine dissolved in 250 mL of dry THF. A mass of 7.0g (42 mmol) of samarium metal was added to the solution. The mixture was allowed to stir overnight under a nitrogen atmosphere during which time the characteristic blue color of SmI_2 appeared indicating the presence of divalent samarium species. The concentration of SmI_2 was determined by iodometric titration to be approximately 0.09M.

Preparation of SmI_2 in CH_3CN :

A typical preparation of SmI_2 in CH_3CN utilized 3.2g (13 mmol) of molecular iodine dissolved in 250 mL of dry acetonitrile. A mass of 5.0g (30 mmol) of samarium metal was added to the solution. The mixture was allowed to stir overnight under a nitrogen atmosphere during which time the characteristic dark green color of SmI_2 in acetonitrile appeared indicating the presence of divalent samarium species. The concentration of SmI_2 was determined by iodometric titration to be approximately 0.05M.

Preparation of SmI_2 in DME:

A typical preparation of SmI_2 in DME utilized 1.3 g (8 mmol) of molecular iodine dissolved in 250 mL of dry DME. A mass of 2.5g (15 mmol) of samarium metal was added to the solution. The mixture was allowed to stir overnight under a nitrogen atmosphere during which time the characteristic blue color of SmI_2 in DME appeared, indicating the presence of divalent samarium species. The concentration of SmI_2 was determined by iodometric titration to be approximately 0.02M.

General Procedure for Aldol Reactions:

LDA (2.0M in heptane/THF) (1.1 equiv.) was dissolved in dry diethyl ether and cooled to -78°C under N_2 . The ketone (1.0 equiv.) was added dropwise to the LDA solution. The reaction mixture was stirred at -78°C until the solution color changed from brown to orange (typically 1 hour). After this color change, the aldehyde (1.1 equiv.) was added dropwise. The reaction mixture was then allowed to stir between 10 minutes to three hours at -78°C under N_2 . The reaction was then placed in an ice bath and 1% HCl was added dropwise. The reaction was allowed to stir for 10 minutes at 0°C . The reaction mixture was poured into water and the organic layer was extracted with ether (3 x 50 mL) and then washed with sodium bicarbonate, brine, and water (3 x 25 mL). The organic layer was dried over MgSO_4 , followed by rotary evaporation of solvent to obtain the crude product. The products were purified utilizing column chromatography or vacuum distillation. Isolated yields ranged from 53-91% for various β -hydroxyketones synthesized.

3-Hydroxy-1-phenyl-butan-1-one:

^1H NMR (500MHz, CDCl_3) δ 1.277 (d, 3H), 2.910-3.202 (m, 2H), 3.294 (s, 1H, exchanges with D_2O), 4.380 (m, 1H), 7.402-7.957 (m, 5H).

^{13}C NMR (125MHz, CDCl_3) δ 22.646, 46.712, 64.243, 128.300, 128.932, 133.801, 136.929, 201.100

Yellow liquid. Yield 68%.

5-Hydroxy-2,2-dimethyl-hexan-3-one:

^1H NMR (500MHz, CDCl_3) δ 1.141 (s, 9H), 1.195 (d, 3H), 2.525 (dd, 1H), 2.691 (dd, 1H), 4.178 (m, 1H).

^{13}C NMR (125MHz, CDCl_3) δ 22.244, 26.229, 44.288, 44.478, 64.007, 217.814.

Light yellow liquid. Yield 74%.

5-Hydroxy-2,2-dimethyl-5-phenyl-pentan-3-one:

^1H NMR (500MHz, CDCl_3) δ 1.144 (s, 9H), 2.88 (dd, 2H), 3.59 (1H, s, exchanges with D_2O), 5.134 (m, 1H), 7.115-7.385 (m, 5H).

^{13}C NMR (125MHz, CDCl_3) δ 26.134, 44.483, 45.453, 70.089, 125.643, 127.562, 128.482, 143.019, 216.910.

Yellow liquid. Yield: 73%.

General Procedure for Proline Catalyzed Aldol Reactions:

(L)-Proline (40 mg, 0.35 mmol) in 10 mL of DMSO/acetone (4:1) was treated with aldehyde (1 mmol). The mixture was stirred at room temperature for a minimum of three days. Aqueous workup with NH_4Cl followed by extraction of the organic layer with ethyl acetate (3 x 50mL) was performed. The organic layer was dried over MgSO_4 . The product was purified by column chromatography utilizing silica gel (3:1 hexanes/ethyl acetate).

(4R)-4-Hydroxy-5,5-dimethyl-hexan-2-one:

^1H NMR (300MHz, CDCl_3) δ 0.912 (s, 9H), 2.207 (s, 3H), 2.458-2.660 (m, 2H), 3.713-3.744 (m, 1H).

^{13}C NMR (125MHz, CDCl_3) δ 25.642, 30.886, 34.175, 45.096, 74.881, 210.474.

Yellow liquid. Yield 59%.

(1R)-Hydroxy-1-phenyl-butan-3-one:

^1H NMR (300MHz, CDCl_3) δ 2.200 (s, 3H), 2.841-2.866 (d, 2H), 5.141-5.183 (dd, 1H), 7.270-7.320 (m, 1H), 7.351 (s, 2H), 7.366 (s, 2H).

Yellow liquid. Yield: 51%.

General Procedure for Reduction of β -Hydroxyketones utilizing SmI_2 in THF, CH_3CN , and DME:

The substrate (β -Hydroxyketone, 1 mmol) was placed in a flame dried round bottom flask and dissolved in 10 mL of anhydrous solvent (THF, DME or acetonitrile depending on the reaction). Methanol (2,5,10,15, or 25 eq.) was placed in the round bottom flask and the reaction was cooled to 0°C . SmI_2 (2.5 equiv.) in THF, DME, or CH_3CN was added dropwise to the reaction mixture. The reaction mixture was allowed to stir at 0°C until the color of the reaction changed from blue (in THF or DME) or dark green (in CH_3CN) to the characteristic yellow color of trivalent samarium. After the reaction was complete, the reaction mixture was diluted with ether and 1% HCl was added to the mixture. The solution was allowed to stir for 15 minutes. The organic layer was extracted with ether (3 x 50 mL). The organic layers were combined washed successively with $\text{Na}_2\text{S}_2\text{O}_3$ (3 x 25 mL), brine (3 x 25 mL), and water (3 x 25 mL). The organic layer was then dried with anhydrous MgSO_4 followed by rotary evaporation of solvent to obtain the products.

Syn: ^1H NMR (300MHz, CDCl_3) δ 1.221-1.242 (d, 3H), 1.802-2.012 (m, 2H), 4.051-4.156 (m, 1H), 5.042-5.120 (m, 1H), 7.242-7.450 (m, 5H).

Anti: ^1H NMR (300 MHz, CDCl_3) δ 1.216 (d, 3H), 1.866 (m, 2H), 2.510 (s, 2H exchanges with D_2O), 4.051 (m, 1H), 5.048 (dd, 1H), 7.332-7.350 (m, 5H).

Syn: ^1H NMR (300MHz, CDCl_3) δ 0.908 (s, 9H), 1.265-1.277 (d, 3H), 1.548 (m, 2H), 3.094 (s, 2H exchanges with D_2O) 3.592-3.610 (dd, 1H), 4.130-4.205 (m, 1H)

Anti: ^1H NMR (300MHz, CDCl_3) δ 0.908 (s, 9H), 1.265-1.277 (d, 3H), 1.548(m, 2H), 2.888 (s, 1H exchanges with D_2O), 3.368 (s, 1H exchanges with D_2O), 3.592-3.610 (dd, 1H), 4.130-4.205 (m, 1H)

^1H NMR (300 MHz, CDCl_3) δ 0.849 (s, 9H), 1.737-1.932 (m, 2H), 3.500 (dd, 1H), 5.045 (dd, 1H), 7.353 (s, 5H).

^1H NMR (300 MHz, CDCl_3) δ 1.216 (d, 3H), 1.866 (m, 2H), 2.510 (s, 2H exchanges with D_2O), 4.051 (m, 1H), 5.048 (dd, 1H), 7.332-7.350 (m, 5H).

^1H NMR (300MHz, CDCl_3) δ 0.908 (s, 9H), 1.265-1.277 (d, 3H), 1.548 (m, 2H), 3.094 (s, 2H exchanges with D_2O) 3.592-3.610 (dd, 1H), 4.130-4.205 (m, 1H)

General Procedure for Determination of Major Diastereomer:

The diastereoselectivity of the major isomers was determined by examining the ^{13}C NMR of the acetal following the elegant procedure by Rychnovsky.¹ The acetals were synthesized by stirring the 1,3 diol in 2,2 dimethoxypropane (2mL), and camphor sulfonic acid (1.1 equiv.). The reaction was poured into H_2O and the organic layer was extracted with ether and dried over MgSO_4 . The crude product was then analyzed by ^{13}C NMR. The *syn*- isomer showed acetal peaks between 19 to 22 and at 31 ppm, while the *anti*- isomer showed two peaks at approximately 25ppm.

1. Rychnovsky, S.; Yang, G.; Powers, J., *J. Org. Chem.*, **1993**, 58, 5251-5255.

Diastereoselection of SmI_2 in THF vs. SmI_2 in CH_3CN in the Reduction of 5-Hydroxy-2,2-dimethyl-hexan-3-one:

GC trace of diol of
5-Hydroxy-2,2-dimethyl-
hexan-3-one after
reduction using
 SmI_2/THF + 25 equiv. of
MeOH
Anti diol -: RT 12.66 min.
Syn diol -: RT 12.77 min

GC trace of diol of
5-Hydroxy-2,2-dimethyl-
hexan-3-one after
reduction using
 $\text{SmI}_2/\text{CH}_3\text{CN}$ + 25 equiv.
of MeOH
Anti diol -: RT 12.6 min.
Syn diol -: RT 12.7 min

Reduction of 5-Hydroxy-2,2-dimethyl-hexan-3-one utilizing SmI_2 in THF and CH_3CN gave opposite diastereoselectivity.

GC Method: Initial temp: 50 °C (hold for 5 min.), rate: 12 °C/min, final temp: 250 °C (hold for 10 min).

Diastereoselection of SmI_2 in THF vs. SmI_2 in DME in the Reduction of 4-Hydroxy-5,5-dimethyl-hexan-2-one:

GC trace of diol of
5-Hydroxy-2,2-dimethyl-
hexan-3-one after
reduction using
 SmI_2 /THF + 25 equiv. of
MeOH
Anti diol:- RT 12.66 min
Syn diol:- RT 12.77 min

GC trace of diol of
5-Hydroxy-2,2-dimethyl-
hexan-3-one after
reduction using
 SmI_2 /DME + 25 equiv.
of MeOH
Syn diol:- RT 12.78 min

Reduction of 4-Hydroxy-5,5-dimethyl-hexan-2-one utilizing SmI_2 in DME gave better diastereoselectivity compared to SmI_2 in THF.

GC Method: Initial temp: 50 °C (hold for 5 min.), rate: 12 °C/min, final temp: 250 °C (hold for 10 min).

Diastereoselectivity in the Reduction of 4-Hydroxy-5,5-dimethyl-hexan-2-one utilizing SmI_2 in THF, CH_3CN , and DME:

Reduction of 4-Hydroxy-5,5-dimethyl-hexan-2-one utilizing SmI_2 in THF and DME gave *syn* diol while SmI_2 in CH_3CN gave *anti* diol as the major isomer.

GC Method: Initial temp: 50 °C (hold for 5 min.), rate: 4 °C/min, final temp: 250 °C (hold for 10 min).

^1H NMR of *Syn*- and *Anti*- 5,6-Dimethyl-hexan-2,4-diol (The $-\text{OH}$ peaks have different δ values):

Diastereoselectivity in the Reduction of (1R)-Hydroxy-1-phenyl-butan-3-one utilizing SmI_2 in THF, CH_3CN , and DME:

GC trace of diol of (1R)-Hydroxy-1-phenyl-butan-3-one after reduction using SmI_2/THF + 25 equiv. of MeOH
anti diol: -RT 18.42 min

GC trace of diol of (1R)-Hydroxy-1-phenyl-butan-3-one after reduction using $\text{SmI}_2/\text{CH}_3\text{CN}$ + 10 equiv. of MeOH
anti diol: -RT 18.42 min

GC trace of diol of (1R)-Hydroxy-1-phenyl-butan-3-one after reduction using SmI_2/DME + 25 equiv. of MeOH
anti diol: -RT 18.41 min

Reduction of (1R)-Hydroxy-1-phenyl-butan-3-one utilizing SmI_2 in THF, CH_3CN , and DME all gave *anti* diol as the major isomer.

GC Method: Initial temp: 50 °C (hold for 5 min.), rate: 12 °C/min till 170 °C (hold for 1 min.), rate: 5 °C/min till 230 °C (hold for 1 min.), rate: 12 °C/min till 250 °C, final temp: 250 °C (hold for 10 min).

NMR Data (crude) for the Reduction of (1R)-Hydroxy-1-phenyl-butan-3-one utilizing SmI_2 in THF, CH_3CN , and DME indicating the anti isomer as the major product in all three solvents:

$^1\text{H-NMR}$ trace of diol of (1R)-Hydroxy-1-phenylbutan-3-one after reduction using $\text{SmI}_2/\text{THF} + 25$ equiv. of MeOH

$^1\text{H-NMR}$ (300MHz):
anti: CDCl_3 , 4.038 (m, 1H), 5.018 (dd, 1H)

$^1\text{H-NMR}$ trace of diol of (1R)-Hydroxy-1-phenylbutan-3-one after reduction using $\text{SmI}_2/\text{CH}_3\text{CN} + 10$ equiv. of MeOH

$^1\text{H-NMR}$ (300MHz):
anti: CDCl_3 , 4.038 (m, 1H), 5.018 (dd, 1H)
syn: CDCl_3 , 4.128 (m, 1H), 4.902 (dd, 1H)

$^1\text{H-NMR}$ trace of diol of (1R)-Hydroxy-1-phenylbutan-3-one after reduction using $\text{SmI}_2/\text{CH}_3\text{CN} + 25$ equiv. of MeOH

$^1\text{H-NMR}$ (300MHz):
anti: CDCl_3 , 4.038 (m, 1H), 5.018 (dd, 1H)
syn: CDCl_3 , 4.128 (m, 1H), 4.902 (dd, 1H)

$^1\text{H-NMR}$ trace of diol of (1R)-Hydroxy-1-phenylbutan-3-one after reduction using $\text{SmI}_2/\text{DME} + 25$ equiv. of MeOH

$^1\text{H-NMR}$ (300MHz):
anti: CDCl_3 , 4.038 (m, 1H), 5.018 (dd, 1H)

Diastereoselectivity decreases going from $\text{SmI}_2/\text{CH}_3\text{CN} + 10$ equiv. of MeOH to $\text{SmI}_2/\text{CH}_3\text{CN} + 25$ equiv. of MeOH.

Diastereoselectivity in the Reduction of 1-phenyl-3-hydroxy-butan-1-one utilizing SmI_2 in THF, CH_3CN , and DME:

GC trace of diol of 3-Hydroxy-1-phenyl-butan-1-one after reduction using SmI_2/THF + 25 equiv. of MeOH

Syn diol: RT 16.50 min
Anti diol: RT 16.61 min

GC trace of diol of 3-Hydroxy-1-phenyl-butan-1-one after reduction using $\text{SmI}_2/\text{CH}_3\text{CN}$ + 10 equiv. of MeOH

Syn diol: RT 16.48 min
Anti diol: RT 16.66 min

GC trace of diol of 3-Hydroxy-1-phenyl-butan-1-one after reduction using SmI_2/DME + 25 equiv. of MeOH

Syn diol: RT 16.50 min
Anti diol: RT 16.65 min

Reduction of 1-Phenyl-3-hydroxy-butan-1-one utilizing SmI_2 in DME gave better diastereoselectivity compared to SmI_2 in THF, SmI_2 in CH_3CN gave *anti* diol as major isomer.

GC Method: Initial temp: 50 °C (hold for 5 min.), rate: 12 °C/min, final temp: 250 °C (hold for 10 min).

^{13}C NMR Trace ($\delta = 21$ to 33) of the *Syn* Acetal of 5,5-Dimethyl hexan-2,4-diol :

^{13}C NMR Trace ($\delta = 21$ to 35) of the *Anti* Acetal of 1-Phenyl-butan-1,3-diol :

^{13}C NMR Trace ($\delta = 20$ to 33) of the *Syn* Acetal of 1-Phenyl-butan-1,3-diol :