SUPPLEMENTARY MATERIAL

Source characterisation and mid-term spatial and temporal distribution of polycyclic aromatic hydrocarbons in molluscs along the Basque coast (northern Spain)

Chemistry and Ecology

O. Solaun*, J.G. Rodríguez, A. Borja, J. Larreta, I. Legorburu and V. Valencia

Marine Research Division, AZTI-Tecnalia Foundation, Pasaia, Spain

Herrera Kaia, Portualdea, z/g, 20110 Pasaia (Spain)

Tel: +34 946574000 Fax: +34 946572555

* Corresponding author, E-mail: osolaun@azti.es

Figure S1. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 1, throughout the study period (2003-2011).

Figure S2. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 2, throughout the study period (2003-2011).

Figure S3. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 3, throughout the study period (2003-2011).

Figure S4. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 4, throughout the study period (2003-2011).

Figure S5. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 5, throughout the study period (2003-2011).

Figure S6. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 6, throughout the study period (2003-2011).

Figure S7. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 7, throughout the study period (2003-2011).

Figure S8. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 8, throughout the study period (2003-2011).

Figure S9. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 9, throughout the study period (2003-2011).

Figure S10. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 10, throughout the study period (2003-2011).

Figure S11. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 11, throughout the study period (2003-2011).

Figure S12. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 12, throughout the study period (2003-2011).

Figure S13. Polycyclic aromatic hydrocarbon (PAH) distribution (%) in molluscs from Site 13, throughout the study period (2003-2011).

Figure S14. Box-plots for concentrations of polycyclic aromatic hydrocarbons (Σ PAH, sum of 18 PAH, in $\mu\text{g kg}^{-1}$, dry weight), for each of the sites considered along the Basque coast, in autumn, over the period 2003-2011.