Supporting Information for:

Sphingolipid synthesis via olefin cross metathesis: Preparation of a differentially protected building block and application to the synthesis of *D-erythro*-ceramide.

Anand Narain Rai and Amit Basu
Chemistry Department, Brown University Box H
Providence, RI 02912
abasu@brown.edu

General Experimental Protocols

Dry solvents (ether/THF/CH₂Cl₂/MeOH) were obtained using commercially available solvent purification system from GlassContour. All other reagent chemicals were purchased from Aldrich. NMR spectra were recorded on a Bruker instrument at 300 or 400 MHz for ¹H and 75 or 100 MHz for ¹³C. ¹H chemical shifts were referenced to TMS at 0 ppm or residual solvent peaks. ¹³C chemical shifts were referenced to the solvent peak, typically CDCl₃ at 77 ppm. The structures of the metathesis catalysts used are shown below. Catalysts **A** and **C** were purchased from Aldrich. Catalyst **B** was prepared using a literature procedure.¹

Experimental Procedures

The azidotriol **2** was prepared from diethyl (D)-tartrate in 3 steps in 31% overall using previously reported methods, shown above.²⁻⁴

¹H NMR (300 MHz, CD₃OD): δ 3.40-3.46 (m, 1H), 3.49-3.55 (m, 1H), 3.58 (m, 3H), 3.69 (dd, J = 7.3, 11.5 Hz, 1H), 3.91 (dd, J = 3.3, 11.5 Hz, 1H).

¹³C NMR (75.5 MHz, CD₃OD): δ 63.1, 64.3, 66.5, 72.6,

HRMS (ESI): Calcd. for C₁₁H₁₃N₃O₄ (M+Na)⁺ 170.0542, found 170.0548.

A mixture of **2** (3.85 g, 26.1 mmol), dibutyltin oxide (7.86 g, 31.4 mmol) in methanol was heated at reflux until the solution was clear. The solution was cooled to 0 °C, and triethylamine (20.2 ml, 144.0 mmol) and benzoyl chloride (15.2 ml, 130.9 mmol) were added. The reaction mixture was stirred for 6h at room temp. Reaction mixture was concentrated, and triturated with ethyl acetate (3x 100 ml). The combined ethyl acetate layers were concentrated to give crude product which was purified by column chromatography (SiO₂, ethyl acetate/hexanes, 3/7 to 1/1) to give 4.75 g (72 %, 18.9 mmol) of **3** as a waxy white solid.

¹H NMR (300 MHz, CDCl₃): δ 2.32 (t, J = 5.9 Hz, 1H), 3.15 (d, J = 5.2 Hz, 1H), 3.63 (m, 1H), 3.89-4.0 (m, 2H), 4.02-4.10 (m, 1H), 4.45 (dd, J = 5.7, 12 Hz, 1H), 4.64 (dd, J = 2.9, 12 Hz, 1H), 7.43 - 7.48 (m, 2H), 7.60 (tt, J = 1.2, 7.4 Hz, 1H), 8.04-8.06 (m, 2H),

¹³C NMR (75.5 MHz, CDCl₃): δ 62.6, 63.8, 66.5, 70.4, 128.5, 129.3, 129.7, 133.5, 167.2.

HRMS (ESI): Calcd. for C₁₁H₁₃N₃O₄ (M+Na)⁺ 274.0803, found 274.0804.

To a cold (0 °C) solution of **3** (4.7 g, 18.7 mmol) and imidazole (2.04 g, 30 mmol) in THF (300 ml) was added *tert*-butyldiphenylsilyl chloride (5.5 ml, 21.5 mmol). The reaction mixture was stirred for 12h at room temperature. Reaction mixture was added to a layer of ethyl acetate (200 ml), and washed with water (50 ml). The aqueous layer was extracted with ethyl acetate (2x100 ml). The combined ethyl acetate layers were dried (Na₂SO₄) and concentrated. The crude reaction mixture was purified by flash column chromatography (SiO₂, ethyl acetate/hexanes, 1/9) to give 6.92 g (75 %, 14.1 mmol) of silylated product as an oil.

¹H NMR (300 MHz, CDCl₃): δ 1.09 (s, 9H), 2.83 (d, *J* = 5 Hz, 1H), 3.56-3.62 (m, 1H), 3.92 (dd, *J* = 6.3, 10.8 Hz, 1H), 4.01 (dd, *J* = 4.1, 10.8 Hz, 1H), 4.39 (dd, *J* = 5.8, 11.9 Hz, 1H), 4.53 (dd, *J* = 3.1, 11.9 Hz, 1H), 7.38-7.43 (m, 8H), 7.55 (m, 1H), 7.68-7.71 (m, 4H), 8.0-8.02 (d, *J* = 7.7 Hz, 2H).

¹³C NMR (75.5 MHz, CDCl₃): δ 19.1, 26.7, 64.1, 64.2, 66.4, 69.7, 127.8, 128.4, 129.5, 129.7, 129.9, 132.6, 133.3, 135.5, 166.9.

HRMS (ESI): Calcd. for C₂₇H₃₁N₃O₄Si (M+H)⁺ 490.2162, found 490.2172.

To a solution of the secondary alcohol prepared above (6.92 g, 14.1 mmol) in toluene (350 ml) was added *p*-methoxybenzyltrichloroacetimidate (5.98 g, 21.2 mmol,) and sodium bicarbonate (1g) followed by addition of lanthanum triflate (828 mg, 1.42 mmol). The reaction mixture was stirred for 2h. The reaction mixture was washed with 200 ml of water and concentrated. The crude product was purified by column chromatography (SiO₂, ethyl acetate /hexanes, 1/19) to give 6.46 g (75 %, 10.6 mmol) of **4** as an oil.

¹H NMR (300 MHz, CDCl₃): δ 1.08 (s, 9H), 3.67-3.85 (m, 2H), 3.76 (s, 3H), 3.94 (dd, *J* = 3.8, 10.6 Hz, 1H), 3.83 (dd, *J* = 6.6, 10.6 Hz, 1H), 4.36 (dd, *J* = 5, 12 Hz), 4.42-4.63 (ab, *J* = 11.1 Hz, 2H), 4.63 (dd, *J* = 3.2, 12 Hz, 1H), 6.75 (d, *J* = 8.6 Hz, 2H), 7.08 (d, *J* = 8.6 Hz, 2H), 7.32-7.45 (m, 8H), 7.54-7.59 (m, 1H), 7.64-7.67 (m, 4H), 7.98-8.0 (m, 2H).

¹³C NMR (100 MHz, CDCl₃): δ 19.1, 26.7, 55.2, 63.4, 63.8, 63.9, 72.3, 75.4, 113.8, 127.8, 128.4, 129.5, 129.59, 129.64, 129.9, 132.8, 132.9, 133.1, 135.60, 135.62, 159.3, 166.2.

HRMS (ESI): Calcd. for C₃₅H₃₉N₃O₅Si (M+Na)⁺ 632.2557, found 632.2544.

To a solution of **4** (6.46 g, 10.6 mmol) in methanol (200 ml) was added sodium methoxide (15.9 mmol, prepared from 365 mg of sodium (15.8 mmol) in methanol (25 ml)). The reaction was stirred at room temperature for 1h. Amberlite (IR-120, H⁺) was added, and the reaction mixture was stirred until the solution was neutral. The reaction mixture was filtered, concentrated and purified by column chromatography (SiO₂, ethyl acetate/hexane, 1/9) to provide 4.3 g (81 %, 8.5 mmol) of the alcohol.

¹H NMR (300 MHz, CDCl₃): δ 1.08 (s, 9H), 3.49 (dt, *J* = 4.0, 7.0 Hz, 1H), 3.63- 3.80 (m, 4H), 3.77 (s, 3H), 3.90 (dd, *J* = 3.6, 10.6 Hz, 1H), 4.39-4.49 (ab, *J* = 11 Hz, 2H), 6.79 (d, *J* = 8.7 Hz, 2H), 7.08 (d, *J* = 8.6 Hz, 2H), 7.35-7.47 (m, 6H), 7.66-7.69 (m, 4H).

¹³C NMR (100 MHz, CDCl₃): δ 19.1, 26.7, 55.2, 61.1, 63.3, 64.1, 72.2, 77.7, 113.9, 127.8, 129.48, 129.54, 129.85, 129.85, 132.8, 132.9, 135.58, 135.63, 159.4.

HRMS (ESI): Calcd. for C₂₈H₃₁N₃O₄Si (M+Na)⁺ 528.2295, found 528.2285.

To a solution of the above alcohol (690 mg, 1.36 mmol) in dichloromethane (5 ml) was added a solution of Dess-Martin periodinane (865 mg, 2.04 mmol) in dichloromethane (15 ml). The reaction mixture was stirred at room temperature for 1.5h. Ether (50 ml) was added to the reaction mixture followed by the addition of 1.3 N NaOH (5 ml), and mixture was stirred till the ether layer became clear. The two layers were separated, and the aqueous layer was washed with ether (2 x10 ml). The combined ether layers were filtered through a plug of silica and concentrated to give 630 mg (92%, 1.25 mmol) of **4** as an oil.

¹H NMR (300 MHz, CDCl₃): δ 1.03 (s, 9H), 3.71-3.89 (m, 3H), 3.80 (s, 3H), 3.94 (dd, *J* = 1.7, 4 Hz, 1H), 4.55-4.64 (ab, *J* = 11.3 Hz, 2H), 6.86 (d, *J* = 8.7 Hz, 2H), 7.23 (d, *J* = 8.6 Hz, 2H), 7.33-7.47 (m, 6H), 7.61-7.66 (m, 4H), 9.63 (d, *J* = 1.8 Hz, 1H).

¹³C NMR (100 MHz, CDCl₃): δ 19.0, 26.7, 55.3, 62.2, 63.5, 73.1, 82.1, 114.0, 127.81, 127.84, 129.87, 129.89, 129.9, 132.5, 135.5, 159.7, 201.1.

HRMS (ESI): Calcd. for C₂₇H₃₁N₃O₄Si (M+Na)⁺ 526.2138, found 526.2136.

To a cold solution (-78 °C) of aldehyde (113 mg, 0.22 mmol) in THF was added cold (-78 °C) solution of ylide (prepared from $\text{Ph}_3\text{P}^+\text{CH}_3\text{Br}^-$ (1 eq.) and BuLi (0.9 eq.) through cannula until the color of the solution containing aldehyde turned yellow. The addition was stopped, and the reaction mixture was warmed to room temperature and stirred. After 30 min reaction mixture was cooled to -78 °C and quenched by addition of saturated methanolic ammonium chloride. The reaction mixture was concentrated to dryness and ether (10 ml) and water (5 ml) was added. The ether layer was separated, and water layer was extracted with ether (3 x 5 ml). The combined ether extracts were concentrated, and was purified by column chromatography (SiO_2 , ethyl acetate/ hexanes, 1/24) to afford 84 mg (76 %, 0.167 mmol) of **6** as an oil.

^1H NMR (300 MHz, CDCl_3): δ 1.05 (s, 9H), 3.49-3.55 (m, 1H), 5.52 (q, $J = 5.6$ Hz, 2H), 3.79 (s, 3H), 3.89-3.94 (m, 1H), 4.23-4.54 (ab, $J = 11.3$ Hz, 2H), 5.28 (m, 2H), 5.77 (ddd, $J = 8, 10.3, 17.0$ Hz), 6.82 (d, $J = 8.5$ Hz, 2H), 7.14 (d, $J = 8.5$ Hz, 2H), 7.34-7.46 (m, 6H), 7.63-7.66 (m, 4H).

^{13}C NMR (100.6 MHz, CDCl_3): δ 19.1, 26.7, 55.2, 63.6, 66.0, 70.0, 78.8, 113.7, 120.2, 127.7, 129.2, 129.9, 133.0, 134.7, 135.6, 159.1

HRMS (ESI): Calcd. for $\text{C}_{29}\text{H}_{35}\text{N}_3\text{O}_3\text{Si}$ ($\text{M}+\text{Na}$) $^+$ 524.2345, found 524.2324.

To a solution of **6** (27 mg, 0.054 mmol) and 1-pentadecene (0.15 ml, 0.55 mmol) in toluene (1 ml) was added **C** (5.5 mg, 0.009 mmol). Reaction mixture was stirred for 48h at room temperature. Reaction mixture was concentrated and directly purified by column chromatography (SiO_2 , ethyl acetate/hexanes, 1/24) to provide 14 mg (38 %, 0.02 mmol) of **6** as a waxy solid.

^1H NMR (300 MHz, CDCl_3): δ 0.88 (app t, $J_{\text{ave}} = 6.8$ Hz, 3H), 1.04 (s, 9H), 1.25-1.39 (m, 22H), 2.07 (q, $J = 7.2$ Hz, 2H), 3.5 (q, $J = 5.6$ Hz, 1H), 3.71- 3.77 (m, 2H), 3.79 (s, 3H), 3.88 (dd, $J = 5.6$ Hz, 8.4 Hz, 1H), 4.21-4.52 (ab, $J = 11.6$ Hz, 2H), 5.36 (dd, $J = 8.5, 15.4$ Hz, 1H), 5.69 (dt, $J = 8.4, 15.4$ Hz, 1H), 6.82 (d, $J = 8.6$ Hz, 2H), 7.14 (d, $J = 8.6$ Hz, 2H), 7.34-7.44 (m, 6H), 7.65 (d, $J = 6.8$ Hz, 4H).

^{13}C NMR (100.6 MHz, CDCl_3): δ 14.1, 19.1, 22.7, 26.7, 29.1, 29.2, 29.3, 29.5, 29.7, 31.9, 32.4, 55.3, 63.7, 66.2, 69.6, 78.6, 113.7, 126.2, 127.7, 129.7, 130.2, 133.1, 135.6, 137.7, 159.1.

HRMS (ESI): Calcd. for $\text{C}_{42}\text{H}_{61}\text{N}_3\text{O}_3\text{Si}$ ($\text{M}+\text{Na}$) $^+$ 706.4380, found 706.4398.

To a solution of **6** (80 mg, 0.16 mmol) in methanol (3 ml) was added NH_4Cl (85 mg, 1.59 mmol), and Zn (dust) (103 mg, 1.59 mmol). The reaction mixture was stirred vigorously for 2h at room temperature. The reaction mixture was concentrated, and the residue was triturated with ether (5 x 10 ml) and the combined ether fractions were filtered through small pad of silica to afford 63 mg (83 %, 0.13 mmol) of **8** as an oil which was used directly in the next step. To a cold (0°C) solution of **8** (52 mg, 0.11 mmol) in toluene (3 ml) was added diisopropylamine (0.094 ml, 0.54 mmol) followed by addition of Fmoc-Cl (84 mg, 0.33 mmol). The reaction mixture was stirred at room temperature for 2h. The reaction mixture was diluted with ethyl acetate (10 ml) and washed with water (5 ml) and brine (5ml). The organic layer was separated, dried (Na_2SO_4), concentrated and purified by column chromatography (ethyl acetate/ hexanes, 1/19), to provide 58 mg (76 %, 0.083 mmol) of **9** as a waxy solid.

^1H NMR (300 MHz, CDCl_3): δ 1.06 (s, 9H), 3.77 (s, 3H), 3.88-4.03 (m, 4H), 4.21-4.39 (m, 3H), 4.22-4.54 (ab, $J=11.1$ Hz, 2H), 5.10 (d, $J=9.1$ Hz, 1H), 5.29-5.35 (m, 2H), 5.78 (m, 1H), 6.81 (d, $J=8.6$ Hz, 2H), 7.15 (d, $J=8.5$ Hz, 2H), 7.28-7.43 (m, 10H), 7.56-7.65 (m, 6H), 7.76 (d, $J=7.4$ Hz, 2H).

^{13}C NMR (100 MHz, CDCl_3): δ 19.3, 26.9, 47.2, 55.2, 62.3, 66.8, 70.3, 79.7, 113.7, 119.5, 125.1, 125.2, 127.0, 127.6, 127.8, 129.3, 129.8, 130.2, 133.2, 135.57, 135.58, 135.8, 141.27, 141.28, 144.0, 144.1, 155.9, 159.1

HRMS (ESI): Calcd. for $\text{C}_{44}\text{H}_{47}\text{NO}_5\text{Si}$ ($\text{M}+\text{Na}$) $^+$ 720.3121, found 720.3135.

To a solution of **9** (23 mg, 0.033 mmol) and 1-pentadecene (0.044 ml, 0.16 mmol) in dichloromethane (1 ml) was added **A** (6 mg, 0.007 mmol). The reaction mixture was heated at reflux for 36h. The reaction mixture was concentrated and purified by column chromatography (SiO_2 , ethyl acetate/hexanes, 1/9) to provide 24 mg (82 %, 0.027 mmol) of **10** as a waxy solid.

^1H NMR (300 MHz, CDCl_3): δ 0.88 (app t, $J_{\text{ave}} = 6.5$ Hz, 3H), 1.06 (s, 9H), 1.22-1.32 (m, 22), 2.02 (q, $J=6.9$ Hz, 2H), 3.77 (s, 3H), 3.87-3.97 (m, 4H), 4.20-4.53 (ab, $J=11.1$ Hz, 2H), 4.23-4.31 (m, 3H), 5.07 (d, $J=9\text{Hz}$, 1H), 5.38 (dd, $J=8.1, 15$ Hz, 1H), 5.71 (dt, 7.2, 15 Hz, 1H), 6.81 (d, $J=8.4$ Hz, 2H), 7.16 (d, $J=8.4$ Hz, 2H), 7.28-7.41 (m, 10H), 7.56-7.66 (m, 6H), 7.76 (d, $J=7.7$ Hz, 2H).

^{13}C NMR (75.5 MHz, CDCl_3): δ 14.1, 19.3, 22.7, 26.9, 29.3, 29.4, 29.5, 29.7, 31.9, 32.3, 47.3, 55.3, 55.5, 62.5, 66.9, 69.9, 79.4, 113.7, 119.9, 125.2, 127.0, 127.3, 127.6, 127.7, 129.2, 129.3, 129.7, 130.5, 133.3, 135.6, 136.8, 141.3, 144.0, 155.9, 159.1.

HRMS (ESI): Calcd. for $\text{C}_{57}\text{H}_{73}\text{NO}_5\text{Si}$ ($\text{M}+\text{Na}$) $^+$ 902.5156, found 902.5135.

11

To a solution of **9** (16 mg, 0.023 mmol) and 1-nonene (0.02 ml, 0.12 mmol) in dichloromethane was added **A** (4 mg, 0.0046 mmol). Reaction mixture was heated at reflux for 24h. Reaction mixture was concentrated and purified by column chromatography (SiO_2 , ethyl acetate/hexane, 1/19) to provide 16 mg (87%, 0.02 mmol) of **11** as an oil.

^1H NMR (300 MHz, CDCl_3): δ 0.83-0.89 (m, 3H), 1.06 (s, 9H), 1.24-1.35 (m, 10H), 2.02 (q, J = 6.7 Hz, 2H), 3.77 (s, 3H), 3.86-3.99 (m, 4H), 4.23-4.31 (m, 3H), 4.21-4.53 (ab, J = 11.3 Hz, 2H), 5.07 (d, J = 9.4 Hz, 1H), 5.38 (dd, J = 7.6, 15.1 Hz, 1H), 5.71 (dt, J = 6.4, 15.1 Hz, 1H), 6.81 (d, J = 8.6 Hz, 2H), 7.15 (d, J = 8.6 Hz, 2H), 7.26-7.41 (m, 10 H), 7.56-7.65 (m, 6H), 7.76 (d, J = 7.5 Hz, 2H).

^{13}C NMR (75.5 MHz, CDCl_3): δ 14.1, 19.3, 22.6, 29.1, 29.18, 29.22, 31.8, 32.3, 47.2, 55.3, 62.5, 66.9, 69.9, 79.4, 113.7, 119.9, 125.2, 127.0, 127.3, 127.6, 127.7, 129.2, 130.5, 133.5, 135.6, 136.8, 141.3, 144.0, 144.1, 155.9, 159.1

HRMS (ESI): Calcd. for $\text{C}_{51}\text{H}_{61}\text{NO}_5\text{Si}$ ($\text{M}+\text{Na}$) $^+$ 818.4217, found 818.4200.

12

To a solution of **9** (10 mg, 0.0144 mmol) and **17** (35 mg, 0.1 mmol) in dichloromethane was added **A** (3 mg, 0.0036 mmol). The reaction mixture was heated at reflux for 24h. The reaction mixture was concentrated and purified by column chromatography (SiO_2 , ethyl acetate/hexanes 1/11 to 1/9) to provide 10 mg (60 %, 0.009 mmol) of **12** as a waxy solid.

^1H NMR (300 MHz, CDCl_3): δ 1.06 (s, 9H), 1.18-1.40 (m, 22 H), 2.02 (m, 2H), 2.12 (t, J = 7.2 Hz, 2H), 3.70-3.97 (m, 4H), 3.77 (s, 3H), 4.21-4.52 (ab, J = 11.3 Hz, 2H), 4.22-4.31 (m, 3H), 5.06 (d, J = 9.1 Hz, 1H), 5.37 (dd, J = 8.0, 15.2 Hz, 1H), 5.71 (dt, J = 6.7, 14.9 Hz, 1H), 6.81 (d, J = 8.5 Hz, 2H), 7.14-7.42 (m, 27H), 7.56-7.65 (m, 6H), 7.75 (d, J = 7.4 Hz, 2H).

^{13}C NMR (100 MHz, CDCl_3): δ 19.3, 26.9, 28.6, 29.0, 29.3, 29.4, 29.5, 29.6, 31.6, 32.0, 32.3, 47.2, 55.2, 62.4, 66.3, 69.9, 79.3, 113.7, 119.9, 125.2, 126.5, 127.0, 127.3, 127.6, 127.8, 129.2, 129.6, 129.7, 130.5, 133.2, 135.6, 136.9, 141.2, 144.0, 144.1, 145.1, 155.9, 159.0.

HRMS (FAB): Calcd. for $C_{76}H_{87}NO_5SSi$ ($M+Na$)⁺ 1176.5972, found 1176.5950.

A solution of **10** (18 mg, 0.02 mmol) and 20% piperidine in DMF (1ml) was stirred for 5 min. After completion of the reaction DMF was removed by azeotrope with heptane (5x5 ml) to provide crude amine. The crude amine was taken up in dichloromethane (3 ml) and triethylamine (0.02 ml, 0.14 mmol), DMAP (1mg, 0.008 mmol) and palmitoyl chloride (0.02 ml, 0.06 mmol) were added. The reaction mixture was stirred at room temperature for 3h. The reaction mixture was diluted with dichloromethane (10 ml) and washed successively with 1N HCl (5 ml), water (5 ml) and brine (5ml). The dichloromethane layer was dried (Na_2SO_4) and concentrated to give crude which was purified by column chromatography (SiO_2 , ethyl acetate/hexanes, 1/19) to provide 12 mg (66 %, 0.013 mmol) of **13** as a waxy solid.

1H NMR (400 MHz, $CDCl_3$): δ 0.88 (app t, J_{ave} = 6.8 Hz, 3H), 1.05 (s, 9H), 1.25 (m, 46H), 1.51-1.54 (m, 2H), 2.02-2.04 (m, 4H), 3.71 (dd, J = 4, 10.4 Hz, 1H), 3.78 (s, 3H), 3.92 (t, J = 7.8 Hz, 1H), 4.01 (dd, J = 4, 10.4 Hz, 1H), 4.10-4.16 (m, 1H), 4.19-4.51 (ab, J = 11.2 Hz, 2H), 5.35 (dd, J = 8.4, 10.2 Hz, 1H), 5.60 (d, J = 9.2 Hz, 1H), 5.66 (dt, J = 6.8, 15.2 Hz, 1H), 6.82 (d, J = 8.4 Hz, 2H), 7.16 (d, J = 8.8 Hz, 2H), 7.25-7.43 (m, 6H), 7.59-7.63 (m, 4H).

^{13}C NMR (75.5 MHz, $CDCl_3$): δ 14.1, 19.3, 22.7, 25.7, 26.8, 26.9, 29.31, 29.36, 29.4, 29.5, 29.7, 31.9, 32.3, 37.0, 53.1, 55.2, 62.4, 69.8, 79.3, 113.7, 127.5, 127.7, 129.17, 129.24, 129.7, 130.6, 133.3, 133.4, 135.6, 136.5, 159.0, 172.2.

HRMS (ESI): Calcd. for $C_{34}H_{58}NO_4Si$ ($M+H$)⁺ 896.6952, found 896.6958.

A solution of **13** (75 mg, 0.083 mmol) in 10% trifluoroacetic acid in dichloromethane (3 ml) was stirred for 2h. Solid potassium carbonate was added and was stirred until effervescence stopped. The suspension was filtered and concentrated to afford 46 mg (72%, 0.06 mmol) of the alcohol as an oil. To a solution of alcohol (40 mg, 0.051 mmol) in THF (3 ml) was added *tert*-butylammonium fluoride (0.1 ml, 0.11 mmol, 1M solution in THF). The reaction mixture was stirred at room temperature for 1h, and was

concentrated and purified by column chromatography (SiO₂, ethyl acetate/hexane, 2/3) to provide 11 mg (40%, 0.02 mmol) of **14** as a white waxy solid. Spectral and optical rotation data of the ceramide were in accord with that reported by Duclos.⁵

To a suspension of lithium aluminum hydride (512 mg, 13.49 mmol) in ether (14 ml) was added solution of *t*-butyl 14-pentadecenoate⁶ (2 g, 6.74 mmol) in ether (10 ml) drop wise at room temperature. The reaction mixture was heated at reflux for 12h. The reaction mixture was cooled (0 °C) and water (10 ml) was added drop wise with stirring. The reaction mixture was stirred for 3h at room temperature. The reaction mixture was diluted with ethyl acetate (50 ml) and, was washed with water (20 ml) and brine. The ethyl acetate layer was dried (Na₂SO₄) and concentrated to provide 1.4 g (91 %, 6.18 mmol) of the alcohol.

¹H NMR (300 MHz, CDCl₃): δ 1.26-1.37 (m, 20 H), 1.56 (quintet, *J* = 6.7 Hz, 2H), 2.0-2.07 (m, 2H), 3.62 (t, *J* = 6.6 Hz, 2H), 4.92 (d, *J* = 10.2 Hz, 1H), 4.99 (d, *J* = 18.4 Hz, 1H), 5.81 (ddt, *J* = 6.6, 10.2, 17.0 Hz, 1H).

¹³C NMR (75.5 MHz, CDCl₃): δ 25.7, 28.8, 29.1, 29.4, 29.6, 32.6, 33.7, 62.5, 113.9, 139.0.

To a cold solution of **15** (520 mg, 2.29 mmol) and pyridine (0.91 ml, 11.45 mmol) in dichloromethane (5 ml) was added methanesulfonyl chloride (0.7 ml, 9.16 mmol) and was stirred for 12h at room temperature. The reaction mixture was diluted with dichloromethane (20 ml) and, was washed with 0.5 N HCl (5 ml x 2), water (10 ml), aq. saturated Sodium bicarbonate (10 ml) and brine. The dichloromethane layer was dried (Na₂SO₄) and concentrated to provide 660 mg (94 %, 2.17 mmol) of the mesylate as a waxy solid, which was used directly in next step. The mesylate (660 mg, 2.16 mmol) and potassium thioacetate (1.23 g, 10.8 mmol) in DMF (5 ml) was stirred for 12h at room temperature. The reaction mixture was directly purified by column chromatography (SiO₂ ethyl acetate/ hexanes, 1/49) to afford 443 mg (72 %, 1.56 mmol) of **16** as yellow waxy solid.

¹H NMR (400 MHz, CDCl₃): δ 1.25-1.35 (m, 20 H), 1.52-1.60 (m, 2H), 2.01-2.06 (m, 2H), 2.86 (t, *J* = 7.32 Hz, 2H), 4.92 (d, *J* = 10.2 Hz, 1H), 4.98 (d, *J* = 17 Hz), 5.81 (tdd, *J* = 6.7, 10.2, 17 Hz, 1H).

¹³C NMR (100 MHz, CDCl₃): δ 28.8, 28.9, 29.07, 29.1, 29.4, 29.46, 29.5, 29.56, 29.58, 30.5, 33.8, 114.0, 139.1, 195.8

HRMS (FAB-MS): Calcd. for C₁₇H₃₂OS (M+H)⁺ 285.2252, found 285.2258.

The mixture of **16** (100 mg, 0.35 mmol), absolute ethanol (5 ml) and conc. HCl (0.5 ml) were heated at reflux for 12h. The reaction mixture was cooled and adjusted to a pH of 7 with 5 % methanolic ammonium hydroxide. The solution was concentrated and diluted with water (5 ml). The aqueous layer was extracted with dichloromethane (5 x 5 ml). The combined organic phases were dried (Na₂SO₄) and concentrated to afford the thiol as a pale yellow oil. To a solution of thiol in dry THF (8 ml) was added pyridine (0.06 ml, 0.70 mmol) followed by the addition of trityl chloride (147 mg, 0.53 mmol). The reaction mixture was allowed to stir for 12h at room temperature. The reaction mixture was concentrated and purified by column chromatography (SiO₂, ethyl acetate/hexanes, 1/99) to afford 98 mg (57 %, 0.20 mmol) of **17** as a waxy solid.

¹H NMR (300 MHz, CDCl₃): δ 1.66-1.43 (m, 22 H), 2.03 (q, J = 7 Hz, 2H), 2.13 (t, J = 7.2 Hz, 2H), 4.90-5.02 (m, 2H), 5.81 (tdd, J = 6.6, 10.2, 17 Hz, 1H), 7.16-7.29 (m, 9H), 7.39-7.43 (m, 6H).

¹³C NMR (100 MHz, CDCl₃): δ 28.6, 28.9, 29.0, 29.1, 29.2, 29.4, 29.49, 29.5, 29.59, 29.6, 32.0, 33.8, 66.4, 114.1, 126.5, 127.8, 129.6, 139.2, 145.1.

HRMS (FAB): Calcd. for C₃₄H₄₄S (M+Na)⁺ 507.3061, found 507.3070

References

- (1) Love, J. A.; Morgan, J. P.; Trnka, T. M.; Grubbs, R. H. *Angew. Chem. Int. Ed. Engl.* **2002**, *41*, 4035-4037.
- (2) Mori, K.; Kinsho, T. *Liebigs Ann. Chem.* **1991**, 1309-1315.
- (3) Metz, K.; Honda, M.; Komori, T. *Liebigs Ann. Chem.* **1993**, 55-60.
- (4) He, L.; Byun, H.-S.; Bittman, R. *Tetrahedron Lett.* **1998**, *39*, 2071-2074.
- (5) Duclos, R. I. *Chem Phys Lipids* **2001**, *111*, 111-138.
- (6) Hostetler, E. D.; Fallis, S.; McCarthy, T. J.; Welch, M. J.; Katzenellenbogen, J. A. *J. Org. Chem.* **1998**, *63*, 1348-1351.