Supporting Information

Stereocontrolled Synthesis of Onchidins

Yungui Peng ^a Heung Wing Pang ^a and Tao Ye* ^{a,b}

^a Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong;

^b Open Laboratory of Chirtechnology of the Institute of Molecular Technology for Drug Discovery & Synthesis and Department of Applied Biology & Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, China

bctaoye@inet.polyu.edu.hk

E-mail: bctaoye@inet.polyu.edu.hk

GENERAL EXPERIMENTAL

All non-aqueous reactions were run under an inert atmosphere (nitrogen or argon) with rigid exclusion of moisture from reagents and all reaction vessels were oven-dried. Solvents were distilled prior to use: THF from Na/benzophenone, dichloromethane, DMF, triethylamine and diisopropylethylamine from CaH₂. NMR spectra were recorded on Bruker Advance DPX 300MHz, AV400MHz or AV600MHz spectrometers. Chemical shifts are reported in parts per million (ppm), relative to either a tetramethylsilane internal standard or the signals due to the solvent. Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad), integration, and coupling constants. Low- and high- resolution EI and FAB mass spectra were obtained using a Finnigan MAT 95 mass spectrometer. IR spectra were recorded neatly or as KBr disk on a Bio-Red FTS 165 Fourier Transform Spectrophotometer. Optical rotations were recorded on a Perkin Elmer 343 Polarimeter. TLC was carried out using pre-coated sheets (Merck silica gel 60-F₂₅₀, 0.2mm) which, after development, were visualized at 254nm, and/or staining in p-anisole, ninhydrin or phosphomolybdic acid solution followed by heating. Flash column chromatography was performed using the indicated solvents (with R_f = 1.5-2.0 for the desired component) on E. Merck silica gel 60 (230-400 mesh ASTM). Melting points were measured on Carl Zeiss melting point apparatus and are uncorrected.

Experimental procedures

Compound (4)

Freshly distilled benzaldehyde (2.9 mL, 28.5 mmol) was added to a solution of L-lysine monohydrochloride (5 g, 27 mmol) in 2N lithium hydroxide solution (14 mL) at 0°C. The reaction flask was shaken until the

benzaldehyde had dissolved and a white solid precipitated. After standing in the refrigerator for several hours, the solid was filtered, washed with cold ethanol and dried over phosphorus pentoxide for one day to give N^ε-benzylidene-L-lysine (4.78 g, 75%) as a white solid, which was used in the next step without purification.

N^ε-benzylidene-L-lysine (4.78 g, 20 mmol) was dissolved in a mixture of 1N sodium hydroxide solution (20 mL) and ethanol (20 mL) below -5°C. A cooled mixture of 1N sodium hydroxide solution and ethanol (1:1, 80 mL) and benzyl chloroformate (3.9 mL, 26 mmol) were added in two portions over 5 to 6 min at -10°C with vigorous stirring. The mixture was stirred at the same temperature for 10 min and then room temperature for a further 10 min. Conc. hydrochloric acid (6 mL) was added and the resulting mixture was heated at 50°C for 10 min. The mixture was then extracted with diethyl ether (2x50 mL) and the aqueous layer was adjusted to pH 6.2. The resulting mixture was concentrated *in vacuo* to a volume of approximately 50 to 60 mL. After standing in the refrigerator for 2 days, the white crystals were filtered off, washed with a small amount of cold water and dried over phosphorus pentoxide for one day to give N^α-carbobenzoxy-L-lysine (3.36 g, 60%) as a white solid, which was used in next step without purification.

Sodium nitrite (2.8 g, 40.1 mmol) was added in three portions over 20 min to a stirred mixture of N^α-carbobenzoxy-L-lysine (3.5 g, 12.5 mmol) and sodium acetate (1.1 g, 13.6 mmol) in acetic acid (90 mL) at 40°C. The resulting mixture was stirred at the same temperature for 6 h. The solvent was removed *in vacuo* as far as possible. Water (100 mL) was added to the resulting residue and the mixture was extracted by dichloromethane (3x100 mL). The combined organic extracts were dried (Na₂SO₄) and concentrated *in vacuo* to give crude *acid* (2.4 g, 60%) as a pale yellow oil. The acid was used in the next reaction without further purification; ¹H NMR (400 MHz, CDCl₃) δ 1.39 – 1.50 (m, 2H), 1.59 – 1.76 (m, 3H), 1.86 – 1.98 (m, 1H), 2.03 (s, 3H), 4.05 (t, J = 6.5 Hz, 2H), 4.36 – 4.41 (m, 1H), 5.11 (s, 2H), 5.49 (d, J = 8.1 Hz, 1H), 7.29 – 7.35 (m, 5H).

Compound (5)

Triethylamine (3.8 mL, 27 mmol) and isobutyl chloroformate (3.7 mL) were added to a stirred solution of acid (**4**) (8.72 g, 27 mmol) in dry THF (135 mL) at -25°C and under an atmosphere of argon.

The resulting mixture was stirred at the same temperature for 30 min then at -10°C for 30 min.

The mixture was then added dropwise to a stirred solution of diazomethane¹¹¹ in diethyl ether at -10°C and under an atmosphere of argon. The resulting mixture was stirred for 1 h then allowed to warm to room temperature and stirred overnight. Excess diazomethane was destroyed by blowing air in the reaction flask for 1 h. The volatile solvents were removed *in vacuo* and the residue was dissolved in ethyl acetate (500 mL) and washed with saturated NaHCO₃ solution (50 mL), saturated NaCl solution (50 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 15% ethyl acetate – hexane as eluent to give the *diazoketone* (6.56 g, 70%) as a pale yellow oil; IR (neat) ν_{max} 3326, 3092, 3033, 2954, 2868, 2107, 1748, 1714, 1637, 1520, 1455, 1361, 1243, 1150, 1044, 909, 847, 753, 698 and 666 cm⁻¹; [α]_D²⁰ -15.6 (c = 1.52, CHCl₃); HRMS(EI) for maximum ion C₁₇H₂₁NO₅ (Loss N₂): Calcd 319.1420; Found 319.1410; ¹H NMR (400 MHz, CDCl₃) δ 1.37 – 1.44 (m, 2H), 1.56 – 1.68 (m, 3H), 1.82 – 1.87 (m, 1H), 2.04 (s, 3H), 4.04 (t, J = 6.5 Hz, 2H), 4.24 (m, 1H), 5.10 (s, 2H), 5.44 (s, 1H), 5.49 (d, J = 7.8 Hz, 1H), 7.31 – 7.36 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 20.89, 21.64, 28.21, 32.06, 54.06, 57.54, 63.86, 67.01, 128.03, 128.19, 128.49, 136.11, 155.91, 171.08, 193.11.

Compound (6)

Silver benzoate (85.4 mg, 0.37 mmol), dissolved in triethylamine (1.38 mL, 9.8 mmol), was added to a stirred solution of diazoketone (**5**) (1.18 g, 3.4 mmol) in dry methanol (15 mL) at

-25°C with the exclusion of light and under an atmosphere of argon. The resulting mixture

was allowed to warm to room temperature and stirred at that temperature overnight. The mixture was then filtered through a pad of celite and the resulting filtrate was evaporated *in vacuo*. The residue was dissolved in ethyl acetate (300 mL) and washed with saturated NaHCO₃ solution (30 mL), saturated NaCl solution (30 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale green oil. The oil was purified by silica gel chromatography using 25% ethyl acetate – hexane as eluent to give the *ester* (1.07 g, 90%) as a colorless oil; IR (neat) $\nu_{\max}$ 3349, 3033, 2952, 2868, 1740, 1698, 1531, 1455, 1438, 1367, 1243, 1169, 1038, 909, 846, 775, 740 and 698 cm⁻¹; [α]_D²⁰ -20.1 (c = 1.29, CHCl₃); HRMS(EI) for C₁₈H₂₅NO₆: Calcd 351.1682; Found 351.1686; ¹H NMR (400 MHz, CDCl₃) δ 1.38 – 1.45 (m, 2H), 1.53 – 1.66 (m, 4H), 2.04 (s, 3H), 2.56 (t, J = 6.2 Hz, 2H), 3.66 (s, 3H), 3.96 – 4.01 (m, 1H), 4.04 (t, J = 6.5 Hz, 2H), 5.09 (s, 2H), 5.22 (d, J = 8.7 Hz, 1H), 7.29 – 7.36 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 20.93, 22.55, 28.22, 33.98, 38.83, 47.95, 51.66, 64.09, 66.63, 128.02, 128.08, 128.48, 136.48, 155.81, 171.13, 171.87.

Compound (7)

Anhydrous K₂CO₃ (1.74 g, 12.6 mmol) was added to a stirred solution of ester (6) (2.22 g, 6.3 mmol) in dry methanol (70 mL) at -5°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 6 h and quenched by the addition of saturated NaCl solution (20 mL). The volatile solvent was removed *in vacuo* and the residue was extracted by ethyl acetate (3x100 mL) and the combined extracts were dried (Na₂SO₄) and concentrated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 50% ethyl acetate – hexane as eluent to give the *alcohol* (1.79 g, 92%) as a colorless oil; IR (neat) $\nu_{\max}$ 3336, 2925, 2856, 1736, 1686, 1542, 1461, 1436, 1375, 1275, 1252, 1221, 1194, 1168, 1116, 1069, 1048, 1033, 1000, 981, 962, 878, 841, 757, 729, 702 and 657 cm⁻¹; [α]_D²⁰ -10.1 (c = 1.01, EA); HRMS(EI) for C₁₆H₂₃NO₅: Calcd 309.1576; Found 309.1572; ¹H NMR (400 MHz, CDCl₃) δ 1.39 – 1.48 (m, 2H), 1.49 – 1.58 (m, 4H), 1.99 (br, 1H), 2.54 (t, J = 5.7 Hz, 2H),

3.61 (t, $J = 6.2$ Hz, 2H), 3.65 (s, 3H), 3.96 – 4.01 (m, 1H), 5.08 (s, 2H), 5.24 (d, $J = 8.8$ Hz, 1H), 7.29 – 7.35 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 22.24, 32.12, 34.10, 38.90, 47.98, 51.62, 62.35, 66.59, 128.05, 128.08, 128.43, 136.47, 155.93, 171.97.

[2-(Trimethylsilyl)ethoxy]methyl chloride SEMCl (1.7 mL, 9.6 mmol) was added dropwise to a stirred mixture of above alcohol (1.48 g, 4.8 mmol) and diisopropylethylamine DIPEA (3.5 mL, 20 mmol) in dry dichloromethane (20 mL) at 0 °C and under an atmosphere of argon. The resulting mixture was stirred at room temperature for 5 h and then evaporated *in vacuo* to leave a brown residue. The residue was dissolved in ethyl acetate (200 mL) and washed with saturated aqueous NH_4Cl (40 mL), saturated aqueous NaCl (40 mL), dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 15% ethyl acetate – hexane as eluent to give the *SEM ether* (1.84 g, 87%) as a colorless oil; IR (neat) ν_{max} 3336, 3065, 3033, 2951, 2871, 1747, 1720, 1653, 1527, 1455, 1438, 1374, 1247, 1110, 1027, 937, 921, 860, 836, 774, 753, 737 and 697 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ -9.9 ($c = 3.56$, EA); HRMS(EI) for $\text{C}_{22}\text{H}_{37}\text{NO}_6\text{Si}$: Calcd 439.2390; Found 439.2427; ^1H NMR (400 MHz, CDCl_3) δ 0.01 (s, 9H), 0.94 (t, $J = 8.5$ Hz, 2H), 1.38 – 1.46 (m, 2H), 1.53 – 1.61 (m, 4H), 2.54 (t, $J = 5.3$ Hz, 2H), 3.51 (t, $J = 6.4$ Hz, 2H), 3.60 (t, $J = 8.4$ Hz, 2H), 3.65 (s, 3H), 3.96 – 3.98 (m, 1H), 4.64 (s, 2H), 5.08 (s, 2H), 5.23 (d, $J = 9.0$ Hz, 1H), 7.28 – 7.35 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ -1.47 (3C), 18.04, 22.83, 29.33, 34.16, 38.77, 48.06, 51.58, 64.91, 66.53, 67.41, 94.72, 127.97, 128.00, 128.42, 136.50, 155.75, 171.87.

Compound (8)

Lithium diisopropylamide (LDA) (4.4 mL, 5.6 mmol) was added dropwise over 10 min to a stirred mixture of anhydrous lithium chloride (400 mg, 9.4 mmol) and SEM ether (7) (1.11 g, 2.53 mmol) in dry THF (20 mL) at -78°C and under an atmosphere at argon. The resulting mixture was stirred at the same temperature for another 1 h. Iodomethane (1 mL, 16 mmol) was added slowly to the reaction mixture at -78°C and under an atmosphere at argon. The resulting

mixture was stirred at the same temperature overnight and then hydrolyzed with the addition of saturated NH_4Cl solution (10 mL) at the same temperature. The mixture was dissolved in ethyl acetate (100 mL) and washed with saturated NH_4Cl solution (20 mL), saturated NaHCO_3 solution (20 mL), saturated NaCl solution (20 mL), dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 15% ethyl acetate – hexane as eluent to give the *ester* (854 mg, 75%) as a colorless oil; IR (neat) ν_{max} 3425, 3338, 3065, 3033, 2951, 2873, 1726, 1508, 1457, 1437, 1382, 1247, 1109, 1029, 936, 921, 860, 836, 753, 737 and 697 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ -6.3 ($c = 1.79$, EA); HRMS(EI) for $\text{C}_{22}\text{H}_{36}\text{NO}_6\text{Si}$ (Loss CH_3): Calcd 438.2312; Found 438.2314; ^1H NMR (400 MHz, CDCl_3) δ 0.01 (s, 9H), 0.94 (t, $J = 8.4$ Hz, 2H), 1.21(d, $J = 7.2$ Hz, 3H), 1.37 – 1.50 (m, 4H), 1.53 – 1.62 (m, 2H), 2.69 (dq, $J = 4.0, 7.2$ Hz, 1H), 3.50 (t, $J = 6.5$ Hz, 2H), 3.60 (t, $J = 8.4$ Hz, 2H), 3.66 (s, 3H), 3.75 – 3.80 (m, 1H), 4.64 (s, 2H), 5.10 (s, 2H), 5.41 (d, $J = 9.9$ Hz, 1H), 7.28 – 7.36 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ -1.44 (3C), 14.80, 18.08, 22.92, 29.42, 33.83, 42.68, 51.64, 53.31, 64.95, 66.57, 67.50, 94.76, 127.94, 128.00, 128.46, 136.66, 156.49, 175.63.

Compound (12)

A catalytic amount of 10% Pd/C was added in one portion to a stirred mixture of ester (**8**) (448 mg, 0.99 mmol) and trifluoroacetic acid (3 mL) in methanol (20 mL). The resulting mixture was stirred at room temperature in an atmosphere of hydrogen overnight and then filtered through a pad of celite and the residue was washed with methanol (2x10 mL). The filtrate was evaporated *in vacuo* to give the corresponding amine as a yellow solid which was used in the next reaction without further purification.

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride EDCI (570 mg, 3 mmol) was added in one portion to a stirred mixture of the crude amine, N-(*tert*-butoxycarbonyl)-L-valine (302 mg, 1.4 mmol), 1-hydroxybenzotriazole HOBt (402 mg, 3 mmol) and N-methylmorpholine NMM (0.8 mL, 8 mmol) in dry dichloromethane (5

mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and at room temperature overnight. The solvent was removed *in vacuo* and the resulting residue was dissolved in ethyl acetate (100 mL) and washed with saturated NaHCO₃ solution (50 mL) and saturated NaCl solution (50 mL). The organic layer was dried (Na₂SO₄) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 30% ethyl acetate – hexane as eluent to give the *amide* (266 mg, 70%) as a colorless viscous oil; IR (neat) ν_{max} 3313, 2960, 2872, 1732, 1681, 1650, 1521, 1457, 1391, 1367, 1302, 1249, 1172, 1046, 1019, 930, 878 and 756 cm⁻¹; [α]_D²⁰ -23.7 (c = 1.14, EA); HRMS(EI) for C₁₉H₃₆N₂O₆: Calcd 388.2573; Found 388.2575; ¹H NMR (400 MHz, CDCl₃) δ 0.90 (d, J = 6.8 Hz, 3H), 0.97 (d, J = 6.8 Hz, 3H), 1.18 (d, J = 7.2 Hz, 3H), 1.36 – 1.60 (m, 6H), 1.45 (s, 9H), 2.14 – 2.18 (m, 1H), 2.58 (br, 1H), 2.68 – 2.71 (m, 1H), 3.59 (t, J = 6.3 Hz, 2H), 3.68 (s, 3H), 3.90 (dd, J = 5.8, 8.5 Hz, 1H), 4.06 – 4.13 (m, 1H), 5.17 (d, J = 8.2 Hz, 1H), 6.88 (d, J = 9.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.88, 17.54, 19.38, 22.13, 28.19 (3C), 30.35, 32.09, 33.24, 42.39, 51.01, 51.63, 60.29, 62.13, 79.72, 155.80, 171.76, 175.79.

Compound (13)

A solution of dimethyl sulfoxide (0.4 mL, 7 mmol) in dry dichloromethane (3 mL) was added slowly to a stirred solution of oxalyl chloride (0.3 mL, 3.5 mmol) in dry dichloromethane (3 mL) at -78°C and under an atmosphere of argon. The resulting mixture

was stirred at the same temperature for 20 min and then a pre-cooled solution of amide (**12**) (313 mg, 0.8 mmol) in dry dichloromethane (3 mL) was added slowly to the stirred mixture at -78°C and under an atmosphere of argon. The resulting mixture was stirred at the same temperature for 2 h. Diisopropylethylamine DIPEA (3 mL, 17 mmol) was added dropwise over 20 min at -78°C and the mixture was stirred at that temperature for 1.5 h, and at -55°C for further 1.5 h. A saturated NH₄Cl solution (5 mL) was added to the reaction mixture and the separated aqueous layer was extracted by dichloromethane (3x15 mL). The combined

organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 25% ethyl acetate – hexane as eluent to give the corresponding aldehyde (**13**) (257 mg, 82%) as a colorless viscous oil; IR (neat) $\nu_{\max}$ 3313, 3070, 2960, 2872, 2821, 2724, 1726, 1679, 1650, 1539, 1520, 1456, 1390, 1366, 1337, 1303, 1249, 1172, 1112, 1046, 1020, 917, 878, 844, 799, 733 and 647 cm⁻¹; [α]_D²⁰ -28.3 (c = 1.55, EA); HRMS(EI) for C₁₉H₃₄N₂O₆: Calcd 386.2417; Found 386.2414; ¹H NMR (400 MHz, CDCl₃) δ 0.91 (d, J = 6.8 Hz, 3H), 0.99 (d, J = 6.8 Hz, 3H), 1.19 (d, J = 7.2 Hz, 3H), 1.45 (s, 9H), 1.48 – 1.50 (m, 2H), 1.62 – 1.67 (m, 2H), 2.04 – 2.23 (m, 1H), 2.42 – 2.48 (m, 2H), 2.67 – 2.69 (m, 1H), 3.69 (s, 3H), 3.92 (dd, J = 5.9, 8.3 Hz, 1H), 4.08 – 4.12 (m, 1H), 5.06 (d, J = 7.5 Hz, 1H), 6.83 (d, J = 9.1 Hz, 1H), 9.72 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 15.08, 17.52, 18.55, 19.39, 28.19 (3C), 30.34, 33.19, 42.28, 43.16, 50.64, 51.69, 60.32, 79.74, 155.71, 171.73, 175.77, 201.96.

Compound (14)

Anhydrous K₂CO₃ (498 mg, 3.6 mmol) was added in one portion to a stirred solution of Ohira-Bestmann reagent (700 mg, 3.6 mmol) in dry methanol (10 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 30 min and a solution of aldehyde (**13**) (257 mg, 0.66 mmol) in dry methanol (5 mL) was added slowly to the stirred reaction mixture at the same temperature. The resulting mixture was stirred at the same temperature for 1 h and allowed to warm to room temperature and stirred at that temperature overnight. A saturated NaCl solution (10 mL) was added to the reaction mixture and the separated aqueous layer was extracted with dichloromethane (3x20 mL). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 15% ethyl acetate – hexane as eluent to give the corresponding alkyne (**14**) (235 mg, 93%) as a colorless oil; IR (neat) $\nu_{\max}$ 3317, 2959, 2873, 2118, 1732, 1681, 1651, 1556, 1504, 1435, 1392, 1367, 1304, 1173, 1075, 1046,

1019, 930, 878, 844, 755 and 666 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ -28.9 ($c = 2.38$, EA); HRMS(EI) for $\text{C}_{20}\text{H}_{34}\text{N}_2\text{O}_5$: Calcd 382.2468; Found 382.2469; ^1H NMR (400 MHz, CDCl_3) δ 0.92 (d, $J = 6.9$ Hz, 3H), 0.98 (d, $J = 6.8$ Hz, 3H), 1.18 (d, $J = 7.2$ Hz, 3H), 1.45 (s, 9H), 1.49 – 1.66 (m, 4H), 1.93 (t, $J = 2.6$ Hz, 1H), 2.18 – 2.22 (m, 3H), 2.68 – 2.73 (m, 1H), 3.69 (s, 3H), 3.90 (dd, $J = 5.7, 8.4$ Hz, 1H), 4.08 – 4.12 (m, 1H), 5.07 (d, $J = 7.9$ Hz, 1H), 6.78 (d, $J = 9.1$ Hz, 1H); ^{13}C NMR (400 MHz, CDCl_3) δ 15.04, 17.52, 17.93, 19.38, 24.81, 28.19 (3C), 30.32, 32.80, 42.43, 50.52, 51.62, 60.30, 68.66, 79.66, 83.75, 155.66, 171.58, 175.78.

Compound (15)

Water (200 mL) was added to a three-necked flask containing L-valine (23.4 g, 0.2 mol). The flask was fitted with two addition funnels, one containing 2N sulphuric acid (150 mL, 0.3 mol) and another a solution of NaNO_2 (20.7 g, 0.3 mol) in water (150 mL). The acid was added dropwise to the stirred reaction mixture at 0°C . After the L-valine had dissolved, the NaNO_2 solution was added dropwise with the same rate of the addition as the acid. After the addition was complete, the resulting mixture was stirred at 0°C for 3 h, allowed to warm to room temperature and stirred at that temperature overnight. The mixture was extracted by ethyl acetate (5x200 mL). The combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give (S)-2-Hydroxy-3-methylbutyric acid (12 g, 50%) as a colourless viscous oil which was used in the next reaction without further purification.

A 1:99 (w/v) solution of I_2 in dichloromethane (1.6 mL) was added to a mixture of (S)-2-hydroxy-3-methyl-butylric acid (1.44 g, 12 mmol) and benzophenone hydrazone (3.59 g, 18.3 mmol) in dry dichloromethane (25 mL) at -10°C . The resulting mixture was stirred for 10 min and iodobenzene diacetate (5.9 g, 18.3 mmol) was added in several portions over 30 min to the stirred reaction mixture at -10°C . The mixture was stirred at that temperature for 2 h and at room temperature for a further 2 h. The solvent was evaporated *in vacuo* to leave a yellow residue. The residue was purified by silica gel chromatography using 10% ethyl acetate –

hexane as eluent to give the (S)-2-hydroxy-3-methyl-butyrac acid benzhydryl ester (2 g, 60%) as a colorless viscous oil; ^1H NMR (400 MHz, CDCl_3) δ 0.77 (d, J = 6.9 Hz, 3H), 1.02 (d, J = 7.0 Hz, 3H), 2.17 (dq, J = 3.3, 6.9 Hz, 1H), 2.68 (d, J = 2.5 Hz, 1H), 4.16 (dd, J = 3.3, 5.7 Hz, 1H), 6.96 (s, 1H), 7.27 – 7.36 (m, 10H); ^{13}C NMR (400 MHz, CDCl_3) δ 15.61, 18.90, 32.14, 74.99, 78.22, 127.02, 127.25, 128.15, 128.22, 128.56, 139.41, 139.45, 174.17.

Dicyclohexylcarbodiimide DCC (9.5 g, 46 mmol) was added in one portion to a stirred mixture of (S)-2-hydroxy-3-methyl-butyrac acid (4.7 g, 40 mmol), 2,2,2-trichloroethanol (3.87 mL, 40 mmol), 4-dimethylaminopyridine DMAP (5.6 g, 46 mmol) and 98% (1R)-(-)-10-camphorsulfonic acid CSA (1.86 g, 8 mmol) in dry dichloromethane (30 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and stirred at room temperature overnight. The mixture was filtered through a pad of celite and the residue was washed with hexane (2 x 20 mL). The combined organic layers were concentrated *in vacuo* to leave a residue which was dissolved in ethyl acetate (300 mL) and washed with saturated NH_4Cl solution (50 mL), saturated NaCl solution (50 mL), dried (Na_2SO_4) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 5% ethyl acetate – hexane as eluent to give (S)-2-hydroxy-3-methylbutyric acid 2,2,2-trichloro-ethyl ester (6.85 g, 69%) as a colorless oil; IR (neat) ν_{max} 3479, 2914, 2878, 1749, 1403, 1369, 1334, 1316, 1271, 1179, 1130, 1061, 1037, 1014, 9553, 891, 816, 760 and 715 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ - 12.50 (c = 1.2, EA); ^1H NMR (400 MHz, CDCl_3) δ 0.94 (d, J = 6.9 Hz, 3H), 1.09 (d, J = 6.9 Hz, 3H), 2.16 – 2.22 (m, 1H), 2.60 (d, J = 6.5 Hz, 1H), 4.21 (dd, J = 3.0, 6.4 Hz, 1H), 4.83 (dd, J = 11.9, 83.6 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.93, 18.79, 32.11, 74.56, 75.10, 94.33, 173.43.

60% Sodium hydride (5 g, 0.125 mol) was added in one portion to a stirred mixture Cbz-L-valine (9.5 g, 0.038 mol) in dry THF (150 mL) at 0°C and stirred for 30min at the same temperature

under an atmosphere of argon. Iodomethane (20 mL, 0.32 mol) was added and the resulting mixture was stirred overnight at room temperature. Iced water (200 mL) was poured into the mixture at 0°C and the resulting mixture was extracted with diethyl ether (2x200 mL). The aqueous layer was acidified to pH 1.5 using dilute HCl solution and extracted with ethyl acetate (3x200 mL). The combined organic extracts were washed with saturated sodium thiosulphate solution (100 mL), dried (Na₂SO₄) and evaporated *in vacuo* to give N-methyl-N-Cbz-L-valine (5 g, 55%) as a colorless viscous oil which was used in the next step without purification.

Dicyclohexylcarbodiimide DCC (1 g, 5 mmol) was added in one portion to a stirred mixture of N-methyl-N-Cbz-L-valine (636 mg, 2.5 mmol), (S)-2-hydroxy-3-methyl-butyric acid benzhydryl ester (502 mg, 2 mmol), 4-dimethylaminopyridine DMAP (610 mg, 5 mmol) and 98% (1R)-(-)-10-camphorsulfonic acid (CSA) (93 mg, 0.4 mmol) in dry dichloromethane (10 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was filtered through a pad of celite and the residue was washed with hexane (2x10 mL). The combined washings were concentrated *in vacuo* to leave a residue which was dissolved in ethyl acetate (100 mL) and washed with saturated NH₄Cl solution (25 mL), saturated NaCl solution (25 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 10% ethyl acetate – hexane as eluent to give (S)-2-(benzyloxy-carbonyl-methyl-amino)-3-methyl-butyric acid 1-benzhydryloxy carbonyl-2-methyl-propyl ester (930 mg, 93%) as a colorless oil; IR (neat) ν_{max} 3472, 3090, 3065, 3033, 2966, 2936, 2877, 1749, 1701, 1604, 1587, 1496, 1455, 1402, 1372, 1305, 1198, 1126, 1024, 996, 912, 869, 844, 754, 697 and 600 cm⁻¹; [α]_D²⁰ -64.6 (c = 8.48, EA); ¹H NMR (400 MHz, CDCl₃) δ 0.77 – 1.05 (m, 12H), 2.16 – 2.27 (m, 2H), 2.89, 2.91 (s, 3H), 4.7, 4.60 (d, J = 10.4 Hz, 1H), 4.94, 4.97 (d, J = 4.2 Hz, 1H), 5.14 (d, J = 4.5 Hz, 2H), 6.92, 6.93 (s, 1H), 7.22 – 7.34 (m, 15H); ¹³C NMR (400 MHz, CDCl₃) δ 16.69, 16.75, 18.56, 18.66, 18.81, 18.84, 19.55, 27.41,

27.48, 29.93, 30.00, 30.47, 30.69, 63.97, 64.30, 67.24, 67.30, 76.94, 76.96, 77.64, 77.01, 126.81, 126.85, 127.36, 127.60, 127.64, 127.80, 128.00, 128.28, 128.33, 136.37, 136.51, 139.20, 139.22, 139.33, 156.02, 156.68, 168.18, 168.26, 170.17, 170.57.

Trifluoroacetic acid (5 mL) was added dropwise to a stirred solution of (S)-2-(benzyloxycarbonyl-methyl-amino)-3-methyl-butyl-1-carboxy-2-methyl-propyl ester (472 mg, 0.95 mmol) in dry dichloromethane (5 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 10 h. The reaction mixture was poured into a mixture of benzene and chloroform (1:1, 10 mL) and evaporated *in vacuo* to give 2-(benzyloxycarbonyl-methyl-amino)-3-methyl-butyl-1-carboxy-2-methyl-propyl ester as a brown residue which was used in the next reaction without further purification.

Dicyclohexylcarbodiimide DCC (488 mg, 2.4 mmol) was added in one portion to a stirred mixture of 2-(benzyloxycarbonylmethylamino)-3-methyl-butyl-1-carboxy-2-methyl-propyl ester, (S)-2-hydroxy-3-methyl-butyl-1-carboxy-2-methyl-propyl ester, (S)-2-hydroxy-3-methyl-butyl-1-carboxy-2-methyl-propyl ester, 2,2,2-trichloro-ethyl ester (236 mg, 0.95 mmol), 4-dimethylaminopyridine DMAP (289 mg, 2.4 mmol) and 98% (1R)-(-)-10-camphorsulfonic acid CSA (44 mg, 0.19 mmol) in dry dichloromethane (4 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h then at room temperature overnight. The mixture was filtered through a pad of celite and the residue was washed with hexane (2x5 mL). The combined filtrate was evaporated *in vacuo* to leave a residue which was dissolved in ethyl acetate (50 mL) and washed with saturated NH₄Cl solution (15 mL), saturated NaCl solution (15 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 10% ethyl acetate – hexane as eluent to give the corresponding ester (397 mg, 70%) as a colorless oil; IR (neat) $\nu_{\max}$ 3464, 3091, 3066, 3034, 2968, 2938, 2878, 1747, 1699, 1497, 1469, 1455, 1392, 1373, 1289, 1241, 1191, 1125, 1029, 984, 910, 821, 768, 718 and 698 cm⁻¹; [α]_D²⁰ -63.3 (c =

3.83, EA); HRMS(EI) for $C_{26}H_{36}NCl_3O_8$: Calcd 595.1507; Found 595.1508; ^1H NMR (400 MHz, $CDCl_3$) δ 0.90 – 1.11 (m, 18H), 2.22 – 2.42 (m, 3H), 2.90 (s, 3H), 4.39, 4.58 (d, J = 10.6 Hz, 1H), 4.69 (d, J = 11.9 Hz, 1H), 4.89 – 4.94 (m, 2H), 5.06 (d, J = 4.0 Hz, 1H), 5.18 (t, J = 4.4 Hz, 2H), 7.29 – 7.38 (m, 5H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 15.93, 16.87, 16.93, 16.99, 18.67, 18.75, 18.77, 18.95, 19.01, 19.65, 27.60, 27.72, 30.02, 30.10, 30.67, 30.75, 32.10, 64.08, 64.33, 67.41, 67.49, 74.24, 74.53, 76.71, 76.83, 94.36, 127.73, 127.81, 127.88, 127.95, 128.44, 136.50, 136.65, 156.20, 156.85, 167.66, 168.87, 168.93, 170.40, 170.80.

A catalytic amount of 10% Pd/C was added in one portion to a stirred mixture of the above ester) and 1N hydrogen chloride in methanol (2 mL) in methanol (8 mL). The resulting mixture was stirred at room temperature in an atmosphere of hydrogen for 4 h and then filtered through a pad of celite and the residue was washed with methanol (2x10 mL). The filtrate was evaporated *in vacuo* to give the corresponding amine (**15**) which was used in the next reaction without further purification.

Compound (16)

Cesium carbonate (3.5 g, 10 mmol) was added to a stirred mixture of N-carbobenzyloxy-L-valine (1.26 g, 5 mmol) and iodomethane (2 mL, 30 mmol) in dry DMF (50 mL) at room temperature and under an atmosphere of argon. The resulting mixture was stirred at the room temperature overnight and then dissolved in a mixture of ethyl acetate and benzene (3:1, 400 mL) and washed with saturated NH_4Cl solution (50 mL), saturated $NaHCO_3$ solution (50 mL), saturated $NaCl$ solution (50 mL), dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 10% ethyl acetate – hexane as eluent to give 2-(benzyloxycarbonyl-methyl-amino)-3-methyl-butanoic acid methyl ester (1.33 g, 95%) as a colorless oil; IR (neat) ν_{max} 3065, 3033, 2965, 2876, 1747, 1702, 1498, 1401, 1371, 1307, 1262, 1206, 1149, 1009, 978, 768, 736 and 697 cm^{-1} ; $[\alpha]_D^{20}$ - 87.8 (c = 2.64, EA); HRMS(EI) for $C_{15}H_{21}NO_4$: Calcd 279.1471; Found 279.1479; ^1H NMR (400 MHz, $CDCl_3$) δ 0.87 (d, J =

6.7 Hz), 0.90 (d, J = 6.7 Hz, 3H), 0.95 (d, J = 6.5 Hz), 0.98 (d, J = 6.5 Hz, 3H), 2.13 – 2.22 (m, 1H), 2.91 (s, 3H), 3.65, 3.69 (s, 3H), 4.31 (d, J = 10.6 Hz), 4.52 (d, J = 10.6 Hz, 1H), 5.15, 5.16 (s, 2H), 7.26 – 7.33 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 18.63, 18.73, 19.39, 19.47, 27.38, 29.86, 30.28, 51.50, 63.63, 64.13, 67.15, 67.23, 127.51, 127.75, 128.24, 136.44, 156.02, 156.74, 170.99, 171.44.

A catalytic amount of 10% Pd/C was added in one portion to a stirred mixture of 2-(benzyloxycarbonyl-methyl-amino)-3-methyl-butyril acid methyl ester (380 mg, 1.36 mmol) and 1N hydrogen chloride in methanol (2 mL) in methanol (8 mL). The resulting mixture was stirred at room temperature in an atmosphere of hydrogen for 4 h and then filtered through a pad of celite and the residue was washed with methanol (2x10 mL). The filtrate was evaporated *in vacuo* to give N-methyl valine methyl ester as a yellow solid which was used in the next reaction without further purification.

Solid lithium hydroxide monohydrate (420 mg, 10.5 mmol) was added in one portion to a stirred solution of ester (**8**) (479 mg, 1.05 mmol) in THF/MeOH/water (2/2/1, 10 mL) at 0°C. After 1 h, the reaction mixture was allowed to warm at room temperature and stirred at that temperature overnight. The mixture was poured into diethyl ether (20 mL) and acidified to pH2 using dilute aqueous KHSO_4 solution at 0°C. The separated aqueous layer was extracted with ethyl acetate (3x30 mL) and the combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give the corresponding acid as a colorless oil which was used in the next reaction without further purification.

Diisopropylethylamine (1.46 mL, 8.4 mmol) was added dropwise to a stirred mixture of above acid, N-methyl valine methyl ester, 1-hydroxy-7-azabenzotriazole (HOAt) (163 mg, 1.26 mmol) and 2-bromo-1-ethyl pyridinium tetrafluoroborate (BEP) (700 mg, 2.6 mmol) in dry dichloromethane (5 mL) at -25°C and under an atmosphere of argon. The resulting mixture was stirred at the same temperature for 1 h and allowed to warm to room temperature

and stirred at that temperature overnight. The solvent was removed *in vacuo* and the resulting residue was dissolved in ethyl acetate (100 mL) and washed with saturated NH₄Cl solution (20 mL), saturated NaHCO₃ solution (20 mL) and saturated NaCl solution (20 mL). The organic layer was dried (Na₂SO₄) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 20% ethyl acetate – hexane as eluent to give the corresponding amide (**16**) (280 mg, 47%) as a colorless viscous oil; IR (neat) $\nu_{\max}$ 3396, 3333, 3065, 3033, 2952, 2874, 1720, 1637, 1500, 1458, 1411, 1373, 1254, 1211, 1061, 1027, 936, 859, 836, 736 and 697 cm⁻¹; HRMS(EI) for C₂₉H₅₀N₂O₇Si: Calcd 566.3387; Found 566.3389; (1S,2S,3S)-isomer (**16**): $[\alpha]_D^{20}$ - 53.2 (c = 3.64, EA); ¹H NMR (400 MHz, CDCl₃) δ -0.01 (s, 9H), 0.80 (d, J = 6.8 Hz, 3H), 0.90 (t, J = 8.5 Hz, 2H), 0.98 (d, J = 6.5 Hz, 3H), 1.16 (d, J = 7.1 Hz, 3H), 1.32 – 1.42 (m, 3H), 1.46 – 1.56 (m, 3H), 2.17 – 2.20 (m, 1H), 2.94 – 2.95 (m, 1H), 2.82, 2.97 (s, 3H), 3.45 (t, J = 10.2 Hz, 2H), 3.56 (t, J = 8.4 Hz, 2H), 3.66, 3.70 (s, 3H), 3.71 – 3.73 (m, 1H), 4.60 (s, 2H), 4.81 (d, J = 10.5 Hz, 1H), 5.06 (s, 2H), 6.36 (d, J = 9.6 Hz, 1H), 7.24 – 7.31 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ -1.53 (3C), 14.89, 15.02, 17.95, 18.54, 18.74, 19.45, 19.82, 22.95, 23.13, 27.23, 27.63, 29.41, 29.44, 31.72, 33.92, 38.07, 38.70, 51.71, 52.04, 54.01, 54.30, 61.06, 64.80, 65.66, 66.14, 67.44, 94.61, 127.57, 127.66, 128.23, 136.83, 156.72, 156.76, 170.14, 171.16, 176.46, 176.48;

The isomer **16a** was synthesized according to the above procedure from **ent-8**, which was in turn synthesized from D-Lysine.

(1R,2R,3S)-isomer (**16a**): $[\alpha]_D^{20}$ - 61.1 (c = 1.30, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 9H), 0.85 (d, J = 6.5 Hz, 3H), 0.93 (t, J = 8.1 Hz, 2H), 1.01 (d,

J = 6.2 Hz, 3H), 1.18 (d, J = 6.9 Hz, 3H), 1.35 – 1.48 (m, 3H), 1.50 – 1.56 (m, 3H), 2.17 – 2.26 (m, 1H), 2.95 – 2.97 (m, 1H), 2.86, 3.02 (s, 3H), 3.48 (t, J = 6.4 Hz, 2H), 3.59 (t, J = 8.2 Hz, 2H), 3.69, 3.72 (s, 3H), 3.71 – 3.73 (m, 1H), 4.63 (s, 2H), 4.82 (d, J = 10.5 Hz, 1H), 5.08 (s, 2H), 6.34 (d, J = 9.6 Hz), 6.39, (d, J = 9.6 Hz, 1H), 7.28 – 7.34 (m, 5H); ¹³C NMR (100

MHz, CDCl₃) δ -1.41 (3C), 14.69, 15.64, 18.11, 18.98, 19.03, 19.63, 19.81, 23.00, 23.18, 27.21, 27.74, 29.57, 29.66, 31.73, 34.14, 38.21, 38.35, 51.87, 52.06, 54.06, 54.51, 61.33, 64.96, 65.88, 66.29, 67.54, 67.63, 94.76, 127.71, 127.79, 128.38, 136.94, 156.83, 171.51, 176.55, 176.65.

Compound (18)

Trifluoroacetic acid (6 mL) was added dropwise to a stirred solution of amide (**16**) (669 mg, 1.18 mmol) in dry dichloromethane (6 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C overnight.

The reaction mixture was poured into a mixture of benzene and chloroform (1:1, 50 mL) and evaporated *in vacuo* to give the corresponding alcohol (464 mg, 90%) as a brown residue which was used in the next reaction without further purification.

A catalytic amount of 10% Pd/C was added in one portion to a stirred mixture of the above alcohol (464 mg, 1.06 mmol) and 1N hydrogen chloride in methanol (5 mL) in methanol (10 mL). The resulting mixture was stirred at room temperature in an atmosphere of hydrogen overnight and then filtered through a pad of celite and the residue was washed with methanol (2x30 mL). The filtrate was evaporated *in vacuo* to give the *amine* (**17**) as a yellow solid which was used in the next reaction without further purification.

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride EDCI (960 mg, 5 mmol) was added in one portion to a stirred mixture of the above amine (**17**), N-(*tert*-butoxycarbonyl)-L-valine (235 mg, 1.1 mmol), 1-hydroxybenzotriazole (HOBt) (674 mg, 5 mmol) and N-methylmorpholine (NMM) (1.1 mL, 10 mmol) in dry dichloromethane (5 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h then at room temperature overnight. The solvent was removed *in vacuo* and the resulting residue was dissolved in ethyl acetate (100 mL) and washed with saturated NaHCO₃ solution (20 mL) and saturated NaCl solution (20 mL). The organic layer was dried (Na₂SO₄) and

evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 1% methanol – chloroform as eluent to give amide (**18**) (266 mg, 50%) as a colorless viscous oil; IR (neat) $\nu_{\max}$ 3379, 2967, 2935, 2875, 1739, 1707, 1651, 1505, 1412, 1391, 1246, 1204, 1171, 1093, 1046, 1015, 928, 872, 755 and 665 cm^{-1} ; HRMS(EI) for $\text{C}_{25}\text{H}_{47}\text{N}_3\text{O}_7$: Calcd 501.3414; Found 501.3413; (1S,2S,3S,4S)-isomer (**18**): $[\alpha]_{\text{D}}^{20} = -76.6$ ($c = 4.28$, EA); ^1H NMR (400 MHz, CDCl_3) δ 0.83 (d, $J = 6.7$ Hz, 3H), 0.93 (d, $J = 7.0$ Hz), 0.90 (d, $J = 6.9$ Hz, 3H), 1.00 (d, $J = 6.9$ Hz, 3H), 1.02 (d, $J = 6.5$ Hz, 3H), 1.19 (d, $J = 7.4$ Hz), 1.16 (d, $J = 7.0$ Hz, 3H), 1.31 – 1.42 (m, 3H), 1.43 (s, 9H), 1.48 – 1.59 (m, 3H), 2.17 – 2.26 (m, 2H), 2.42 (br, 1H), 2.98 – 3.04 (m, 1H), 2.85, 3.02 (s, 3H), 3.58 (t, $J = 6.2$ Hz, 2H), 3.72, 3.73 (s, 3H), 3.96 – 3.99 (m, 1H), 4.02 – 4.09 (m, 1H), 4.90 (d, $J = 10.5$ Hz, 1H), 5.09 (d, $J = 8.2$ Hz, 1H), 7.80 (d, $J = 9.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.17, 15.34, 17.16, 17.24, 18.60, 18.82, 19.49, 19.83, 22.24, 22.43, 27.25, 27.65, 28.19, 28.23 (3C), 28.64, 30.52, 30.61, 31.62, 32.23, 33.68, 33.89, 37.69, 38.50, 51.85, 51.92, 52.10, 52.22, 60.03, 60.28, 60.85, 62.10, 62.25, 65.74, 79.48, 155.64, 170.13, 171.31, 171.71, 171.82, 176.71, 176.79

The isomer **18a** was synthesized according to the above procedure from **16a**

(1S,2R,3R,4S)-isomer (**18a**): $[\alpha]_{\text{D}}^{20} = -39.5$ ($c = 1.30$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 0.86 (d, $J = 6.7$ Hz, 3H), 0.91 (d, $J = 6.9$ Hz, 3H), 0.98 (d, $J = 6.8$ Hz, 3H), 1.02 (d, $J = 6.6$ Hz, 3H), 1.16 (d, $J = 7.0$ Hz, 3H), 1.35 – 1.41 (m,

2H), 1.45 (s, 9H), 1.50 – 1.52 (m, 2H), 1.55 – 1.60 (m, 2H), 2.17 (br, 1H), 2.18 – 2.26 (m, 2H), 2.97 (dq, $J = 3.4, 7.1$ Hz, 1H), 2.89, 3.04 (s, 3H), 3.59 (t, $J = 6.1$ Hz, 2H), 3.72, 3.75 (s, 3H), 3.94 – 4.03 (m, 1H), 4.04 – 4.08 (m, 1H), 4.84 (d, $J = 10.5$ Hz, 1H), 5.17 (d, $J = 7.4$ Hz, 1H), 7.76 (d, $J = 9.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.68, 15.76, 17.44, 18.92, 18.98, 19.16, 19.33, 19.50, 19.74, 22.20, 22.39, 27.10, 27.68, 28.20 (3C), 28.31, 30.53, 30.63, 31.61, 31.77, 32.19, 32.29, 33.76, 37.48, 37.70, 51.77, 51.81, 52.06, 52.25, 60.00, 60.13, 61.29, 62.09, 62.16, 65.89, 79.47, 155.67, 170.39, 171.31, 171.67, 176.60, 176.81.

Compound (19)

A solution of dimethyl sulfoxide (228 μ L, 3.88 mmol) in dry dichloromethane (5 mL) was added slowly to a stirred solution of oxalyl chloride (170 μ L, 1.94 mmol) in dry dichloromethane (5 mL) at -78°C and under an atmosphere

of argon. The resulting mixture was stirred at the same temperature for 20 min and then a solution of amide (**18**) (243 mg, 0.485 mmol) in dry dichloromethane (5 mL) was added slowly to the stirred mixture at -78°C and under an atmosphere of argon. The resulting mixture was stirred at the same temperature for 2 h. Diisopropylethylamine (1.69 mL, 9.7 mmol) was added dropwise over 20 min at -78°C and the mixture was stirred at that temperature for 1.5 h, then at -55°C for a further 1.5 h. Saturated NH_4Cl solution (10 mL) was added to the reaction mixture and the separated aqueous layer was extracted with dichloromethane (3x50 mL). The combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 50% ethyl acetate – hexane as eluent to give aldehyde (**19**) (212 mg, 87%) as a colorless viscous oil; (1S,2S,3S,4S)-(**19**) IR ν_{max} 3365, 2966, 2935, 2875, 2724, 1715, 1651, 1634, 1504, 1412, 1391, 1367, 1245, 1171, 1092, 1044, 1014, 929, 872, 755 and 665 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ -84.876 ($c = 0.88$, EA); HRMS(EI) for $\text{C}_{25}\text{H}_{45}\text{N}_3\text{O}_7$: Calcd 499.3258; Found 499.3254; ^1H NMR (400 MHz, CDCl_3) δ 0.82 (d, $J = 6.7$ Hz, 3H), 0.90 (d, $J = 6.9$ Hz, 3H), 1.00 (d, $J = 7.1$ Hz, 3H), 1.02 (d, $J = 6.7$ Hz, 3H), 1.15 (d, $J = 7.2$ Hz), 1.12 (d, $J = 7.1$ Hz, 3H), 1.41 (s, 9H), 1.34 – 1.48 (m, 2H), 1.53 – 1.60 (m, 2H), 2.14 – 2.21 (m, 2H), 2.41 (t, $J = 6.9$ Hz, 2H), 2.93 (dq, $J = 3.3, 7.1$ Hz, 1H), 2.80, 2.97 (s, 3H), 3.67, 2.69 (s, 3H), 3.92 – 3.97 (m, 1H), 4.00 – 4.02 (m, 1H), 4.84 (d, $J = 10.6$ Hz, 1H), 5.01 (d, $J = 8.2$ Hz, 1H), 7.77 (d, $J = 9.2$ Hz, 1H), 9.72 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.22, 15.35, 17.18, 17.26, 18.53, 18.64, 18.78, 19.45, 19.79, 27.24, 27.62, 28.16 (3C), 28.20, 28.64, 30.46, 30.57, 31.57, 33.41, 33.47, 37.84, 38.53, 43.12, 43.19, 51.49, 51.74, 51.84, 52.10, 60.03, 60.80, 65.76, 79.47, 155.57, 170.03,

171.12, 171.75, 171.81, 176.55, 176.65, 202.10.

Compound (20)

Anhydrous K_2CO_3 (300 mg, 2.18 mmol) was added in one portion to a stirred solution of Ohira-Bestmann reagent (418 mg, 2.18 mmol) in dry methanol (5 mL) at $0^\circ C$ and under an atmosphere of argon. The resulting mixture was stirred at $0^\circ C$

for 30 min and a solution of aldehyde (**19**) (217 mg, 0.44 mmol) in dry methanol (5 mL) was added slowly to the stirred reaction mixture at the same temperature. The resulting mixture was stirred for 1 h, allowed to warm to room temperature and stirred at that temperature overnight. A saturated NaCl solution (10 mL) was added to the reaction mixture and the separated aqueous layer was extracted by dichloromethane (4x50 mL). The combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 30% ethyl acetate – hexane as eluent to give alkyne (**20**) (176 mg, 80%) as a colorless viscous oil; IR (neat) ν_{max} 3311, 2966, 2933, 2875, 2118, 1739, 1714, 1651, 1514, 1412, 1391, 1367, 1245, 1172, 1089, 1042, 1014, 928, 871, 755 and 665 cm^{-1} ; HRMS(EI) for $C_{26}H_{45}N_3O_6$: Calcd 495.3308; Found 495.3297; (1S,2S,3S,4S)-isomer (**20**): $[\alpha]_D^{20}$ - 86.68 ($c = 1.79$, EA): ^1H NMR (400 MHz, $CDCl_3$) δ 0.81 (d, $J = 6.7$ Hz, 3H), 0.88 (d, $J = 6.9$ Hz, 3H), 0.89 (d, $J = 6.8$ Hz), 0.92 (d, $J = 6.7$ Hz), 0.98 (d, $J = 6.9$ Hz, 3H), 1.00 (d, $J = 6.5$ Hz, 3H), 1.18 (d, $J = 7.1$ Hz), 1.15 (d, $J = 7.1$ Hz, 3H), 1.43 (s, 9H), 1.47 – 1.53 (m, 4H), 1.90 (t, $J = 2.6$ Hz, 1H), 2.14 – 2.23 (m, 4H), 2.96 (dq, $J = 3.3, 7.4$ Hz, 1H), 2.84, 3.00 (s, 3H), 3.71, 3.72 (s, 3H), 3.95 – 4.00 (m, 1H), 4.04 – 4.08 (m, 1H), 4.89 (d, $J = 10.6$ Hz, 1H), 5.03 (d, $J = 8.2$ Hz, 1H), 7.75 (d, $J = 9.1$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 15.35, 15.53, 17.24, 17.33, 17.93, 18.05, 18.58, 18.94, 19.55, 19.87, 25.03, 25.06, 27.28, 27.72, 28.26, 28.30 (3C), 28.71, 30.60, 30.73, 31.65, 33.26, 33.42, 38.09, 38.77, 51.53, 51.78, 51.91, 52.18, 60.09, 60.85, 65.91, 68.59, 79.55, 83.99, 155.66, 170.15, 171.24, 171.73, 176.80, 176.82

The isomer **20a** was synthesized according to the above procedures from **18a**.

(1S,2R,3R,4S)-isomer (**20a**): $[\alpha]_D^{20} - 57.1$ ($c = 1.90$, CHCl_3);

^1H NMR (400 MHz, CDCl_3) δ 0.88 (d, $J = 6.7$ Hz, 3H), 0.91 (d, $J = 6.9$ Hz, 3H), 0.98 (d, $J = 6.9$ Hz, 3H), 1.03 (d, $J = 6.5$ Hz, 3H), 1.17 (d, $J = 7.1$ Hz, 3H), 1.44 (s, 9H), 1.48 – 1.60

(m, 4H), 1.91 (t, $J = 2.6$ Hz, 1H), 2.10 – 2.26 (m, 4H), 2.96 (dq, $J = 3.4, 7.0$ Hz, 1H), 2.89, 3.05 (s, 3H), 3.72, 3.77 (s, 3H), 3.96 – 4.01 (m, 1H), 4.06 – 4.10 (m, 1H), 4.85 (d, $J = 10.6$ Hz, 1H), 5.16 (d, $J = 7.9$ Hz, 1H), 7.68 (d, $J = 9.2$ Hz, 1H), 7.73 (d, $J = 9.7$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 14.79, 15.84, 17.48, 17.53, 18.01, 19.00, 19.10, 19.40, 19.57, 19.85, 19.78, 24.95, 25.13, 27.18, 27.67, 28.26 (3C), 30.86, 31.04, 31.79, 33.05, 33.23, 37.62, 38.07, 51.08, 51.36, 51.86, 52.16, 59.98, 60.12, 61.32, 65.89, 68.50, 68.65, 79.46, 83.78, 155.67, 170.31, 171.36, 171.67, 176.62, 176.84.

Compound (23)

Benzyl bromide (5.95 mL, 50 mmol) was added dropwise to a stirred mixture of (S)-2-hydroxy-3-methyl-butyl benzoate (1.18 g, 10 mmol) and anhydrous NaHCO_3 (1.68 g, 20 mmol) in dry DMF (30 mL) under an atmosphere of argon. The resulting mixture was stirred at the same temperature overnight and the mixture was dissolved in ethyl acetate (300 mL) and washed with water (2x50 mL), saturated NH_4Cl solution (50 mL) and saturated NaCl solution (50 mL). The organic layer was dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 10% ethyl acetate – hexane as eluent to give (S)-2-hydroxy-3-methyl-butyl benzoate (1.64 g, 80%) as a colorless oil; IR (neat) ν_{max} 3504, 3066, 3034, 2964, 2933, 2875, 1735, 1498, 1465, 1456, 1388, 1370, 1261, 1137, 1106, 1070, 1029, 990, 913, 830, 751 and 698 cm^{-1} ; $[\alpha]_D^{20} - 14.4$ ($c = 1.17$, EA); HRMS(EI) for $\text{C}_{12}\text{H}_{16}\text{O}_3$: Calcd 208.1099; Found 208.1093; ^1H NMR (300 MHz, CDCl_3) δ 0.83 (d, $J = 6.9$ Hz, 3H), 0.99 (d, $J = 6.9$ Hz, 3H), 2.01 – 2.11 (m, 1H), 2.91 (d, $J = 6.1$ Hz, 1H), 4.07 (dd, $J = 3.6, 6.2$ Hz, 1H), 5.20 (d, $J = 12.2, 18.3$ Hz, 2H), 7.34 (s,

5H); ^{13}C NMR (75 MHz, CDCl_3) δ 15.82, 18.65, 32.05, 67.06, 74.94, 128.28, 128.39, 128.49, 135.15, 174.65.

[2-(Trimethylsilyl)ethoxy]methyl chloride (1.84 mL, 10.4 mmol) was added dropwise to a stirred mixture of (S)-2-hydroxy-3-methyl-butanoic acid benzyl ester (1.442 g, 6.93 mmol) and diisopropylethylamine (5.2 mL, 30 mmol) in dry dichloromethane (15 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at room temperature overnight and then evaporated *in vacuo* to leave a brown residue. The residue was dissolved in ethyl acetate (200 mL) and washed with saturated NH_4Cl solution (50 mL), saturated NaCl solution (50 mL), dried (Na_2SO_4) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 5% ethyl acetate – hexane as eluent to give (S)-3-methyl-2-(2-trimethylsilyl-ethoxymethoxy)-butanoic acid benzyl ester (2.07 g, 88%) as a colorless oil; IR (neat) ν_{max} 3066, 3035, 2957, 2893, 1749, 1498, 1456, 1385, 1368, 1249, 1183, 1156, 1131, 1119, 1057, 1039, 936, 918, 860, 835, 750 and 696 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ - 56.7 ($c = 2.72$, EA); ^1H NMR (400 MHz, CDCl_3) δ 0.01 (s, 9H), 0.89 (dd, $J = 8.2, 8.9\text{ Hz}$, 2H), 0.94 (d, $J = 6.4\text{ Hz}$, 3H), 0.96 (d, $J = 6.4\text{ Hz}$, 3H), 2.09 – 2.14 (m, 1H), 3.59 – 3.69 (m, 2H), 3.96 (d, $J = 5.4\text{ Hz}$, 1H), 4.71 (s, 2H), 5.18 (dd, $J = 12.3, 14.0\text{ Hz}$, 2H), 7.30 – 7.37 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ -1.51 (3C), 17.48, 17.92, 18.72, 31.38, 65.57, 66.30, 80.31, 94.42, 128.23 (2C), 128.47, 135.63, 172.14.

A catalytic amount of 10% Pd/C was added in one portion to a stirred mixture of (S)-3-methyl-2-(2-trimethylsilyl-ethoxymethoxy)-butanoic acid benzyl ester (1.43 g, 3.97 mmol) in methanol (30 mL). The resulting mixture was stirred at room temperature in an atmosphere of hydrogen for 4 h and then filtered through a pad of celite and the residue was washed with MeOH (2x30 mL). The combined filtrates were evaporated *in vacuo* to give (S)-3-methyl-2-(2-trimethylsilyl-ethoxymethoxy)-butanoic acid as a yellow oil which was used in the next reaction without further purification.

Dicyclohexylcarbodiimide (1.69 g, 8.2 mmol) was added in one portion to a stirred mixture of (S)-3-methyl-2-(2-trimethylsilyl-ethoxymethoxy)-butyric acid, (S)-2-hydroxy-3-methyl-butyl-2,2,2-trichloro-ethyl ester (1.02 g, 4.1 mmol), 4-dimethylaminopyridine (1.0 g, 8.2 mmol) and (1R)-(-)-10-camphorsulfonic acid (0.18 g, 0.8 mmol) in dry dichloromethane (15 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and stirred at room temperature overnight. The mixture was filtered through a pad of celite and the residue was washed with hexanes (2 x 20 mL). The filtrate was evaporated *in vacuo* to leave a residue which was dissolved in ethyl acetate (200 mL) and washed with saturated NH₄Cl solution (50 mL), saturated NaCl solution (50 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 2% ethyl acetate – hexane as eluent to give the corresponding ester (**23**) (1.54 g, 81%) as a colorless viscous oil; IR (neat) ν_{max} 2962, 2893, 1756, 1467, 1391, 1372, 1328, 1276, 1248, 1179, 1058, 989, 936, 919, 860, 835, 768, 718, 694, 665 and 572 cm⁻¹; [α]_D²⁰ - 68.2 (c = 1.8, EA); HRMS(FAB) for C₁₈H₃₂ClO₆Si (Loss HCl₂): Calcd 407.1657; Found 407.1667; ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 9H), 0.88 – 0.93 (m, 2H), 1.02 (d, J = 6.8 Hz, 3H), 1.05 (d, J = 6.7 Hz, 3H), 1.06 (d, J = 6.1 Hz, 3H), 1.08 (d, J = 6.2 Hz, 3H), 2.19 – 2.24 (m, 1H), 2.33 – 2.38 (m, 1H), 3.56 – 3.63 (m, 1H), 3.68 – 3.75 (m, 1H), 4.10 (d, J = 4.7 Hz, 1H), 4.75 (s, 2H), 4.81 (dd, J = 12.0, 67.2 Hz, 1H), 5.01 (d, J = 4.1 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ -1.46 (3C), 17.03, 17.16, 18.03, 18.83, 18.92, 30.07, 31.47, 65.66, 74.22, 76.47, 79.40, 94.25, 94.40, 167.90, 172.00.

Compound (22)

Trifluoroacetic acid (6 mL) was added dropwise to a stirred solution of (S, S)-3-methyl-2-(2-trimethylsilyl-ethoxymethoxy)-butyric acid 2-methyl-1-(2,2,2-trichloro-ethoxycarbonyl)-propyl ester (1 g, 2.1 mmol) in dry dichloromethane (6 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 5h. The reaction mixture

was poured into a mixture of benzene and chloroform (1:1, 30 mL) and evaporated *in vacuo* to leave a brown residue. The residue was purified by silica gel chromatography using 12% ethyl acetate – hexane as eluent to give the corresponding alcohol (**22**) (624 mg, 85%) as a colorless oil; IR (neat) $\nu_{\max}$ 3522, 2970, 2936, 2879, 1771, 1746, 1469, 1393, 1373, 1179, 1129, 1033, 964, 905, 821, 769, 719 and 571 cm^{-1} ; $[\alpha]_{\text{D}}^{20}$ – 19.0 ($c = 3.5$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 0.98 (d, $J = 6.9$ Hz, 3H), 1.05 (d, $J = 6.9$ Hz, 3H), 1.08 (d, $J = 6.9$ Hz, 3H), 1.09 (d, $J = 6.9$ Hz, 3H), 2.20 (dq, $J = 3.6, 6.9$ Hz, 1H), 2.39 (dq, $J = 4.0, 6.9$ Hz, 1H), 2.46 (br, 1H), 4.16 (d, $J = 3.6$ Hz, 1H), 4.82 (dd, $J = 12.0, 72.9$ Hz, 1H), 5.06 (d, $J = 4.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.86, 16.91, 18.65, 18.74, 30.06, 32.28, 74.23, 74.90, 77.06, 94.32, 167.60, 174.56.

Compound (**24**)

Solid lithium hydroxide monohydrate (148 mg, 3.54 mmol) was added in one portion to a stirred solution of acetylenic amide (**20**) (175 mg, 0.35 mmol) in THF/MeOH/water (2/2/1, 5 mL) at 0°C. After 1 h, the reaction mixture was warmed to room temperature and stirred at that temperature for 5h. The volatile solvents were removed *in vacuo* and the aqueous phase was poured into diethyl ether (15 mL) and acidified to pH2 using dilute aqueous KHSO_4 solution. The separated aqueous layer was extracted with ethyl acetate (3x50 mL) and the combined organic extracts were washed with saturated NaCl solution (20 mL), dried (Na_2SO_4) and evaporated *in vacuo* to give the corresponding acid (**21**) as a pale yellow oil which was used in the next reaction without further purification.

Dicyclohexylcarbodiimide (365 mg, 1.77 mmol) was added in one portion to a stirred mixture of the above acid, alcohol (**22**) (245 mg, 0.7 mmol), 4-dimethylaminopyridine (216 mg, 1.8 mmol) and (1R)-(-)-10-camphorsulfonic acid (16.4 mg, 0.07 mmol) in dry

dichloromethane (4 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was filtered through a pad of celite and the residue was washed with a mixture of hexanes and ethyl acetate (1:1, 3x5 mL). The filtrate was evaporated *in vacuo* to leave a residue which was dissolved in ethyl acetate (50 mL) and washed with saturated NH₄Cl solution (5 mL), saturated NaCl solution (5 mL), dried (Na₂SO₄) and evaporated *in vacuo* to leave a pale yellow oil. The oil was purified by silica gel chromatography using 25% ethyl acetate – hexane as eluent to give the corresponding ester (**24**) (205 mg, 70%) as a colorless viscous oil; IR (neat) $\nu_{\max}$ 3311, 2967, 2935, 2877, 2126, 1760, 1714, 1659, 1500, 1392, 1368, 1260, 1177, 1125, 1027, 870, 801, 757, 718 and 667 cm⁻¹; HRMS(EI) for C₃₇H₆₀Cl₃N₃O₁₀: Calcd 811.3344; Found 811.3337; (**24**): ¹H NMR (400 MHz, CDCl₃) δ 0.85 (d, J = 6.7 Hz, 3H), 0.90 (d, J = 6.9 Hz, 3H), 1.00 (d, J = 6.8 Hz, 3H), 1.04 – 1.10 (m, 12H), 1.17 (d, J = 7.1 Hz, 3H), 1.26 – 1.23 (m, 3H), 1.43 (s, 9H), 1.52 – 1.53 (m, 4H), 1.91 (t, J = 2.5 Hz, 1H), 2.14 – 2.28 (m, 4H), 2.32 – 2.39 (m, 2H), 2.94 – 2.97 (m, 1H), 2.87, 3.05 (s, 3H), 3.97 – 4.07 (m, 1H), 4.07 – 4.11 (m, 1H), 4.81 (dd, J = 12.0, 96.4 Hz, 2H), 4.96 (d, J = 4.0 Hz, 1H), 5.00 (d, J = 10.5 Hz, 1H), 5.06 (d, J = 4.0 Hz, 2H), 7.75 (d, J = 9.5 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 15.38, 15.63, 16.92, 16.97, 17.22, 17.28, 17.89, 17.93, 18.65, 18.70, 18.76, 19.03, 19.53, 19.77, 19.96, 24.99, 25.18, 27.50, 28.25 (3C), 29.94, 30.01, 30.04, 30.66, 31.95, 32.30, 33.32, 33.40, 38.31, 38.81, 51.47, 51.71, 60.01, 60.31, 61.03, 65.82, 68.57, 68.66, 74.18, 76.76, 76.79, 76.87, 79.48, 83.92, 84.07, 94.29, 155.64, 167.50, 167.60, 168.44, 168.74, 169.44, 170.25, 171.67, 176.67, 176.71

The isomer **24a** was synthesized according to the above procedures from **20a**

(2R,3R)-(**24a**): $[\alpha]_D^{20}$ - 58.4 (c = 1.46, CHCl₃);

¹H NMR (400 MHz, CDCl₃) δ 0.91 (d, J = 6.9 Hz, 6H), 0.98 (d, J = 6.8 Hz, 3H), 1.04 – 1.12

(m, 15H), 1.16 (d, $J = 7.1$ Hz, 3H), 1.44 (s, 9H), 1.49 – 1.60 (m, 4H), 1.90 (t, $J = 2.6$ Hz, 1H), 2.09 – 2.29 (m, 4H), 2.33 – 2.41 (m, 2H), 2.96 (dq, $J = 3.4, 7.0$ Hz, 1H), 2.95, 3.06 (s, 3H), 3.99 – 4.04 (m, 1H), 4.06 – 4.08 (m, 1H), 4.80 (dd, $J = 12.0, 92.7$ Hz, 2H), 4.95 (d, $J = 4.1$ Hz, 1H), 5.05 (d, $J = 4.0$ Hz, 1H), 5.06 (d, $J = 10.3$ Hz, 1H), 5.16 (d, $J = 8.3$ Hz, 1H), 7.67 (d, $J = 9.1$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.93, 15.99, 17.02, 17.22, 17.52, 17.98, 18.05, 18.66, 18.69, 18.77, 19.27, 19.33, 19.43, 19.57, 19.73, 25.17, 25.23, 27.29, 28.21, 28.31 (3C), 30.03, 30.10, 30.13, 31.05, 31.15, 31.81, 33.18, 33.37, 38.07, 38.24, 51.41, 51.91, 59.96, 61.08, 66.17, 68.54, 68.68, 74.27, 76.89, 76.93, 79.48, 83.82, 94.36, 155.74, 167.52, 167.66, 168.34, 168.81, 169.72, 170.44, 171.68, 176.68, 176.86.

Compound (25)

Zinc powder (81 mg, 1.24 mmol) was added in three portions to a stirred solution of ester (**24**) (26.8 mg, 0.034 mmol) in THF (1.5 mL) under an atmosphere of argon and then 1N NH_4OAc solution (0.2 mL) was added. The reaction mixture was stirred at room

temperature overnight and filtered through a pad of celite. The residue was washed with ethyl acetate (3 x 5 mL) and the combined filtrates were diluted with ethyl acetate (25 mL), washed with dilute KHSO_4 solution (2x5 mL), saturated NaCl solution (10 mL), dried (Na_2SO_4) and evaporated *in vacuo* to give acid (**3**) (19 mg, 81%) as a pale yellow oil which was used in the next reaction without further purification.

Trifluoroacetic acid (2 mL) was added dropwise to a stirred solution of ester (**24**) (25.7 mg, 0.03 mmol) in dry dichloromethane (2 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 16h. The reaction mixture was poured into a mixture of benzene and chloroform (1:1, 10 mL) and evaporated *in vacuo* to give amine (**2**) as a brown residue which was used in the next reaction without further purification.

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (132 mg, 0.69 mmol) was added in one portion to a stirred mixture of acid (**3**) (19 mg, 0.027 mmol), amine (**2**), 1-hydroxybenzotriazole (93 mg, 0.69 mmol) and diisopropylethylamine (0.26 ml, 15 mmol) in dry dichloromethane (1 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 2 h and stirred at room temperature overnight. The mixture was dissolved in ethyl acetate (20 mL) and washed with saturated NH₄Cl solution (5 mL) and saturated NaCl solution (5 mL). The organic layer was dried (Na₂SO₄) and evaporated *in vacuo* to leave a yellow oil. The oil was purified by silica gel chromatography using 1% methanol – chloroform as eluent to give the *amide* (21 mg, 53%) as a colorless viscous oil; IR (neat); ν_{max} 3311, 2967, 2935, 2876, 2118, 1743, 1650, 1505, 1468, 1411, 1391, 1370, 1279, 1181, 1126, 1026, 756, 718 and 665 cm⁻¹; Precursor (**25**): $[\alpha]_{\text{D}}^{20}$ - 71.7 (c = 1.41, EA); ¹H NMR (400 MHz, CDCl₃) δ 0.85 (d, J = 6.9 Hz, 3H), 0.87 (d, J = 6.8 Hz, 3H), 0.90 (d, J = 6.9 Hz, 3H), 0.93 – 1.00 (m, 18H), 1.04 – 1.10 (m, 18H), 1.16 – 1.18 (m, 6H), 1.24 (d, J = 7.5 Hz, 3H), 1.43 (s, 9H), 1.47 – 1.58 (m, 8H), 1.89 – 1.91 (m, 1H), 1.94 (t, J = 2.5 Hz, 1H), 2.10 – 2.41 (m, 12H), 2.93 – 3.00 (m, 2H), 2.86, 3.03 (s, 3H), 2.87, 3.06 (s, 3H), 3.97 – 4.12 (m, 3H), 4.34 (dd, J = 5.3, 8.4 Hz, 1H), 4.69 (d, J = 12.0 Hz, 1H), 4.90 – 4.96 (m, 4H), 5.01 – 5.08 (m, 4H), 6.70 (d, J = 8.6 Hz, 1H), 7.56 (d, J = 9.4 Hz, 1H), 7.74 (d, J = 9.3 Hz, 1H), 7.80 (d, J = 9.4 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 15.36, 15.43, 15.53, 15.62, 16.97, 17.00, 17.15, 17.26, 17.66, 17.68, 17.89, 17.93, 17.97, 18.62, 18.68, 18.76, 18.80, 18.92, 19.06, 19.25, 19.39, 19.56, 19.64, 19.76, 19.95, 22.76, 24.94, 25.04, 25.11, 25.19, 27.46, 27.57, 27.94, 28.29, 28.32 (3C), 29.07, 29.66, 29.92, 29.97, 30.06, 30.08, 30.64, 30.68, 31.23, 31.27, 31.57, 31.69, 31.96, 32.10, 33.13, 33.25, 33.32, 33.40, 38.19, 38.24, 38.45, 38.67, 51.12, 51.50, 51.61, 51.73, 51.78, 51.82, 58.27, 58.40, 60.06, 60.98, 61.29, 61.84, 65.88, 68.60, 68.66, 68.69, 74.22, 76.81, 76.84, 77.20, 77.84, 79.20, 79.29, 79.55, 83.88, 83.94, 84.02, 84.12, 94.32, 155.69, 167.55, 167.64, 168.00, 168.26, 168.37, 168.50, 168.79, 169.44, 169.70, 170.22, 170.26, 170.38, 170.44, 170.68, 171.74, 176.52, 176.59, 176.77

The isomer **25a** was synthesized according to the above procedures from **24a**

Precursor (**25a**): $[\alpha]_D^{20} - 70.7$ ($c = 0.40$, CHCl_3); ^1H NMR (600 MHz, CDCl_3) δ 0.91 (d, $J = 6.5$ Hz, 3H), 0.92 (d, $J = 6.7$ Hz, 3H), 0.99 – 0.99 (m, 18H), 1.03 – 1.11 (m, 27H), 1.16 (d, $J = 7.0$ Hz, 3H), 1.44 (s, 9H), 1.51 – 1.58 (m, 8H), 1.90 (t, $J = 2.6$ Hz, 2H), 1.97 (t, J

= 2.6 Hz), 2.12 – 2.42 (m, 12H), 2.94 – 2.98 (m, 2H), 2.92, 3.04 (s, 3H), 2.93, 3.05 (s, 3H), 3.95 – 4.10 (m, 3H), 4.33 (dd, $J = 5.6, 8.4$ Hz, 1H), 4.80 (dd, $J = 12.0, 134.4$ Hz, 1H), 4.94 (d, $J = 4.1$ Hz, 1H), 4.96 (d, $J = 4.0$ Hz, 1H), 5.01 – 5.04 (m, 3H), 5.09 (d, $J = 10.5$ Hz, 1H), 5.14 (d, $J = 7.6$ Hz, 1H), 6.70 (d, $J = 8.4$ Hz, 1H), 7.52 (d, $J = 9.3$ Hz, 1H), 7.57 (d, $J = 9.4$ Hz), 7.60 (d, $J = 9.5$ Hz), 7.65 (d, $J = 9.1$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ 14.65, 14.82, 14.95, 15.75, 16.93, 17.01, 17.04, 17.21, 17.23, 17.52, 17.98, 18.05, 18.54, 18.67, 18.70, 18.76, 18.96, 19.04, 19.19, 19.23, 19.28, 19.34, 19.44, 19.56, 19.70, 19.89, 23.82, 25.17, 25.25, 27.17, 27.29, 28.31 (3C), 29.33, 29.68, 29.92, 30.01, 30.10, 30.13, 30.55, 31.17, 31.41, 31.77, 33.05, 33.36, 38.04, 38.10, 51.42, 51.48, 58.38, 58.56, 59.95, 61.00, 61.07, 66.15, 68.69, 68.72, 74.26, 76.88, 76.95, 77.39, 79.33, 79.47, 83.74, 83.82, 94.36, 155.71, 167.38, 167.66, 168.32, 168.40, 168.47, 168.81, 169.70, 170.42, 170.46, 170.85, 171.66, 176.50, 176.59.

Onchidin (**1**) & (**1a**)

Zinc powder (81 mg, 1.24 mmol) was added in three portions to a stirred solution of the amide (**25**) (21 mg, 0.0147 mmol) in THF (1.5 mL) under an atmosphere of argon and then 1N NH_4OAc (0.2 mL) was added. The reaction

mixture was stirred at room temperature overnight and filtered through a pad of celite. The

residue was washed with ethyl acetate (3x5 mL) and the combined filtrates were diluted by ethyl acetate (25 mL), washed by dilute KHSO₄ solution (2x5 mL), saturated NaCl solution (5 mL), dried (Na₂SO₄) and evaporated *in vacuo* to give the *crude acid* as a pale yellow oil which was used in the next reaction without further purification.

Trifluoroacetic acid (2 mL) was added dropwise to a stirred solution of the crude acid in dry dichloromethane (2 mL) at 0°C and under an atmosphere of argon. The resulting mixture was stirred at 0°C for 16h. The reaction mixture was poured into a mixture of benzene and chloroform (1:1, 10 mL) and evaporated *in vacuo*. The crude product was dissolved in dry DMF (30 mL). To this solution was added diisopropylethylamine (83 µL, 0.48 mmol) and pentafluorophenyl diphenylphosphinate (101.4 mg, 0.24 mmol) at room temperature and under an atmosphere of argon. The resulting mixture was stirred at room temperature for 4 days. The solvent was evaporated *in vacuo* to leave a yellow residue, which was dissolved in ethyl acetate (30 mL) and washed with saturated NH₄Cl solution (5 mL), saturated NaHCO₃ solution (5 mL) and saturated NaCl solution (5 mL). The organic layer was dried (Na₂SO₄) and evaporated *in vacuo* to leave a brown residue. The residue was purified by silica gel chromatography using 1% methanol – chloroform as eluent to give (**1**) (8.7 mg, 62.4%) as a colorless oil; IR (neat) ν_{max} 3449, 3313, 2966, 2876, 2126, 1776, 1744, 1678, 1633, 1519, 1500, 1473, 1414, 1392, 1372, 1302, 1189, 1125, 1030, 1002, 755 and 665 cm⁻¹; HRMS(FAB) for C₆₀H₉₉N₆O₁₄ (M+1): Calcd 1127.722; Found 1127.720; (**1**): $[\alpha]_{\text{D}}^{20}$ - 95.5 (c = 0.58, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 0.81 (d, J = 6.7 Hz, 3H), 0.91 (d, J = 7.0 Hz, 3H), 0.93 (d, J = 6.9 Hz, 3H), 0.97 (d, J = 7.0 Hz, 3H), 1.01 (d, J = 6.8 Hz, 3H), 1.04 (d, J 6.9 Hz, 6H), 1.07 (d, J = 6.4 Hz, 3H), 1.15 (d, J = 7.2 Hz, 3H), 1.37 – 1.42 (m, 1H), 1.47 – 1.52 (m, 2H), 1.53 – 1.57 (m, 1H), 1.88 (t, J = 2.6 Hz, 1H), 2.07 – 2.12 (m, 1H), 2.14 – 2.20 (m, 2H), 2.40 (dq, J = 2.8, 6.8 Hz, 1H), 2.46 (dq, J = 2.9, 6.8 Hz, 1H), 2.63 (dq, J = 3.3, 7.0 Hz, 1H), 2.76 (dq, J = 2.5, 7.2 Hz, 1H), 2.86 (s, 3H), 3.92 (t, J = 9.4 Hz, 1H), 4.42 (dd, J = 3.3, 8.2 Hz, 1H), 5.22 (d, J = 2.9 Hz, 1H), 5.33 (d, J = 11.2 Hz, 1H), 5.60 (d, J = 2.8 Hz, 1H), 6.37 (d, J = 8.3

Hz, 1H), 8.98 (d, $J = 9.2$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 15.37, 16.14, 16.45, 16.50, 17.64, 18.08, 18.75, 19.06, 19.52, 20.01, 25.55, 26.64, 28.09, 30.05, 30.16, 31.18, 34.05, 39.00, 52.33, 58.69, 59.72, 68.71, 75.34, 78.33, 83.83, 165.27, 169.74, 170.54, 170.66, 176.37

The isomer **1a** was synthesized according to the above procedures from **25a**

(**1a**): $[\alpha]_{\text{D}}^{20} - 81.5$ ($c = 0.40$, CHCl_3); ^1H

NMR (600 MHz, CDCl_3) δ 0.91 (d, $J = 6.7$ Hz, 6H), 0.95 (d, $J = 6.9$ Hz, 3H), 0.96 – 0.98

(m, 6H), 0.99 (d, $J = 7.0$ Hz, 3H), 1.04 (d, $J = 6.9$ Hz, 3H), 1.10 (d, $J = 6.4$ Hz, 3H), 1.11 (d,

$J = 6.9$ Hz, 3H), 1.48 (dq, $J = 3.6, 6.7$ Hz, 1H), 1.50 – 1.58 (m, 3H), 1.90 (t, $J = 2.6$ Hz, 1H), 2.12 – 2.18 (m, 1H), 2.20 – 2.27 (m, 2H), 2.30 – 2.38 (m, 2H), 2.38 – 2.44 (m, 1H), 2.89 (dq, $J = 3.2, 7.0$ Hz, 1H), 2.97 (s, 3H), 4.03 – 4.06 (m, 1H), 4.39 (dd, $J = 5.4, 8.6$ Hz, 1H), 5.13 (d, $J = 4.6$ Hz, 1H), 5.15 (d, $J = 10.6$ Hz, 1H), 5.32 (d, $J = 3.1$ Hz, 1H), 6.68 (d, $J = 8.5$ Hz, 1H), 7.43 – 7.45 (br, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ 14.50, 16.73, 17.20, 17.62, 17.91, 18.67, 19.10, 19.27, 19.53, 19.94, 24.97, 26.66, 29.25, 30.34 (2C), 31.19, 32.08, 38.61, 51.09, 58.88, 60.87, 68.62, 76.93, 79.70, 84.08, 168.27, 169.60, 170.55, 170.79, 175.70.

Current Data Parameters

NAME D10-3560
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20090710
Time 16.05
INSTRUM dr-500
PROBHD BBO 5mm
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 0
DS 1
SWH 7485.030 Hz
FIDRES 0.228425 Hz
AQ 2.198924 sec
RG 812.7
DW 66.990 usec
DE 4.50 usec
TE 300.2 K
YC 0.000000 sec
DT 1.0000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.00 usec
PL1 -6.00 dB
SFO1 500.132506 MHz
F2 - Processing parameters
SI 32768
SF 500.1300131 MHz
WDW EN
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

10 MHz plot parameters

CX 25.00 cm
F1P 10.000 ppm
F1 5001.30 Hz
F2P -1.000 ppm
F2 -500.13 Hz
P1PCOR 0.44000 ppm/cm
NUC2 2D1.03721 Hz/cm

1

```

Current Data Parameters
NAME      PYG-356p
EXPNO     2
PROCNO    1

F2 - Acquisition Parameters
Date_     20080711
Time      15.14
INSTRUM   spect
PROBHD    BBO 5mm
PULPROG   zgpg30
TD         32768
SOLVENT   DMSO-d6
NS         1813
DS         1
SH         31446.541 Hz
AQ         0.965672 sec
RG         0.5210612 sec
DM         15.900 usec
DE         4.00 usec
TE         300.0 K
D1         3.00000000 sec
d11        0.03000000 sec

===== CHANNEL f1 =====
NUC1       13C
P1         8.20 usec
PL1        -1.50 dB
SFO1       125.771524 MHz

===== CHANNEL f2 =====
COPPR2     waltz16
NUC2       1H
PCPD2      100.00 usec
PL2        -1.50 dB
PL12       17.00 dB
SFO2       500.1312000 MHz

F2 - Processing parameters
SI         32768
SF         125.7577867 MHz
RG         64
WDW         EM
SSB         0
LB         1.00 Hz
GB         0
PC         0.70

1D NMR D101 parameters
CR         25.00 cm
CP         2149.00 ppm
F1         2640.13 ppm
F2         -10.000 ppm
F3         -1257.96 Hz
P1PCN      8.80000 ppm/cm
NUC1CN     1106.66846 Hz/cm
  
```

Current Data Parameters
NAME onchidin-wing
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20020911
Time 9.37
INSTRUM av600
PROBHD 5 mm TBO BB-1H
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 8
DS 0
SWH 8992.805 Hz
FIDRES 0.274439 Hz
AQ 1.8220063 sec
RG 256
DM 55.600 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

***** CHANNEL f1 *****

NUC1 1H
P1 12.50 usec
PL1 -6.00 dB
SF01 600.1330006 MHz

F2 - Processing parameters

SI 32768
SF 600.1300079 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 25.00 cm
CY 12.97 cm
F1P 10.000 ppm
F1 6001.30 Hz
F2P -1.000 ppm
F2 -600.13 Hz
PPMCM 0.44000 ppm/cm
HZCM 264.05719 Hz/cm

1a

Current Data Parameters
NAME onchidiin-wing
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20020911
Time 9.47
INSTRUM av600
PROBHD 5 mm TBO BB-1H
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 1987
DS 2
SWH 37593.984 Hz
FIDRES 1.147277 Hz
AQ 0.4369777 sec
RG 9195.2
DM 13.300 usec
DE 10.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 7.00 usec
PL1 -3.00 dB
SF01 150.9193483 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 -6.00 dB
PL12 9.00 dB
SF02 600.1350006 MHz

F2 - Processing parameters
SI 32768
SF 150.9028057 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 25.00 cm
CY 55.68 cm
FIP 210.000 ppm
F1 31689.59 Hz
F2 -10.000 ppm
F2 -1569.03 Hz
PPhCN 8.80000 ppm/cm
HZCN 1327.94458 Hz/cm

1a

Current Data Parameters
NAME P15-178
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010714
Time 10:56
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 107
DS 2
SWH 8012.820 Hz
FIDRES 0.244632 Hz
AQ 2.047731 sec
RG 101.6
DM 62.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.1324008 MHz

F2 - Processing parameters
SI 32768
SF 400.1300051 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 25.00 cm
CY 20.70 cm
FIP 10.000 ppm
F1 4000.30 Hz
F2 -1.000 ppm
F2 400.13 Hz
PRACH 0.44000 ppm/cm
HZCN 176.05121 Hz/cm

5

Current Data Parameters

NAME pyg-175
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010714
Time 11.03
INSTRUM 4000
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 203
DS 2
SWH 25125.829 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 1.4696.5
DW 19.900 usec
DE 18.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SF01 100.6227864 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SF02 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127755 MHz
WOM EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 11.78 cm
FIP 210.000 ppm
F1 21.26 68 Hz
F2 -10.000 ppm
F2 -1006.13 Hz
P0MCM 8.80000 ppm/cm
HZCM 865.39240 Hz/cm

171.081
155.914
136.108
128.489
128.186
128.026
77.318
77.000
76.682
67.011
63.857
57.536
54.064
32.062
28.208
21.644
20.888

5

Current Data Parameters

NMR
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010628
Time 6.06
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 271
DS 2
SWH 251.25 629 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 11565.2
DM 19.900 usec
DE 18.00 usec
TE 300.0 K
D1 3.0000000 sec
d11 0.0300000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SFO1 100.6237564 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.612728 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 8.70 cm
F1P 210.000 ppm
F1 21126.68 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 8.60000 ppm/cm
HZCM 885.39233 Hz/cm

Current Data Parameters

NAME pyg-3052
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011130
Time 21:16
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 35
DS 2
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 64
DW 62.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.1324008 MHz

F2 - Processing parameters

SI 32768
SF 400.1300046 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

10 NMR plot parameters

CX 25.00 cm
CY 53.87 cm
F1P 10.000 ppm
F1 4001.30 Hz
F2P -1.000 ppm
F2 -400.13 Hz
PPMCK 0.44000 ppm/cm
HZCM 176.05721 Hz/cm

7

Current Data Parameters
 NAME PY9-305Z
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20011130
 Time 21:18
 INSTR av400
 PROBHD 5 mm QNP 1H/13
 PULPROG zgpg
 TD 32768
 SOLVENT CDCl3
 NS 111
 DS 2
 SWH 25125.629 Hz
 FIDRES 0.766773 Hz
 AQ 0.6521332 sec
 RG 11595.2
 DW 19.900 usec
 DE 18.00 usec
 TE 300.0 K
 D1 3.00000000 sec
 d11 0.03000000 sec
 d12 0.0002000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 6.20 usec
 PL1 -5.00 dB
 SF01 100.6237954 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 P2 100.00 usec
 PL2 120.00 dB
 PL12 14.00 dB
 PL13 14.00 dB
 SF02 400.1316005 MHz

F2 - Processing parameters
 SI 32768
 SF 100.6127729 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 0.70

1D NMR plot parameters
 CX 25.00 cm
 CY 9.86 cm
 F1P 210.000 ppm
 F1 21128.68 Hz
 F2P -15.000 ppm
 F2 -1505.19 Hz
 PPM0H 9.00000 ppm/cm
 HZ0M 905.51501 Hz/cm

136.499
 128.455
 128.422
 127.998
 127.972
 94.716
 77.314
 76.996
 76.679
 67.408
 66.534
 64.914
 51.580
 48.061
 38.765
 34.155
 29.332
 22.833
 18.036
 -1.474

7

ppm

Current Data Parameters
 NAME pyg-307
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20011204
 Time 17:58
 INSTRUM av400
 PROBHD 5 mm QNP 1H/13
 PULPROG zgpg30
 TD 32768
 SOLVENT CDCl3
 NS 67
 DS 2
 SWH 8012.820 Hz
 FIDRES 0.244532 Hz
 AQ 2.0447731 sec
 RG 90.5
 DM 62.400 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 10.50 usec
 PL1 -4.00 dB
 SFO1 400.1324008 MHz

F2 - Processing parameters
 SI 32768
 SF 400.1300071 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 25.00 cm
 CT 31.95 cm
 F1P 10.000 ppm
 F1 400.130 Hz
 F2P -1.000 ppm
 F2 -400.13 Hz
 PPHC 0.44000 ppm/cm
 HZCN 176.05121 Hz/cm

8

Current Data Parameters
NAME pyg-307
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011204
Time 18.08
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TO 32768
SOLVENT CDCl3
NS 104
DS 2
SWH 25125.625 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 20642.5
DM 19.900 usec
DE 18.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec
d12 0.00022000 sec

===== CHANNEL f1 =====

NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SFO1 100.6237964 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters

SI 32768
SF 100.6127726 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters

CX 25.00 cm
CY 10.27 cm
F1P 210.000 ppm
F1 21126.68 Hz
F2P -15.000 ppm
F2 -1505.19 Hz
PRMCM 9.00000 ppm/cm
HZCM 905.51489 Hz/cm

8

Current Data Parameters
NAME PYS-313
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 2001213
Time 11.54
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 65
DS 2
SWH 8012.850 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 2813
AQ 62.441000 usec
DE 5.00 usec
TE 300.1 K
D1 1.0000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.1324008 MHz

F2 - Processing parameters

SI 32768
SF 400.1300071 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 25.00 cm
CY 34.36 cm
F1P 10.000 ppm
F1 4001.30 Hz
F2P -1.000 ppm
F2 -400.13 Hz
PPHMC 0.44000 ppm/cm
HZCN 176.05721 Hz/cm

16

Current Data Parameters
 NAME Ne-1ys-Va1 (N)
 EXPNO 12
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20020621
 Time 21.06
 INSTRUM av400
 PROBHD 5 mm QNP 1H/13
 PULPROG zg30
 TO 32768
 SOLVENT CDCl3
 NS 60
 DS 2
 SWH 8012.820 Hz
 FIDRES 0.244532 Hz
 AQ 2.0447731 sec
 RG 228.1
 DM 62.400 usec
 DE 5.00 usec
 TE 300.0 K
 D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
 P1 10.40 usec
 PL1 -4.00 dB
 SFO1 400.1324008 MHz

F2 - Processing parameters

SF 32768
 SF 400.1300084 MHz
 MDW rg
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters

CX 25.00 cm
 CY 34.42 cm
 F1 10.000 ppm
 F2 400.130 Hz
 F3 10.000 ppm
 F4 -100.13 Hz
 PPMH 0.44000 ppm/cm
 HZCM 176.05721 Hz/cm

16a

Current Data Parameters
NAME PYG-324r
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011216
Time 23.27
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 600
DS 2
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.044731 sec
RG 45.3
DM 62.400 usec
DE 5.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.1324008 MHz

F2 - Processing parameters

SF 32768
SF 400.1599870 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 25.00 cm
CY 25.09 cm
F1 10.000 ppm
F2 4001.30 Hz
F3 -1.000 ppm
F4 0.000 Hz
PRACH 0.00000000 ppm/cm
HZCN 175.05719 Hz/cm

18

Current Data Parameters
NAME pyg-324r
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011217
Time 5:02
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
DS 5000
NS 2
DS 25125.629 Hz
FIDRES 0.76573 Hz
AQ 0.652132 sec
RG 8193
RW 19.600 usec
DE 16.00 usec
TE 300.0 K
D1 3.0000000 sec
d11 0.0300000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====

NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SF01 100.623766 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SF02 400.1315005 MHz

F2 - Processing parameters

SF 32768
SF 100.6127766 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters

CX 25.00 cm
CY 19.69 cm
F1P 210.000 ppm
F1 21126.68 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 8.80000 ppm/cm
HZCM 885.39240 Hz/cm

Current Data Parameters
NAME Boc-Val-Lys-Val
EXPNO 6
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020711
Time 16.04
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 220
DS 2
SWH 25125.629 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 1.4596.5
DM 19.500 usec
DE 18.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 6.10 usec
PL1 -5.00 dB
SF01 100.6237864 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 -4.00 dB
PL12 13.50 dB
PL13 14.00 dB
SF02 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127759 MHz
WDW EM
SSE 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 34.02 cm
F1P 210.000 ppm
F1 21128.68 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 8.80000 ppm/cm
HZCM 885.35240 Hz/cm

Current Data Parameters
NAME DVS-267
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20011123
Time 18.00
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 102
DS 2
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 50.5
DM 62.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.132408 MHz

F2 - Processing parameters
SI 32768
SF 400.130071 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 25.00 cm
CY 27.32 cm
F1F 10.000 ppm
F1 400.130 Hz
F2F -1.000 ppm
F2 -400.13 Hz
P1F0CN 0.44000 ppm/cm
H1CN 176.05721 Hz/cm

20

Current Data Parameters	
NAME	pyg-297
EXPNO	2
PROCNO	1

F2 - Acquisition Parameters

Date_	Time	Inst	Temp	Pressure	PULPHOG	TD	SOLVENT	COC13	NS	DW	FLOW	FLOW	DE	TE	D1	D1	g11	d12				
20061123	16:07	E	av.000	5 mm	QNP	14113	z90g	32768	CCO13	187	2	25125.629 Hz	0.766773 Hz	0.6521332 sec	18390.4	19.900 uSec	18.00 uSec	300.0 K	3.0000000 sec	0.0300000 sec	0.0002000 sec	0.0002000 sec

CHANNEL f1

NUC1	13C
P1	6.20 usec
PL1	-5.00 dB
SFO1	100.6237954 MHZ

===== CHANNEL f2 =====

Channel 12	WJZZ15	1H
CDPH62		
NUC2	100.00	usec
PCP02	120.00	dB
PL2	14.00	dB
PL12	14.00	dB
PL13	14.00	dB
SF02	400.1316005	MHz

F2 - Processing parameters

PC - processing parameters	
SI	32768
SF	100.6127728 MHz
WDW	EM
SSB	0
LB	1.00 Hz
GB	0
PC	0.70

1D NMR plot parameters

ID	NMR	proton parameters
CX		25.00 cm
CY		14.90 cm
F1P		210.000 ppm
F1		21128.68 Hz
F2P		-10.000 ppm
F2		-1006.13 Hz
PPMCM		8.80000 ppm/cm
HZCM		885.39233 Hz/cm

20

Current Data Parameters
NAME Boc-Val-Tyr-DCCl-Val (N)
EXPNO 4
PROCNO 1
F2 - Acquisition Parameters
Date_ 20020730
Time 15:55
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 2
DS 2
SWH 8012.560 Hz
FIDRES 0.244630 Hz
AQ 2.0447731 sec
RG 64
DM 62.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.0000000 sec
***** CHANNEL f1 *****
NUC1 1H
P1 10.40 usec
PL1 -4.00 dB
SF01 400.1324000 MHz
F2 - Processing parameters
SI 400.1299975 MHz
SF 400.1299975 MHz
NUC2 13C
LB 0.00 Hz
GB 0
PC 1.00
1D NMR plot parameters
CX 25.00 cm
CY 22.15 cm
F1P 10.000 ppm
F2P 400.130 Hz
F3P 100.628 MHz
F2 -400.13 Hz
PPH0 0.44000 ppm/cm
HZCN 176.05721 Hz/cm

Current Data Parameters
NAME Boc-Val-lys (CCl₄-V01 IN)
EXPNO 5
PROCNO 1

F2 - Acquisition Parameters

Date_ 20070721
Time 12:05
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TD 32768
SOLVENT CDCl₃
DS 2
SWH 25125.659 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 14596.5
DM 19.900 usec
DE 19.900 usec
TE 300.0
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 ¹³C
PL1 6.10 usec
PL1 -5.00 dB
SF01 100.6279864 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 ¹H
PL2 6.00 usec
PL2 -4.00 dB
PL12 13.60 dB
PL13 14.00 dB
SF02 400.138605 MHz

F2 - Processing parameters
SI 32768
SF 100.6127726 MHz
HOM EN
SSB 0
GB 1.00 Hz
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 29.34 cm
F1 200.000000 MHz
F2 100.000000 MHz
F2 -1006.13 Hz
APPCN 8.60000 ppm/cm
HZCN 885.36233 Hz/cm

20a

Current Data Parameters
 NAME pyg-283
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20011112
 Time 19.22
 INSTRUM av400
 PROBHD 5 mm QNP 1H/13
 PULPROG zg30
 TO 32768
 SOLVENT CDCl3
 NS 174
 DS 2
 SWH 8012.820 Hz
 FIDRES 0.244632 Hz
 AQ 2.0447731 sec
 RG 114
 DW 62.400 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
 P1 10.50 usec
 PL1 -4.00 dB
 SFO1 400.1324008 MHz

F2 - Processing parameters

SI 32768
 SF 400.1300058 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters

CX 25.00 cm
 CY 34.67 cm
 F1P 10.000 ppm
 F1 4001.30 Hz
 F2P -1.000 ppm
 F2 -400.13 Hz
 PPHCM 0.44000 ppm/cm
 HZCM 176.05721 Hz/cm

23

Integral

Current Data Parameters
NAME py9-283
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011112
Time 19.31
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 168
DS 2
SWH 25125.629 Hz
FIDRES 0.766773 Hz
AQ 0.6521332 sec
RG 14596.5
DM 19.900 usec
DE 18.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SFO1 100.6237964 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127713 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 9.69 cm
F1P 210.000 ppm
F1 21128.68 Hz
F2P -15.000 ppm
F2 -1509.19 Hz
PPMCH 9.00000 ppm/cm
HZCM 905.51488 Hz/cm

172.004
167.900

94.404
94.252
79.396
77.316
76.999
76.681
76.595
76.471
74.219
65.660

31.469
30.070
18.922
18.834
18.030
17.161
17.032

-1.458

23

Current Data Parameters
 NAME p99-531
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20011225
 Time 1143
 INSTRUM spect
 PROBRD 5 mm QNP 1H/13
 PULPROG zgpg30
 TD 32768
 SOLVENT CDCl3
 NS 572
 DS 2
 SWH 8012.826 Hz
 FIDRES 0.144532 Hz
 AQ 2.0447731 sec
 RG 181
 DQ 62.400 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
 P1 10.50 usec
 PL1 -4.00 dB
 SFO1 400.1324008 MHz

F2 - Processing parameters

SI 32768
 SF 400.1300556 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 25.00 cm
 CY 21.52 cm
 F1P 10.000 ppm
 F1 400.130 Hz
 F2P -1.000 ppm
 F2 -400.13 Hz
 PPMCH 0.44000 ppm/cm
 HZCM 176.05721 Hz/cm

24

Current Data Parameters
NAME p19-33f
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20011222
Time 20.38
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 299
DS 2
SWH 25125.629 Hz
FIDRES 0.766773 Hz
AQ 0.652132 sec
RG 16384
DM 19.500 usec
DE 18.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 6.20 usec
PL1 -5.00 dB
SFO1 100.6237954 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.612751 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70

1D NMR plot parameters
CX 25.00 cm
CY 17.79 cm
F1P 210.000 ppm
F1 21428.68 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
RFMW 8.80000 ppm/cm
HZCM 885.39240 Hz/cm

Cl₃CH₂CO

24

Current Data Parameters
NAME VALU@VAL-TYPE@CD-VOL INI
EXPNO 3
PROCNO 1
F2 - Acquisition Parameters
Date_ 20080824
Time 15 15
INSTRUM av400
PROBHD 5 mm BBO H-13
PULPROG zgpg30
CROSSPO 2DZG30
SOLVENT CDCl3
NS 2272
DS 2
SWH 25125.626 Hz
FIDRES 0.750773 Hz
AQ 0.0618556 SEC
RG 658.5
DE 19.800 uSBC
TE 300.0 K
D1 2.0000000 SEC
d11 0.0500000 SEC
d12 0.0000000 SEC
***** CHANNEL f1 *****
NUC1 13C
P1 6.10 uSBC
PL1 0.00 dB
SFO1 100.627764 MHz
***** CHANNEL f2 *****
CPDPRG2 MOLD16
NUC2 1H
P2 60.00 uSBC
PL2 -4.00 dB
PL12 13.60 dB
PL13 14.00 dB
SFO2 400.1316005 MHz
F2 - Processing parameters
SI 32768
SF 100.6127697 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 0.70
1D NMR plot parameters
CX 25.00 cm
CY 65.00 cm
FID 210.00 cm
F1 21126.66 Hz
F2 -10.000 ppm
P2 -1006.13 Hz
PRQW 8.80000 ppm/cm
NUC# 865.39233 Hz/cm

24a

Current Data Parameters
NAME dyg-334r
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20011227
Time 22.38
INSTRUM av400
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TO 32768
SOLVENT CDCl3
NS 500
DS 2
SWH 8012.820 Hz
FIDRES 0.244532 Hz
AQ 2.0447731 sec
RG 90.5
CW 62.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 10.50 usec
PL1 -4.00 dB
SFO1 400.1324008 MHz

F2 - Processing parameters

SF 400.1300039 MHz
SE 32768
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 25.00 cm
CY 17.73 cm
F1P 10.000 dBm
F2P 400.36 Hz
F3P 10.000 dBm
F2P -400.13 Hz
PFRCH 0.46000 dBm/cm
HZCR 176.05721 Hz/cm

25

Current Data Parameters
NAME linear-wing
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020827
Time 12.28
INSTRUM av600
PROBHD 5 mm TBO BB-1H
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 8
DS 0
SWH 8992.806 Hz
FIDRES 0.274439 Hz
AQ 1.822063 sec
RG 161
DM 55.600 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 12.50 usec
PL1 -6.00 dB
SF01 600.1330006 MHz

F2 - Processing parameters
SI 32768
SF 600.1300079 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 25.00 cm
CY 21.86 cm
F1P 10.000 ppm
F1 6001.30 Hz
F2 -1.000 ppm
F2 -600.13 Hz
PPMCM 0.44000 ppm/cm
HZCM 264.05719 Hz/cm

25a

Current Data Parameters
NAME linear-wing
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020827
Time 12.38
INSTRUM av600
PROBHD 5 mm TGD BB-1H
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 2026
DS 2
SWH 37593.984 Hz
FIDRES 1.147277 Hz
AQ 0.4338777 sec
RG 9195.2
DM 13.300 usec
DE 10.00 usec
TE 300.0 K
D1 3.00000000 sec
d11 0.03000000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 7.00 usec
PL1 -3.00 dB
SF01 150.9193483 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 -6.00 dB
PL12 9.00 dB
SF02 600.1330005 MHz

F2 - Processing parameters
SI 150.9028665 MHz
SF 32765
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 25.00 cm
CY 125.16 cm
F1P 210.000 ppm
F1 31685.35 Hz
F2P -10.000 ppm
F2 -1595.03 Hz
PFGCH 8.50000 ppm/cm
HZCH 1327.94458 Hz/cm

25a